NEET QUESTION PAPER (2018) BOOKLET CODE - ACHLA (CC)

| - 14 | I. Niche is | | |
|------|--|------|--|
| | (1) the | | Offsets are produced by |
| | the functional role played by the organism the range of tem. | 17. | Offsets are proceed (1) Parthenogenesis |
| | (2) where it lives role played by the organism | | (1) Parthenocarpy (2) Parthenocarpy |
| | (2) the range of temperature that the organism (3) the physical space | | |
| | (3) heeds to live (4) the physical space where an organism lives environment. | | an intic division a comiconservative |
| | (4) the physical | 10.1 | (4) Melotic and proof for semiconverter The experimental proof for semiconverter replication of DNA was first shown in a |
| | (4) all the biological space where an organism lives environment Which of the second sec | 8. | The experiment of DNA was first showing |
| 2 | environment factors in the organism's | | replication of 2- |
| 4 | Which of the are | | |
| | Which of the following is a secondary pollutant ? (1) O₃ (2) and | | |
| | (2) SO | | |
| | SO_2 | 2 | t atch ' |
| | ⁽³⁾ CO ₂ | 9. | Select the correct match - Lac operon |
| | (4) CO ² | | (1) Francois Jacob and |
| 3. | | | Jacques Monod Jacques Monod – Pisum sativum |
| | III Stratogal | | (2) Matthew Meselson |
| | In stratosphere, which of the following elements acts as a catalyst in degradation of game and release of male | | and F. Stahl |
| | | | (3) Alfred Hersney and |
| | (1) Oxygen | 100 | (4) Alec Jeffreys - Streptococcus |
| | (2) Fe | | (4) Alec Jeffreys – Streptococcus pneumoniae |
| | (3) Cl | 1 | |
| | (4) Carbon | 10. | Which of the following has proved helpful in |
| 4. | World O | | preserving pollen as fossils ? |
| | World Ozone Day is celebrated on | | (1) Sporopollenin |
| | (1) 22 nd April | | (2) Oil content |
| | | 1 | (3) Cellulosic intine |
| | Deptember | 12 | (4) Pollenkitt |
| | (3) 21 st April | 11. | Which of the following pairs is <i>wrongly</i> matched? |
| | (4) 5^{th} June | | (1) T.H. Morgan : Linkage |
| 5. | What type of ecological pyramid would be | | (9) VOL |
| | obtained with the following data? | | determination : Grasshopper |
| | Secondary consumer : 120 g | | (3) ABO blood |
| | Primary consumer : 120 g | | (4) Stouch and the store and a store and the |
| | Primary consumer : 60 g | 12. | Which of the following and the |
| | Primary producer : 10 g | | Which of the following flowers only once in its life-time? |
| | (1) Upright pyramid of biomass | 1. | (1) Papaya . |
| | (2) Upright pyramid of numbers | | (2) Mango |
| | (3) Pyramid of energy | | (3) Jackfruit |
| | (4) Inverted pyramid of biomass | | (4) Bamboo species |
| · | ry-mind of biolinusy | 13. | Select the correct statement ; |
| 6. | Natality refers to | | Transduction was discovered by S. Altman. Splitscore and the second second |
| | (1) Number of individuals entering a habitat | | (2) Spliceosomes take part in translation. |
| | (2) Number of individuals leaving the habitat | | (3) Punnett square was developed by a British |
| | (3) Birth rate | - | scientist. |
| * * | (4) Death rate | | (4) Franklin Stahl coined the term "linkage". |

ACHLA/CC/Page 2

SPACE FOR ROUGH WORK

English

| 14. The c | |
|--|--|
| - ne Goloria | |
| Activation of amino acid Formation in back | |
| (2) Respiration of amino acid (3) Formation of second (4) Foto | Match the items given in Column I with those in II and select the correct option given |
| (3) Formation in bact | 22. Match the items given in Column I with those in Column II and select the correct option given |
| (4) Fatter of second | |
| 15. The store acid break a | below: Column II |
| (4) Fatty acid breakdown 15. The stage during which separation of the paired (1) Zygotene (2) Diakinesis (3) Diplote | Column I It is a place having a |
| (1) Zu chromos | <i>Column</i> Herbarium i. It is a place having a collection of preserved |
| (2) Discovere and somes begins of the | plants and animals. |
| (3) Diakinesis | t list that any set |
| | |
| 16. Stomatel | b. Key methodically all the |
| 16. Stomatal movement is not affected by (1) CO_2 concentration (2) O_2 concentration | species found in an area |
| (1) CO comment is not | with brief description |
| (2) O 2 concentration affected by | aiding identification. |
| (2) O₂ concentration (3) Light | iii. Is a place where dried and |
| (3) Light (4) Te | Museum in. Is a pressed plant specimens |
| Tom . | mounted on sheets are |
| | kept. |
| (1) Barrel shaped (2) Becta | A healthat containing a list |
| (2) Rectangel | d. Catalogue iv. A bookiet containing a list of characters and their |
| (3) Kidney shaped (4) Dumb I shaped | alternates which are |
| (4) Dumb ball | |
| (4) Dumb-bell shaped 18. Which of the following is not a product of light reaction of photosynthesis? (1) Oxygen | various taxa. |
| reaction of the following is not | various taxa. |
| (1) Oxygen | a b c d |
| (2) NADPH | -(1) iii iv i ii |
| (3) NADU | (2) ii iv iii i |
| $\begin{array}{c} (3) \text{NADH} \\ (4) \text{ATP} \end{array}$ | (3) - iii ii i iv |
| · • • • • • • • • • • • • • • • • • • • | $\begin{array}{cccccccccccccccccccccccccccccccccccc$ |
| which of the following is t | |
| (1) It is a site for anti- | 23) Which one is wrongly matched? |
| synthesis. | (1) Unicellular organism – Chlorella |
| 41 It takes part in spindle c | (2) Gemma cups – Marchantia |
| (3) It is a membrane-bound structure. (4) Larger pucket: | (3) Biflagellate zoospores – Brown algae |
| (4) Larger pugloal: | |
| (4) Larger nucleoli are present in dividing cells.20. Which among the following in the foll | |
| | Alter karyogamy followed by meiosis, spores are |
| | produced exogenously in |
| | (1) Saccharomyces |
| (3) Mycobacterium | (2) Agaricus |
| (4) Saccharomyces | (3) Alternaria |
| 21. The two functional groups characteristic of | (4) Neurospora |
| sugars are | |
| (1) carbonyl and hydroxyl | grands are present in |
| (2) carbonyl and phosphate | (1) Pinus (2) Manage |
| | (2) Mango |
| (3) carbonyl and methyl | (3) Cycas |
| (4) hydroxyl and methyl | (4) Mustard |
| ACHLA/CC/Page 3 SPACE FOR | ROUGH WORK |



| | | | | hat is the role of NAD ⁺ in cellula |
|---------------------|--|---|--|--|
| Whi | ich of the following is commonly used as a tor for introducing a DNA fragment in b | - | | at is the role of NAD anaerobi piration? It is the final electron acceptor for anaerobi respiration. |
| vect | tor for introducing a DNA fragment in as a | 32. | TIN | hat is the loctron acceptor for thesis. |
| 1ym | phocytes ? | 1 02. | Wh | piration? biration? |
| (1) | pBR 322 | | Ni | It is the final of for ATP synta |
| (2) | λ phage | | (4) | is the piration? It is the final electron acceptor respiration. It is a nucleotide source for ATP synthesis. It is a nucleotide source for ATP synthesis. It functions as an electron carrier. |
| (3) | Ti plasmid | | (2) | It is a nucleotide electron ct |
| (4) | Retrovirus | | (3) | it tunes as an energy lents shows where |
| A 'n com pres | ew' variety of rice was patented by a foreign pany, though such varieties have been then in India for a long time. This | 33, | Whi clos | It is a hud It functions as an electric It functions as an enzyme. It functions as an enzyme. ich one of the following plants shows a ver, ich one of the following plants shows where ich one of the following plants of moth, where ich one of the two can complete its life cycle without other ? |
| as | Basmati | | the | other? |
| • | | | (1) | Viola |
| | | | (2) | Banana |
| | | | (3) | Yucca |
| | | | (4) | Hydrilla vears in |
| | - Other of the Willing will all homeotics - P | 34. | Poll liqu | Hydrilla en grains can be stored for several years in id nitrogen having a temperature of |
| (1) | Bioexploitation | | (1) | – 160°C |
| 1999 | | 12 | (2) | – 196°C |
| | | - | (3) | – 80°C |
| | | | (4) | – 120°C |
| | | 35. | | which of the following forms is iron absorbed lants? |
| 1 | A 14 14 | | (1) | Both ferric and ferrous |
| 1 | | | (2) | Free element |
| C | | | (3) | Ferrous |
| | 신경 이번에 여름 것이 집중 것이 가지 않는 것이 많이 많을 것 같아요. 것이 같아. | | . (4)- | Ferric |
| (4) | Ribozyme – Nucleic acid | 36. | Doul | ble fertilization is |
| | | | | Syngamy and triple fusion |
| | | 105 | | Fusion of two male gametes with one egg Fusion of one male gamete with two polar |
| | | | | |
| | | 1 | (4) | Fusion of two male gametes of a pollen tube |
| | | | | |
| 1000 | 성경 성상 전기가 많은 영상성장은 일을 가지 않는 것 같아. | 37. | Oxyg | gen is <i>not</i> produced during photosynthesis by |
| n In | idia, the organisation responsible for | | (1) | Onara |
| CODCC | ing the safety of introducing genetically | | | Cycas |
| nodifi | ed organisms for public use is | in the | | Nostoc |
|) G | enetic Engineering Appraisal Committee | 5 | (4) | Green sulphur bacteria |
| ((| GEAC) | 38 | Whi | ch of the following elements is responsible for |
| R | esearch Committee on Genetic | | mai | intaining turgor in cens . |
| M | Inimulation (RCGM) | | (1) | |
| | a und Industrial | | (2) | Potassium |
| D | accorch (CSIR) | | .(3) | Sodium |
| R | and a second of Medical Research (ICMR) | | (4) | Magnesium English |
| | (1) (2) (3) (4) A 'n com pres (2) (2) (3) (4) Use and conc (1) (2) (3) (4) Selec (1) (1) (2) (3) (4) (1) (1) (2) (1) (2) (1) (1) (1) (1) (2) (1) (1) (1) (1) (1) (1) (1) (1) (1) (1 | (1) pBR 322 (2) λ phage (3) Ti plasmid (4) Retrovirus A 'new' variety of rice was patented by a foreign present in India for a long time. This is related to (2) Basmati (2) Lerma Rojo (3) Sharbati Sonora (4) Co-667 Use of bioresources by multinational companies and organisations without authorisation from the concerned country and its people is called (1) Bioexploitation (2) Biodegradation (3) Biopiracy (4) Bio-infringement Select the correct match: (1) G. Mendel – Transformation (2) T.H. Morgan – Transduction (3) $F_2 \times \text{Recessive parent}$ – Dihybrid cross (4) Ribozyme – Nucleic acid The correct order of steps in Polymerase Chain Reaction (PCR) is (1) Denaturation, Annealing, Extension 2) Denaturation, Extension, Annealing 3) Annealing, Extension, Denaturation 4) Extension, Denaturation, Annealing in India, the organisation responsible for ssessing the safety of introducing genetically nodified organisms for public use is (2) Genetic Engineering Appraisal Committee (GEAC) (Council for Scientific and Industrial Research (CSIR) I redian Council of Medical Research (ICMR) | (1) pBR 322 (2) λ phage (3) Ti plasmid (4) Retrovirus A 'new' variety of rice was patented by a foreign present in India for a long time. This is related to (4) Retrovirus (5) Basmati (2) Lerma Rojo (3) Sharbati Sonora (4) Co-667 Use of bioresources by multinational companies and organisations without authorisation from the concerned country and its people is called (1) Bioexploitation (2) Biodegradation (3) Biopiracy (4) Bio-infringement Select the correct match: (1) G. Mendel – Transformation (2) T.H. Morgan – Transduction (3) F₂ × Recessive parent – Dihybrid cross (4) Ribozyme – Nucleic acid The correct order of steps in Polymerase Chain Reaction (PCR) is (1) Denaturation, Annealing, Extension (2) Denaturation, Annealing (3) Annealing, Extension, Denaturation (4) Extension, Denaturation (5) Genetic Engineering Appraisal Committee (GEAC) (6) Research Committee on Genetic Manipulation (RCGM) (7) Council for Scientific and Industrial Research (CSIR) (2) Indian Council of Medical Research (ICMR) | (1)pBR 322(1)(2) λ phage(2)(2) λ phage(2)(3)Ti plasmid(2)(4)Retrovirus(3)A 'new' variety of rice was patented by a foreign present in India for a long time. This is related to(3)(4)Basmati(2)(2)Lerma Rojo(3)(3)Sharbati Sonora(4)(4)Co-667(2)Use of bioresources by multinational companies and organisations without authorisation from the concerned country and its people is called(4)(1)Bioexploitation(2)(2)Biodegradation(3)(3)Biopiracy(4)(4)Bio-infringement(4)Select the correct match : (1)(1)(2)T.H. Morgan- Transformation (3)(3)F ₂ × Recessive parentDihybrid cross(4)Ribozyme- Nucleic acid (4)The correct order of steps in Polymerase Chain Reaction (PCR) is(3)(1)Denaturation, Annealing, Extension (2)(3)(3)Annealing, Extension, Denaturation (4)(4)(4)Extension, Denaturation, Annealing (3)(4)(4)Seessing the safety of introducing genetically (3)(4)(4)Genetic Engineering Appraisal Committee (GEAC)(3)(4)Genetic Engineering Appraisal Committee (GEAC)(3)(4)Genetic Engineering Appraisal Committee (GEAC)(3)(4)Genetic Engineeri |

HLA/CC/Page 4

Scanned by CamScanner

÷

| | | and the second se |
|-----|--|---|
| 39 | | and the second of |
| | (1) Submerged ball | 46. Nissl bodies are mainly composed of |
| | (2) Carnivorous plants | (1) Free ribosomes and rear |
| | to a locating has | (2) Nucleic acids and SER |
| | (4) Halophytes | (3) DNA and RNA |
| 40. | and three | The proteing and linids |
| | Mitochondria are the powerhouse of the cell Pseudopodia are l | 47. Which of these statements is <i>incorrect</i> ? |
| | (2) Pseudo (2) Pseudo (2) (2) (2) (2) (2) (2) (2) (2) (2) (2) | 47. Which of these statements is the statements is the statements of the statements is the statement of the |
| | Structures . and locomet | (1) Oxidative phosphorylation tendor i outer mitochondrial membrane. |
| | (3) Mushrooms hele | (2) Glycolysis operates as long as it is supplied |
| | (3) Mushrooms belong to Basidiomycetes. (4) Cell wall is present in members of Fungi Secondary xylem and the second second | (2) Glycolysis operates as long and with NAD that can pick up hydrogen atoms. |
| 122 | and Plantae. | (3) Glycolysis occurs in cytosol. |
| 41. | Secondary xylem and phloem in dicot stem are (1) Axillary merister | (4) Enzymes of TCA cycle are present in |
| | produced by | mitochondrial matrix. |
| | Axillary meristems Phellogram | 48. Many ribosomes may associate with a single |
| | (2) Phellogen | mRNA to form multiple copies of a polypeptide |
| ÷., | (2) Vascular cambium (4) Anicol | simultaneously. Such strings of ribosomes are |
| | -pical meristem- | termed as |
| 42. | Sweet potato is a modified | (1) Nucleosome |
| | (1) Rhizome | (2) Plastidome |
| | (2) Tap root | (3) Polyhedral bodies |
| | (3) Adventitious root | (A) Polysome |
| | (4) Stem | 49. Which of the following torms describe human |
| 43. | Which of the car | 49. Which of the following terms describe human dentition? |
| | Which of the following statements is correct? (1) Stems are usually and | (1) Pleurodont, Diphyodont, Heterodont |
| | Stems are usually unbranched in both Cycas and Cedrus. | (2) Pleurodont, Monophyodont, Homodont |
| | (2) Horsetails and Cedrus. | (3) Thecodont Diphysdant II (|
| | | Heterodont, Diphyodont, Heterodont |
| | (3) Selaginella is heterosporous, while Salvinia is homosporous. | (4) Thecodont, Diphyodont, Homodont50. Which of the fall |
| | | |
| | (4) Ovules are not enclosed by ovary wall in gymnosperms. | |
| 44. | | synthesis |
| | Casparian strips occur in (1) / Endodermis | and a signal peptide |
| | | a storm grycosylation |
| | (2) Cortex | (4) Protein folding |
| | (3) Pericycle | 51. Select the <i>incorrect</i> match : |
| | (4) Epidermis | L Dolate |
| 45. | Plants having little or no secondary growth are | chromosomes - Oocytes of amphibians |
| | (1) Cycads | (2) Submetacentric $-$ L-shaped -) |
| | (2) Conifers | (2) Submetacentric - L-shaped chromososmes (2) Allo |
| ÷ ' | (3) Deciduous angiosperms | (3) Allosomes - Sex chromos |
| | (4) Grasses | (4) Lampbrush – Diplotene bivalents |
| | A/CC/Page 5 SPACE FOR | chromosomes - piotene bivalents |

English



| | | | | | | | | a 1 | |
|-----|---------|---------------------------------|---------------------------------------|----------|--|----------------------------|-------------------|----------|-------------------------|
| 152 | 1. W.L. | i | | | | populatio | n of a cou | intry, | loce than |
| C | hor | ich of the following i mone? | is an amino acid derived | 56. In a | growing | populatio | individua | als are | less than |
| | | | | (1) | pre-repr | oductive oductive i | ndividua | ls. | aductive |
| | (1) | Estriol | 11 | | the repl | 04 | | nre-ror | oroductive |
| | (2) | Estradiol | | (2) | | | | | than the |
| | (3) | Ecdysone | | (3) | 1 | ative mu | | -19 | than the |
| | (4) | Epinephrine | 1 - 1 - 1 - 1 | | post-rep | ctive indi | individua | ls are r | nore than |
| | 2.05 | -pmephrine | · · · · · · · · · · · · · · · · · · · | (4) | pre-repr | oductive oductive i | ndividua | ls. | nore than |
| 53 | Whi | ich of the follow: | 4 | Ŭ | the repr | oaucuro | | T wit | h those in |
| | inc | orrectly paired with | structures or regions is | 57. Mat | ch the ite | ms given | in Colun | rect opt | h those in tion give |
| | (1) | | its function ? | Con | umn II a | nd select | the corr | 200 -1 | |
| | | Corpus callosum | : band of fibers | belo | w: | | C | olumn I | I . |
| | | | connecting left and | S | Column | | i. UV | B radia | ation |
| | | | right cerebral hemispheres. | a. | Eutroph | | 1. De | forestat | ion |
| | (2) | Hypothalamus | | b. | | y landfill | ii. De iii. Nu | | |
| | | 4 | : production of releasing hormones | c. | Snow bl | indness | | ichmen | t |
| | | | and regulation of | d. | | | | ste disp | X |
| | | | temperature, | α. | A.S. 1993. | ultivation | d | ste uisp | |
| | (3) | Limbia | hunger and thirst. | (1) | a t | | a iii | | * |
| | (0) | Limbic system | : consists of fibre | (1) | i i iii i | 0 | | | 1 |
| | | 5 S | tracts that interconnect | | | v i ii iv | ii | - *(| |
| | 1 | 19 | different regions of | (4) | i i | | ii | 9 | |
| | | 19 g - 1 | brain; controls | 1.900 | | | iv | | |
| | (18 | | movement. | 58. Whi | ich part o | f poppy p | lant is u | sed to o | btain the |
| | (4) | Medulla oblongata | | and sh | g Smack | ~? | | | |
| | | | and cardiovascular reflexes. | (2) | Leaves Roots | 1.1 | | | |
| ~ | | 8 | renexes. | (3) | Latex | 1 | ÷ | | |
| 4.) | Whi | ch of the following | g hormones can play a | | Flowers | A | | | £ |
| | sign | ificant role in osteop | porosis? | | | • | | | 1. |
| | (1) | Parathyroid hormo | one and Prolactin | 59. Wh | nich one | of th | e follov | ving p | opulation |
| | (2) | Estrogen and Para | | th | e producti | is widely i on of antib | icod in m | edical s | opulation cience for |
| | (3) | Progesterone and A | | (1) | | | lotics ? | | |
| | (4) | Aldosterone and Pr | | (2) | | 212.1 | | 5 | |
| | | | | (3 | 1. | Sec. 4. 16 | | | |
| 5. | The | transparent lens in | the human eye is held in | | | nensalism | | | |
| | its p | lace by | the numan eye is held in | | | | | | |
| 3 | (1) | | | 60) A | ll of the | following | g are incl | uded in | Ex-situ |
| .+ | 1.16 | | tached to the ciliary body | | | on oncept | | | -a-siti |
| | (2) | smooth muscles att | | (1 | | banks | | | |
| | (3) | ligaments attached | to the iris | (2 | | nical gard | ens | | |
| | (4) | ligaments attached | to the ciliary body | (3 | 3) Sacre | ed groves | | | |
| _ | - | /Page 6 | 0, - , | (4 | 4). Wild | life safari | parks | | |

| 61 | | |
|------|---|---|
| 91, | Which of the following gastric cells indirectly help in erythropoiesis ? | 65. AGGTATCGCAT is a sequence from the coding what will be the corresponding |
| | (1) a guarde cells indirectly | 65. AGGTATCGCAT is a sequence from the coding strand of a gene. What will be the corresponding strand of the transcribed mRNA? |
| - | (o) cells | 65. AGGTATO sequence of the transcribed mRNA? |
| | Goblet coll | (1) UCCAUAGCGUA |
| | (3) Mucous cells | a contal Good |
| | | UCCTUTCGOAL |
| 62. | (4) Chief cells | TICGCAU |
| 1 | Match | |
| | Match the items given in Column I with those in Column II and select the <i>correct</i> option given Column I | (4) AGGUAUCCOULT 66. A woman has an X-linked condition on one of her X chromosomes. This chromosome can be X chromosome can be a chromosome. |
| | below: | X chromosomes. |
| | | inherited by (1) Both sons and daughters |
| | | (1) Both sons and date |
| | a. Fibrinogen i. Osmail | (2) Only grandchildren |
| | b. Globulin i. Osmotic balance | (a) Only sons |
| | c. Albumin | 1 archters |
| | iii. Defense | Column with those I |
| | a h | 67. Match the items given in Column 1 when Column II and select the <i>correct</i> option give |
| | (1) ji c | Column II and select the |
| | $(2) i \qquad i$ | below: Column II |
| | (3) i ii | Column |
| | ii) | a. Proliferative Phase i. Breakdown of |
| N. | ii ii | endometrial |
| 3) | | lining |
| / | Calcium is important in skeletal | b. Secretory Phase ii. Follicular Phase |
| | contraction because it | iii Luteal Phase |
| | (1) prevents the | c. Menstruation III. Latera - |
| | (1) prevents the formation of bonds between the myosin cross bridges and the | a b c |
| \$ | the myosin cross bridges and the actin filament. | (1) jii i ii |
| | (2) detaches 4 | (2) ii iii i |
| | (2) detaches the myosin head from the actin filament. | (3) i iii ii |
| 1811 | filament. | |
| | (8) activates the muccin Am | (4) iii ii i |
| | (3) activates the myosin ATPase by binding to it. | 68. According to Hugo de Vries, the mechanism |
| | | avalution is |
| | (4) binds to troponin to remove the masking of | (1) Minor mutations |
| | active sites on actin for myosin. | |
| | | |
| | Which of the following is an occupational | (3) Saltation |
| | respiratory disorder ? | (4) Multiple step mutations |
| | 승규가 가장님께서 가장을 즐기 같다. 것이 같아요. 이 것이 같아요. | 69. All of the following are part of an operon except |
| | (1) Emphysema | (1) a promoter |
| | (2) Botulism | (2) an enhancer |
| 9 | (3) Silicosis | (3) structural genes |
| | (4) Anthracis | (4) an operator |

| To wait a cut out out on the section of the | 73. Hormones secreted by the placenta to maintain |
|--|--|
| 70. Which of the following options correctly represents the lung conditions in asthma and emphysema, respectively? | pregnancy are |
| (1) Decreased respiratory surface; Inflammation of bronchioles | hCG, progestogens, estrogens, ¥ glucocorticoids |
| (2) Increased respiratory surface; Inflammation of bronchioles (3) Increased number of bronchioles; Increased respiratory surface (4) Inflammation of bronchioles; Decreased | (2) hCG, hPL, progestogens, estrogens (3) hCG, hPL, estrogens, relaxin, oxytocin (4) hCG, hPL, progestogens, prolactin |
| respiratory surface71. Match the items given in Column I with those in Column II and select the <i>correct</i> option given below : | 74. The contraceptive 'SAHELI' (1) is a post-coital contraceptive. (2) is an IUD. (3) increases the concentration of estrogen an |
| Column IColumn IIa.Tricuspid valvei.Between left atrium and left ventricleb.Bicuspid valveii.Between right ventricle and | prevents ovulation in females. (4) blocks estrogen receptors in the uterus preventing eggs from getting implanted. |
| c. Semilunar valve iii. Between right atrium and right ventricle (1) ii i iii | The difference between spermiogenesis an spermiation is (1) In spermiogenesis spermatozoa are formed while in spermiation spermatozoa ar released from sertoli cells into the cavity or in the sertoli cells into the sertoli celli |
| (2) i ii iii (3) i iii ii • (4) iii i ii • c_{1} column I with those in | seminiferous tubules. (2) In spermiogenesis spermatozoa from sertol cells are released into the cavity of seminiferous tubules, while in spermiation spermatozoa are formed. |
| Column II and select the correct option g below: Column I i., 2500 - 3000 mL | (3) In spermiogenesis spermatozoa are formed while in spermiation spermatids are formed. |
| a. Tidal volume b. Inspiratory Reserve . ii. 1100 - 1200 mL volume c. Expiratory Reserve iii. 500 - 550 mL | (4) In spermiogenesis spermatids are formed while in spermiation spermatozoa are formed. |
| volume d. Residual volume \mathbf{a} \mathbf{b} \mathbf{c} \mathbf{d} | from |
| (1) iv iii ii i ● (2) i iv ii iii | (1) ectoderm and endoderm (2) mesoderm and trophoblast (2) definition of the second arm |
| (3) iii. i iv ii (4) iii ii i iv | (3) endoderm and mesoderm(4) ectoderm and mesoderm |

| 1.00 | | - | | | | | | - 10 00 | |
|---------------------|---|---------------|-------------|-------------|-----------------|----------|-------------|------------------------|----------------------------|
| Which | of the following animals does <i>not</i> undergo morphosis ? | | | | - | - | | | |
| (1) | norphosis ? | 83. | Mato | h the | items | river ! | - Col | umn I w | ith those i |
| 101 | Starfish | 11- | Colu | mn II | and s | select 4 | ho c | orrect 0 | ption give |
| 101 | Moth | 1.1 | below | w : | und c | beleet (| the c | onneer | |
| | Tunicate | | | Colun | nn I | | Co | lumn II | |
| | Earthworm | í | | | | | | | furic |
| • Which home | h one of these animals is not a | | a. | Glyco | suria | i. | | umulation in joints | |
| (1) ' | Psittacula | | þ. | Gout | | ii. | | s of cryst s within | tallised the kidney |
| (2) | Camelus | | c. | Renal | calcul | : ::: | Infl | ammatio | n in |
| • (3) | Chelone | | | nena | calcul | 1 111. | | neruli | |
| (4) | Macropus | 4 2 | d. ' | Glom | erular ritis | iv. | Pre urir | | glucose in |
| ** 111 | ch of the following features is used to identify ale cockroach from a female cockroach 2 | | | a | b | . c | | | |
| (1) | ale cockroach from a female cockroach ? Presence of another | | (1) | iv | i | ii | | 3 | |
| (2) | - of allal Cerci | | (2) | ii | | | | ii | |
| ~ (3) | Forewings with darker tegmina | 10. | (3) | | iii | i | i | v | |
| (4) | caudal styles | | 1.1 | i | ii | iii | i | v | |
| | Presence of a boat shaped sternum on the 9 th abdominal segment | | (4) | iii | ii , | iv | i | | |
| 0. Wh chi (1) | ich of the following organisms are known a ef producers in the oceans ? Euglenoids | s | Col belo | ow: | and | select | the c | correct o | ith those i ption give |
| (2) | Cyanobacteria | 1 | | Colu | | | | Colum | n II " |
| • (3) | Diatoms · | 1 | | (Fun | ection) | | | | Excretory |
| (4) | Dinoflagellates | 4 | a. | Πtr | afiltrat | | | System) | |
| 1. Cil | iates differ from all other protozoans in | | | | | | i. | Henle's | loop |
| (1) | naving two types of nuclei | | b. | Con of u | centrat rine | tion | / ii. | Ureter | |
| e (2) | using pseudopodia for capturing prey | 1 | c. | Tra | nsport | of | | G | 6 |
| (3) | having a contractile vacuale for more | | | uri | ne | | (m. | Urinary | bladder |
| | encess water | ng | đ | . St | orage o | f urine | | | |
| (4) | using flagella for locomotion | | | | 0.1 | · | 10 | Malpight | le . |
| cn | entify the vertebrate group of animaracterized by crop and gizzard in its dige stem. | mals stive | | | | | | . Proxima convolu | u ^{ted} tubule |
| sy (1) | | | 30 | 1.22 | a | b | c | d | 100 |
| | , | 1 14 | | (1) | v | iv | i | iii≯ | |
| (2) | | | | (2) | v | iv | i | ii x | |
| (3 |) Reptilia | | 1 | • (3) | iv | i | | | |
| • (4 |) Amphibia | | 1 | | | | ü | iii | |
| | CC/Page 9 SPAC | | | (4) | iv | v | ii | iii | |

13

I THE O

2.1

| Complete for 01 | A carbon resistor of (47 ± 4.7) k Ω is to be marked |
|---|--|
| | with rings of different colours for its |
| divergent evolution, select the mean of the | identification. The colour code sequence will be |
| | |
| (7) Dian or set | (1) Green - Orange - Violet - Gold |
| (3) Heart of bat, man and cheetah | (2) Yellow - Green - Violet - Gold |
| | (3) Yellow - Violet - Orange - Silver |
| Conversion of milk to curd improves its | (4) Violet - Yellow - Orange - Silver |
| 92 | A set of 'n' equal resistors, of value 'R' each, are battery of emf 'E' and |
| | A set of 'n' equal resistors, of value connected in series to a battery of emf 'E' and D' The current drawn is I. |
| | connected in series to a battery of drawn is I. internal resistance 'R'. The current drawn is I. |
| (3) Vitamin A | internal resistance 'R'. The current in parallel to Now, the 'n' resistors are connected in parallel to the same battery. Then the current drawn from the same battery. Then the current drawn from |
| • (4) Vitamin D | the same bottomy Then the ourse |
| 87. Which of the following characteristics represent 'Inheritance of blood groups' in humans ? | battery becomes 10 I. The value of 'n' is (1) 9 $V \leq 16$ |
| a. Dominance | (2) 20 $\left(\frac{\partial}{\gamma_{TR}}\right)R$ |
| b. Co-dominance 🖌 | (3) 11 |
| c. Multiple allele | * (4) 10 |
| a. incompiete dominance | |
| e. Polygenic inheritance 93. | A battery consists of a variable number 'n' of |
| (1) a, c and e \succ (2) b, d and e | identical cells (having internal resistance r |
| (3) a, b and c \times | each) which are connected in series. The |
| • (4) b, c and e | terminals of the battery are short-circuited and |
| | the current I is measured. Which of the graphs shows the correct relationship between I and n? |
| 88. Which of the following is <i>not</i> an autoimmune disease? | shows the correct relationship between 1 and 1 |
| (1) Vitiligo | |
| (2) Alzheimer's disease | (1) |
| (3) Rheumatoid arthritis | $0 \rightarrow n$ |
| • (4) Psoriasis | 241 |
| | τΙ |
| 89. The similarity of bone structure in the forelimbs of many vertebrates is an example of | (2) |
| (1) Adaptive radiation | $0 \rightarrow n$ |
| (2) Convergent evolution | |
| (3) Analogy | a menor of the second |
| (4) Homology | I |
| 90. In which disease does mosquito transmitted | (3) 1 |
| pathogen cause chronic inflammation of lymphatic vessels ? | $0 \rightarrow n$ |
| (1) Amoebiasis | |
| (2) Ringworm disease | I |
| (3) Ascariasis | |
| (d) Elephantiasia | |

- •(4) Elephantiasis
- ACHLA/CC/Page 10

SPACE FOR ROUGH WORK

51 Roy. 45 VALO

0

>n

English


The moment of the force, $\overrightarrow{F} = 4\hat{i} + 5\hat{j} - 6\hat{k}$ at (2, 0, -3), about the point (2, -2, -2), is given by -GThe power radiated by a black body is P and it 98. radiates maximum energy at wavelength, λ_0 . If (1) $-7\hat{i} - 4\hat{j} - 8\hat{k}$ the temperature of the black body is now (2) $-7\hat{i} - 8\hat{j} - 4\hat{k}$ 121 changed so that it radiates maximum energy at (3) $-4\hat{i} - \hat{j} - 8\hat{k}$ wavelength $\frac{3}{4}\lambda_0$, the power radiated by it (4) $-8\hat{i} - 4\hat{j} - 7\hat{k}$ becomes nP. The value of n is P= wxt. A student measured the diameter of a small steel 81 (1) $\mathcal{E} = \frac{\omega}{F}$ 99. ball using a screw gauge of least count 256 0.001 cm. The main scale reading is 5 mm and hu E=wxt2 zero of circular scale division coincides with 256(2)25 divisions above the reference level. If screw 81 gauge has a zero error of -0.004 cm, the correct E=pt. (3)diameter of the ball is he = P At Pxi 0.529 cm (1)0.053 cm (2)• (4) 0.525 cm (3)0.521 cm (4)95. Two wires are made of the same material and A toy car with charge q moves on a frictionless 100. have the same volume. The first wire has horizontal plane surface under the influence of a cross-sectional area A and the second wire has uniform electric field E . Due to the force q E, cross-sectional area 3A. If the length of the first its velocity increases from 0 to 6 m/s in one wire is increased by Δl on applying a force F, how much force is needed to stretch the second second duration. At that instant the direction of wire by the same amount ? the field is reversed. The car continues to move . (1) F for two more seconds under the influence of this (2)4 Ffield. The average velocity and the average speed of the toy car between 0 to 3 seconds are (3) 6 FF= M a = 9 9 F respectively (4)(1)1.5 m/s, 3 m/s $= \frac{dv}{dt} = 2E_{c}$ = dv =A sample of 0.1 g of water at 100°C and normal 96. (2)1 m/s, 3.5 m/s pressure $(1.013 \times 10^5 \text{ Nm}^{-2})$ requires 54 cal of (3)1 m/s, 3 m/s (4)2 m/s, 4 m/sheat energy to convert to steam at 100°C. If the A block of mass m is placed on a smooth inclined 101. volume of the steam produced is 167.1 cc, the wedge ABC of inclination θ as shown in the figure. The wedge is given an acceleration 'a' change in internal energy of the sample, is towards the right. The relation between a and θ (1)84.5 J for the block to remain stationary on the wedge (2)42.2 J is (3)208.7 J (4)104·3 J A small sphere of radius 'r' falls from rest in a 97. viscous liquid. As a result, heat is produced due to viscous force. The rate of production of heat when the sphere attains its terminal velocity, is $a = g \tan \theta$ (1)proportional to • (2) $a = g \cos \theta$ (1) r^4 $a = \frac{g}{\sin \theta}$ (3)r⁵ (2)(3) $a = \frac{g}{cosec 0}$ (4)'(4)

ACHLA/CC/Page 11

SPACE FOR ROUGH WORK

 $\frac{AC}{AA} = t \qquad \Rightarrow \frac{P}{P} = \frac{Y}{3} p$

English



- 106. The ratio of kinetic energy to the total energy of 102. An em wave is propagating in a medium with a velocity $\overrightarrow{V} = V \overrightarrow{i}$. The instantaneous oscillating electric field of this em wave is along +y axis. Then the direction of oscillating magnetic field of the em wave will be along
 - x direction (1)
 - y direction (2)
 - + z direction (3)
 - z direction . (4)

103. The refractive index of the material of a prism is $\sqrt{2}$ and the angle of the prism is 30°. One of the

two refracting surfaces of the prism is made a mirror inwards, by silver coating. A beam of monochromatic light entering the prism from the other face will retrace its path (after reflection from the silvered surface) if its angle of incidence ADDES=1+e

- on the prism is
- zero (1)
- 30° (2)
- 45° (3)
- 60° (4)

104. The magnetic potential energy stored in a certain inductor is 25 mJ, when the current in the inductor is 60 mA. This inductor is of inductance

- 13·89 H (1)
- 1.389 H (2)
- 138·88 H (3)
- 0.138 H (4)


105. An object is placed at a distance of 40 cm from a concave mirror of focal length 15 cm. If the object is displaced through a distance of 20 cm towards the mirror, the displacement of the image will be

- 36 cm towards the mirror (1)
- 30 cm towards the mirror (2)
- 36 cm away from the mirror • (3)
 - 30 cm away from the mirror (4)

an electron in a Bohr orbit of the hydrogen atom, is 1:-2(1)

- 2:-1(2)
- 1:-1(3)
- 1:1 (4)

107. An electron of mass m with an initial velocity $\vec{V} = V_0 \hat{i} (V_0 > 0)$ enters an electric field $\overrightarrow{E} = -E_0 \overrightarrow{i}$ (E₀ = constant > 0) at t = 0. If λ_0 is its de-Broglie wavelength initially, then its de-Broglie wavelength at time t is



108. When the light of frequency $2v_0$ (where v_0 is threshold frequency), is incident on a metal plate, the maximum velocity of electrons emitted is v_1 . When the frequency of the incident radiation is increased to $5v_0$, the maximum velocity of electrons emitted from the same plate is v2. The ratio of v1 to v2 is

- (1) 2:1
- 4:1 (2)
- (3)1:4
- (4)1:2
- 109. a. radioactive For material, half-life is 10 minutes. If initially there are 600 number of nuclei, the time taken (in minutes) for the disintegration of 450 nuclei is

(1) 15
$$10^{-} = 60^{-}$$

- (2)30 450.
- 10 (3)
- 20 (4)

VI = VI tat 45 SPACE FOR ROUGH WORK ACHLA/CC/Page 12 mv= -

The volume (V) of a monatomic gas varies with 110) its temperature (T), as shown in the graph. The ratio of work done by the gas, to the heat absorbed by it, when it undergoes a change from state A to state B, is

0 27 (1)1 (2)3 2 ;(3) 3 The fundamental frequency in an open organ 2 5 pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is 111. 16 cm (1) 12.5 cm The efficiency of an ideal heat engine working The efficiency of an inear near engine working between the freezing point and boiling point of 112. water, is 12.5% 6.25% (1) what temperature will the rms speed of molecules become just sufficient . (2) At what temperature will the rms speed of molecules become just sufficient for oxygen from the Earth's atmosphere 2 oxygen from the Earth's atmosphere? 113.

(Given . Mass of oxygen molecule (m) = 2.76×10^{-26} kg Mass Boltzmann's constant $k_B = 1.38 \times 10^{-23} J K^{-1}$

 $1.254 \times 10^4 \,\mathrm{K}$

 $5.016 \times 10^4 \text{ K}$ (1) $8.360 \times 10^4 \text{ K}$ (2)

 $2.508 \times 10^4 \text{ K}$ (3)

(4) INCC/Page 13 062

- 114. Unpolarised light is incident from air on a plane surface of a material of refractive index 'µ'. At a particular angle of incidence 'i', it is found that rays are refracted and reflected the perpendicular to each other. Which of the following options is correct for this situation ?
- (1) $i = \tan^{-1}\left(\frac{1}{n}\right)$ • (2) $i = \sin^{-1}\left(\frac{1}{n}\right)$
 - Reflected light is polarised with its electric vector perpendicular to the plane of (3)incidence
 - Reflected light is polarised with its electric
- vector parallel to the plane of incidence (4)


115. In Young's double slit experiment the separation d between the slits is 2 mm, the wavelength λ of the light used is 5896 Å and distance D between the screen and slits is 100 cm. It is found that the angular width of the fringes is 0.20° . To increase the fringe angular width to 0.21° (with same λ and D) the separation between the slits needs to

- be changed to
- 1.7 mm (1)
- 2.1 mm (2)
- 1.9 mm (3)
- 1.8 mm (4)

SPACE FOR ROUGH WORK

DY

- 116. An astronomical refracting telescope will have large angular magnification and high angular resolution, when it has an objective lens of small focal length and small diameter
 - (1) large focal length and large diameter
 - (2)large focal length and small diameter
 - (3) small focal length and large diameter • (4)
 - English 40 21 ×1 / 40



A metallic rod of mass per unit length 0.5 kg m⁻¹ is lying horizontally on a smooth inclined plane which makes an angle of 30° with the horizontal. The rod is not allowed to slide down by flowing a current through it when a magnetic field of induction 0.25 T is acting on it in the vertical direction. The current flowing in the rod to keep it stationary is

- 11.32 A
- 14.76 A
- 5.98 A
- 7.14 A
- 121. An inductor 20 mH, a capacitor 100 µF and a resistor 50 Ω are connected in series across a source of emf, $V = 10 \sin 314 t$. The power loss in the circuit is
 - 1.13 W
 - (2) 2·74 W
 - 0.43 W
 - 0:79 W
 - A thin diamagnetic rod is placed vertically between the poles of an electromagnet. When the current in the electromagnet is switched on, then the diamagnetic rod is pushed up, out of the horizontal magnetic field. Hence the rod gains gravitational potential energy. required to do this comes from The work
 - the induced electric field due to the changing magnetic field
 - the lattice structure of the material of the
 - the magnetic field
 - the current source

123. Current sensitivity of a moving coil galvanometer is 5 div/mA and its voltage sensitivity (angular deflection per unit voltage applied) is 20 div/V. The resistance of the galvanometer is

500 Ω 250 Ω 25 Q 40 Ω

Scanned by CamScanner

124. A tuning fork is used to produce resonance in a 128. glass tube. The length of the air column in this tube can be adjusted by a variable piston. At room temperature of 27°C two successive resonances are produced at 20 cm and 73 cm of column length. If the frequency of the tuning fork is 320 Hz, the velocity of sound in air at 27°C is

- (2)350 m/s
- (3) 339 m/s
- (4) 330 m/s

125. The electrostatic force between the metal plates of an isolated parallel plate capacitor C having a charge Q and area A, is .

- (1)inversely proportional to the distance between the plates.
- 2(2) proportional to the square root of the distance between the plates.
 - (3)linearly proportional to the distance between the plates.
 - (4) independent of the distance between the plates.


126. A pendulum is hung from the roof of a sufficiently high building and is moving freely to and fro like a simple harmonic oscillator. The acceleration of the bob of the pendulum is 20 m/s² at a distance of 5 m from the mean position. The time period of oscillation is

- (1)1s
- 2s(2)
- (3)πs
- $2\pi s$ (4)
- 127. An electron falls from rest through a vertical distance h in a uniform and vertically upward directed electric field E. The direction of electric field is now reversed, keeping its magnitude the same. A proton is allowed to fall from rest in it through the same vertical distance h. The time of fall of the electron, in comparison to the time of fall of the proton is
 - equal • (1)
 - 10 times greater (2)
 - (3)5 times greater
 - (4)smaller

ACHLA/CC/Page 15

The kinetic energies of a planet in an elliptical orbit about the Sun, at positions A, B and C are K_A , K_B and K_C , respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure.





- (1) $K_B > K_A > K_C$
- (2) $K_B < K_A < K_C$
- $K_A > K_B > K_C$ (3)
- $K_A < K_B < K_C$ (4)

129. A solid sphere is in rolling motion. In rolling motion a body possesses translational kinetic energy (K_t) as well as rotational kinetic energy (K_r) simultaneously. The ratio $K_t : (K_t + K_r)$ for

the sphere is

- 2:5 (1)
- 10:7 (2)
- 5:7 (3)
- (4)7:10

130. If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is not correct ?

- 'g' on the Earth will not change. (1)
- Time period of a simple pendulum on the 6(2) Earth would decrease.
 - Walking on the ground would become more (3)difficult.
- Raindrops will fall faster. A (4)
- 131. A solid sphere is rotating freely about its symmetry axis in free space. The radius of the sphere is increased keeping its mass same. Which of the following physical quantities would remain constant for the sphere?
 - (1) Angular momentum
 - Rotational kinetic energy (2)
 - Moment of inertia (3)
 - Angular velocity •(4)

English

SPACE FOR ROUGH WORK

| 132. A body initially at rest and sliding along a frictionless track from a height h (main along a | |
|--|---|
| the figure) just completes a vertical circle | (1) dinuclear |
| $\mathcal{U} = \mathbf{D}. \text{ The height h is equal to} \\ \mathcal{U} = \mathbf{O}$ | (2) trinuclear (3) mononuclear |
| h B u=0 | (4) tetranuclear |
| | 137. Match the metal ions given in Column I with spin magnetic moments of the ions given Column II and assign the correct code. |
| (1) $\frac{5}{4}$ D | Column II and assign the correct code : Column I Column II |
| 2 | a. Co^{3+} i. $\sqrt{8}$ B.M. |
| $(2) \frac{7}{5} D \qquad $ | b. Cr^{3+} ii. $\sqrt{35}$ B.M. |
| (3) D | c. Fe^{3+} iii. $\sqrt{3}$ B.M. |
| (4) $\frac{3}{2}$ D | d. Ni ²⁺ iv. $\sqrt{24}$ B.M. |
| 133. Three objects, A : (a solid sphere), B : (a thin | · · · √15 B M |
| and USK) and C: (a circular ring) each have | |
| the same mass M and radius R. They all spin with the same angular speed ω about their own | |
| symmetry axes. The amounts of work (W) | |
| required to bring them to rest, would satisfy the relation | |
| (1) $\dot{W}_{A} > W_{C} > W_{B}$ | |
| $(2) W_{\rm B} > W_{\rm A} > W_{\rm C}$ | |
| • (3) $W_A > W_B > W_C$ | 138. Which one of the following ions exh d-d transition and paramagnetism as well? |
| $(4) W_{C} > W_{B} > W_{A}$ | 0 (1) MnO ₄ ²⁻ + 2 24 S |
| | x (2) MO-45 X-8=-2 |
| 134. Which one of the following statements is <i>incorrect</i> ? | $(3) \operatorname{Cr}_2 O_7^{2-} 2$ |
| (1) Coefficient of sliding friction has dimensions of length. | (4) CrO_4^{2-} |
| • (2) Frictional force opposes the relative motion. | |
| (3) Limiting value of static friction is directly proportional to normal reaction. | 139. The geometry and magnetic behaviour of complex $[Ni(CO)_4]$ are |
| (4) Rolling friction is smaller than sliding | (1) tetrahedral geometry and paramagnetic |
| friction. 135. A moving block having mass m, collides with | (2) square planar geometry and paramagnetic (3) tetrahedral geometry and diamagnetic (4) square planar geometry and diamagnetic |
| another stationary block having mass 4m. The | square planar geometry and diamaged |
| lighter block comes to rest after collision. When the initial velocity of the lighter block is v, then the value of coefficient of restitution (e) will be | [$CoCl_2(en)_2$] is |
| (1) 0.4 | (1) Linkage isomerism |
| (2) 0.8 | (2) Ionization isomerism |
| (3) 0.25 | soordination isomerism |
| (4) 0·5 | acometrical isomerism |
| ACHLA/CC/Page 16 SPACE FOR R | $eq - Cl = 2x - 14 = -2^{En}$ 7 = 0 $x = 6$ |
| 6+2) x-8 =-2 Ro | - co - Cl 2x-14 = -2 -" |

141. Identify the major products P, Q and R in the 143. For the redox reaction following contract following contract for the redox reaction for the redox red following sequence of reactions :

$$\begin{array}{c} \begin{array}{c} \begin{array}{c} \text{Anhydrous} \\ \text{AlCl}_3 \end{array} \\ \end{array} \\ \begin{array}{c} \text{AlCl}_3 \end{array} \end{array} \\ \begin{array}{c} P \xrightarrow{\text{(i) } O_2} \\ \hline \text{(ii) } H_3 O^{+/\Delta} \end{array} Q + R \end{array}$$

R

(1)
$$(1)$$
 (1)

໑

(2)
$$(2)$$
 (2)

(3)
$$(3)$$
 (3)

(4)
$$(4)$$
 (4)

142. Which of the following compounds can form a zwitterion ?

Glycine (1)

P

- Benzoic acid (2)
- Acetanilide (3)
- Aniline (4)

SPACE FOR ROUGH WORK

ACHLA/CC/Page 17

 $q MnO_4^- + C_2O_4^{2-} + H^+$ the correct coefficients of the reactants for the balanced equation are $C_2 O_4^{2-}$ H^{+} MnO₄ 2 16 5 (1) 5 16 . 2 (2)16 5 2 (3)2 5 144. The correction factor 'a' to the ideal gas equation 16 gas corresponds to attraction between the forces of (1) molecules gas the electric field present between (2)molecules volume of the gas molecules (3)density of the gas molecules (4) 145. Which one of the following conditions will favour maximum formation of the product in the reaction, $A_2(g) + B_2(g) \rightleftharpoons X_2(g) \Delta_r H = -X kJ?$ High temperature and low pressure (1)High temperature and high pressure (2)Low temperature and low pressure (3)Low temperature and high pressure (4)146. The bond dissociation energies of X_2 , Y_2 and XYare in the ratio of 1:0.5:1. ΔH for the formation of XY is -200 kJ mol^{-1} . The bond dissociation energy of X2 will be

5


 $\xrightarrow{d} Mn^{2+} \stackrel{e}{+} CO_2 + H_2O$

- 400 kJ mol⁻¹ (1)
- 800 kJ mol⁻¹ (2)
- 100 kJ mol^{-1} (3)200 kJ mol⁻¹

(4)

- 147. When initial concentration of the reactant is doubled, the half-life period of a zero order reaction
 - (1)remains unchanged
 - (2)is tripled
 - (3)is doubled
 - .(4) is halved

- 148. Which of the following molecules represents the order of hybridisation sp^2 , sp^2 , sp, sp from left to right atoms ?
 - (1) $CH_3 CH = CH CH_3$
 - (2) $CH_2 = CH CH = CH_2$
 - (3) $CH_2 = CH - C \equiv CH$
 - $HC \equiv C C \equiv CH$ (4)
- 149. Which of the following carbocations is expected to be most stable ?



- (4)
- 150. Which of the following is correct with respect to - I effect of the substituents ? (R = alkyl)
 - (1) $-NR_2 > -OR > -F$
 - (2) $-NH_2 > -OR > -F$
 - (3) $-NR_2 < -OR < -F$
 - **b** (4) $\mathrm{NH}_2 < \mathrm{OR} < -\mathrm{F}^+$

ACHLA/CC/Page 18

correct difference between first-151. The and second-order reactions is that

- the rate of a first-order reaction does (1)depend on reactant concentrations; the rate of a second-order reaction does not depend on reactant concentrations
- a first-order reaction can be catalyzed; a (2)second-order reaction cannot be catalyzed
- the half-life of a first-order reaction does not X(3) depend on [A]0; the half-life of a second-order reaction does depend on [A]0
 - the rate of a first-order reaction does not (4) depend on reactant concentrations; the rate of a second-order reaction does depend on reactant concentrations.
- 152. Among CaH₂, BeH₂, BaH₂, the order of ionic character is
 - (1) $BaH_2 < BeH_2 < CaH_2$
 - (2) $BeH_2 < BaH_2 < CaH_2$
 - (3) $CaH_2 < BeH_2 < BaH_2$
 - (4) $BeH_2 < CaH_2 < BaH_2$
- 153. Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below :

Then the species undergoing disproportionation

- (1)HBrO
- (2) Br₂
- (3)Bro-
- (4) BrO-

154. In which case is the number of molecules of water

- 10⁻³ mol of water (1)
- (2)
- 0.00224 L of water vapours at 1 atm and
- (3)0.18 g of water
- 18 mL of water (4)

SPACE FOR ROUGH WORK

155. and with PCl₅ gives C. B and C react together to give diethyl ether. A, B and C are in the order A(1) $\mathrm{C_{2}H_{5}OH,\,C_{2}H_{5}ONa,\,C_{2}H_{5}Cl}$ (2)

(3) C H
$$_{2}H_{5}Cl, C_{2}H_{6}, C_{2}H_{5}OH$$

- $\mathrm{C_{2}H_{5}OH,\,C_{2}H_{5}Cl,\,C_{2}H_{5}ONa}$
- (4) C_2H_5OH , C_2H_6 , C_2H_5Cl

156. Hydrocarbon (A) reacts with bromine substitution to form an alkyl bromide which by reaction is converted hydrocarbon containing less than four carbon

- (1)CH4
- $CH_3 CH_3$ D(2)

 $CH_2 = CH_2$ (3)

- (4) $CH \equiv CH$

157. The compound C_7H_8 undergoes the following

 $C_7H_8 \xrightarrow{3 Cl_2/\Delta} A \xrightarrow{Br_2/Fe} B \xrightarrow{Zn/HCl} C$ The product 'C' is

- (1)*p*-bromotoluene
- 3-bromo-2,4,6-trichlorotoluene (2)
- (3)o-bromotoluene
- (4)*m*-bromotoluene

- 158. Which oxide of nitrogen is not a common 163. Regarding cross-linked or network polymers, which of the following statements is incommon is in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statements is incommon in the statement of the following statement of the foll due to natural and human activity?
 - (1) NO. (2) N_2O
 - (3)NO2
 - N_2O_5 (4)

ACHLA/CC/Page 19

SPACE FOR ROUGH WORK

The compound A on treatment with Na gives B, 159. A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc. H2SO4. The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be

- (1)4.4
- (2)2.8
- (3)3.0
- (4)1.4
- 160. The difference between amylose and amylopectin is
 - Amylose is made up of glucose and (1)galactose
 - (2) Amylopectin have $1 \rightarrow 4 \alpha$ -linkage and $1 \rightarrow 6 \beta$ -linkage
 - (3)Amylose have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6 \beta$ -linkage
 - (4) Amylopectin have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6 \alpha$ -linkage
- 161. Which of the following oxides is most acidic in nature?
 - (1)CaO
 - (2)BaO
 - (3)BeO (4)
 - MgO

162. Nitration of aniline in strong acidic medium also gives m-nitroaniline because (1)

- In acidic (strong) medium aniline is present as anilinium ion. (2)
- In absence of substituents nitro group always goes to m-position. (3)In
- electrophilic substitution amino group is meta directive. reactions (4)
- In spite of substituents nitro group always

- which of the following statements is *incorrect*? • (1)
 - They contain strong covalent bonds in their (2)
 - Examples are bakelite and melamine. (3)
 - They are formed from bi- and tri-functional
 - They contain covalent bonds (4)between various linear polymer chains.

- 164. Following solutions were prepared by mixing 168. Magnesium reacts with an element (X) to form an different volumes of NaOH and HCl of different concentrations :
 - $60 \text{ mL} \frac{\text{M}}{10} \text{ HCl} + 40 \text{ mL} \frac{\text{M}}{10} \text{ NaOH}$ a. 55 mL $\frac{M}{10}$ HCl + 45 mL $\frac{M}{10}$ NaOH b.
 - 75 mL $\frac{M}{5}$ HCl + 25 mL $\frac{M}{5}$ NaOH c.
 - 100 mL $\frac{M}{10}$ HCl + 100 mL $\frac{M}{10}$ NaOH d.
 - pH of which one of them will be equal to 1?
 - (1)C.
 - d (2)
 - (3)a
 - (4)b
 - 165. On which of the following properties does the coagulating power of an ion depend ?
 - The sign of charge on the ion alone (1)
 - Both magnitude and sign of the charge on (2)the ion
 - Size of the ion alone (3)
 - The magnitude of the charge on the ion (4)alone
 - solubility of BaSO₄ in water is 166. The 2.42×10^{-3} gL⁻¹ at 298 K. The value of its solubility product (K_{sp}) will be (Given molar mass of $BaSO_4 = 233 \text{ g mol}^{-1}$)
 - $1.08 imes 10^{-8} \, \mathrm{mol}^2 \, \mathrm{L}^{-2}$ (1)
 - $1.08 imes 10^{-14} ext{ mol}^2 ext{ L}^{-2}$ (2)
 - (3) $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$
 - $1.08 imes 10^{-10} ext{ mol}^2 ext{ L}^{-2}$. (4)

167. Given van der Waals constant for NH_3 , H_2 , O_2 and CO_2 are respectively 4.17, 0.244, 1.36 and 3.59, which one of the following gases is most


- easily liquefied ? NH3, CO2, O2, H2 CO_2 (1)02 (2)(3) H₂
 - (4) NH₃

ACHLA/CC/Page 20

\$40-23 + 230 + 23 × 10 SPACE FOR ROUGH WORK

ionic compound. If the ground state electronic configuration of (X) is $1s^2 2s^2 2p^3$, the simplest formula for this compound is

- Mg₃X₂ (1)
- Mg₂X (2)
- MgX₂ (3)
- Mg₂X₃ (4)
- 169. Iron exhibits bcc structure at room temperature. Above 900°C, it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature)



Which one is a *wrong* statement? 170.

The value of m for d_{z^2} is zero. (1)

- The electronic configuration of N atom is (2)
 - $2s^2 = \frac{2p_x^1 2p_y^1 2p_z^1}{2p_x^1 2p_y^1 2p_z^1}$ 1↓
- An orbital is designated by three quantum (3)numbers while an electron in an atom is designated by four quantum numbers.
- Total orbital angular momentum of electron (4)in 's' orbital is equal to zero.

171. Consider the following species :

CN⁺, CN⁻, NO and CN

Which one of these will have the highest bond order ?

- ·(1) CN (2) CN^+
- (3)CN⁻
- (4)NO

4210-19

English

172. Which of the following statements is not true for 178. In the reaction O-Na⁺ Chlorine has the highest electron-gain CHO OH + CHCl3 + NaOH (2)All but fluorine show positive oxidation 0 the electrophile involved is (3)All are oxidizing agents. e(4) All form monobasic oxyacids. dichlorocarbene (:CCl₂) 173. Which one of the following elements is unable to (1)dichloromethyl anion (CHCl₂) form MF₆³⁻ion? (2)•(1) . In (3) formyl cation (CHO) (2)B dichloromethyl cation (CHCl₂) (3)(4)Al (4)Ga 179. Carboxylic acids have higher boiling points than 174. In the structure of ClF_3 , the number of lone pairs aldehydes, ketones and even alcohols of of electrons on central atom 'Cl' is comparable molecular mass. It is due to their (1)three formation of intermolecular H-bonding (1) (2)four (2)more extensive association of carboxylic L(3) two acid via van der Waals force of attraction (4)one (3)formation of carboxylate ion 175. Considering Ellingham diagram, which of the formation of intramolecular H-bonding (4)following metals can be used to reduce alumina? (1)Cu 180. Compound A, C₈H₁₀O, is found to react with •(2) Mg NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic Zn (3)* (4) Fe A and Y are respectively 176. The corréct order of atomic radii in group 13 elements is (1) B < Ga < Al < In < Tl(1) CH₃ OH and I2 B < Ga < Al < Tl < In(2)(3) B < Al < Ga < In < TlB < Al < In < Ga < Tl(2) $H - CH_3$ and I_2 (4)177. The correct order of N-compounds in its OH decreasing order of oxidation states is (3) $CH_2 - CH_2 - OH$ and I_2 NH4Cl, N2, NO; HNO3 (1)HNO3, NH4Cl, NO, N2 (2)H₃C HNO3, NO, NH4Cl, N2 (4) $CH_2 - OH and I_2$ (3)HNO3, NO, N2, NH4Cl SPACE FOR ROUGH WORK (4)ACHLA/CC/Page 21 English