
NEET QUESTION PAPER (2018) BOOKLET CODE - HLAAC (KK)

orbit about the Sun, at positions A, B and C are K_A, K_B and K_C, respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure. Then



- (1) $K_A > K_B > K_C$
- (2) $K_B < K_A < K_C$
- (3) $K_A < K_B < K_C$
- (4) $K_B > K_A > K_C$
- 2. A solid sphere is in rolling motion. In rolling motion a body possesses translational kinetic energy (K_t) as well as rotational kinetic energy (K_r) simultaneously. The ratio $K_t:(K_t+K_r)$ for the sphere is
 - (1) 5:7 ,
 - (2) 10:7
 - (3) 7:10
 - (4) 2:5
- 3. A solid sphere is rotating freely about its symmetry axis in free space. The radius of the sphere is increased keeping its mass same. Which of the following physical quantities would remain constant for the sphere?
 - (1) Moment of inertia
 - (2) Rotational kinetic energy
 - (3) Angular velocity
 - (4) Angular momentum.
- If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is **not** correct?
 - Walking on the ground would become more difficult.
 - (2) Time period of a simple pendulum on the Earth would decrease.
 - (3) Raindrops will fall faster.
 - (4) 'g' on the Earth will not change.

- A toy car with charge q moves on a frictionless horizontal plane surface under the influence of a uniform electric field E. Due to the force qE, its velocity increases from 0 to 6 m/s in one second duration. At that instant the direction of the field is reversed. The car continues to move for two more seconds under the influence of this field. The average velocity and the average speed of the toy car between 0 to 3 seconds are respectively
- (1) 1 m/s, 3 m/s
- (2) 1 m/s, 3·5 m/s
- (3) 2 m/s, 4 m/s
- (4) 1.5 m/s, 3 m/s
- 6. A block of mass m is placed on a smooth inclined wedge ABC of inclination θ as shown in the figure. The wedge is given an acceleration 'a' towards the right. The relation between a and 0 for the block to remain stationary on the wedge is




- (1) $a = \frac{g}{\sin \theta}$
- (2) $a = g \cos \theta$
- (3) $a = \frac{g}{\csc \theta}$
- (4) $a = g \tan \theta$ The moment of the force, $\overrightarrow{F} = 4 \hat{i} + 5 \hat{j} - 6 \hat{k}$ at (2, 0, -3), about the point (2, -2, -2), is given by
 - (1) $-4\hat{i} \hat{j} 8\hat{k}$
 - (2) $-7\hat{i} 8\hat{j} 4\hat{k}$
 - (3) $-8\hat{i} 4\hat{j} 7\hat{k}$
 - (4) $-7\hat{i} 4\hat{j} 8\hat{k}$

A student measured the diameter of a small steel ball using a screw gauge of least count 0.001 cm. The main scale reading is 5 mm and 'zero of circular scale division coincides with 25 divisions above the reference level. If screw gauge has a zero error of -0.004 cm, the correct diameter of the ball is

- (1) 0.525 cm
- (2) 0.053 cm
- (3) 0.521 cm
- (4) 0.529 cm

9. The volume (V) of a monatomic gas varies with its temperature (T), as shown in the graph. The ratio of work done by the gas, to the heat absorbed by it, when it undergoes a change from state A to state B, is





- (1) $\frac{2}{3}$
- (2) $\frac{1}{3}$
- (3) $\frac{2}{5}$
- $(4) \frac{2}{7}$
- 10. The fundamental frequency in an open organ pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is
 - (1) 8 cm
 - (2) 12·5 cm
 - (3) 13·2 cm
 - (4) 16 cm
- At what temperature will the rms speed of oxygen molecules become just sufficient for escaping from the Earth's atmosphere?

Mass of oxygen molecule (m) = 2.76×10^{-26} kg Boltzmann's constant $k_B = 1.38 \times 10^{-23}$ J K⁻¹)


- (1) 8·360 × 10⁴ K
- (2) $5.016 \times 10^4 \text{ K}$
- (3) $2.508 \times 10^4 \text{ K}$
- (4) 1·254 × 10⁴ K
- The efficiency of an ideal heat engine working between the freezing point and boiling point of water, is
 - (1) 20%
 - (2) 6.25%
 - (3) 26.8%
 - (4) 12.5%


- 13. A carbon resistor of $(47\pm4.7)~k\Omega$ is to be marked with rings of different colours for its identification. The colour code sequence will be
 - (1) Yellow Violet Orange Silver
 - (2) Yellow Green Violet Gold
 - (3) Violet Yellow Orange Silver
 - (4) Green Orange Violet Gold
- 14. A set of 'n' equal resistors, of value 'R' each, are connected in <u>series</u> to a battery of emf 'E' and internal resistance 'R'. The current drawn is I. Now, the 'n' resistors are connected in parallel to the same battery. Then the current drawn from battery becomes 10 I. The value of 'n' is
 - (1) 11
 - (2) 20
 - (3) 10°
 - (4) 9
- A battery consists of a variable number 'n' of identical cells (having internal resistance 'r' each) which are connected in series. The terminals of the battery are short-circuited and the current I is measured. Which of the graphs shows the correct relationship between I and n?






B








- 16. An em wave is propagating in a medium with a velocity $\overrightarrow{V} = \overrightarrow{V}$ i. The instantaneous oscillating electric field of this em wave is along +y axis.

 Then the direction of oscillating magnetic field of the em wave will be along
 - (1) + z direction -
 - (2) y direction
 - (3) z direction
 - (4) x direction
- 17. The refractive index of the material of a prism is √2 and the angle of the prism is 30°. One of the two refracting surfaces of the prism is made a mirror inwards, by silver coating. A beam of monochromatic light entering the prism from the other face will retrace its path (after reflection from the silvered surface) if its angle of incidence on the prism is
 - (1) 45°
 - (2) 30°
 - (3) 60°
 - (4) zero
- 18. An object is placed at a distance of 40 cm from a concave mirror of focal length 15 cm. If the object is displaced through a distance of 20 cm towards the mirror, the displacement of the image will be
 - (1) 36 cm away from the mirror.
 - (2) 30 cm towards the mirror
 - (3) 30 cm away from the mirror
 - (4) 36 cm towards the mirror
- 19. The magnetic potential energy stored in a certain inductor is 25 mJ, when the current in the inductor is 60 mA. This inductor is of inductance
 - (1) 138·88 H
 - (2) 1·389 H
 - (3) 0·138 H
 - (4) 13·89 H

- 20. An electron of mass m with an initial velocity $\overrightarrow{V} = \overrightarrow{V_0} \, \hat{i} \, (V_0 > 0)$ enters an electric field $\overrightarrow{E} = \, E_0 \, \hat{i} \, (E_0 = \text{constant} > 0)$ at t = 0. If λ_0 is its de-Broglie wavelength initially, then its de-Broglie wavelength at time t is
 - $(1) \quad \ \, \lambda_0 \left(1 + \frac{e E_0}{m V_0} \, t \, \right)$
 - (2) $\lambda_0 t$
 - $(3) \qquad \frac{\lambda_0}{\left(1 + \frac{eE_0}{mV_0}t\right)}$
 - (4) λ₀
- 21. For a radioactive material, half-life is 10 minutes. If initially there are 600 number of nuclei, the time taken (in minutes) for the disintegration of 450 nuclei is
 - (1) 10
 - (2) 30
 - (3) 20,
 - (4) 15
- 22. The ratio of kinetic energy to the total energy o an electron in a Bohr orbit of the hydrogen atom
 - (1) 1:-1,
 - (2) 2:-1
 - (3) 1:1
 - (4) 1:-2
- 23. When the light of frequency $2v_0$ (where v_0 i threshold frequency), is incident on a meta plate, the maximum velocity of electrons emitted is v_1 . When the frequency of the inciden radiation is increased to $5v_0$, the maximum velocity of electrons emitted from the same plat is v_2 . The ratio of v_1 to v_2 is
 - (1) 1:4
 - (2) 4:1
 - (3) 1:2
 - (4) 2:1

- 24. Unpolarised light is incident from air on a plane surface of a material of refractive index '\u03c4'. At a particular angle of incidence 'i', it is found that the reflected and refracted rays are perpendicular to each other. Which of the following options is correct for this situation?
 - (1) Reflected light is polarised with its electric vector perpendicular to the plane of incidence
 - $(2) \quad i = \sin^{-1}\left(\frac{1}{\mu}\right)$
 - (3) Reflected light is polarised with its electric vector parallel to the plane of incidence
 - $(4) \quad i = tan^{-1} \left(\frac{1}{\mu}\right)$
- 25. An astronomical refracting telescope will have large angular magnification and high angular resolution, when it has an objective lens of
 - (1) large focal length and small diameter
 - (2) large focal length and large diameter
 - (3) small focal length and large diameter
 - (4) small focal length and small diameter
- 26. In Young's double slit experiment the separation d between the slits is 2 mm, the wavelength λ of the light used is 5896 Å and distance D between the screen and slits is 100 cm. It is found that the angular width of the fringes is 0.20°. To increase the fringe angular width to 0.21° (with same λ and D) the separation between the slits needs to be changed to
 - (1) 1.9 mm
 - (2) 2·1 mm
 - (3) 1.8 mm
 - (4) 1.7 mm


27. In the circuit shown in the figure, the input voltage V_i is 20 V, V_{BE} = 0 and V_{CE} = 0. The values of I_B , I_C and β are given by



- (1) $I_B = 25 \mu A$, $I_C = 5 mA$, $\beta = 200$
- (2) $I_B = 20 \mu A$, $I_C = 5 mA$, $\beta = 250$
- (3) $I_B = 40 \mu A$, $I_C = 10 \text{ mA}$, $\beta = 250$
- (4) $I_B = 40 \,\mu\text{A}, \ I_C = 5 \,\text{mA}, \ \beta = 125$

In a p-n junction diode, change in temperature due to heating

- (1) affects only forward resistance
- (2) does not affect resistance of p-n junction
- (3) affects only reverse resistance
- (4) affects the overall V I characteristics of p-n junction
- 29. In the combination of the following gates the output Y can be written in terms of inputs A and B as



- (1) $A \cdot \overline{B} + \overline{A} \cdot B \cdot$
- (2) $(A \cdot B + A \cdot B)$
- (3) A.B
- (4) $\overline{A+B}$

SPACE FOR ROUGH WORK

=> 0.20 = 12 2

English

down by flowing a current through it when a column length. If the frequency of the tuning fork magnetic field of induction 0.25 T is acting on it is 320 Hz, the velocity of sound in air at 27°C is in the vertical direction. The current flowing in (1)339 m/s the rod to keep it stationary is \$ = 0.5 (2)350 m/s 5.98 A 330 m/s (3)(2) 14.76 A (4) 300 m/s 7.14 A The electrostatic force between the metal plates 11.32 A (4) 31. of an isolated parallel plate capacitor C having a Current sensitivity of a moving coil galvanometer 35. charge Q and area A, is is 5 div/mA and its voltage sensitivity (angular distance linearly proportional to the deflection per unit voltage applied) is 20 div/V. between the plates. The resistance of the galvanometer is proportional to the square root of the distance between the plates. independent of the distance between the 250Ω SV = NABXI plates. * 40 Ω inversely proportional to the distance (4) 500 Ω between the plates. A thin diamagnetic rod is placed vertically An electron falls from rest through a vertical 36 (32) between the poles of an electromagnet. When the distance h in a uniform and vertically upward current in the electromagnet is switched on, then directed electric field E. The direction of electric field is now reversed, keeping its magnitude the the diamagnetic rod is pushed up, out of the same. A proton is allowed to fall from rest in it horizontal magnetic field. Hence the rod gains through the same vertical distance h. The time of gravitational potential energy. The work fall of the electron, in comparison to the time of required to do this comes from fall of the proton is the magnetic field (1)5 times greater the lattice structure of the material of the (2)10 times greater rod (3) smaller the current source (4) equal the induced electric field due to the changing magnetic field 33. A pendulum is hung from the roof of a sufficiently high building and is moving freely to An inductor 20 mH, a capacitor 100 µF and a and fro like a simple harmonic oscillator. The resistor 50 Ω are connected in series across a acceleration of the bob of the pendulum is source of emf, V = 10 sin 314 t. The power loss in 20 m/s2 at a distance of 5 m from the mean the circuit is position. The time period of oscillation is 0.43 W (1) T S 2.74 W (2)25 0.79 W (3) $2\pi s$ (4)1.13 W (4) 1s SPACE FOR ROUGH WORK English HLAAC/KK/Page 6

A tuning fork is used to produce resonance in a

glass tube. The length of the air column in this tube can be adjusted by a variable piston. At

room temperature of 27°C two successive

resonances are produced at 20 cm and 73 cm of

A metallic rod of mass per unit length


0.5 kg m-1 is lying horizontally on a smooth

inclined plane which makes an angle of 30° with

the horizontal. The rod is not allowed to slide

- 38. The power radiated by a black body is P and it radiates maximum energy at wavelength, λ_0 . If the temperature of the black body is now changed so that it radiates maximum energy at wavelength $\frac{3}{4}\lambda_0$, the power radiated by it becomes nP. The value of n is
 - (1) $\frac{4}{3}$
 - (2) $\frac{256}{81}$,
 - (3) $\frac{3}{4}$
 - (4) $\frac{81}{256}$
- 39. Two wires are made of the same material and have the same volume. The first wire has cross-sectional area A and the second wire has cross-sectional area 3A. If the length of the first wire is increased by Δl on applying a force F, how much force is needed to stretch the second wire by the same amount?
 - (1) 6F 4, 2 12
 - (2) 4F R
 - (3) 9 F
 - (4) 1
- F A A A F 2 = YA
- 40. A small sphere of radius 'r' falls from rest in a viscous liquid. As a result, heat is produced due to viscous force. The rate of production of heat when the sphere attains its terminal velocity, is proportional to
 - (1) r^2
 - (2) r⁵
 - (3) r^3
 - (4) r⁴
- 41. A sample of 0·1 g of water at 100°C and normal pressure (1·013 × 10⁵ Nm⁻²) requires 54 cal of heat energy to convert to steam at 100°C. If the volume of the steam produced is 167·1 cc, the change in internal energy of the sample, is
 - (1) 208·7 J
 - (2) 42·2 J
 - (3) 104·3 J
 - (4) 84·5 J


A body initially at rest and sliding along a frictionless track from a height h (as shown in the figure) just completes a vertical circle of diameter AB = D. The height h is equal to



(2) $\frac{7}{5}$ D

(1)

- $(3) \quad \frac{3}{2} D$
- (4) $\frac{5}{4}$
- 43. Three objects, A: (a solid sphere), B: (a thin circular disk) and C: (a circular ring), each have the same mass M and radius R. They all spin with the same angular speed ω about their own symmetry axes. The amounts of work (W) required to bring them to rest, would satisfy the relation
 - (1) $W_A > W_B > W_C$
 - $(2) \quad \mathbf{W}_{\mathrm{B}} > \mathbf{W}_{\mathrm{A}} > \mathbf{W}_{\mathrm{C}}$
 - (3) $W_C > W_B > W_A$
 - $(4) \quad \mathbf{W_A} > \mathbf{W_C} > \mathbf{W_B}$
- 44. A moving block having mass m, collides with another stationary block having mass 4m. The lighter block comes to rest after collision. When the initial velocity of the lighter block is v, then the value of coefficient of restitution (e) will be
 - (1) 0.25 *
 - (2) 0.8
 - (3) 0.5
 - (4) 0.4
- 45. Which one of the following statements is incorrect?
 - Limiting value of static friction is directly proportional to normal reaction.
 - (2) Frictional force opposes the relative motion.
 - (3) Rolling friction is smaller than sliding friction.
 - (4) Coefficient of sliding friction has dimensions of length. •



46. Match the metal ions given in Column I with the spin magnetic moments of the ions given in Column II and assign the *correct* code:

	$Column\ I$		Column II
a.	Co^{3+}	i/	$\sqrt{8}$ B.M.
b.	Cr ³⁺	Ti.	$\sqrt{35}$ B.M.
c.	Fe ³⁺	iii.	$\sqrt{3}$ B.M.
d.	Ni ²⁺	iv.	$\sqrt{24}$ B.M.
		v.	$\sqrt{15}$ B.M.
		7	

	a	ь	c	d
(1)	i	ii	iii	iv

- (2) iv i ii iii
- (3) iv v ii i •
- (4) iii v i ii
- 47. Iron carbonyl, Fe(CO)₅ is
 - (1) mononuclear
 - (2) trinuclear
 - (3) tetranuclear
 - (4) dinuclear
- 48. The type of isomerism shown by the complex $[CoCl_2(en)_2]$ is
 - (1) Coordination isomerism
 - (2) Ionization isomerism
 - (3) Geometrical isomerism
 - (4) Linkage isomerism
- **49.** Which one of the following ions exhibits d-d transition and paramagnetism as well?
 - (1) $Cr_2O_7^{2-}$
 - (2) MnO₄
 - (3) CrO₄²⁻
 - (4) MnO_{4}^{2}
- 50. The geometry and magnetic behaviour of the complex [Ni(CO)₄] are
 - (1) tetrahedral geometry and diamagnetic:
 - (2) square planar geometry and paramagnetic
 - (3) square planar geometry and diamagnetic
 - (4) tetrahedral geometry and paramagnetic

- 51. A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc. H₂SO₄. The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be
 - (1) 3.0
 - (2) 2.8
 - (3) 1.4
 - (4) 4.4

53.

- 52. The difference between amylose and amylopectin is
 - (1) Amylose have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6$ β -linkage
 - (2) Amylopectin have 1 \rightarrow 4 α -linkage and 1 \rightarrow 6 β -linkage
 - (3) Amylopectin have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6$ α -linkage .
 - (4) Amylose is made up of glucose and galactose
 - Regarding cross-linked or network polymers, which of the following statements is *incorrect*?
 - They are formed from bi- and tri-functional monomers.
 - (2) Examples are bakelite and melamine.
 - (3) They contain covalent bonds between various linear polymer chains.
 - (4) They contain strong covalent bonds in their polymer chains.
- 54. Nitration of aniline in strong acidic medium also gives m-nitroaniline because
 - In electrophilic substitution reactions amino group is meta directive.
 - (2) In absence of substituents nitro group always goes to m-position. ⊀
 - (3) In spite of substituents nitro group always goes to only m-position.
 - (4) In acidic (strong) medium aniline is present as anilinium ion.
- **55.** Which of the following oxides is most acidic in nature?
 - (1) BeO -
 - (2) BaO
 - (3) MgO
 - (4) CaO

56. In the reaction

$$\begin{array}{c}
\text{OH} \\
\hline
\text{O} \\
\text{+ CHCl}_3 + \text{NaOH}
\end{array}$$

the electrophile involved is

- (1) formyl cation (CHO)
- (2) dichloromethyl anion (CHCl₂)
- (3) dichloromethyl cation (CHCl₂)
- (4) dichlorocarbene (:CCl2) ,
- 57. Carboxylic acids have higher boiling points than aldehydes, ketones and even alcohols of comparable molecular mass. It is due to their
 - (1) formation of carboxylate ion
 - (2) more extensive association of carboxylic acid via van der Waals force of attraction
 - (3) formation of intramolecular H-bonding
 - (4) formation of intermolecular H-bonding
- 58. Compound A, C₈H₁₀O, is found to react with NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell.

A and Y are respectively

(1)
$$\sim$$
 CH₂ – CH₂ – OH and I₂

(2)
$$\leftarrow$$
 CH – CH₃ and I₂ OH

(3)
$$H_3C \longrightarrow CH_2 - OH \text{ and } I_2$$

$$^{
m CH_3}$$
 $^{
m CH_3}$ $^{
m CH_3}$ OH and $^{
m I}_2$

- 59. The compound A on treatment with Na gives B, and with PCl₅ gives C. B and C react together to give diethyl ether. A, B and C are in the order
 - (1) C_2H_5OH , C_2H_5Cl , C_2H_5ONa
 - (2) C_2H_5Cl , C_2H_6 , C_2H_5OH
 - (3) C_2H_5OH , C_2H_6 , C_2H_5CI
 - (4) C₂H₅OH, C₂H₅ONa, C₂H₅Cl
 - Which oxide of nitrogen is **not** a common pollutant introduced into the atmosphere both due to natural and human activity?
 - (1) NO₂
 - (2) N₂O
 - (3) N₂O₅
 - (4) NO
- 61. Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms. (A) is
 - (1) $CH_2 = CH_2$
 - (2) $CH_3 CH_3$
 - (3) CH ≡ CH
 - (4) CH₄
- **62.** The compound C₇H₈ undergoes the following reactions:

$$C_7H_8 \xrightarrow{3 \operatorname{Cl}_2/\Delta} A \xrightarrow{\operatorname{Br}_2/\operatorname{Fe}} B \xrightarrow{\operatorname{Zn}/\operatorname{HCl}} C$$

The product 'C' is

- (1) o-bromotoluene
- (2) 3-bromo-2,4,6-trichlorotoluene
- (3) m-bromotoluene
- (4) p-bromotoluene.

Which of the following carbocations is expected to be most stable?

$$(4) \qquad \stackrel{\text{NO}_2}{Y}$$

64. Which of the following is correct with respect to - I effect of the substituents? (R = alkyl)

(1)
$$-NR_2 < -OR < -F$$

(2)
$$-NH_2 > -OR > -F$$

$$(3) - NH_2 < -OR < -F$$

$$(4)$$
 $-NR_2 > -OR > -F$

65. Which of the following molecules represents the order of hybridisation sp², sp², sp, sp from left to right atoms?

(1)
$$CH_2 = CH - C \equiv CH$$

$$(2) \quad \operatorname{CH}_2 = \operatorname{CH} - \operatorname{CH} = \operatorname{CH}_2$$

(3)
$$HC \equiv C - C \equiv CH$$

(4)
$$CH_3 - CH = CH - CH_3$$

66. Identify the major products P, Q and R in the following sequence of reactions:

$$\begin{array}{c} \text{Anhydrous} \\ & \text{AlCl}_3 \\ \\ \text{P} \xrightarrow{\text{(i) O}_2} \\ & \text{(ii) H}_3\text{O}^+\!/\!\Delta} \neq \text{Q} + \end{array}$$

P Q R
$$\begin{array}{cccc} \text{CH}_2\text{CH}_2\text{CH}_3 & \text{CHO} & \text{COOH} \\ & & & & & \\ \text{(1)} & & & & & \\ \end{array}$$

(2)
$$CH(CH_3)_2$$
 OH $CH_3CH(OH)CH$

(3)
$$CH_2CH_2CH_3$$
 CHO , $CH_3CH_2 - OH$

(4)
$$CH(CH_3)_2$$
 $CH_3 - CO - CH_2$

- 67. Which of the following compounds can form zwitterion?
 - (1) Acetanilide
 - (2) Benzoic acid
 - (3) Aniline
 - (4) Glycine ,

- 68. Following solutions were prepared by mixing different volumes of NaOH and HCl of different concentrations:
 - a. $60 \text{ mL } \frac{\text{M}}{10} \text{ HCl} + 40 \text{ mL } \frac{\text{M}}{10} \text{ NaOH}$
 - b. $55 \text{ mL } \frac{\text{M}}{10} \text{ HCl} + 45 \text{ mL } \frac{\text{M}}{10} \text{ NaOH}$
 - c. $75 \text{ mL } \frac{\text{M}}{5} \text{ HCl} + 25 \text{ mL } \frac{\text{M}}{5} \text{ NaOH}$
 - d. 100 mL $\frac{M}{10}$ HCl + 100 mL $\frac{M}{10}$ NaOH

pH of which one of them will be equal to 1?

- (1) a
- (2) d
- (3) b
- (4) Ce
- 69. On which of the following properties does the coagulating power of an ion depend?
 - (1) Size of the ion alone
 - (2) Both magnitude and sign of the charge on the ion
 - (3) The magnitude of the charge on the ion alone
 - (4) The sign of charge on the ion alone
- Given van der Waals constant for NH₃, H₂, O₂ and CO₂ are respectively 4·17, 0·244, 1·36 and 3·59, which one of the following gases is most easily liquefied?
 - (1) H₂
 - (2) O₂
 - (3) NH2
 - (4) CO₂
- 71. The solubility of BaSO₄ in water is $2.42 \times 10^{-3} \text{ gL}^{-1}$ at 298 K. The value of its solubility product (K_{sp}) will be

(Given molar mass of $BaSO_4 = 233 \text{ g mol}^{-1}$)

- $(1) \quad 1.08 \times 10^{-12} \ mol^2 \ L^{-2}$
- (2) $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$
- (3) $1.08 \times 10^{-10} \, \text{mol}^2 \, \text{L}^{-2}$,
- $(4) \quad \ 1 \cdot 08 \times 10^{-8} \ \text{mol}^2 \ L^{-2}$

- The bond dissociation energies of X₂, Y₂ and XY are in the ratio of 1:0.5:1. ΔH for the formation of XY is -200 kJ mol⁻¹. The bond dissociation
 - energy of X_2 will be (1) 100 kJ mol⁻¹
 - (2) 800 kJ mol-1.
 - (3) 200 kJ mol⁻¹
 - (4) 400 kJ mol⁻¹
- 73. When initial concentration of the reactant is doubled, the half-life period of a zero order reaction
 - (1) is doubled
 - (2) is tripled
 - (3) is halved
 - (4) remains unchanged
- 74. For the redox reaction

$$\mathrm{MnO_4^-} + \mathrm{C_2O_4^{2-}} + \mathrm{H^+} {\longrightarrow} \mathrm{Mn^{2+}} + \mathrm{CO_2} + \mathrm{H_2O}$$

the correct coefficients of the reactants for the balanced equation are

	MnO_4^-	$C_2O_4^{2-}$	H ⁺		
1)	2	5	16		
2)	2	16	5		
a	10	-	0		

- (3) 16 5 2 (4) 5 16 2
- 75. Which one of the following conditions will favour maximum formation of the product in the reaction,

$$\mathbf{A_{2}}\left(\mathbf{g}\right)+\mathbf{B_{2}}\left(\mathbf{g}\right)\rightleftarrows\mathbf{X_{2}}\left(\mathbf{g}\right)\ \Delta_{\mathbf{r}}\mathbf{H}=-\mathbf{X}\;\mathbf{kJ}\;?$$

- (1) Low temperature and low pressure
- (2) High temperature and high pressure
- (3) Low temperature and high pressure ,
- (4) High temperature and low pressure
- 76. The correction factor 'a' to the ideal gas equation corresponds to
 - (1) volume of the gas molecules
 - (2) electric field present between the gas molecules
 - (3) density of the gas molecules
 - (4) forces of attraction between the gas molecules.

- 77. The correct order of N-compounds in its decreasing order of oxidation states is
 - (1) HNO₃, NO, NH₄Cl, N₂
 - (2) HNO3, NH4Cl, NO, N2
 - (3) HNO_3 , NO, N_2 , NH_4Cl
 - (4) NH₄Cl, N₂, NO, HNO₃
- 78. Which one of the following elements is unable to form MF₆³⁻ ion?
 - (1) Al
 - (2) B •
 - (3) Ga
 - (4) In
- 79. Considering Ellingham diagram, which of the following metals can be used to reduce alumina?
 - (1) Zn
 - (2) Mg
 - (3) Fe
 - (4) Cu
- 80. The correct order of atomic radii in group 13 elements is
 - (1) B < Al < Ga < In < Tl
 - (2) B < Ga < Al < Tl < In
 - (3) B < Al < In < Ga < Tl
 - (4) B < Ga < Al < In < Tl.
- Which of the following statements is **not** true for 86. halogens?
 - All are oxidizing agents.
 - (2) All but fluorine show positive oxidation states.
 - (3) All form monobasic oxyacids.
 - (4) Chlorine has the highest electron-gain enthalpy./
- 82. In the structure of ClF₃, the number of lone pairs of electrons on central atom 'Cl' is
 - (1) two
 - (2) four
 - (3) one
 - (4) three

- 83. The correct difference between first- and second-order reactions is that
 - (1) the half-life of a first-order reaction does not depend on [A]₀; the half-life of a second-order reaction does depend on [A]₀
 - a first-order reaction can be catalyzed; a second-order reaction cannot be catalyzed
 - (3) the rate of a first-order reaction does not depend on reactant concentrations; the rate of a second-order reaction does depend on reactant concentrations
 - (4) the rate of a first-order reaction does depend on reactant concentrations; the rate of a second-order reaction does not depend on reactant concentrations
- Among CaH₂, BeH₂, BaH₂, the order of ionic character is
 - (1) $CaH_2 < BeH_2 < BaH_2$
 - (2) $BeH_2 < BaH_2 < CaH_2$
 - (3) BeH2 < CaH2 < BaH2 F
 - (4) BaH₂ < BeH₂ < CaH₂
 - In which case is the number of molecules of water maximum?
 - (1) 0.18 g of water
 - (2) 0.00224 L of water vapours at 1 atm and 273 K
 - (3) 18 mL of water
 - (4) 10^{-3} mol of water

Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below:

$$BrO_4^- \xrightarrow{1.82 \text{ V}} BrO_3^- \xrightarrow{1.5 \text{ V}} HBrO$$

$$Br^- \xleftarrow{1.0652 \text{ V}} Br_2 \xleftarrow{1.595 \text{ V}}$$

Then the species undergoing disproportionation is

- (1) BrO-
- (2) Br₂
- (3) BrO₂
- (4) HBrO

87. Consider the following species:

Which one of these will have the highest bond order?

- (1) CN-
- (2) CN+
- (3) NO
- (4) CN
- 88. Which one is a wrong statement?
 - An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers.
 - (2) The electronic configuration of N atom is

$1s^2$	$2s^2$	$2p_x^1 \ 2p_y^1 \ 2p_z^1$				
$\uparrow \downarrow$	↑ ↓	1	1	1		

- (3) Total orbital angular momentum of electron in 's' orbital is equal to zero.
- (4) The value of m for d,2 is zero.
- 89. Iron exhibits bcc structure at room temperature.

 Above 900°C, it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature) is
 - $(1) \quad \frac{4\sqrt{3}}{3\sqrt{2}}$
 - $(2) \qquad \frac{3\sqrt{3}}{4\sqrt{2}}$
 - $(3) \qquad \frac{\sqrt{3}}{\sqrt{2}}$
 - (4) $\frac{1}{2}$
- 90. Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is 1s² 2s² 2p³, the simplest formula for this compound is
 - (1) MgX₂
 - (2) Mg_2X
 - (3) Mg_2X_3
 - (4) Mg₃X₂ •

- **91.** Pollen grains can be stored for several years in liquid nitrogen having a temperature of
 - (1) -80°C
 - (2) 196°C
 - (3) 120°C
 - (4) 160°C
- 92. Oxygen is not produced during photosynthesis by
 - (1) Nostoc
 - (2) Cycas
 - (3) Green sulphur bacteria.
 - (4) Chara
- 93. What is the role of NAD⁺ in cellular respiration?
 - (1) It functions as an electron carrier.
 - (2) It is a nucleotide source for ATP synthesis.
 - (3) It functions as an enzyme.
 - (4) It is the final electron acceptor for anaerobic respiration.
- **94.** Which of the following elements is responsible for maintaining turgor in cells?
 - (1) Sodium
 - (2) Potassium.
 - (3) Magnesium
 - (4) Calcium
- 95. Which one of the following plants shows a very close relationship with a species of moth, where none of the two can complete its life cycle without the other?
 - (1) Yucca.
 - (2) Banana
 - (3) Hydrilla
 - (4) Viola
- In which of the following forms is iron absorbed by plants?
 - (1) Ferrous
 - (2) Free element
 - (3) Ferric
 - (4) Both ferric and ferrous
- 97. Double fertilization is
 - Fusion of one male gamete with two polar nuclei
 - (2) Fusion of two male gametes with one egg
 - (3) Fusion of two male gametes of a pollen tube with two different eggs
 - (4) Syngamy and triple fusion e

- A 'new' variety of rice was patented by a foreign (10). company, though such varieties have been 98. present in India for a long time. This is related to
 - Sharbati Sonora (1)
 - Lerma Rojo (2)
 - Co-667 (3)
 - Basmati.
- India, the organisation responsible for assessing the safety of introducing genetically (99) modified organisms for public use is

(CAPELOS

- Industrial Council for Scientific and Research (CSIR)
- Genetic on Committee Research (2)Manipulation (RCGM)
- Indian Council of Medical Research (ICMR) (3)
- Genetic Engineering Appraisal Committee (4) (GEAC)
- 100. Which of the following is commonly used as a vector for introducing a DNA fragment in human lymphocytes?
 - Ti plasmid
 - λ phage
 - Retrovirus.
 - (4) pBR 322
 - 101. Use of bioresources by multinational companies and organisations without authorisation from the concerned country and its people is called
 - Biopiracy . (1)
 - Biodegradation (2)
 - Bio-infringement (3)
 - Bioexploitation (4)
 - 102. The correct order of steps in Polymerase Chain Reaction (PCR) is
 - Annealing, Extension, Denaturation (1)
 - Denaturation, Extension, Annealing (2)
 - Extension, Denaturation, Annealing (3)
 - Denaturation, Annealing, Extension.
 - (03) Select the correct match :
 - Dihybrid cross $\mathbf{F}_2 \times \mathbf{Recessive \ parent}$ (1) Transduction
 - T.H. Morgan Nucleic acid,
 - Ribozyme (3) G. Mendel (4)
 - Transformation

- - the physical space where an organism lives Niche is (1)
 - the range of temperature that the organism (2)needs to live
 - all the biological factors in the organism's (3) environment
 - the functional role played by the organism (4) where it lives .
- 105. Which of the following is a secondary pollutant?
 - CO_2 (1)
 - SO2 (2)
 - CO (3)
 - (4) O3,
 - 106. Natality refers to
 - Birth rate * (1)
 - Number of individuals leaving the habitat (2)
 - Death rate
 - Number of individuals entering a habitat (3)(4)
 - 107. World Ozone Day is celebrated on
 - (1) 21st April
 - 16th September
 - (3) 5th June
 - 22nd April
 - 108. What type of ecological pyramid would obtained with the following data?

Secondary consumer: 120 g

Primary consumer: 60 g

Primary producer: 10 g

- Pyramid of energy (1)
- Upright pyramid of numbers (2)
- Inverted pyramid of biomass . (3)
- Upright pyramid of biomass (4)
- 109. In stratosphere, which of the following elem acts as a catalyst in degradation of ozone release of molecular oxygen?
 - (1) Cl.
 - Fe (2)
 - Carbon (3)
 - Oxygen (4)

£10)	Which of the following pairs is wrongly matched?	117.	Secondary xylem and phloem in dicot stem are produced by
	(1) ABO blood grouping : Co-dominance	1	(1) Vascular cambium-
	(2) XO type sex : Grasshopper		(2) Phellogen
	determination		(3) Apical meristems
	(3) Starch synthesis in pea : Multiple alleles		(4) Axillary meristems
	(4) T.H. Morgan : Linkage/	118	Plants having little or no secondary growth are
111.	Select the correct statement:	110.	(1) Deciduous angiosperms
	(1) Punnett square was developed by a British		(2) Conifers
	scientist.		(3) Grasses
	(2) Spliceosomes take part in translation.		
	(3) Franklin Stahl coined the term "linkage".		*
	(4) Transduction was discovered by S. Altman.	119.	Sweet potato is a modified
112	The experimental proof for semiconservative		(1) Adventitious root
0	replication of DNA was first shown in a		(2) Tap root •
	(1) Bacterium.		(3) Stem
	(2) Plant		(4) Rhizome
	(3) Fungus	120.	Pneumatophores occur in
	(4) Virus		(1) Free-floating hydrophytes
113.	Select the correct match:		(2) Carnivorous plants
	(1) Alfred Hershey and - TMV		(3) Halophytes.
	Martha Chase		(4) Submerged hydrophytes
	(2) Matthew Meselson - Pisum sativum and F. Stahl	121.	Casparian strips occur in
	(3) Alec Jeffreys – Streptococcus		(1) Pericycle
	pneumoniae		(2) Cortex,
	(4) Francois Jacob and – Lac operon		(3) Epidermis
	Jacques Monod		(4) Endodermis,
114.	Offsets are produced by	122.	Which of the following statements is correct ?
	(1) Mitotic divisions	1225	(1) Selaginella is heterosporous, while Salvinia
	(2) Parthenocarpy		is homosporous. ×
	(3) Meiotic divisions		(2) Horsetails are gymnosperms. ×
115.	(4) Parthenogenesis Which of the following flowers only once in its		(3) Ovules are not enclosed by ovary wall in gymnosperms.
	life-time? (1) Jackfruit (2) Manage		(4) Stems are usually unbranched in both Cycas and Cedrus.
	(2) Mango (3) Bamboo species	123	Select the wrong statement:
	(4) Papaya		(1) Mushrooms belong to Basidiomycetes.
116.	Which of the following has proved helpful in preserving pollen as fossils?		(2) Pseudopodia are locomotory and feeding structures in Sporozoans.
	(1) Cellulosic intine (2) Oil content		(3) Cell wall is present in members of Fungi and Plantae.
	(3) Pollenkitt		(4) Mitochondria are the powerhouse of the cell
	(4) Sporopollenin •	- 00	in all kingdoms except Monera.

124.	Mate Colu belo	ımn II and	giver select	in Column I with those in the correct option given	128.	The sugar (1) (2)	two functional groups characteristic of carbonyl and methyl carbonyl and phosphate	
	a. Herbarium i.		i.	It is a place having a collection of preserved		(3) (4)	hydroxyl and methyl carbonyl and hydroxyl •	
	b.	Key	ii.	plants and animals. A list that enumerates methodically all the species found in an area with brief description aiding identification.	(129.		ch of the following is not a product of light tion of photosynthesis? NADH NADPH • ATP Oxygen	
	c.	Museum	iii.	Is a place where dried and pressed plant specimens mounted on sheets are kept.	130.	(1) (2) (3)	ch among the following is <u>not</u> a prokaryote? Mycobacterium Nostoc Saccharomyces.	
	d.	Catalogue	iv.	A booklet containing a list of characters and their alternates which are helpful in identification of various taxa.	131.	(4) Storr (1) (2) (3)	Oscillatoria natal movement is not affected by Light O_2 concentration,	
		a b	c	d ·		(4)	CO_2 concentration	
	(1)	iii ii	i	iv	132.	The (Golgi complex participates in	
	(2)	ii iv	iii	i		(1)	Formation of secretory vesicles.	
	(3)	i iv	iii	ii		(2)	Respiration in bacteria	
	(4)	iii iv	i	ii •		(3)	Fatty acid breakdown	
			0.11	34 State - 1		(4)	Activation of amino acid	
125.				wed by meiosis, spores are	133.	Which of the following is true for nucleolus?		
		luced exogen		in		(1)	It is a membrane-bound structure.	
	(1) (2)	Alternaria ;	((2)	It takes part in spindle formation.	
	(3)	Neurospora	A				Larger nucleoli are present in dividing cells.	
	(4)	Saccharomy				(4)	It is a site for active ribosomal RNA synthesis.	
126.			ains a	are present in	134.		stage during which separation of the paired	
	(1)	Cycas					ologous chromosomes begins is	
	(2)	Mango				(1)	Diplotene	
	(3)	Mustard				(2)	Diakinesis .	
	(4)	Pinus				(3)	Pachytene	
127.	Whic	ch one is wro	ngly	matched?	105	(4)	Zygotene	
	(1)	Biflagellate	4.		135.		ata in grass leaf are	
	(2)	Gemma cup		- Marchantia/		(1) (2)	Kidney shaped	
	(3)	TT 'C 11		netes - Polysiphonia & for	m	(4)	Rectangular	
	 (3) Uniflagellate gametes - Polysiphonia 6 (4) Unicellular organism - Chlorella , 					(3)	Dumb-bell shaped,	

136.	Niss	al bodies are mainly composed of	142.	All	of the	followin	g are pa	art o	f an operon except
	(1)	DNA and RNA		(1)		ctural g			***************************************
	(2)	Nucleic acids and SER		(2)		nhancer			
	(3)	Proteins and lipids		(3)	an o	perator			4
	(4)	Free ribosomes and RER,		(4)	a pr	omoter			
137.	Whie	ch of these statements is <i>incorrect</i> ? Glycolysis occurs in cytosol.	143.			has an			ndition on one of her romosome can be
	(2) (3) (4)	Glycolysis operates as long as it is supplied with NAD that can pick up hydrogen atoms. Enzymes of TCA cycle are present in mitochondrial matrix. Oxidative phosphorylation takes place in outer mitochondrial membrane.		(1) (2) (3) (4)	Only	sons grando daught	children ters nd daug		x x x x x x x x x x x x x x x x x x x
138.	(1) (2) (3) (4)	ch of the following terms describe human tition?		evol (1) (2) (3) (4) AGC stra	Salt Pher Mul Min GTAT nd of uence UGO	is ation, notypic tiple ste or muta CGCAT a gene.	variations is a see What w	ns tions que	nce from the coding be the corresponding
	(4)	chromosomes Polytene – Oocytes of amphibians chromosomes	146.	(3) (4) Mat	UCC	GUAUC CAUAGO items	CGUA	Col	umn I with those in
(140.)		ich of the following events does not occur in gh endoplasmic reticulum?	11.72		ımn l				orrect option given
	(1)	Protein glycosylation			Colu	imn I			Column II
	(2) (3)	Cleavage of signal peptide Protein folding		a.	Prol	iferative	e Phase	i.	Breakdown of endometrial lining
	(4)	Phospholipid synthesis				TO	e di con		
141	Mar	ny ribosomes may associate with a single		b.		etory P		ii.	Follicular Phase
1421	mR.	NA to form multiple copies of a polypeptide ultaneously. Such strings of ribosomes are ned as		c. (1)	Mer a i	struatio b iii	on c ii	111.	Luteal Phase
	(1)	Polyhedral bodies							
	(2)	Plastidome		(2)	ii 	iii	i'.		
	(3)	Polysome .		(3)	iii	ii	i		
	(4)	Nucleosome		(4)	iii	i	ii		

the p (1) (2) (3) (4) 148. All (actions is widely used in medical science for roduction of antibiotics? Mutualism Parasitism Commensalism	152. Among the following sets of examples for divergent evolution, select the incorrect option: (1) Heart of bat, man and cheetah (2) Brain of bat, man and cheetah (3) Forelimbs of man, bat and cheetah (4) Eye of octopus, bat and man. 153. Which of the following is not an autoimmune disease? (1) Rheumatoid arthritis (2) Alzheimer's disease. We ace fylcholime (3) Psoriasis (4) Vitiligo
(3) (4)	Wildlife safari parks Seed banks th the items given in Column I with those in	154. In which disease does mosquito transmitted pathogen cause chronic inflammation of lymphatic vessels?
Colubelov a. b. c. d. (1) (2) (3) (4) 150. In a (1)	mm II and select the correct option given w: Column I Eutrophication i. UV-B radiation Sanitary landfill ii. Deforestation Snow blindness iii. Nutrient enrichment Jhum cultivation iv. Waste disposal a b c d i iii iv ii iii iv ii iii iv ii ii iii i	(1) Ascariasis (2) Ringworm disease (3) Elephantiasis (4) Amoebiasis 155. Conversion of milk to curd improves its nutritional value by increasing the amount of (1) Vitamin A (2) Vitamin B ₁₂ (3) Vitamin D (4) Vitamin E 156. The similarity of bone structure in the forelimbs of many vertebrates is an example of (1) Analogy (2) Convergent evolution (3) Homology (4) Adaptive radiation
(2)	reproductive and pre-reproductive individuals are equal in number. pre-reproductive individuals are more than	157. Which of the following characteristics represent 'Inheritance of blood groups' in humans?
(4)	the reproductive individuals. pre-reproductive individuals are less than the reproductive individuals.	c. Multiple allele
151. White dru (1) (2) (3)	ich part of poppy plant is used to obtain the g "Smack"? Latex . Roots Flowers	d. Incomplete dominance e. Polygenic inheritance, (1) a, b and c (2) b, d and e (3) b, c and e (4) a, c and e
(4)	Leaves	POUGH WORK English

- 158. Hormones secreted by the placenta to maintain pregnancy are
 - (1) hCG, hPL, estrogens, relaxin, oxytocin
 - (2) hCG, hPL, progestogens, estrogens.
 - (3) hCG, hPL, progestogens, prolactin
 - (4) hCG, progestogens, estrogens, glucocorticoids →
- 159. The contraceptive 'SAHELI'
 - increases the concentration of estrogen and prevents ovulation in females.
 - (2) is an IUD.
 - (3) blocks estrogen receptors in the uterus, preventing eggs from getting implanted.
 - (4) is a post-coital contraceptive.
- 160. The amnion of mammalian embryo is derived from
 - (1) endoderm and mesoderm
 - (2) mesoderm and trophoblast
 - (3) ectoderm and mesoderm ,
 - (4) ectoderm and endoderm
- 161. The difference between spermiogenesis and spermiation is
 - (1) In spermiogenesis spermatozoa are formed, while in spermiation spermatids are formed.
 - (2) In spermiogenesis spermatozoa from sertoli cells are released into the cavity of seminiferous tubules, while in spermiation spermatozoa are formed.
 - (3) In spermiogenesis spermatids are formed, while in spermiation spermatozoa are formed.
 - (4) In spermiogenesis spermatozoa are formed, while in spermiation spermatozoa are released from sertoli cells into the cavity of seminiferous tubules.

- 162. Which of the following is an amino acid derived hormone?
 - (1) Ecdysone
 - (2) Estradiol
 - (3) Epinephrine •
 - (4) Estriol
- 163. Which of the following structures or regions is incorrectly paired with its function?
 - (1) Limbic system

consists of fibre

tracts that interconnect

different regions of brain; controls

movement.

(2) Hypothalamus

production of

releasing hormones and regulation of

temperature, hunger and thirst.

(3) Medulla oblongata:

controls respiration

and cardiovascular

reflexes.

(4) Corpus callosum

band of fibers

connecting left and right cerebral

hemispheres.

- The transparent lens in the human eye is held in its place by
 - (1) ligaments attached to the iris
 - (2) smooth muscles attached to the iris
 - (3) ligaments attached to the ciliary body
 - (4) smooth muscles attached to the ciliary body
- 165. Which of the following hormones can play a significant role in osteoporosis?
 - (1) Progesterone and Aldosterone ⋈
 - (2) Estrogen and Parathyroid hormone .
 - (3) Aldosterone and Prolactin x
 - (4) Parathyroid hormone and Prolactin >

166. Which of the following options correctly 69. Which of the following gastric cells indire represents the lung conditions in asthma and help in erythropoiesis? emphysema, respectively? (1) Mucous cells Increased number of bronchioles; Increased respiratory surface (2)Goblet cells (2)Increased respiratory surface: Chief cells Inflammation of bronchioles Inflammation of bronchioles; Decreased (3) Parietal cells, (4) respiratory surface, 170. Match the items given in Column I with thos (4) Decreased respiratory surface; Inflammation of bronchioles Column II and select the correct option gi 167. Match the items given in Column I with those in below: Column II and select the correct option given Column I Column II below: Fibrinogen î. Osmotic balance Column I Column II Tricuspid valve i. Between left atrium ii. a. Globulin Blood clotting b. and left ventricle Albumin Defence mechanis iii. b. Bicuspid valve ii. Between right ventricle and pulmonary artery ii (1) iii Semilunar valve Between right (2)iii ii atrium and right ventricle (3) iii ii (4) ii iii iii (1) i ii 171. Which of the following is an occupation ii (2)iii respiratory disorder? (3)ii . (4) ii Silicosis . iii Botulism 168. Match the items given in Column I with those in Column II and select the correct option given (3)Anthracis below: (4) Emphysema Column I Column II i 2500 - 3000 mL a. Tidal volume 172. Calcium is important in skeletal mu Inspiratory Reserve b. ii. 1100 - 1200 mL contraction because it volume activates the myosin ATPase by binding Expiratory Reserve C. iii. 500 - 550 mL volume detaches the myosin head from the acd. Residual volume iv. 1000 - 1100 mL filament. b d a binds to troponin to remove the masking (1) iii iv ii . active sites on actin for myosin. (2)iv ii iii prevents the formation of bonds betw (3) iii ii i iv the myosin cross bridges and the ac (4) iv ii i filament.

			Prince	III C	Column I	with th	ose in
lumn	II	and	select	the	correct	option	given
ow:							
							lumn II and select the ${\it correct}$ option ow:

		Colu	mnI		Column II		
	a.	Glyc	osuria	i.	Accumulation of uric acid in joints		
	b.	Gout	t \	ii.	Mass of crystallised salts within the kidney		
	c.	Rena	al calculi	iii.	Inflammation in glomeruli		
d.			nerular nritis	\ iv.	Presence of glucose in urine		
		a	b	c	d		
	(1)	i	ii	iii	iv		
	(2)	ii	iii	i	iv		
	(3)	iii	ii	iv	i -		
	(4)	iv	í	ii -	iii •		

174. Match the items given in Column I with those in Column II and select the correct option given below:

	Colu	mnI		9	Column II
	(Fun	ction)			Part of Excretory (ystem)
a.	Ultr	afiltrati	ion	i. H	Ienle's loop
b.	Cone of ur	centrati	on	ii. U	reter
c.	Tran	nsport o e	f	iii. U	rinary bladder
d.	Stor	age of u	rine		Ialpighian orpuscle
	-6				roximal 'onvoluted tubule
	a	b	c	d	
(1)	iv	i	ii	iii 🗸	
(2)	v	iv	i	ii	
(3)	iv	V	ii	iii	(*)
(4)	v	iv	i	iii	

- **175.** Which of the following features is used to identify a male cockroach from a female cockroach?
 - (1) Presence of caudal styles
 - (2) Forewings with darker tegmina
 - (3) Presence of a boat shaped sternum on the 9th abdominal segment
 - (4) Presence of anal cerci
- 176. Identify the vertebrate group of animals characterized by crop and gizzard in its digestive system.
 - (1) Reptilia
 - (2) Aves.
 - (3) Amphibia
 - (4) Osteichthyes
- 177. Which one of these animals is not a homeotherm?
 - (1) Chelone .
 - (2) Camelus
 - (3) Macropus,
 - (4) Psittacula,
- .178. Which of the following organisms are known as chief producers in the oceans?
 - (1) Diatoms
 - (2) Cyanobacteria
 - (3) Dinoflagellates
 - (4) Euglenoids
- 179. Which of the following animals does <u>not</u> undergo metamorphosis?
 - (1) Tunicate
 - (2), Moth
 - (3) Earthworm,
 - (4) Starfish
- 180. Ciliates differ from all other protozoans in
 - having a contractile vacuole for removing excess water
 - (2) using pseudopodia for capturing prey
 - (3) using flagella for locomotion
 - (4) having two types of nuclei