# ICSE Question Paper(2014) CHEMISTRY

## (Two hour)

Answers to this Paper must be written on the paper provided separately. You will **not** be allowed to write during the first **15** minutes. This time is to be spent in reading the Question Paper. The time given at the head of this Paper is the time allowed for writing the answers.

Section I is compulsory. Attempt any four questions from Section II. The intended marks for questions or parts of questions are given in brackets [].

## **SECTION-I** (40 Marks)

Attempt all questions from this Section

#### **Question 1.**

- (a) Choose the correct answer from the options given below :
  - (i) Ionisation Potential increases over a period from left to right because the :
    - (A) Atomic radius increases and nuclear charge increases
    - (B) Atomic radius decreases and nuclear charge decreases
    - (C) Atomic radius increases and nuclear charge decreases
    - (D) Atomic radius decreases and nuclear charge increases.
    - A compound X consists of only molecules. Hence X will have :
    - (A) A crystalline hard structure
    - (B) A low melting point and low boiling point
    - (C) An ionic bond
    - (D) A strong force of attraction between its molecules.
  - (iii) When fused lead bromide is electrolysed we observe :
    - (A) a silver grey deposit at anode and a reddish brown deposit at cathode
    - (B) a silver grey deposit at cathode and a reddish brown deposit at anode
    - (C) a silver grey deposit at cathode and reddish brown fumes at anode
    - (D) silver grey fumes at anode and reddish brown fumes at cathode.
  - (iv) The main ore used for the extraction of iron is :
    - (A) Haematite (B) Calamine
    - (C) Bauxite (D) Cryolite
  - (v) Heating an ore in a limited supply of air or in the absence of air at a temperature just below its melting point is known as :
    - (A) smelting (B) ore dressing
    - (C) calcination (D) bessemerisation
  - (vi) If an element A belongs to Period 3 and Group II then it will have :

		<ul> <li>(A) 3 shells and 2</li> <li>(B) 2 shells and 3</li> <li>(C) 3 shells and 3</li> <li>(D) 2 shells and 2</li> <li>(D) 2 shells and 2</li> <li>(D) 2 shells and 2</li> <li>(A) ammonia</li> <li>(C) water</li> <li>(C) water</li> <li>(C) water nitrate s</li> <li>(C) sodium argent</li> </ul>	valence valence valence ining a ining	electrons electrons electrons triple co-valent (B) (D) troplating an o (B) e solution (D)	methane nitrogen article with silver silver cyanide s nickel sulphate	olution		
	(ix)	Aluminium powder (A) it is a strong re			it is a strong ox	idising agent		
		(C) it is corrosion	-	0	it is a good cond			
	(x) <sub>.</sub>	The I.U.P.A.C. nam	ne of ace		C			
		(A) propane		(B)	propyne			
		(C) ethene		(D)	ethyne	[10]		
<b>(b)</b>	) Fill in the blanks from the choices given within brackets :							
	(i)	The basicity of Acetic Acid is (3, 1, 4)						
	<ul> <li>(ii) The compound formed when ethanol reacts with sodium (sodium ethanoate, sodium ethoxide, sodium propanoate).</li> </ul>							
	(iii)	Quicklime is not used to dry HCl gas because (CaO is alkaline, CaO is acidic, CaO is neutral).						
	(iv)	Ammonia gas is collected by (an upward displacement of air, a downward displacement of water, a downward displacement of air)						
	(v) -							
(c)	Give one word or phrase for the following :							
	<ul> <li>(i) The ratio of the mass of a certain volume of gas to the mass of an equal volume of hydrogen under the same conditions of temperature and pressure.</li> </ul>							
	*(ii)	Formation of ions from molecules.						
	(iii)	Electrolytic deposition of a superior metal on a baser metal.						
	0 							
	(iv)	Hydrocarbons containing a —C— functional group.						
	( <b>v</b> )	The amount of energy released when an atom in the gaseous state accepts						
		an electron to form an anion. [5]						
(d)	Match the options A to E with the statements (i) to (v) :							
	A	alkynes	(i)	No. of molec dioxide at s.t.	ules in 22.4 dm p.	<sup>3</sup> of carbon		
	B	alkane	(ii)	An element i 2, 8, 8, 3	vith electronic co	onfiguration		
			I	1				

(iii)

 $C_n H_{2n+2}$ 

C

iron

D	$6.023 \times 10^{23}$	(iv)	$C_n H_{2n-2}$
E	metal	(v)	The metal that forms two types of ions

[5]

[5]

- (e) Write balanced equations for the following :
  - (i) Action of heat on a mixture of copper and concentrated nitric acid.
  - (ii) Action of warm water on magnesium nitride.
  - (iii) Action of concentrated sulphuric acid on carbon.
  - (iv) Action of dilute hydrochloric acid on sodium sulphide.
  - (v) Preparation of ethane from sodium propionate.
- (f) Distinguish between the following pairs of compounds using the test given within brackets :
  - (i) Iron (II) sulphate and iron(III) sulphate (using ammonium hydroxide)
  - (ii) A lead salt and a zinc salt (using excess ammonium hydroxide)
  - (iii) Sodium nitrate and sodium sulphite (using dilute sulphuric acid)
  - (iv) Dilute sulphuric acid and dilute hydrochloric acid (using barium chloride solution)
  - (v) Ethane and ethene (using alkaline potassium permanganate solution.

[5]

(g) (i) Oxygen oxidises ethyne to carbon dioxide and water as shown by the equation:

$$2C_2H_2 + 5O_2 \longrightarrow 4CO_2 + 2H_2O$$

What volume of ethyne gas at s.t.p. is required to produce 8.4  $dm^3$  of carbon dioxide at s.t.p. ? [H = 1, C = 12, O = 16]

(ii) A compound made up of two elements X and Y has an empirical formula  $X_2$  Y. If the atomic weight of X is 10 and that of Y is 5 and the compound has a vapour density 25, find its molecular formula. [5]

## Answer.

- (a) (i) (D) Atomic radius decreases and nuclear charge increases.
  - (ii) (B) A low melting point and low boiling point.
  - (iii) (C) A silver grey deposit at cathode and reddish brown fumes at anode.
  - (iv) (A) Haematite
  - (v) (C) Calcination
  - (vi) (A) 3 shells and 2 valence electrons
  - (vii) (D) Nitrogen
  - (viii) (C) Sodium argentocyanide solution
  - (ix) (A) It is a strong reducing agent.
  - (x) (D) Ethyne

1

- **(b)** (i)
  - (ii) Sodium ethoxide
  - (iii) CaO is alkaline
  - (iv) a downward displacement of air
  - (v) nitric oxide.

- (c) (i) Vapour density
  - (ii) Ionisation
  - (iii) Electroplating
  - (iv) Ketone or Carbonyl compound
  - (v) Electron affinity
- (d) A (iv)  $C_n H_{2n-2}$ 
  - B (iii)  $C_n H_{2n+2}$
  - C (v) The metal that forms two types of ions
  - D (i) No. of molecules in  $22.4 \text{ dm}^3$  of carbon dioxide at s.t.p.
  - E (ii) An element with electronic configuration 2, 8, 8, 3
- (e) (i)  $Cu + 4HNO_3 \longrightarrow Cu(NO_3)_2 + 2NO_2 + 2H_2O$ 
  - (ii)  $Mg_3N_2 + 6H_2O \longrightarrow 3Mg(OH)_2 + 2NH_3$
  - (iii)  $C + 2H_2SO_4 \longrightarrow CO_2 + 2H_2O + 2SO_2$
  - (iv)  $Na_2S + 2HCl \longrightarrow 2NaCl + H_2S$
  - (v)  $C_2H_5COONa + NaOH \xrightarrow{CaO} Na_2CO_3 + C_2H_6$
- (f) (i) **Iron II sulphate :** Gives dirty green ppt with ammonium hydroxide insoluble in excess.

**Iron III sulphate :** Gives reddish brown ppt with ammonium hydroxide insoluble in excess.

(ii) Lead salt : Gives white ppt with ammonium hydroxide which is insoluble in excess.

**Zinc salt :** Gives gelatenous white ppt which is soluble in excess ammonium hydroxide.

(iii) **Sodium nitrate :** Colourless vapours of nitric acid which condenses to form nitric acid.

**Sodium sulphite :** Colourless, gas with smell of burning sulphur, acidic in nature that is sulphur di oxide is released.

- (iv) With dil. HCl,  $BaCl_2$  gives no ppt with dil.  $H_2SO_4$ ,  $BaCl_2$  gives a white insoluble ppt of  $BaSO_4$ .
- (v) With ethane, purple colour of potassium permanganate remains unfaded with ethene the purple colour gets decolourised.


(g) (i) 
$$2C_2H_2 + 5O_2 \longrightarrow 4CO_2 + 2H_2O_2$$
  
 $2vol. : 5vol. 4vol. : 2vol.$   
If 4 vol. of CO<sub>2</sub> is produced by 2 vol. of C<sub>2</sub>H<sub>2</sub> at STP  
Then 8·4 dm<sup>3</sup> of CO<sub>2</sub> is produced by  $\frac{2}{4} \times 8 \cdot 4 = 4 \cdot 2$  dm<sup>3</sup>. **Ans.**  
(ii) Emp. formula = X<sub>2</sub>Y, At. wt. of X = 10, At. wt. of Y = 5  
Empirical formula mass = 2 × 10 + 5 = 25  
If Vapour density V.D. = 25  
Mol. Mass = V.D. × 2 = 25× 2  
= 50u


$$n = \frac{\text{Mol. Mass}}{\text{Emp. formula mass}} = \frac{50}{25} = 2$$
  
Mol. formula = Emp. formula × 2  
= X<sub>2</sub>Y · × 2  
= X<sub>4</sub> Y<sub>2</sub>. Ans.

## (Answer any **four** questions from this section)

#### **Question 2.**

- (a) State your observation in each of the following cases :
  - (i) When dilute hydrochloric acid is added to sodium carbonate crystals.
  - (ii) When excess sodium hydroxide is added to calcium nitrate solution.
  - (iii) At the cathode when acidified aqueous copper sulphate solution is electrolyzed with copper electrodes.
  - (iv) When calcium hydroxide is heated with ammonium chloride crystals.
  - (v) When moist starch iodide paper is introduced into chlorine gas. [5]
- (b) Study the figure given below and answer the questions that follow :





- (i) Identify the gas Y.
- (ii) What property of gas Y does this experiment demonstrate ?
- (iii) Name another gas which has the same property and can be demonstrated through this experiment. [3]
- (c) (i) Name the other ion formed when ammonia dissolves in water.
  - (ii) Give one test that can be used to detect the presence of the ion produced.

[2]

#### Answer.

- (a) (i) Brisk effervescence with the release of a colourless odourless gas that extinguish a glowing splint and turns lime water milky *i.e.*,  $CO_2$  gas is released.
  - (ii) A white ppt of Ca(OH)<sub>2</sub> is obtained that remains insoluble in excess of NaOH.
  - (iii) The blue colour of aq.CuSO<sub>4</sub> remains unchanged.
  - (iv) A colourless pungent smelling basic gas *i.e.*, Ammonia is obtained.
  - (v) Moist starch iodide paper turns blue black.

- (b) (i) Hydrogen chloride gas (HCl).
  - (ii) Y Gas *i.e.*, HCl gas is highly soluble and acidic in nature.
  - (iii) Ammonia gas.
- (c) (i) Hydroxyl ion (OH<sup>-</sup>) other than Ammonium ion.
  - (ii) Red litmus turns blue/Methyl orange yellow/Phenolphthalein turns pink.

## **Question 3.**

(a) State the conditions required for the following reactions to take place :

- (i) Catalytic hydrogenation of ethyne.
- (ii) Preparation of ethyne from ethylene dibromide.
- (iii) Catalytic oxidation of ammonia to nitric oxide.
- (iv) Any two conditions for the conversion of sulphur dioxide to sulphur trioxide. [5]
- (b) State the main components of the following alloys :
  - (i) Brass.
  - (ii) Duralumin.
  - (iii) Bronze.
- (c) Give balanced equations for the following :
  - (i) Laboratory preparation of nitric acid.
  - (ii) Preparation of ethanol from monochloroethane and aq. sodium hydroxide. [2]

[3]

[4]

## Answer.

(a) (i) In presence of Catalyst like Ni/Pt/Pd etc.

- (ii) Heating of ethylene dibromide by using alcoholic KOH.
- (iii) In presence of Platinum catalyst at 800 °C.
- (iv) In presence of vanadium pentaoxide ( $V_2O_5$ ) or Pt as catalyst at 450 °C.
- (b) (i) **Brass**: Cu + Zn.
  - (ii) **Duralumin :** Al + Cu + Mg + Mn
  - (iii) **Bronze :** Cu + Sn.

(c) (i) 
$$NaNO_3 + H_2SO_4 \xrightarrow{< 200 \ C} NaHSO_4 + HNO_3$$
  
Nitric Acid

(ii)  $C_2H_5Cl + NaOH_{(aq)} \longrightarrow C_2H_5OH + NaCl$ Chloroethane Sod. hydroxide Ethanol

000 00

#### **Question 4.**

- (a) Give the structural formula of the following :
  - (i) *ethanol*.
  - (ii) 1-propanal
  - (iii) ethanoic acid
  - (iv) 1, 2, dichloroethane.
- (b) Draw the structure of the stable positive ion formed when an acid dissolves in water.
   [2]
- (c) State the inference drawn from the following observations :
  - (i) On carrying out the flame test with a salt P a brick red flame was obtained. What is the cation in P?

- (ii) A gas Q turns moist lead acetate paper silvery black. Identify the gas Q.
- (iii) pH of liquid R is 10. What kind of substance is R?
- (iv) Salt S is prepared by reacting dilute sulphuric acid with copper oxide. Identify S. [4]



- нн нно ł -C--С--ОН (ii) H--C **(a) (i)** H-ΗH Н Н H O н H 11 L (iv) H - C - H(iii) H\_C\_OH Η Cl Cl
- **(b)** Hydronium ion  $(H_3O^+)$

$$\mathbf{H}^{+} + \mathbf{H}_{2}\mathbf{O} \longrightarrow \mathbf{H}_{3}\mathbf{O}^{+} = \begin{bmatrix} \mathbf{H} \\ \uparrow \\ \mathbf{H} \longrightarrow \mathbf{O} \longrightarrow \mathbf{H} \\ \cdots \end{bmatrix}^{+}$$

- (c) (i)  $P \rightarrow Calcium ion (Ca^{++})$ 
  - (ii)  $\mathbf{Q} \rightarrow \mathbf{Hydrogen} \text{ sulphide gas } (\mathbf{H}_2 \mathbf{S})$
  - (iii)  $\mathbf{R} \rightarrow \mathbf{Base}$ .
  - (iv)  $S \rightarrow Copper Sulphate (CuSO_4).$

#### **Question 5.**

- (a) Name the following :
  - (i) The property possessed by metals by which they can be beaten into sheets.
  - (ii) A compound added to lower the fusion temperature of electrolytic bath in the extraction of aluminium.
  - (iii) The ore of zinc containing its sulphide. [3]

[3]

- (b) Give one equation each to show the following properties of sulphuric acid: [3]
  - (i) Dehydrating property.
  - (ii) Acidic nature.
  - (iii) As a non-volatile acid.
- (c) Give balanced chemical equations to prepare the following salts :
  - (i) Lead sulphate from lead carbonate.
  - (ii) Sodium sulphate using dilute sulphuric acid.
  - (iii) Copper chloride using copper carbonate. [4]

## Answer.

- (a) (i) Malleability.
  - (ii) Cryolite  $(Na_3AlF_6)$
  - (iii) Zinc Sulphide/Zinc blende (ZnS)

(b) (i) 
$$C_{12}H_{22}O_{11} \xrightarrow{Conc. H_2SO_4} 12C + 11H_2O$$
  
(ii)  $NaOH + H_2SO_4 \xrightarrow{<200 \ ^\circ C} NaHSO_4 + H_2O$   
(dil)  
(iii)  $KNO_3 + H_2SO_4 \xrightarrow{<200 \ ^\circ C} KHSO_4 + HNO_3$   
(c) (i)  $PbCO_3 + 2HNO_3 \longrightarrow Pb(NO_3)_2 + H_2O + CO_2$   
lead  
carbonate  
 $Pb(NO_3)_2 + H_2SO_4 \longrightarrow PbSO_4 + 2HNO_3$   
lead  
sulphate  
(ii)  $2NaOH + H_2SO_4 \xrightarrow{>200 \ ^\circ C} Na_2SO_4 + 2H_2O$   
(iii)  $CuCO_3 + 2HCI \longrightarrow CuCl_2 + H_2O + CO_2^{\uparrow}$   
Copper carbonate Copper chloride

#### **Question 6.**

- (a) (i) State Avogadro's law.
  - (ii) A cylinder contains 68g of ammonia gas at s.t.p.
    - (1) What is the volume occupied by this gas ?
    - (2) How many moles of ammonia are present in the cylinder ?
    - (3) How many molecules of ammonia are present in the cylinder ? [N-14, H-1]

[4]

[3]

Ans.

- (b) (i) Why do covalent compounds exist as gases, liquids or soft solids ?
  - (ii) Which electrode : anode or cathode is the oxidising electrode ? Why ? [3]
- (c) Name the kind of particles present in :
  - (i) Sodium Hydroxide solution.
    - (ii) Carbonic acid.
    - (iii) Sugar solution.

#### Answer.

- (a) (i) Under the similar conditions of temperature and pressure, equal volumes of all gases contains equal number of molecules.
  - (ii) (1)  $NH_3 = 14 + 3 = 17$ If 17 gm of  $NH_3$  contains  $22 \cdot 4l$  at STP Then 68 gm of  $NH_3$  contains  $\frac{22 \cdot 4}{17} \times 68 = 89 \cdot 60 l$ .
    - (2) No. of moles =  $\frac{\text{Mass in gm}}{\text{Gram molecular mass}}$ =  $\frac{68}{17}$  = 4 moles. Ans.
    - (3) One mole of NH<sub>3</sub> contains =  $6.022 \times 10^{23}$  molecules 4 moles of NH<sub>3</sub> contains =  $4 \times 6.022 \times 10^{23}$ =  $2.4088 \times 10^{24}$  molecules.
- (b) (i) Because the particles/atoms are held by weak Wander Vaal's forces.
  - (ii) Anode. Because anode is the oxidising electrode, there is loss of electrons.

- (c) (i) Ions *i.e.*,  $Na^+$  and  $OH^-$ 
  - (ii) Ions *i.e.*,  $H^+$  and  $CO_3^{2-}$

(iii) Molecules  $C_{12}H_{22}O_{11}$ .

## **Question 7.**

(a) An element Z has atomic number 16. Answer the following questions on Z:

- (i) State the period and group to which Z belongs.
- (ii) Is Z a metal or a non-metal?
- (iii) State the formula between Z and Hydrogen.
- (iv) What kind of a compound is this?

(b) M is a metal above hydrogen in the activity series and its oxide has the formula  $M_2O$ . This oxide when dissolved in water forms the corresponding hydroxide which is a good conductor of electricity. In the above context answer the following :

[5]

[5]

- (i) What kind of combination exists between M and O?
- (ii) How many electrons are there in the outermost shell of M?
- (iii) Name the group to which M belongs.
- (iv) State the reaction taking place at the cathode.
- (v) Name the product at the anode.

## Answer.

(a)

- Z = 16 = 2, 8, 6.
- (i) Period No. = 3 Group No. = VI A/16
- (ii) Non metal.
- (iii) Z's valency = -2

H = +1 So formula  $H_2Z$ 

(iv) Polar Covalent compound.

- (b) (i) Electrovalent/Ionic compound is formed.
  - (ii) 1
  - (iii) M belong to alkali metal group *i.e.*, Group-1.
  - (iv)  $M^+ + e^- \longrightarrow M$  Reduction.
  - (v) Oxygen gas.