




## Learning Objectives

**CHARLES ADOLPHE WURTZ**

He is a French organic Chemist. He discovered phosphoryl chlorides. He showed that alkylhalides react with sodium to produce higher alkanes. This reaction was later named as Wurtz reaction. He is also known for his discoveries of ethylamine glycol and aldol condensation reactions.

After studying this unit, the students will be able to

- classify hydrocarbons according to nature of bond between the carbon atoms.
- name hydrocarbons according to IUPAC system of nomenclature.
- describe various methods of preparing hydrocarbon.
- interpret and use the following terminology in organic chemistry like free radicals, initiation, propagation and termination.
- describe chemistry of alkanes by the following reaction of halogenations, combustion, cracking.
- predict the formation of addition products of unsymmetrical alkenes which applies both Markovnikov's rule and peroxide effect.
- deduce acidic hydrogen in alkynes.
- comprehend the structure of benzene, explain aromaticity and understand mechanism of electrophilic substitutions reactions.
- predict the directive influence of substituents on mono substituted benzene.
- recognize the toxicity and carcinogenic nature of aromatic hydrocarbons.

## Introduction

The hydrocarbons are compounds composed entirely of Carbon and Hydrogen. They occur widely in nature. The crude oil consists of complex mixtures of hydrocarbons, mangoes contain cyclohexane, a cyclic hydrocarbon, cockroaches secrete a hydrocarbon, undecane which attract opposite gender of its species. Hydrocarbons are primarily used as fuel. For example, Liquefied mixture of propane and butane is used as Liquefied petroleum gas (LPG). They also find many applications in industries such as solvents etc. In this unit we will study the classification, preparation, properties and uses of aliphatic and aromatic hydrocarbons.

### 13.1 Introduction and classification of alkanes:

Depending upon the characteristic pattern of bonding between the carbon atoms, hydrocarbons are divided into two main classes: aliphatic and aromatic. The word aliphatic was derived from the Greek word 'aleiphar' meaning fat. Important sources of aliphatic hydrocarbons are oils and fats. The word 'aroma' means odour, which is obtained by chemical treatment of pleasant-smelling plant extracts.

Aliphatic hydrocarbons include three major groups: alkanes, alkenes and alkynes. Alkanes are saturated hydrocarbons in which all the bonds between the carbon atoms are single bond, alkenes consist of at least one carbon-carbon double bond, and alkynes have at least one carbon-carbon triple bond. Hydrocarbons having localised carbon - carbon multiple bonds are called unsaturated hydrocarbons.

Aromatic hydrocarbons are cyclic compounds which contain characteristic benzene ring or its derivatives. The classification of hydrocarbons is as shown below.




Fig.13.1. Classification of Hydrocarbons

### 13.2. Alkanes:

Alkanes are saturated hydrocarbons represented by the general formula  $C_nH_{2n+2}$  where 'n' is the number of carbon atoms in the molecule. Methane  $CH_4$ , is the first member of alkane family. The successive members are ethane  $C_2H_6$ , propane  $C_3H_8$ , butane  $C_4H_{10}$ , pentane  $C_5H_{12}$  and so on. It is evident that each member differs from its preceding or succeeding member by a  $-CH_2$  group.


**DO  
YOU  
KNOW?**

Methane is the major component of the atmosphere of Jupiter, Saturn, Uranus and Neptune but only minor component of earth atmosphere. However, decomposition of plant and animal matter in an oxygen deficient environment like swamps, marshes, bogs and the sediments of lakes produces methane gas. So, it is otherwise known as marsh gas. It is also associated with coal and petroleum deposits.

**DO  
YOU  
KNOW?**

#### "FLAMMABLE ICE"

This catchy phrase describes a frozen mixture of water and methane gas chemically known as methane clathrates. The methane molecule which is produced by biological process under the deep-ocean, (at  $4^\circ C$  & 50 atm) doesn't simply reach the surface, instead each molecule is trapped inside clusters of 6 to 18 water molecules forming methane clathrates. Many countries are working on how to tap out these vast resources of natural gas but mining and extracting are very difficult.





### Nomenclature and isomerism:

We have already discussed the nomenclature of organic compound in Unit:11. Let us understand the nomenclature and isomerism in few examples. The first three members methane  $CH_4$ , ethane  $C_2H_6$  and propane  $C_3H_8$  have only one structure.

IUPAC Name	Molecular Formula	Condensed Structural formula
Methane	CH <sub>4</sub>	CH <sub>4</sub>
Ethane	C <sub>2</sub> H <sub>6</sub>	CH <sub>3</sub> -CH <sub>3</sub>
Propane	C <sub>3</sub> H <sub>8</sub>	CH <sub>3</sub> -CH <sub>2</sub> -CH <sub>3</sub>




However, higher members can have more than one structure leading to constitutional isomers (differ in connectivity) or structural isomers. For example, an alkane with molecular formula C<sub>4</sub>H<sub>10</sub> can have two structures. They are n-butane and iso-butane. In n-butane, all the four carbon atoms are arranged in a continuous chain. The 'n' in n-butane stand for 'normal' and means that the carbon chain is unbranched. The second isomer iso-butane has a branched carbon chain. The word iso indicates it is an isomer of butane.

Molecular Formula	Structural Formula	Bond line Formula	IUPAC name	Common Name
C <sub>4</sub> H <sub>10</sub>	CH <sub>3</sub> -CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>3</sub> <div style="text-align: center;">       CH<sub>3</sub>                 CH<sub>3</sub>-CH-CH<sub>3</sub> </div>	 	butane  2-methylpropane	n-butane  iso-butane

Though both the structures have same molecular formula but their carbon chains differ leading to chain isomerism

Let us understand the chain isomerism by writing the isomers of pentane C<sub>5</sub>H<sub>12</sub>

### Solution:

To begin, draw the carbon backbone of the straight chain-isomer	C—C—C—C—C
Determine the carbon backbone structure of the other isomers by arranging the carbon atoms in the other way	<div style="display: flex; align-items: center; justify-content: space-around;"> <div style="text-align: center;">       C—C—C—C                 C     </div> <div style="text-align: center;">       C—C—C                 C     </div> </div>
Fill in all the hydrogen atoms so that each carbon forms four bonds	<div style="display: flex; justify-content: space-around;"> <div>CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>3</sub></div> <div>       CH<sub>3</sub>-CH-CH<sub>2</sub>-CH<sub>3</sub>                 CH<sub>3</sub> </div> <div>       CH<sub>3</sub>                 CH<sub>3</sub>-C-CH<sub>3</sub>                 CH<sub>3</sub> </div> </div>
	<div style="display: flex; justify-content: space-around; align-items: center;">    </div>

## Evaluate Yourself

1) Write the structural formula and carbon skeleton formula for all possible chain isomers of  $C_6H_{14}$  (Hexane)

### IUPAC name for some branched alkanes

Let us write the IUPAC name for the below mentioned alkanes by applying the general rules of nomenclature that we already discussed in unit No.11




S.NO	COMPOUND	IUPAC NAME
1	$  \begin{array}{ccccccc}  & 5 & 4 & 3 & 2 & 1 \\  & CH_3 & -CH_2- & CH_2- & CH- & CH_3 \\  & & & &   \\  & & & & CH_3  \end{array}  $	2- Methyl pentane
2	$  \begin{array}{ccccccc}  & CH_3 & -CH- & CH_2- & CH- & CH_3 \\  & 1 & 2 & 3 & 4 & 5 \\  & &   & &   \\  & & CH_3 & & CH_3  \end{array}  $	2,4- Dimethyl pentane
3	$  \begin{array}{ccccccc}  & & & CH_3 & & & \\  & & &   & & & \\  & CH_3 & -CH_2- & C- & CH_2- & CH_3 \\  & 12 & & 3 & 4 & 5 \\  & & &   \\  & & & CH_3  \end{array}  $	3,3- Dimethyl pentane
4	$  \begin{array}{ccccccc}  & CH_3 & -CH- & CH_2- & CH_2- & CH_3 \\  & 1 & 2 & 3 & 4 & 5 \\  & &   & & & \\  & & CH_3 & & & \\  & &   & & & \\  & & CH_2 & & & \\  & &   & & & \\  & & CH_3 & & &  \end{array}  $	3-Ethyl-2-methylpentane
5	$  \begin{array}{ccccccc}  & CH_3 & -CH_2- & CH- & CH- & CH- & CH_2- & CH_2- & CH_3 \\  & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\  & & &   &   &   \\  & & & CH_2 & CH_2 & CH_2 \\  & & &   &   &   \\  & & & CH_3 & CH_2 & CH_2 \\  & & & &   &   \\  & & & & CH_3 & CH_3  \end{array}  $	3-Ethyl-4,5-dipropyl octane
6	$  \begin{array}{ccccccc}  & CH_3 & -CH- & CH- & CH_3 \\  & &   &   \\  & & CH_2 & CH_3 \\  & &   \\  & & CH_3  \end{array}  $	2,3 - Dimethylpentane



## 1. Preparation of alkanes from catalytic reduction of unsaturated hydrocarbons.

When a mixture hydrogen gas with alkene or alkyne gas is passed over a catalysts such as platinum or palladium at room temperature, an alkane is produced. This process of addition of  $H_2$  to unsaturated compounds is known as hydrogenation. The above process can be catalysed by nickel at 298K. This reaction is known as Sabatier-Sendersens reaction

for example:




## 2. Preparation of alkanes from carboxylic acids:

### i) Decarboxylation of sodium salt of carboxylic acid


When a mixture of sodium salt of carboxylic acid and soda lime (sodium hydroxide + calcium oxide) is heated, alkane is formed. The alkane formed has one carbon atom less than carboxylic acid. This process of eliminating carboxylic group is known as decarboxylation.

for example:



### ii) Kolbe's Electrolytic method

When sodium or potassium salt of carboxylic acid is electrolyzed, a higher alkane is formed. The decarboxylative dimerization of two carboxylic acid occurs. This method is suitable for preparing symmetrical alkanes (R-R).




## 3. Preparation of alkanes using alkyl halides (or) halo alkanes

### i) By reduction with nascent hydrogen

Except alkyl fluorides, other alkyl halides can be converted to alkanes by reduction with nascent hydrogen. The hydrogen for reduction may be obtained by using any of the following reducing agents:  $\text{Zn} + \text{HCl}$ ,  $\text{Zn} + \text{CH}_3\text{COOH}$ ,  $\text{Zn-Cu}$  couple in ethanol,  $\text{LiAlH}_4$  etc.,

for example:



### ii) Wurtz reaction

When a solution of halo alkanes in dry ether is treated with sodium metal,





higher boiling point compared to branch chain isomers. The boiling point decreases with increase in branching as the molecule becomes compact and the area of the contact decreases.

## 2) Solubility and density

Water molecules are polar and alkanes are non-polar. The insolubility of alkanes in water makes them good water repellent for metals which protects the metal surface from corrosion. Because of their lower density than water, they form two layers and occupy top layer. The density difference between alkanes and water explains why oil spills in aqueous environment spread so quickly.



Apples produce its own wax which contains long unbranched alkanes ( $C_{27}H_{56}$  and  $C_{29}H_{60}$ ). This natural wax helps the apple to resist moisture lost, enhances the fruit firmness and slows down its natural degradation. The consumption of apple with artificial wax coating is harmful to the digestive system




infinite number of readily interconvertible three dimensional arrangements. Such readily interconvertible three dimensional arrangement of a molecule is called conformations.

### (i) Conformations of ethane:

The two tetrahedral methyl groups can rotate about the carbon – carbon bond axis yielding several arrangements called conformers. The extreme conformations are staggered and eclipsed conformation. There can be number of other arrangements between staggered and eclipsed forms and their arrangements are known as skew forms.

#### Eclipsed conformation:



In this conformation, the hydrogen's of one carbon are directly behind those of the other. The repulsion between the atoms is maximum and it is the least stable conformer.

### 13.2.3 Conformations of alkane:

Each carbon in alkanes is  $sp^3$  hybridized and the four groups or atoms around the carbon are tetrahedrally bonded. In alkanes having two or more carbons, there exists free rotation about C-C single bond. Such rotation leaves all the groups or atoms bonded to each carbon into an

### Staggered conformation:



In this conformation, the hydrogens of both the carbon atoms are far apart from each other. The repulsion between the atoms is minimum and it is the most stable conformer.


### Skew Conformation :

The infinite numbers of possible intermediate conformations between the two extreme conformations are referred as skew conformations.

The stabilities of various conformations of ethane are

Staggered > Skew > Eclipsed

The potential energy difference between the staggered and eclipsed conformation of ethane is around  $12.5 \text{ KJmol}^{-1}$ . The various conformations can be represented by new man projection formula.



Newman projection formula for Ethane

### Conformations of n-Butane:

n-Butane may be considered as a

derivative of ethane, as one hydrogen on each carbon is replaced by a methyl group


### Eclipsed conformation:

In this conformation, the distance between the two methyl group is minimum. So there is maximum repulsion between them and it is the least stable conformer.

### Anti or staggered form

In this conformation, the distance between the two methyl groups is maximum and so there is minimum repulsion between them. And it is the most stable conformer.

The following potentially energy diagram shows the relative stabilities of various conformers of n-butane.



### 13.2.4 Chemical properties:

Alkanes are quite unreactive towards most reagents. However under favorable conditions, alkanes undergo the following type of reaction.


Paraffin is the older name for the alkane group family of compounds. This name comes from the Latin which means 'little activity'

#### 1) Combustion:


A combustion reaction is a chemical reaction between a substance and oxygen with evolution of heat and light (usually as a flame). In the presence of sufficient oxygen, alkanes undergo combustion when ignited and produce carbon dioxide and water.

The combustion reaction is expressed as follows

for example:



When alkanes burn in insufficient supply of oxygen, they form carbon monoxide and carbon black.



#### Evaluate Yourself


6) Write down the combustion reaction of propane whose  $\Delta H^\circ = -2220 \text{ kJ}$

#### 2) Halogenation:

A halogenation reaction is the chemical reaction between an alkane and

halogen in which one or more hydrogen atoms are substituted by the halogens.

Chlorination and Bromination are two widely used halogenation reactions. Fluorination is too quick and iodination is too slow. Methane reacts with chlorine in the presence of light or when heated as follows.



#### Mechanism:

The reaction proceeds through the free radical chain mechanism. This mechanism is characterized by three steps: initiation, propagation and termination.


i) **CHAIN INITIATION:** The chain is initiated by UV light leading to homolytic fission of chlorine molecules into free radicals (chlorine atoms).




Here we choose Cl-Cl bond for fission because C-C & C-H bonds are stronger than Cl-Cl.

ii) **PROPAGATION:** It proceeds as follows,

(a) Chlorine free radical attacks the methane molecule and breaks the C-H bond resulting in the generation of methyl free radical




(b) The methyl free radical thus obtained attacks the second molecule of chlorine to give chloromethane ( $\text{CH}_3\text{Cl}$ ) and a chlorine free radical as follows.



(c) This chlorine free radical then cycles back to step (a) and both step (a) and (b) are repeated many times and thus chain of reaction is set up.

### iii) Chain termination:

After sometimes, the reactions stops due to consumption of reactant and the chain is terminated by the combination of free radicals.



### Evaluate Yourself


7) Why ethane is produced in chlorination of methane?

### 3) Aromatisation

Alkanes with six to ten carbon atoms are converted into homologous of benzene at high temperature and in the presence of catalyst. This process is known

as aromatization. It occurs by simultaneous cyclisation followed by dehydrogenation of alkanes.

n-Hexane passed over  $\text{Cr}_2\text{O}_3$  supported on alumina at 873 K gives benzene.




### Evaluate Yourself

8) How toluene can be prepared by this method?

### 4) Reaction With Steam:

Methane reacts with steam at 1273K in the presence of Nickel and decomposes to form carbon monoxide and hydrogen gas.




Production of  $\text{H}_2$  gas from methane is known as steam reforming process and it is a well-established industrial process for the production of  $\text{H}_2$  gas from hydrocarbons.

### 5) Pyrolysis

Pyrolysis is defined as the thermal decomposition of organic compound into smaller fragments in the absence of air through the application of heat. 'Pyro' means 'fire' and 'lysis' means 'separating'. Pyrolysis of alkanes also named as cracking.

In the absence of air, when alkane vapours are passed through red-hot metal it


breaks down into simpler hydrocarbons.



The products depends upon the nature of alkane, temperature, pressure and presence or absence of catalyst. The ease of cracking in alkanes increases with increase in molecular weight and branching in alkanes. Cracking plays an important role in petroleum industry.

### 6) Isomerisation:

Isomerisation is a chemical process by which a compound is transformed into any its isomeric forms. Normal alkanes can be converted into branched alkanes in the presence of  $\text{AlCl}_3$  and  $\text{HCl}$  at 298 K.



This process is of great industrial importance. The quality of gasoline is improved by isomerising its components.

### Uses

The exothermic nature of alkane combustion reaction explains the extensive use of alkanes as fuels. Methane present in natural gas is used in home heating. Mixture of propane and butane are known as LPG gas which is used for domestic cooking purpose. GASOLINE is a complex mixture of many hydrocarbons used as a fuel for internal-

combustion engines.

Carbon black is used in the manufacture of ink, printer ink and black pigments. It is also used as fillers.

No of Carbon Atoms	State at room temperature	Major uses
1-4	Gas	Heating fuel, Cooking fuel
5-7	Low boiling liquid	Solvents, Gasoline
6-12	Liquid	Gasoline
12-24	Liquid	Jet fuel- portable stove fuel
18-50	High boiling liquid	Diesel fuel, lubricant, heating oil
50+	Solid	Petroleum jelly and paraffin wax

## 13.3. Alkenes:

Alkenes are unsaturated hydrocarbons that contain carbon-carbon double bond. They are represented by the general formulae  $\text{C}_n\text{H}_{2n}$  where 'n' stands for number of carbon atoms in the molecule. Alkenes are also known as olefins (in Latin - oil maker) because the first member ethene combines with chlorine gas to form an oily liquid as a product.

### (I) Nomenclature Of Alkenes:

Let us write the IUPAC name for the below mentioned alkenes by applying the general rules of nomenclature that we

already discussed in unit No.11





TAB 13.4 IUPAC NAMES FOR ALKENES

S No	Structure	IUPAC name
1	$\text{CH}_3-\text{CH}=\text{CH}_2$	propene
2	$\text{CH}_3-\text{CH}=\text{CH}-\text{CH}_3$	2-butene
3	$\begin{array}{c} \text{CH}_2=\text{C}-\text{CH}_3 \\   \\ \text{CH}_3 \end{array}$	2-methyl-1-propene
4	$\begin{array}{c} \text{CH}_2=\text{CH}-\text{CH}-\text{CH}_2-\text{CH}_3 \\   \\ \text{CH}_3 \end{array}$	3-methyl-1-pentene
5	$\begin{array}{ccccccc} \text{CH}_2 & = & \text{CH} & - & \text{CH} & - & \text{CH}_2 - \text{CH}_3 \\ 1 & & 2 & & 3 & & 4 & & 5 \\ & & & &   & & & & \\ & & & & \text{CH}_3 & & & & \end{array}$	3-methyl-1-pentene

### Evaluate Yourself



1. Write the IUPAC names for the following alkenes.



2. Draw the structures for the following alkenes.



- 6 - Bromo - 2,3 - dimethyl - 2 - hexene
- 5 - Bromo - 4 - chloro - 1 - heptene
- 2,5 - Methyl - 4 - octene
- 4 - Methyl - 2 pentene

### (ii) Isomerism:

Presence of double bond in Alkene provides the possibility of both structural and geometrical isomerism.

#### Structural Isomerism:

The first two member's ethene  $\text{C}_2\text{H}_4$  and propene  $\text{C}_3\text{H}_6$  do not have isomers because the carbon atoms in the molecules can be arranged only one distinct way. However from the third member of alkene family butene  $\text{C}_4\text{H}_{10}$ , structural isomerism exists.



structures (i) & (ii) are position isomers. structures (i) & (iii), (ii) & (iii) are chain isomers.

### Evaluate Yourself



9) Draw the structure and write down the IUPAC name for the isomerism exhibited by the molecular formulae:



#### Geometrical isomerism:


It is a type of stereoisomerism and it is also called cis-trans isomerism. Such type of isomerism results due to the restricted rotation of doubly bounded carbon atoms.

If the similar groups lie on the same side, then the geometrical isomers are called Cis-isomers. When the similar groups lie on



the opposite side, it is called a Trans isomer.

for example: the geometrical isomers of 2-Butane is expressed as follows



### Evaluate Yourself

10) Determine whether each of the following alkenes can exist in cis-trans isomers?

- (a) 1 - Chloro propene
- (b) 2 - Chloro propene


11) Draw cis-trans isomers for the following compounds

- (a) 2- chloro - 2 - butene
- (b)  $\text{CH}_3 - \text{CCl} = \text{CH} - \text{CH}_2\text{CH}_3$


### 13.3.1: General methods of preparation of alkenes:

#### (1) Preparation of alkene by dehydration of alcohol:

When an alcohol is heated at 430-440 K with excess of concentrated sulphuric acid, a molecule of water from alcohol is removed and an alkene is formed. This reaction is called elimination reaction.




Ethene can also be prepared in laboratory by catalytic dehydration of alcohol.




#### (2) Preparation of alkenes from alkynes:

Alkynes can be reduced to cis-alkenes using Lindlar's catalyst ( $\text{CaCO}_3$  supported in palladium partially deactivated with sulphur (or) gasoline). This reaction is stereo specific giving only the cis- alkene.




Alkynes can also be reduced to trans-alkenes using sodium in liquid ammonia. This reaction is stereospecific giving only the trans-alkene.




#### (3) Preparation of alkenes by dehydrohalogenation of halo alkanes.

Halo alkanes react with alcoholic KOH and eliminate hydrohalide resulting in the formation of alkene.



#### (4) Preparation of alkenes from vicinal dihalogen derivative of alkanes or vicinal dihalides

The compound in which two halogen atoms are attached to adjacent carbon-atoms are called as vicinal dihalides. When vicinal dihalides are warmed with granulated zinc in methanol, they lose a molecule of  $\text{ZnX}_2$  to form an alkene.




#### Evaluate Yourself


12) How propene is prepared from 1, 2-dichloro propane?

#### (5) Preparation of ethene by Kolbe's electrolytic method:

When an aqueous solution of potassium succinate is electrolyzed between two platinum electrodes, ethene is produced at the anode.



At anode



#### 13.3.2. Physical properties of alkenes:

The first three members (Ethene, Propene and Butene) are gases, next fourteen members are liquids and the higher alkenes are waxy solids. They are all colourless and odourless except ethene which has a sweet smell.

1. The melting and boiling point of alkenes increases along the homologous series. Like alkanes, straight chain alkenes have high boiling point compared to its isomeric branched alkenes.
2. Alkenes are slightly soluble in water but readily in organic solvents.

#### 13.3.3. Chemical properties of alkenes:

Alkenes are more reactive than alkanes due to the presence of a double bond. The  $\sigma$ -bond is strong but the  $\pi$ -bond is weak. The typical reactions of alkenes involve addition of an electrophile across the double bonds proceeding through ionic mechanism. However addition reactions proceed through free-radical mechanism. Ozonolysis and polymerization are some of the characteristic reactions of alkenes.

##### (i) Addition Reactions

##### (ii) Addition of hydrogen: (Hydrogenation of alkenes)


Hydrogen adds on to alkenes in the presence of a metal catalyst (Ni, Pd (or) Pt) to yield corresponding alkanes. This is known as catalytic hydrogenation. This process is




of great importance in the manufacture of vanaspathi from vegetable oil. This helps to prevent rancidity of vegetable oils.

## (ii) Addition of halogens: (Halogenation of alkenes)

When alkene is treated with halogens like chlorine or bromine, addition takes place rapidly and forms 1, 2- dihalo alkane (or) vicinal dihalide.



Iodine reacts very slowly to form 1, 2 – diiodo alkane which are unstable and regenerate the original alkene by elimination of iodine.



## TEST FOR ALKENE:




Bromine in water is reddish brown colour. When small amount of bromine water is added to an alkene, the solution is decolourised as it forms dibromo compound. So, this is the characteristic test for unsaturated compounds.

## Markovnikoff's rule:

“When an unsymmetrical alkene reacts with hydrogen halide, the hydrogen adds to the carbon that has more number of hydrogen and halogen adds to the carbon having fewer hydrogen”. This rule can also be stated as in the addition reaction of alkene / alkyne, the most electro negative part of the reagent adds on to the least hydrogen attached doubly bonded carbon.

## (iii) Addition of water:- (Hydration of alkenes)

Normally, water does not react with alkenes. In the presence of concentrated sulphuric acid, alkenes react with water to form alcohols. This reaction follows carbocation mechanism and Markovnikoff's rule.





## (iv) Addition of hydrohalides: (Hydrohalogenation of Alkenes)

Hydrogen halides (HCl, HBr and HI) add to alkene to yield alkyl halides. The order of reactivity of different hydrogen halides is  $\text{HI} > \text{HBr} > \text{HCl}$ . It is an example for electrophilic addition.


### (a) Addition of HBr to symmetrical alkene:

Addition of HBr to symmetrical alkene (similar groups are attached to double bond) yields alkyl halides (haloalkanes)



**(b) Addition HBr to unsymmetrical alkene:**

In the addition of hydrogen halide to an unsymmetrical alkene, two products are obtained.




### Mechanism:

Consider addition of HBr to propene

**Step: 1** Formation of electrophile:


In H-Br, Br is more electronegative than H. When bonded electron moves toward Br, polarity is developed and creates an electrophile  $H^+$  which attaches the double bond to form carbocation, as shown below.




**Step:2** Secondary carbocation is more stable than primary carbocation and it predominates over a the primary carbocation.

**Step:3** The Br<sup>-</sup>ion attack the 2° carbocation to form 2-Bromobutane, the major product.

Consider addition of HBr to 3-methyl-1-butene. Here the expected product according to markovnikoff's rule is 2-bromo-3-methyl butane but the actual major product is 2-Bromo-2-methyl butane. This is because, the secondary carbocation formed during the reaction rearranged to more stable tertiary carbocation. Attack of  $\text{Br}^-$  on this tertiary carbocation gives the major product 2-bromo-2-methyl butane.



The addition of HBr to an alkene in the presence of organic peroxide, gives the anti Markovnikoff's product. This effect is called peroxide effect.




### Mechanism:

The reaction proceeds via free radical mechanism.

#### Step:1

The weak O-O single bond linkages of peroxides undergoes homolytic cleavage to generate free radical.



#### Step:2


The radicals abstracts a hydrogen from HBr thus generating bromine radical.




#### Step:3

The Bromine radical adds to the double bond in the way to form more stable alkyl free radical.

### Carbocation rearrangement




### Anti-Markovnikoff's Rule (Or) Peroxide Effect (Or) Kharasch Addition



#### Step:4

Addition of HBr to secondary free radical



The H-Cl bond is stronger (430.5


kJmol<sup>-1</sup>) than H-Br bond (363.7 kJmol<sup>-1</sup>), thus H-Cl is not cleaved by the free radical. The H-I bond is weaker (296.8 kJ mol<sup>-1</sup>), than H-Cl bond. Thus H-I bond breaks easily but iodine free radicals combine to form iodine molecules instead of adding to the double bond and hence peroxide effect is not observed in HCl& HI.

#### Kharasch Addition

Metal catalysed free radical addition of CXCl<sub>3</sub> Compounds to alkene is called Kharash addition reaction

#### (v) Addition of sulphuric acid to alkenes



Alkenes react with cold and concentrated sulphuric acid to form alkyl hydrogen sulphate accordance with Markownikoff 's rule. Further hydrolysis yields alcohol.



#### (2) Oxidation:



##### (i) With cold dilute alkaline KMnO<sub>4</sub> solution (Baeyer's Reagent)

Alkenes react with Baeyer's reagent to form vicinal diols. The purple solution (Mn<sup>2+</sup>) becomes dark green (Mn<sup>6+</sup>), and then produces a dark brown precipitate (Mn<sup>4+</sup>).




## (ii) With acidified KMnO<sub>4</sub> Solution:

Alkenes react with acidified KMnO<sub>4</sub> solution and are oxidised to ketones or carboxylic acid depends on the substituent at the olefinic carbon atom.. The purple solution becomes colourless. This is one of the test for unsaturation.



## (iii) Ozonolysis:

Ozonolysis is a method of oxidative cleavage of alkenes or alkynes using ozone and forms two carbonyl compounds. Alkenes react with ozone to form Ozonide and it is cleaved by Zn/H<sub>2</sub>O to form smaller molecules. This reaction is often used to identify the structure of unknown alkene or alkyne by detecting the position of double or triple bond.



## Evaluate Yourself

13) How ozone reacts with 2-methyl propene?


14) An organic compound (A) on ozonolysis gives only acetaldehyde. (A) reacts with Br<sub>2</sub>/CCl<sub>4</sub> to give compound (B) Identify the compound (A) and (B). Write the IUPAC name of (A) and (B). Give the Geometrical isomers of (A)

15) An organic compound (A) C<sub>2</sub>H<sub>4</sub> decolourises bromine water. (A) on reaction with chlorine gives (B) A reacts with HBr to give (C). identify (A), (B), (C), Explain the reactions.

#### (iv) Polymerisation:

A polymer is a large molecule formed by the combination of larger number of small molecules. The process is known as polymerisation. Alkenes undergo polymerisation at high temperature and pressure, in the presence of a catalyst.

for example










#### Recycling plastics

Extensive use of polymers clogs up landfills and pollute the environment. Because of diversity of polymers in consumer products, recycling requires sorting the polymers into various sub-types, labels with codes and symbols, which are then recycled separately.

**Table shows the codes and symbols used in recycling of ethene-based addition polymers.**

(Lower the number, greater the ease of recycling the material)

CODE	TYPE	NAME	EXAMPLES
PET		Polyethylene terephthalate	Soft drinks bottles, jars, vegetable oil bottle.
HDPE		High-density polyethylene	Milk, water and juice containers
PVC		Polyvinyl chloride	Shampoo bottles, plastic pipes
LDPE		Low density polyethylene	Sandwich bags, grocery bags
PP		Polypropylene	Straws, diaper, toys
PS		Polystyrene	Disposable utensil, foam cups
Other		Multilayer plastics	Various flexible item.

#### 13.3.4. Uses of Alkenes

1) Alkenes find many diverse applications in industry. They are used as starting materials in the synthesis of alcohols, plastics, liquors, detergents and fuels

2) Ethene is the most important organic feed stock in the polymer industry. E.g. PVC, Sarans and polyethylene. These polymer are used in the manufacture of floor tiles, shoe soles, synthetic fibres, raincoats, pipes etc.,

### 13.4. Alkenes:

Alkynes are unsaturated hydrocarbons that contain carbon-carbon triple bonds in their molecules. Their general formula is  $C_n H_{2n-2}$ . The first member of alkyne series is Ethyne popularly known as acetylene. Oxyacetylene torch is used in welding.

#### Nomenclature of alkynes:

Let us write the IUPAC name for the below mentioned alkynes by applying the general rules of nomenclature that we already discussed in unit No.11


Structure	IUPAC name	Carbon Skeleton Formula
$CH_3-C\equiv CH$	Propyne	
$CH_3-CH_2-C\equiv CH$	1-butyne	
$CH_3-C\equiv C-CH_3$	2-butyne	
$CH_3-\underset{\substack{  \\ CH_3}}{CH}-C\equiv CH$	2-methyl-1-butyne	

#### 13.4.1. General Methods Of Preparation Of Alkynes

##### 1. Preparation of alkynes from alkenes:


This process involves two steps:

- Halogenation of alkenes to form vicinal dihalides
- Dehalogenation of vicinal dihalides to form alkynes.




##### 2. Preparation of alkene from gem dihalides:

A compound containing two halogen atoms on the same carbon atom is called gem dihalide (Latin word 'Gemini' means twins). On heating with alcoholic KOH, gem dihalides give alkynes.




### 3. Preparation Of Alkenes From Electrolysis Of Salts Of Unsaturated Dicarboxylic Acids. (Kolbe's Electrolytic Method)

Electrolysis of sodium or potassium salt of maleic or fumaric acid yields alkynes.



### 4. Industrial preparation of ethyne:

Ethyne can be manufactured in large scale by action of calcium carbide with water.




Calcium carbide required for this reaction is prepared by heating quick lime and coke in an electric furnace at 3273 K



### Evaluate Yourself

- 16) Prepare propyne from its corresponding alkene.
- 17) Write the products A & B for the following reaction.



### 13.4.2. Physical properties of alkynes:

1. The first three members are gases, next eight are liquids and the higher alkynes are solids. They are all colourless and odourless except acetylene which has a garlic odour.
2. They are slightly soluble in water but dissolve readily in organic solvents like benzene, acetone and ethyl alcohol

### 13.4.3. Chemical properties of alkynes


Terminal Alkynes are acidic in nature. It undergoes polymerization and addition reaction.

#### 1. Acidic nature of alkynes:

An alkyne shows acidic nature only if it contains terminal hydrogen. This can be explained by considering sp hybrid orbitals of carbon atom in alkynes. The percentage of s-character of sp hybrid orbital (50%) is more than sp<sup>2</sup> hybrid orbital of alkene (33%) and sp<sup>3</sup> hybrid orbital of alkane (25%). Because of this, Carbon becomes more electronegative facilitating donation of H<sup>+</sup>




ions to bases. So hydrogen attached to triply bonded carbon atoms is acidic.




## 2. Addition reactions of alkynes

### i) addition of hydrogen




### ii) Addition Of Halogens:

When Br<sub>2</sub> in CCl<sub>4</sub> (Reddishbrown) is added to an alkyne, the bromine solution is decolourised. This is the test for unsaturation.




### iii) Addition Of Hydrogen Halides:

Reaction of hydrogen halides to symmetrical alkynes is electrophilic addition reaction. This reaction also follows Markovnikoff's rule.




Addition of HBr to unsymmetrical alkene follows Markownikoff's rule.





#### iv) Addition Of Water:

Alkynes undergo hydration on warming with mercuric sulphate and dilute  $\text{H}_2\text{SO}_4$  at 333K to form carbonyl compounds.



### 3. Ozonolysis:

Ozone adds to carbon-carbon triple bond of alkynes to form ozonides. The ozonides are hydrolyzed by water to form carbonyl compounds. The hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) formed in the reaction may oxidise the carbonyl compound to carboxylic acid.




## 4. Polymerisation:

Alkyne undergoes two types of polymerisation reaction


### (i) Linear Polymerisation:

Ethyne forms linear polymer, when passed into a solution of cuprous chloride and ammonium chloride.



### (ii) Cyclic Polymerisation:


Ethyne undergoes cyclic polymerisation on passing through red hot iron tube. Three molecules of ethynepolymerise to benzene.



## 13.4.4 Uses of Alkynes

- 1) Acetylene is used in oxy acetylene torch used for welding and cutting metals.
- 2) It is used for manufacture of PVC, polyvinyl acetate, polyvinyl ether, orlon and neoprene rubbers.

### Evaluate Yourself




## 13.5. Aromatic Hydrocarbons

Take a moment and think of substances that have a strong fragrance. What kind of things come to your mind?

Perfume, Vanilla or cinnamon? They smell differently, they have something in common. These substances are made of aromatic compounds [Greek: Aroma-Pleasant smelling]. However, some compounds are chemically aromatic but do not have distinct smell. The aromatic hydrocarbons are classified depending upon number of rings present in it.


(i) Monocyclic aromatic hydrocarbon (MAH)

(Ex) Benzene ( $\text{C}_6\text{H}_6$ ) and Toluene ( $\text{C}_7\text{H}_8$ )



(ii) Polycyclic aromatic hydrocarbon (PAH)

(Ex) Naphthalene ( $\text{C}_{10}\text{H}_8$ ) and Anthracene ( $\text{C}_{14}\text{H}_{10}$ )



### Evaluate Yourself


19) Calculate the number of rings present in  $\text{C}_{18}\text{H}_{12}$ .

### 13.5.1. Nomenclature and Isomerism


- We have already discussed about nomenclature of aromatic hydrocarbons in Unit:11. The first member of aromatic hydrocarbon is benzene ( $\text{C}_6\text{H}_6$ )

represented by a regular hexagon with a circle inscribed in it.

- Since, all the six hydrogen atom in benzene are equivalent, it can give only one mono-substituted compound (Ex) methyl benzene ( $C_6H_5-CH_3$ ) which named as toluene.



- When di substitution occurs either by a similar monovalent atom or two different atoms or groups in benzene, then three different position isomers are possible. Their relative positions are indicated as ortho (1,2), meta (1,3) and para (1,4). For example, consider dimethyl benzene which is named as xylene.



### Evaluate Yourself

- 20) write all possible isomers for an aromatic benzenoid compound having the molecular formula  $C_8H_{10}$
- 21) write all possible isomers for a monosubstituted aromatic benzenoid compound having the molecular formula  $C_9H_{12}$







#### 13.5.2. Aromaticity

Huckel proposed that aromaticity is a function of electronic structure. A compound may be aromatic, if it obeys the following rules

- The molecule must be co-planar
- Complete delocalization of  $\pi$  electron in the ring
- Presence of  $(4n+2)$   $\pi$  electrons in the ring where  $n$  is an integer ( $n=0,1,2,\dots$ )

This is known as Huckel's rule.

Some of the examples for Huckel rule

1.	 <p><b>Benzene</b></p>	<p>(i) The benzene is a planar molecule  (ii) It has six delocalised <math>\pi</math> electrons  (iii) <math>4n + 2 = 6</math>  <math>4n = 6 - 2</math>  <math>4n = 4</math>  <math>n = 1</math>  it obeys Huckel's <math>(4n+2)</math> <math>\pi</math> electron rule with <math>n = 1</math>  hence, benzene is aromatic.</p>
2.	 <p><b>Naphthalene</b></p>	<p>(i) Naphthalene has a planar ring structure  (ii) It has 10 delocalised <math>\pi</math> electrons  (ii) <math>4n + 2 = 10</math>  <math>4n = 10 - 2</math>  <math>4n = 8</math>  <math>n = 8/4 = 2</math>  hence, naphthalene is aromatic.</p>
3.	 <p><b>Anthracene</b></p>	<p>It has a planar ring structure with 14 delocalised <math>\pi</math> electrons.  Applying Huckel's rule <math>4n+2=14\pi</math> electrons.  <math>n=3</math> Hence it is an aromatic compound</p>
4.	 <p><b>Cyclo penta diene</b></p>	<p>(i) It has planar structure  (ii) It has four <math>\pi</math> electron but the <math>\pi</math> electrons are not delocalised and hence it is not an aromatic compound</p>
5.	 <p><b>Cyclooctatetraene</b></p>	<p>molecule is non planar  and hence it is not an aromatic compound</p>
6	 <p><b>Cyclopropenylcation</b></p>	<p>(i) cyclopropenyl cation has planar structure  (ii) It has 2 delocalised <math>\pi</math> electron.  (ii) <math>4n + 2 = 2</math>  <math>4n = 0</math>  <math>n = 0</math> (an interger)  and hence it is aromatic compound.</p>

### 13.5.3. Structure of benzene:

#### 1. Molecular formula:

Elemental Analysis and molecular weight determination have proved that the molecular formula of benzene is  $C_6H_6$ . This indicates that benzene is a highly unsaturated compound.


#### 2. Straight chain structure not possible:

Benzene could be constructed as a straight chain or ring compound but it not feasible since it does not show the properties of alkenes or alkynes. for example, it did not decolourise bromine in carbon tetrachloride or acidified  $KMnO_4$ . It did not react with water in the presence of acid.

#### 3. Evidence of cyclic structure:

##### I) substitution of benzene:

Benzene reacts with bromine in the presence of  $AlCl_3$  to form monobromobenzene.



Formation of only one monobromo compound indicates that all the six hydrogen atoms in benzene were identical. This is possible only if it has a cyclic structure of six carbons each containing one hydrogen.

##### Ii) addition of hydrogen:

Benzene can add on to three moles of hydrogen in the presence of nickel catalyst to give cyclohexane.




This confirms cyclic structure of benzene and the presence of three carbon-carbon double bond.

#### 4. Kekule's structure of benzene:


In 1865, August Kekule suggested that benzene consists of a cyclic planar structure of six carbon with alternate single and double bonds.

There were two objections:

- (i) Benzene forms only one orthodisubstituted products whereas the Kekule's structure predicts two o-di substituted products as shown below.




- (ii) Kekule's structure failed to explain why benzene with three double bonds did not give addition reactions like other alkenes. To overcome this objection, Kekule suggested that benzene was mixture of two forms (1 and 2) which are in rapid equilibrium.



#### 5. Resonance description of benzene:

The phenomenon in which two or more structures can be written for a substance which has identical position

of atoms is called resonance. The actual structure of the molecule is said to be resonance hybrid of various possible alternative structures. In benzene, Kekule's structures I & II represented the resonance structure, and structure III is the resonance hybrid of structure I & II



The structures 1 and 2 exist only in theory. The actual structure of benzene is the hybrid of two hypothetical resonance structures.

## 6. Spectroscopic measurements:

Spectroscopic measurements show that benzene is planar and all of its carbon-carbon bonds are of equal length  $1.40\text{\AA}$ . This value lies between carbon-carbon single bond length  $1.54\text{\AA}$  and carbon-carbon double bond length  $1.34\text{\AA}$ .

## 7. Molecular orbital structure:

The structure of benzene is best described in terms of the molecular orbital theory. All the six carbon atoms of benzene are  $sp^2$  hybridized. Six  $sp^2$  hybrid orbitals of carbon overlap with six  $1s$  orbitals of hydrogen atoms to form six C - H sigma bonds. Overlap between the remaining  $sp^2$  hybrid orbitals of carbon forms six C-C sigma bonds.




Figure 13.6. Formation of Sigma bond in benzene

All the  $\sigma$  bonds in benzene lie in one plane with bond angle  $120^\circ$ . Each carbon atom in benzene possesses an unhybridized p-orbital containing one electron. The lateral overlap of their p-orbitals produces 3  $\pi$ -bond. The six electrons of the p-orbitals cover all the six carbon atoms and are said to be delocalised. Due to delocalization, a strong  $\pi$ -bond is formed which makes the molecule stable. Hence unlike alkenes and alkynes, benzene undergoes substitution reactions rather than addition reactions under normal conditions.





Figure 13.7. All carbon atoms have The delocalized  $\pi$  Mo is formed by p orbitals the overlap of six p-orbitals

## 8. Representation of benzene:

Hence, there are three ways in which benzene can be represented.



### Benzene and its homologous series

Benzene and its homologous series are colorless liquids with pleasant odour. They are lighter than water and insoluble in it. Their vapours are highly flammable, and volatile and toxic in nature.

#### 13.5.4. Sources of aromatic compound:

- Benzene and other aromatic compound are obtained from coal tar and petroleum
- It can also be prepared in laboratory using some simple aliphatic compounds

#### 1. Preparation of benzene

##### (i) industrial preparation of benzene from coal tar :

Coal tar is a viscous liquid obtained by the pyrolysis of coal. During fractional distillation, coal tar is heated and distills away its volatile compounds namely benzene, toluene, xylene in the temperature range of 350 to 443 K. These vapours are collected at the upper part of the fractionating column (Table 13.5.)

TABLE.13.5COMPENENTS OF DISTILLATION OF COAL TAR


NAME OF THE FRACTION	TEMPERATURE RANGE	NAME OF THE COMPENENTS
1. Crude light oil	350 - 443 K	Benzene, Toluene, Xylenes
2. Middle oil	443 - 503 K	Phenol, Naphthalene
3. Heavy oil	503 - 543 K	Naphthalene, Cresol
4. Green oil	543 - 633 K	Anthracene
5. Pitch	Alone 633 K	Residue

##### (ii) from acetylene.

Acetylene on passing through a red -hot tube trimerises to give benzene. We have




already studied this concept in polymerization of alkynes.



### (iii) Laboratory Methods Of Preparing Benzene And Toluene


#### (a) Decarboxylation Of Aromatic Acid.

When sodium benzoate is heated with sodalime, benzene vapours distil over.




#### (b) Preparation Of Benzene From Phenol

When phenol vapours are passed over zinc dust, then it is reduced to benzene.




#### (c) Wurtz – Fittig Reaction:

When a solution of bromo benzene and iodo methane in dry ether is treated with metallic sodium, toluene is formed.



#### (d) Friedel Craft's Reaction:

When benzene is treated with methyl chloride in the presence of anhydrous aluminium chloride, toluene is formed.



### Evaluate Yourself

22) How benzene can be prepared by Grignard Reagent

### 13.4. Physical Properties:

- Benzene is a colourless liquid, insoluble in water and solution alcohol, ether and chloroform.

- It burns with luminous sooty flame in contrast to alkanes and alkenes which usually burn with bluish flame.
- Their vapours are highly toxic which on inhalation produce loss of consciousness.


### 13.5. Chemical Properties:

1. Benzene contains delocalized  $\pi$ -electrons which make the ring to act as an electron rich centre. So electrophilic substitution reaction occurs in benzene.
2. Benzene ring is stabilized by delocalized  $\pi$  electrons. Though it is highly stable, it undergoes addition and oxidation reaction under specific conditions.

#### 1. Electrophilic Substitution Reaction

##### (a) Nitration:


When benzene is heated at 330K with a nitrating mixture (Con.  $\text{HNO}_3$  + Con.  $\text{H}_2\text{SO}_4$ ), nitro benzene is formed by replacing a hydrogen atom by a nitronium ion  $\text{NO}_2^+$  (electrophile)



Concentrated  $\text{H}_2\text{SO}_4$  is added to produce nitronium ion  $\text{NO}_2^+$


##### (b) Halogenation:

Benzene reacts with halogens ( $\text{X}_2 = \text{Cl}_2, \text{Br}_2$ ) in the presence of Lewis acid such as  $\text{FeCl}_3$ ,  $\text{FeBr}_3$  or  $\text{AlCl}_3$  and give corresponding halo benzene. In the absence of catalyst, Fluorine reacts vigorously with benzene even in the absence of catalyst. However iodine is very inactive even in the presence of catalyst




##### (c) Sulphonation:

Benzene reacts with fuming sulphuric acid (Con  $\text{H}_2\text{SO}_4$  +  $\text{SO}_3$ ) and gives benzene sulphonic acid. The electrophile  $\text{SO}_3$  is a molecule. Although it does not have positive charge, it is a strong electrophile. This is because the octet of electron around the sulphur atom is not reached. The reaction is reversible and desulphonation occurs readily in aqueous medium.




#### (d) Friedel Craft's Alkylation: (Methylation)

When benzene is treated with an alkyl halide in the presence of only  $\text{AlCl}_3$ , alkyl benzene is formed.




#### (e) Friedel Craft's Acylation : Acetylation

When benzene is treated with acetyl chloride in the presence of  $\text{AlCl}_3$ , acyl benzene is formed.



#### (f) Electrophilic Substitution Reactions: Mechanism

Benzene undergoes electrophilic substitution reaction because it is an electron-rich system due to delocalised  $\pi$  electron. So it is easily attacked by electrophiles and gives substituted products.



## Mechanism:


### Step: 1

Formulation of the electrophile




### Step:2

The electrophile attacks the aromatic ring to form a carbocation intermediate which is stabilized by resonance.


















### Step: 3

Loss of proton gives the substitution product.



## Evaluate Yourself


21) Why benzene undergoes electrophilic substitution reaction where as alkenes undergoes addition reaction?

Types of reaction	NITRATION	HALOGENATION	SULPHONATION	FRIEDEL CRAFT'S ALKYLATION	FRIEDEL CRAFT'S ACYLATION
Reagents	Con $\text{HNO}_3$ + Con $\text{H}_2\text{SO}_4$	$\text{X}_2 / \text{AlX}_3$	Fuming $\text{H}_2\text{SO}_4$ $\text{CH}_2\text{SO}_4 + \text{SO}_3$	$\text{CH}_3\text{Cl}$ + anhydrous $\text{AlCl}_3$	$\text{CH}_3\text{COCl}$ + anhydrous $\text{AlCl}_3$
Electro- phile	$-\text{NO}_2^+$	$-\text{X}^+ (\text{X} = \text{Cl}, \text{Br})$	$-\text{SO}_3$ (Neutral electrophile)	$-\text{CH}_2^+$	$-\text{COCH}_3^+$
Over all reaction					
Mechanism Step 1	$\text{NO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{NO}_2^+ + \text{HSO}_4^- + \text{H}_2\text{O}$	$\text{AlCl}_3 + \text{Cl}_2 \rightarrow \text{Cl}^+ + \text{AlCl}_4^-$	$2\text{H}_2\text{SO}_4 \rightarrow \text{H}_3\text{O}^+ + \text{SO}_3 + \text{HSO}_4^-$	$\text{AlCl}_3 + \text{CH}_3\text{Cl} \rightarrow \text{CH}_3^+ + \text{AlCl}_4^-$	$\text{AlCl}_3 + \text{CH}_3\text{COCl} \rightarrow \text{CH}_3\text{CO}^+ + \text{AlCl}_4^-$
Step 2					
STEP 3					

## (ii) Addition Reaction:


### a. Hydrogenation of benzene:

Benzene reacts with hydrogen in the presence of Platinum or Palladium to yield Cyclohexane. This is known as hydrogenation.



### b. Chlorination of Benzene:


Benzene reacts with three molecules of  $\text{Cl}_2$  in the presence of sun light or UV light to yield Benzene Hexa Chloride (BHC)  $\text{C}_6\text{H}_6\text{Cl}_6$ . This is known as gammaxane or Lindane which is a powerful insecticide.



## (Iii) Oxidation:


### a. Vapour – phase oxidation:-

Although benzene is very stable to strong oxidizing agents, it quickly undergoes vapour phase oxidation by passing its vapour mixed with oxygen over  $\text{V}_2\text{O}_5$  at 773K. The ring breaks to give maleic anhydride.



### b. Birch reduction:

Benzene can be reduced to 1,4-cyclohexadiene by treatment with Na or Li in a mixture of liquid ammonia and alcohol. It is the convenient method to prepare cyclic dienes.



## Evaluate Yourself

24) Convert Ethyne to Benzene and name the process.


### 13.5.1 directive influence of a functional group in monosubstituted benzene:

When mono substituted benzene undergoes an electrophilic substitution reaction, the rate of the reaction and the site of attack of the incoming electrophile depends on the functional group already attached to it. Some groups increase the reactivity of benzene ring and are known as activating groups. While others which decrease the reactivity are known as deactivating groups. We further divide

these groups into two categories depending on the way they influence the orientation of attack by the incoming groups. Those which increase electron density at 'ortho' and 'para' position are known as ortho-para directors while those which increase electron density at 'meta' position is known as meta-directors. Some examples of directive influence of functional groups in mono-substituted benzene are explained below.

### Ortho and para directing groups

All the activating groups are 'ortho-para' directors. Example  $-\text{OH}$ ,  $-\text{NH}_2$ ,  $-\text{NHR}$ ,  $-\text{NHCOCH}_3$ ,  $-\text{OCH}_3$ ,  $-\text{CH}_3$ ,  $-\text{C}_2\text{H}_5$  etc. Let us consider the directive influences of phenolic ( $-\text{OH}$ ) group. Phenol is the resonance hybrid of following structures.




In these resonance structures, the  $(-)$  charge residue is present on ortho and para position of ring structure. It is quite evident that the lone pair of electron on the atom which is attached to the ring involves in resonance and makes the ring more electron rich than benzene. The electron

density at ortho and para positions increases as compared to the meta position. Therefore phenolic group activates the benzene ring for electrophilic attack at 'ortho' and 'para' positions and hence  $-\text{OH}$  group is an ortho-para director and activator.

In aryl halides, the strong  $-I$  effect of the halogens (electron withdrawing tendency) decreases the electron density of benzene ring, thereby deactivating for electrophilic attack. However the presence of lone pair on halogens involved in the resonance with pi electrons of benzene ring, increases electron density at ortho and para position. Hence the halogen group is an ortho-para director and deactivator.

### META DIRECTING GROUPS

Generally all deactivating groups are meta-directors. For example  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{CHO}$ ,  $-\text{COR}$ ,  $-\text{COOH}$ ,  $-\text{COOR}$ ,  $-\text{SO}_3\text{H}$  etc. Let us consider the directive influence of aldehyde ( $-\text{CHO}$ ) group. Benz aldehyde is the resonance hybrid of following structures.



In these resonance structures, the (+) charge residues is present on the ring structure. It is quite evident that resonance delocalizes the positive charge on the atoms of the ring, making the ring less electron rich than benzene. Here overall density of benzene ring decreases due to –I effect of –CHO group there by deactivating the benzene for electrophilic attack. However resonating structure shows that electron density is more in meta position. Compared to o & p-position. Hence –CHO group is a meta-director and deactivator.


### Evaluate Yourself

25) Toluene undergoes nitration easily than benzene. Why?


### 13.5.2. Carcinogenity and toxicity

Benzene and polycyclic aromatic hydrocarbons (PAH) are ubiquitous environmental pollutants generated during incomplete combustion of coal oil, petrol and wood. Some PAH originate from open burning, natural seepage of petroleum and coal deposits and volcanic activities. They are toxic, mutagenic and carcinogenic. It has hematological, immunological and neurological effect on humans. They are radiomimetic and prolonged exposure leads to genetic damage. Some of the examples of PAH are.

“L” shaped polynuclear hydrocarbons are much more toxic & carcinogenic




1,3- Benzpyrene



1,3- Benzanthrachine

Found in cigarette smoke Found in tobacco and cigarette and charcoal boiled food





1,2,5,6 dibenzanthracene



Found in gasoline exhaust and barbaccued food




# FLOWCHART AND REACTION SUMMARY OF HYDROCARBON




## Methods of Preparation





## Chemical Properties





3. An alkyl bromide (A) reacts with sodium in ether to form 4, 5- diethyloctane, the compound (A) is





- 
CC1CCCC1.BrBr>>CC1(Br)CCCC1

c) 


d) 

- d) none of of these

7. The compounds formed at anode in the electrolysis of an aqueous solution of potassium acetate are
- $\text{CH}_4$  and  $\text{H}_2$
  - $\text{CH}_4$  and  $\text{CO}_2$
  - $\text{C}_2\text{H}_6$  and  $\text{CO}_2$
  - $\text{C}_2\text{H}_4$  and  $\text{Cl}_2$
8. The general formula for cyclo alkanes
- $\text{C}_n\text{H}_n$
  - $\text{C}_n\text{H}_{2n}$
  - $\text{C}_n\text{H}_{2n-2}$
  - $\text{C}_n\text{H}_{2n+2}$
9. The compound that will react most readily with gaseous bromine has the formula (NEET)
- $\text{C}_3\text{H}_6$
  - $\text{C}_2\text{H}_2$
  - $\text{C}_4\text{H}_{10}$
  - $\text{C}_2\text{H}_4$
10. Which of the following compounds shall not produce propene by reaction with  $\text{HBr}$  followed by elimination (or) only direct elimination reaction (NEET)
- $\triangle$
  - $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{OH}$
  - $\text{H}_2\text{C} = \text{C} = \text{O}$
  - $\text{CH}_3 - \text{CH}_2 - \text{CH}_2\text{Br}$
11. Which among the following alkenes on reductive ozonolysis produces only propanone ?
- 2 - Methyl propene
  - 2 - Methyl but - 2 - ene
  - 2, 3 - Dimethyl but - 1 - ene
  - 2, 3 - Dimethyl but - 2 - ene
12. The major product formed when 2 - bromo - 2 - methyl butane is refluxed with ethanolic  $\text{KOH}$  is
- 2 - methylbut - 2 - ene
  - 2 - methyl butan - 1 - ol
  - 2 - methyl but - 1 - ene
  - 2 - methyl butan - 2 - ol
13. Major product of the below mentioned reaction is,
- $$(\text{CH}_3)_2\text{C} = \text{CH}_2 \xrightarrow{\text{ICl}}$$
- 2-chloro - 1- iodo - 2 - methyl propane
  - 1-chloro-2-iodo-2-methylpropane
  - 1,2 - dichloro - 2 - methyl propane
  - 1, 2 - diiodo - 2 - methyl propane
14. The IUPAC name of the following compound is
- 
- trans-2-chloro-3-iodo - 2 - pentane
  - cis-3 - iodo - 4 - chloro - 3 - pentane
  - trans-3-iodo-4-chloro - 3 - pentene
  - cis-2 - chloro - 3 - iodo - 2 - pentene
15. Cis - 2 - butene and trans - 2 - butene are
- conformational isomers
  - structural isomers
  - configurational isomers
  - optical isomers
16. Identify the compound (A) in the following reaction
- 



17.  $\text{CH}_2 - \text{CH}_2 \xrightarrow{\text{(A)}} \text{CH} \equiv \text{CH}$ , where A is,



a) Zn

b) Conc  $\text{H}_2\text{SO}_4$

c) alc. KOH

d) dil  $\text{H}_2\text{SO}_4$

18. Consider the nitration of benzene using mixed con  $\text{H}_2\text{SO}_4$  and  $\text{HNO}_3$  if a large quantity of  $\text{KHSO}_4$  is added to the mixture, the rate of nitration will be




a) unchanged

b) doubled

c) faster




d) slower

19. In which of the following molecules, all atoms are co-planar






d) both (a) and (b)


20. Propyne on passing through red hot iron tube gives







d) none of these



c) both (a) and (b)




22. Which one of the following is non aromatic ?



23. Which of the following compounds will not undergo Friedal - crafts reaction easily ? (NEET)

a) Nitro benzene

- b) Toluene  
c) Cumene  
d) Xylene
24. Some meta-directing substituents in aromatic substitution are given. Which one is most deactivating ?  
a)  $-\text{COOH}$                       b)  $-\text{NO}_2$   
c)  $-\text{C}\equiv\text{N}$                       d)  $-\text{SO}_3\text{H}$
25. Which of the following can be used as the halide component for Friedel-Crafts reaction ?  
a) Chloro benzene  
b) Bromo benzene  
c) chloro ethene  
d) isopropyl chloride
26. An alkane is obtained by decarboxylation of sodium propionate. Same alkane can be prepared by  
a) Catalytic hydrogenation of propene  
b) action of sodium metal on iodomethane  
c) reduction of 1-chloro propane  
d) reduction of bromomethane
27. Which of the following is aliphatic saturated hydrocarbon  
a)  $\text{C}_8\text{H}_{18}$                       b)  $\text{C}_9\text{H}_{18}$   
c)  $\text{C}_8\text{H}_{14}$                       d) All of these
28. Identify the compound 'Z' in the following reaction
- $$\text{C}_2\text{H}_6\text{O} \xrightarrow[623\text{ K}]{\text{Al}_2\text{O}_3} \text{X} \xrightarrow{\text{O}_3} \text{Y} \xrightarrow{\text{Zn}/\text{H}_2\text{O}} (\text{Z})$$
- a) Formaldehyde  
b) Acetaldehyde  
c) Formic acid                      d) none of these
29. Peroxide effect (Kharasch effect) can be studied in case of  
a) Oct-4-ene                      b) hex-3-ene  
c) pent-1-ene                      d) but-2-ene
30. 2-butyne on chlorination gives  
a) 1-chloro butane  
b) 1,2-dichloro butane  
c) 1,1,2,2-tetrachlorobutane  
d) 2,2,3,3-tetrachloro butane
31. Give IUPAC names for the following compounds
- 1)  $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}=\text{CH}-\text{C}\equiv\text{C}-\text{CH}_3$
- 2) 
$$\begin{array}{c} \text{C}_2\text{H}_5 \quad \text{CH}_3 \\ | \quad | \\ \text{CH}_3-\text{C}-\text{C}-\text{C}\equiv\text{C}-\text{CH}_3 \\ | \quad | \\ \text{CH}_3 \quad \text{H} \end{array}$$
- 3)  $(\text{CH}_3)_3\text{C}-\text{C}\equiv\text{C}-\text{CH}(\text{CH}_3)_2$
- 4) ethyl isopropyl acetylene
- 5)  $\text{CH}\equiv\text{C}-\text{C}\equiv\text{C}-\text{C}\equiv\text{CH}$
32. Identify the compound A, B, C and D in the following series of reactions



Write short notes on ortho, para directors in aromatic electrophilic substitution reactions.

33. How is propyne prepared from an alkyne dihalide ?

34. An alkylhalide with molecular formula  $C_6H_{13}Br$  on dehydro halogenation gave two isomeric alkenes X and Y with molecular formula  $C_6H_{12}$ . On reductive ozonolysis, X and Y gave four compounds  $CH_3COCH_3$ ,  $CH_3CHO$ ,  $CH_3CH_2CHO$  and  $(CH_3)_2CHCHO$ . Find the alkylhalide.

35. Describe the mechanism of Nitration of benzene.

36. How does Huckel rule help to decide the aromatic character of a compound.

37. Suggest the route for the preparation of the following from benzene.

1) 3 - chloro nitrobenzene

2) 4 - chlorotoluene

3) Bromo benzene

4) m - dinitro benzene

38. Suggest a simple chemical test to distinguish propane and propene.

39. What happens when isobutylene is treated with acidified potassium permanganate ?

40. How will you convert ethyl chloride in to

i) ethane            ii) n - butane

41. Describe the conformers of n - butane.

42. Write the chemical equations for combustion of propane.

43. Explain Markov nikoff's rule with suitable example.

44. What happens when ethylene is passed through cold dilute alkaline potassium permanganate.

45. Write the structures of following alkanes.

1) 2, 3 - Dimethyl - 6 - (2 - methyl propyl) decane

2) 5 - (2 - Ethyl butyl) - 3, 3 - dimethyldecane

3) 5 - (1, 2 - Dimethyl propyl) - 2 - methylnonane

46. How will you prepare propane from a sodium salt of fatty acid ?

47.  $CH_3 - CH(CH_3) - CH(OH) - CH_3$



$H^+/heat$

(A) major product  $\xrightarrow{HBr}$  (B) major product


Identify A and B

48. Complete the following :

i) 2 - butyne  $\xrightarrow{\text{Lindlar Catalyst}}$

ii)  $CH_2 = CH_2 \xrightarrow{I_2}$

iii)  $CH_2 - CH_2 \xrightarrow{Zn/C_2H_5OH}$



iv)  $CaC_2 \xrightarrow{H_2O}$

49. How will you distinguish 1 - butyne and 2 - butyne?