

Respiration

Chapter Outline

- 6.1 Respiratory function
- 6.2 Respiratory organs in various organisms
- 6.3 Mechanism of breathing
- 6.4 Exchange of gases
- 6.5 Transport of gases
- 6.6 Regulation of respiration
- 6.7 Problems in oxygen transport
- 6.8 Disorders of respiratory system
- 6.9 Effects of smoking



Exercise increases the rate and depth of breathing and supplies extra oxygen to the muscles and removes more CO₂ from the tissues.

Learning Objectives:

- *Learns to describe the gross structure of the human gaseous exchange system*
- *Observes and draws the tissues and organs associated with the respiratory system*
- *Understands the process of gaseous exchange and transport of respiratory gases*
- *Knows the problems associated with oxygen transport*
- *Gains knowledge on the ill-effects of smoking.*



We inhale and exhale air. Why is breathing so important for life? What happens when we breathe? Why energy is required for the body to perform various life processes? Where does the energy come from? We eat food for energy. Though the above raised questions look disconnected, we should know that the process of breathing is connected to the process of release of energy from food. Oxygen is utilized by the organisms to breakdown the biomolecules like glucose and to derive energy. During this breakdown carbondioxide, which is a harmful gas is also released. It is very obvious that oxygen has to be provided to cells continuously and the CO₂ to be released immediately by the cells. So the need of a respiratory system is essential for life.

We have discussed in the previous chapter how food provides energy for growth and repair of tissues. As mentioned earlier along with food, oxygen is necessary for breakdown of glucose to energy. In this chapter we shall discuss the respiratory organs of human, the mechanism of breathing, exchange and transport of gases and a few respiratory disorders.

The term respiration refers to the exchange of oxygen and carbon dioxide between environment and cells of our body where organic nutrients are broken down enzymatically to release energy.

6.1 Respiratory functions

The five primary functions of the respiratory system are –

- i. To exchange O_2 and CO_2 between the atmosphere and the blood.
- ii. To maintain homeostatic regulation of body pH.
- iii. To protect us from inhaled pathogens and pollutants.
- iv. To maintain the vocal cords for normal communication (vocalization).
- v. To remove the heat produced during cellular respiration through breathing.

6.2 Respiratory organs in various organisms.

Different animals have different organs for exchange of gases, depending upon their habitats and levels of organization. The amount of dissolved oxygen is very low in water compared to the amount of oxygen in the air. So the rate of breathing in aquatic organisms is much faster than land animals.

In animals like sponges, coelenterates and flatworms exchange of gases takes place through the body surface by simple diffusion. Earthworms use their moist skin, whereas insects have tracheal tubes. Gills are used as respiratory organs in most of the aquatic Arthropods and Molluscs. Among vertebrates, fishes use gills whereas amphibians, reptiles, birds and mammals have well vascularised lungs. Frogs spend most of their time in water and also use their moist skin for respiration along with lungs.

6.2.1 Human Respiratory System

The respiratory system includes the external nostrils, nasal cavity, the pharynx, the larynx, the trachea, the bronchi and bronchioles and the lungs which contain the alveoli (Figure 6.1). The parts starting from the external nostrils up to the terminal bronchioles constitute the conducting zone, whereas the alveoli and the ducts are called the respiratory zone. The parts of the conducting zone, humidifies and warms the incoming air.

In human beings, air enters the upper respiratory tract through the external nostrils. The air passing through the nostrils is filtered by fine hairs and mucus lining the passage. The external nostrils lead to the nasal chamber which opens into the nasopharynx which opens through the glottis of the larynx region into the trachea. The ciliated epithelial cells lining the trachea, bronchi and bronchioles secrete mucus. Mucus membrane lining the airway contains goblet cells which secrete mucus, a slimy material rich in glycoprotein. Microorganisms and dust particles attach in the mucus films and




Figure 6.1 The Human respiratory system

are carried upwards to pass down the gullet during normal swallowing. During swallowing a thin elastic flap called epiglottis prevents the food from entering into the larynx and avoids choking of food.

The trachea is semiflexible tube supported by multiple cartilaginous rings which extends up to the midthoracic cavity and at the level of the 5th thoracic vertebra where it divides into right and left primary bronchi, one bronchus to each lung. Within the lungs the bronchi divides repeatedly into secondary and tertiary bronchi and further divides into terminal bronchioles and respiratory bronchioles.

It is advised not to talk or laugh louder while eating. Can you give the reason?

Bronchi have 'C' shaped curved cartilage plates to ensure that the air passage does not collapse or burst as the air pressure changes during breathing. The bronchioles are without cartilaginous rings and have rigidity that prevent them from collapsing but are surrounded by smooth muscle which contracts or relaxes to adjust the diameter of these airways.

The fine respiratory bronchioles terminate into highly vascularised thin walled pouch like air sacs called alveoli meant for gaseous exchange (Figure 6.2, 6.3). The diffusion membrane of alveolus is made up of three layers – the thin squamous epithelial cells of the alveoli, the endothelium of the alveolar capillaries and the basement substance found in between them. The thin squamous epithelial cells of the alveoli are composed of Type I and




Figure 6.2
The Human Respiratory tract




Figure 6.3 Structure of alveoli

Type II cells. Type I cells are very thin so that gases can diffuse rapidly through them. Type II cells are thicker, synthesize and secrete a substance called **Surfactant**.

The lungs are light spongy tissues enclosed in the thoracic cavity surrounded by an airtight space. The thoracic cavity is bound dorsally by the vertebral column and ventrally by the sternum, laterally by the ribs and on the lower side by the dome shaped diaphragm.

The lungs are covered by double walled pleural membrane containing a several layers of elastic connective tissues and capillaries, which encloses the pleural fluid. Pleural fluid reduces friction when the lungs expand and contract.

Characteristic features of respiratory surface:

- surface area must be very large and richly supplied with blood vessels
- should be extremely thin and kept moist
- should be in direct contact with the environment
- should be permeable to respiratory gases

The steps involved in respiration are

- i. The exchange of air between the atmosphere and the lungs.
- ii. The exchange of O_2 and CO_2 between the lungs and the blood.
- iii. Transport of O_2 and CO_2 by the blood.
- iv. Exchange of gases between the blood and the cells.
- v. Uptake of O_2 by the cells for various activities and the release of CO_2 .



SURFACTANTS are the thin non-cellular films made of protein and phospholipids covering the alveolar membrane. The surfactant lowers the surface tension in the alveoli and prevents the lungs from collapsing. It also prevents pulmonary oedema. Premature Babies have low levels of surfactant in the alveoli may develop the new born respiratory distress syndrome (NRDS) because the synthesis of surfactants begins only after the 25th week of gestation.

6.3. Mechanism of breathing

The movement of air between the atmosphere and the lungs is known as ventilation or breathing. Inspiration and expiration are the two phases of breathing. Inspiration is the movement of atmospheric air into the lungs and expiration is the movement of alveolar air that diffuse out of the lungs. (Figure 6.4)

Lungs do not contain muscle fibres but expands and contracts by the movement of the ribs and diaphragm. The diaphragm is a sheet of tissue which separates the thorax from the abdomen. In a relaxed state, the diaphragm is domed shaped.

Observe a live fish and find out how many times it beats the operculum per minute. Now check your rate of breathing for a minute. The rate of breathing will be more in fish than you – Give reasons.

Ribs are moved by the intercostal muscles. External and internal intercostal muscles found between the ribs and the diaphragm helps in creating pressure gradients. Inspiration occurs if the pressure inside

the lungs (intrapulmonary pressure) is less than the atmospheric pressure likewise expiration takes place when the pressure within the lungs is higher than the atmospheric pressure.




Figure 6.4 Mechanism of breathing

Why do some people snore? – Breathing with a hoarse sound during sleep is caused by the vibration of the soft palate. Snoring is caused by a partially closed upper air way (nose and throat) which becomes too narrow for enough air to travel through the lungs. This makes the surrounding tissues to vibrate and produces the snoring sound.

Inspiration is initiated by the contraction of the diaphragm muscles and external intercostal muscles, which pulls the ribs and sternum upwards and outwards and increases the volume of the thoracic chamber in the dorso-ventral axis, forcing the lungs to expand the pulmonary volume. The increase in pulmonary volume and

You are at high level in a mountain above the sea level. Suddenly you get palpitation and nausea. What condition are you suffering from? What are the other symptoms for this disease and how can it be reduced?

decrease in the intrapulmonary pressure forces the fresh air from outside to enter the air passages into the lungs to equalize the pressure. This process is called **inspiration**.

Relaxation of the diaphragm allows the diaphragm and sternum to return to its dome shape and the internal intercostal muscles contract, pulling the ribs downward reducing the thoracic volume and pulmonary volume. This results in an increase in the intrapulmonary pressure slightly above the atmospheric pressure causing the expulsion of air from the lungs. This process is called **expiration**.

On an average, a healthy human breathes 12–16 times/minute. An instrument called Spirometer is used to measure the volume of air involved in breathing movements for

clinical assessment of a person's pulmonary function.

6.3.1 Respiratory volumes and capacities

The volume of air present in various phases of respiration is denoted as

Respiratory volumes: (Figure 6.5)

- **Tidal Volume (TV)** Tidal volume is the amount of air inspired or expired with each normal breath. It is approximately 500 mL., i.e. a normal human adult can inspire or expire approximately 6000 to 8000mL of air per minute. During vigorous exercise, the tidal volume is about 4–10 times higher.




Figure 6.5 Lung volumes and capacity



Healthy lungs contain large amounts of elastic connective tissue around the alveoli, containing elastin, which makes the lung tissue elastic. People with emphysema and bronchitis have difficulty in exhaling because the enzyme elastase destroys the elastin around the alveoli and reduces the elasticity of the lungs.

- **Inspiratory Reserve volume (IRV)** Additional volume of air a person can inspire by forceful inspiration is called Inspiratory Reserve Volume. The normal value is 2500–3000 mL.
- **Expiratory Reserve volume (ERV)** Additional volume of air a person can forcefully exhale by forceful expiration is called Expiratory Reserve Volume. The normal value is 1000–1100 mL.
- **Residual Volume (RV)** The volume of air remaining in the lungs after a forceful expiration. It is approximately 1100–1200 mL.

Respiratory capacities:

- **Vital capacity (VC)** the maximum volume of air that can be moved out during a single breath following a maximal inspiration. A person first inspires maximally then expires maximally. $VC=ERV+TV+IRV$
- **Inspiratory capacity (IC)** The total volume of air a person can inhale after normal expiration. It includes tidal volume and inspiratory reserve volume. $IC=TV+IRV$

- **Expiratory capacity (EC)** The total volume of air a person can exhale after normal inspiration. It includes tidal volume and expiratory reserve volume. $EC=TV+ERV$
- **Total Lung Capacity (TLC)** The total volume of air which the lungs can accommodate after forced inspiration is called Total Lung Capacity. This includes the vital capacity and the residual volume. It is approximately 6000mL. $TLC=VC+RV$
- **Minute Respiratory Volume** The amount of air that moves into the respiratory passage per minute is called minute respiratory volume.
Normal $TV = 500\text{mL}$; Normal respiratory rate = 12 times/minute
Therefore, minute respiratory volume = 6 Litres/minute (for a normal healthy man).

Dead space

Some of the inspired air never reaches the gas exchange areas but fills the respiratory passages where exchange of gases does not occur. This air is called dead space.

Dead space is not involved in gaseous exchange. It amounts to approximately 150mL.

6.4 Exchange of gases

The primary site for the exchange of gases is the alveoli. The uptake of O_2 and the release of CO_2 occur between the blood and tissues by simple diffusion driven by partial pressure gradient of O_2 and CO_2 . Partial pressure is the pressure contributed by an individual gas in a mixture of gases. It is represented as pO_2 for oxygen and pCO_2 for carbon-

Breathing through nose is healthier than through mouth- Why?




Figure 6.6 Exchange of gases at the alveolus and the tissue with blood and transport of oxygen and carbondioxide

dioxide. Due to pressure gradients, O_2 from the alveoli enters into the blood and reaches the tissues. CO_2 enters into the blood from the tissues and reaches alveoli for elimination. As the solubility of CO_2 is 20–25 times higher than that of O_2 , the partial pressure of CO_2 is much higher than that of O_2 (Tab.6.1 and Figure 6.6).

Respiratory pigments

Haemoglobin

Haemoglobin belongs to the class of conjugated protein. The iron containing pigment portion haem constitutes only 4% and the rest colourless protein of the histone class globin. Haemoglobin has a molecular weight of 68,000 and contains four atoms of iron, each of which can combine with a molecule of oxygen.

Methaemoglobin

If the iron component of the haem moieties is in the ferric state, than the normal ferrous state, it is called methaemoglobin. Methaemoglobin

Respiratory gases	Partial pressure mm Hg				
	Atmospheric air	Alveoli	Deoxygenated Blood	Oxygenated blood	Tissues
O_2	159	104	40	95	40
CO_2	0.3	40	45	40	45

Table 6.1 Partial pressure of Oxygen and Carbon dioxide (in mmHg) in comparison to those gases in the atmosphere

does not bind O₂. Normally RBC contains less than 1% methaemoglobin.

6.5 Transport of gases

6.5.1 Transport of oxygen

Molecular oxygen is carried in blood in two ways: bound to haemoglobin within the red blood cells and dissolved in plasma. Oxygen is poorly soluble in water, so only 3% of the oxygen is transported in the dissolved form. 97% of oxygen binds with haemoglobin in a reversible manner to form oxyhaemoglobin (HbO₂). The rate at which haemoglobin binds with O₂ is regulated by the partial pressure of O₂. Each haemoglobin carries maximum of four molecules of oxygen. In the alveoli high pO₂, low pCO₂, low temperature and less H⁺ concentration, favours the formation of oxyhaemoglobin, whereas in the tissues low pO₂, high pCO₂, high H⁺ and high




Figure 6.7 Oxygen dissociation curve

temperature favours the dissociation of oxygen from oxyhaemoglobin.

A sigmoid curve (S-shaped) is obtained when percentage saturation of haemoglobin with oxygen is plotted against pO₂. This curve is called oxygen haemoglobin dissociation curve (Figure 6.7). This S-shaped curve has a steep slope for pO₂ values between 10 and 50 mmHg and then flattens between 70 and 100 mm Hg.

Under normal physiological conditions, every 100 mL of oxygenated blood can deliver about 5 mL of O₂ to the tissues.

6.5.2 Transport of Carbon-dioxide

Blood transports CO₂ from the tissue cells to the lungs in three ways


- i. **Dissolved in plasma** About 7 – 10% of CO₂ is transported in a dissolved form in the plasma.
 - ii. **Bound to haemoglobin** About 20 – 25% of dissolved CO₂ is bound and carried in the RBCs as carbamino haemoglobin (Hb CO₂)

$$\text{CO}_2 + \text{Hb} \rightleftharpoons \text{Hb CO}_2$$
 - iii. **As bicarbonate ions in plasma** about 70% of CO₂ is transported as bicarbonate ions
- This is influenced by pCO₂ and the degree of haemoglobin oxygenation. RBCs contain a high concentration of the enzyme, carbonic anhydrase, whereas small amounts of carbonic anhydrase is present in the plasma.

At the tissues the pCO₂ is high due to catabolism and diffuses into the blood to form HCO₃⁻ and H⁺

ions. When CO_2 diffuses into the RBCs, it combines with water forming carbonic acid (H_2CO_3) catalyzed by carbonic anhydrase. Carbonic acid is unstable and dissociates into hydrogen and bicarbonate ions.

Carbonic anhydrase facilitates the reaction in both directions.



The HCO_3^- moves quickly from the RBCs into the plasma, where it is carried to the lungs. At the alveolar site where pCO_2 is low, the reaction is reversed leading to the formation of CO_2 and water. Thus CO_2 trapped as HCO_3^- at the tissue level it is

Events in inspiration and expiration

Inspiration	Expiration
Respiratory centre initiates the stimuli during inspiration.	Respiratory centre terminates the stimuli during expiration.
↓	↓
Impulses are carried to the inspiratory muscles through nerves.	The diaphragm and inspiratory muscles relax.
↓	↓
Diaphragm and inspiratory muscles contract.	Chest wall contracts and the thoracic volume gets reduced.
↓	↓
The thoracic volume increases as the chest wall expands.	The intra pulmonary pressure is reduced.
↓	↓
The intra pulmonary pressure is reduced.	The alveolar pressure increases than the atmospheric pressure.
↓	↓
The alveolar pressure decreases than the atmospheric pressure	Air is sent out due to the contraction of alveoli.
↓	↓
Air flows into the alveoli until the alveolar pressure equalizes the atmospheric pressure and the alveoli get inflated.	Air flows out of the alveoli until the alveolar pressure equalizes the atmospheric pressure and the alveoli get deflated.

transported to the alveoli and released out as CO_2 . Every 100mL of deoxygenated blood delivers 4mL of CO_2 to the alveoli for elimination.

6.6 Regulation of Respiration

A specialised respiratory centre present in the medulla oblongata of the hind brain called respiratory rhythm centre is responsible for this regulation. Pneumotaxic centre present in pons varoli region of the brain moderates the function of the respiratory rhythm centre to ensure normal breathing. The chemosensitive area found close to the rhythm centre is highly sensitive to CO_2 and H^+ . And H^+ are eliminated out by respiratory process. Receptors associated with the aortic arch and carotid artery send necessary signals to the rhythm centre for remedial action. The role of O_2 is insignificant in the regulation of respiratory rhythm.

Particulate matter PM 2.5 in the air is increasing day by day which causes respiratory illness. Central Pollution Control Board (CPCB) reports that the quality of air is not good due to soot and smoke. So some cities in India are using CNG (Compressed Natural Gas) as fuel.

6.7 Problems in Oxygen transport

When a person travels quickly from sea level to elevations above 8000ft, where the atmospheric pressure and partial pressure of oxygen are lowered, the individual responds with symptoms of acute mountain sickness (AMS)–

headache, shortness of breath, nausea and dizziness due to poor binding of O_2 with haemoglobin. When the person moves on a long-term basis to mountains from sea level is body begins to make respiratory and haematopoietic adjustments. To overcome this situation kidneys accelerate production of the hormone erythropoietin, which stimulates the bone marrow to produce more RBCs.

When a person descends deep into the sea, the pressure in the surrounding water increases which causes the lungs to decrease in volume. This decrease in volume increases the partial pressure of the gases within the lungs. This effect can be beneficial, because it tends to drive additional oxygen into the circulation, but this benefit also has a risk, the increased pressure can also drive nitrogen gas into the circulation. This increase in blood nitrogen content can lead to a condition called **nitrogen narcosis**. When the diver ascends to the surface too quickly a condition called ‘bends’ or decompression sickness occurs and nitrogen comes out of solution while still in the blood forming bubbles. Small bubbles in the blood are not harmful, but large bubbles can lodge in small capillaries, blocking



Allergy is caused by allergens. When we enter a polluted area, immediately we start sneezing and coughing. The allergens in that place affect our respiratory tracts and the responses to the allergens start within minutes. Allergens provoke an inflammatory response. A common manifestation of allergy is Asthma.

blood flow or can press on nerve endings. Decompression sickness is associated with pain in joints and muscles and neurological problems including stroke. The risk of nitrogen narcosis and bends is common in scuba divers.

During carbon-dioxide poisoning, the demand for oxygen increases. As the O₂ level in the blood decreases it leads to suffocation and the skin turns bluish black.

6.8 Disorders of the Respiratory system

Respiratory system is highly affected by environmental, occupational, personal and social factors. These factors may be responsible for a number of respiratory disorders. Some of the disorders are discussed here.

Asthma – It is characterized by narrowing and inflammation of bronchi and bronchioles and difficulty in breathing. Common allergens for asthma are dust, drugs, pollen grains, certain food items like fish, prawn and certain fruits etc.

Emphysema– Emphysema is chronic breathlessness caused by gradual breakdown of the thin walls of the alveoli decreasing the total surface area of a gaseous exchange. i.e., widening of the alveoli is called emphysema. The major cause for this disease is cigarette smoking, which reduces the respiratory surface of the alveolar walls.

Bronchitis– The bronchi when it gets inflated due to pollution smoke and cigarette smoking, causes bronchitis. The symptoms are cough, shortness of breath and sputum in the lungs.

Pneumonia– Inflammation of the lungs due to infection caused by bacteria or virus is called pneumonia. The common symptoms are sputum production, nasal congestion, shortness of breath, sore throat, etc.

Tuberculosis– Tuberculosis is caused by *Mycobacterium tuberculosis*. This infection mainly occurs in the lungs and bones. Collection of fluid between the lungs and the chest wall is the main complication of this disease.

Occupational respiratory disorders– The disorders due to one's occupation of working in industries like grinding or stone breaking, construction sites, cotton industries, etc. Dust produced affects the respiratory tracts.

Long exposure can give rise to inflammation leading to fibrosis. Silicosis and asbestosis are occupational respiratory diseases resulting from inhalation of particle of silica from sand grinding and asbestos into the respiratory tract. Workers, working in such industries must wear protective masks.


6.9 Effects of Smoking

Today due to curiosity, excitement or adventure youngsters start to smoke and later get addicted to smoking. Research says about 80% of the lung cancer is due to cigarette smoking.

Smoking is inhaling the smoke from burning tobacco. There are thousands of known chemicals which includes nicotine, tar, carbon monoxide, ammonia, sulphur-dioxide and even small quantities of arsenic. Carbon monoxide and nicotine damage the cardiovascular system and tar damages

Disorders of Respiratory System

Respiratory system is highly affected by environmental, occupational, personal and social factors. These factors may be responsible for a number of respiratory disorders.



- **Pulmonary embolism** is a blood clot that occurs in the lungs.
- **Bronchitis** is an inflammation of the lining of your bronchial tubes.
- **Asthma** is a condition in which airways narrow and swell and produce extra mucus.
- **Lung cancer** -The number one cause of cancer deaths. Smoking is the risk factor for lung cancer.
- **Pneumonia** is an inflammatory condition of the lung affecting primarily the small air sacs known as alveoli.
- **Pulmonary edema** is a fluid accumulation in the tissue and air spaces of the lungs.
- **Emphysema** A lung condition that causes shortness of breath due to widening of alveoli.
- **Atelectasis** is a collapse of a lung or lobe of a lung develops when alveoli within the lung become deflated.
- **Tuberculosis** is an infectious disease caused due to *Mycobacterium tuberculae*.
- **Pleurisy** is a condition in which the pleura becomes inflamed

the gaseous exchange system. Nicotine is the chemical that causes addiction and is a stimulant which makes the heart beat faster and the narrowing of blood vessels results in raised blood pressure and coronary heart diseases. Presence of carbon monoxide reduces oxygen supply. Lung cancer, cancer of the mouth and larynx is more common in

Sumanan noticed that his close friend was addicted to cigarette smoking. He advised his friend and explained the ill-effects of smoking. As a Biology student, explain what advice he might have given to his friend regarding the ill-effects of smoking.

smokers than non-smokers. Smoking also causes cancer of the stomach, pancreas and bladder and lowers sperm count in men.

Smoking can cause lung diseases by damaging the airways and alveoli and results in emphysema and chronic bronchitis. These two diseases along with asthma are often referred as Chronic Obstructive Pulmonary Disease (COPD). When a person smokes, nearly 85% of the smoke released is inhaled by the smoker himself and others in the vicinity, called passive smokers, are also affected. Guidance or counselling should be done in such users to withdraw this habit.

AMAZING FACTS


- The World TB Day is March 24.
- Direct Observation Therapy (DOTs) can treat about 95% of the TB patients.
- The surface area of the lungs is roughly the same size as a tennis court (525 feet long).
- It is possible to live with one lung.
- The highest recorded 'sneeze speed' is 165 km per hour.
- Adults breathe around 12 – 16 times per minute where as new borns breathe around 30–60 times per minute.
- Yawning helps us to breathe more oxygen to the lungs. When our brain senses the shortage of O_2 , it send a message to CNS to imbalance to O_2 demand and trigger us to yawn.
- Breathing through mouth results in bladder shrinkage and creates an urge to urinate in the middle of the night.
- Most people can hold their breath between 30 seconds to one minute.
- Hiccups are due to eating too fast or having occasional spasms of the diaphragm.

Activity


To test the presence of CO_2 in exhaled air Take two test tubes A and B with few mL of clear lime water. Blow exhaled air into A with a help of a straw and pass normal air into B with a help of a syringe for about 15 times and observe the changes that occur in the tubes A and B. The lime water (Calcium Hydroxide) in the test tube A turns milky.



Respire



Let's explore the anatomy and function of the **Respiratory system.**



Step – 1

Use the URL to reach the 'Respiratory System' page. In the grid select 'Nasal cavity' and explore its structure and the functions.

Step – 2


Now click back button on the top of the window or use the 'Backspace' key. Select 'Pharynx' from the grid and explore its anatomical regions.

Step – 3

Follow the above steps to explore each part and its functions.

Step – 4

Use the reference given below the page to acquire additional details.



Step 1



Step 2



Step 3



Step 4

Respiratory System's URL:

<https://www.getbodysmart.com/respiratory-system>

Schematics of Gas exchange:


<https://www.wisc-online.com/learn/general-education/anatomy-and-physiology2/ap2404/respiratory-system-gas-exchange>




B167_STD_11_ZOOLOGY_EM

* Pictures are indicative only

Concept map





Summary:

The process of intake of oxygen rich air and giving out of air rich in carbon dioxide is generally called respiration. Pollutants and micro organism are filtered from the inspired air by the hair and mucus present in the nostrils. The two main steps in the mechanism of respiration are inspiration and expiration which takes place due to pressure gradient in the atmosphere and lungs.

O₂ is transported in blood in dissolved form and is also bound to haemoglobin. One molecule of haemoglobin can bind four molecules of O₂. The Sigmoid shape of the O₂ haemoglobin dissociative curve shows increased affinity for each O₂ molecule.

CO₂ is transported in blood in dissolved form as carbamino haemoglobin

and as HCO₃. HCO₃ is produced in RBCs from CO₂ and water catalysed by carbonic anhydrase. Breathing is controlled by medullary respiratory centre.

Respiratory volumes and capacities indicate the amount of air inspired and expired during normal respiration. Our respiratory system can be affected by pollutants, pathogens and other chemical substances found in air. Lung cancer and emphysema cannot be cured and these diseases are common among cigarette smokers.

People at higher level than the sea level are prompted to altitude sickness as the barometric pressure is low in those regions. Surfactant, emphysema, Asthma and Dead space have been discussed. During vigorous exercise the rate of respiration increases.

Glossary

Apnoea – Temporary stopping of respiration.

Book gills – Respiratory organs in aquatic Limulus.

Book lungs – Respiratory organs of Scorpions and most spiders.

COLD – Chronic Obstructive Lung Disease.

Dyspnoea – painful respiration.

Epiglottis – a thin elastic cartilaginous flap which covers the glottis and prevents the entry of food into the larynx.

Haemoglobin – iron containing red pigment of RBCs of vertebrates, gives red colour to blood.

Herring-Breuer reflex – a defensive mechanism against over dilation of lungs.

Hypoxia – the failure of tissues for any reason to receive an adequate supply of oxygen.

Pneumothorax – presence of air in the pleural cavity which causes collapsing of lungs.

Vocal cords – sound regulating cords also called larynx or voice box.

Yawning – prolonged inspiration due to increase in CO₂ concentration.

Evaluation

- Breathing is controlled by
 - cerebrum
 - medulla oblongata
 - cerebellum
 - pons
- Intercostal muscles are found between the
 - vertebral column
 - sternum
 - ribs
 - glottis
- The respiratory structures of insects are
 - tracheal tubes
 - gills
 - green glands
 - lungs
- Asthma is caused due to
 - bleeding in pleural cavity.
 - infection of nose
 - damage of diaphragm.
 - infection of lungs
- The Oxygen Dissociation Curve is
 - sigmoid
 - straight line
 - curved
 - rectangular hyperbola
- The Tidal Volume of a normal person is
 - 800 mL
 - 1200 mL
 - 500 mL
 - 1100 – 1200 mL
- During inspiration, the diaphragm
 - expands.
 - unchanged
 - relaxes to become domed-shaped.
 - contracts and flattens
- CO₂ is transported through blood to lungs as
 - carbonic acid
 - oxyhaemoglobin
 - carbamino haemoglobin
 - carboxy haemoglobin
- When 1500 mL air is in the lungs, it is called
 - vital capacity
 - tidal volume
 - residual volume
 - inspiratory reserve volume
- Vital capacity is
 - TV + IRV
 - TV + ERV
 - RV + ERV
 - TV + TRV + ERV
- After a long deep breath, we do not respire for some seconds due to
 - more CO₂ in the blood
 - more O₂ in the blood
 - less CO₂ in the blood
 - less O₂ in the blood
- Which of the following substances in tobacco smoke damage the gas exchange system?
 - carbon monoxide and carcinogens
 - carbon monoxide and nicotine
 - carcinogens and tar
 - nicotine and tar

13. Column I represents diseases and column II represents their symptoms. Choose the correctly paired option

Column I	Column II
----------	-----------

- | | |
|---------------|---|
| (P) Asthma | (i) Recurring of bronchitis |
| (Q) Emphysema | (ii) Accumulation of W.B.CS in alveolus |
| (R) Pneumonia | (iii) Allergy |

- a. P = iii, Q = ii, R = i
- b. P = iii, Q = i, R = ii
- c. P = ii, Q = iii, R = i
- d. P = ii, Q = i, R = iii

14. Which of the following best describes the process of gas exchange in the lungs?

- a. Air moves in and out of the alveoli during breathing.
- b. Carbon dioxide diffuses from deoxygenated blood in capillaries into the alveolar air.
- c. Oxygen and carbon dioxide diffuse down their concentration gradients between blood and alveolar air.
- d. Oxygen diffuses from alveolar air into deoxygenated blood.

15. Make the correct pairs.

Column-I	Column-II
----------	-----------

- | | |
|--------|--|
| (P) IC | i. maximum volume of air breathe in after forced. |
| (Q) EC | ii. Volume of air present after expiration in lungs. |
| (R) VC | iii. Volume of air inhaled after expiration. |

- (S) FRC iv. Volume of air exhaled after inspiration.

- (a) P – i , Q – ii , R – iii , S – iv
- (b) P – ii , Q – iii , R – iv , S – i
- (c) P – ii , Q – iii , R – i , S – iv
- (d) P – iii , Q – iv , R – i , S – ii

16. Make the correct pairs.

Column-I	Column-II
----------	-----------

- | | |
|--------------------------------|----------------------|
| (P) Tidal volume | i. 1000 to 1100 ml |
| (Q) Residual volume | ii. 500 ml |
| (R) Expiratory reserve volume | iii. 2500 to 3000 ml |
| (S) Inspiratory reserve volume | iv. 1100 to 1200 ml |

- (a) P – ii , Q – iv , R – i , S – iii
- (b) P – iii , Q – ii , R – iv , S – i
- (c) P – ii , Q – iv , R – iii , S – i
- (d) P – iii , Q – iv , R – i , S – ii

17. Name the respiratory organs of flatworm, earthworm, fish, prawn, cockroach and cat.

18. Name the enzyme that catalyses the bicarbonate formation in RBCs.

19. Air moving from the nose to the trachea passes through a number of structures. List in order of the structures.

20. Which structure seals the larynx when we swallow?

21. Resistance in the airways is typically low. Why? Give two reasons.

22. How the body makes long-term adjustments when living in high altitude.

23. Diffusion of gases occurs in the alveolar region only and not in any other part of the respiratory system. Discuss.
24. Sketch a flow chart to show the path way of air flow during respiration.

Competitive Exam Corner

Sarojini's father has congestion of the lungs. His doctor advised him to take bedrest and prescribed him an inhaler. What disease is he suffering from? List the symptoms for the disease.

A villager who came to the city was affected by severe respiratory illness due to the inhalation of particulate pollutants. Suggest the reason for his illness and how does particulate pollutants affect him.

Kumar's mother works in a stone grinding factory. Suddenly she faints and taken to the hospital. The doctor notices fibres in the lungs. What kind of disease is she affected with? How can it be rectified?

Web links

http://kidshealth.org/kid/closet/movies/how_the_body_works_interim.html

25. Why is pneumonia considered a dangerous disease?
26. Explain the conditions which creates problems in oxygen transport.

References

1. Guyton A.C. and Hall J.E, (2006) Textbook of Medical Physiology– Eleventh Edition Elsevier Saunders, International Edition ISBN 0–8089–2317–X.
2. Mackean D.G. and Hayward D (2014). AS and A level biology book, Cambridge International, 3rd edition, Hodder Education, An Hachette UK company, London NW1 3BH.
3. Dee Unglaub Silverthron, Human physiology –an integrated approach – 7th Edition – Pearson Global edition.
4. Elaine N. Marieb and Katja Hoehn (2010). Human Anatomy and Physiology Eighth Edition, Benjamin Cummings, Pearson. New York.
5. Lauralee Sherwood and Robert Kell. (2007). Human physiology from cells to systems. First Canadian Edition Nelson Education Ltd, Toronto, Ontario MIK SG4.
6. Moyes and Schulte, 2016 Principles of Animal Physiology– 2nd edition, Pearson
7. Muthayya N.M., 2010 Human physiology– 4th edition, Jaypee brother medical publishers.