WEST BENGAL BOARD CLASS 12 MATHS SAMPLE PAPER SOLUTIONS

ANSWERS & EXPLANATION

SECTION – A

1. Solution

Option: b

 $x^2 - 6x + 10$ is is the equation to form equal and real roots

We will subtract it by 1 to form $x^2 - 6x + 9$

$$x^2 - 3x - 3x + 9$$

x(x-3) - 3(x-3)

(x-3)(x-3)

2. Solution

120x + 20y + 40z = 0

When x = 2, y = 1

120(2) + 20(1) + 40z = 0

40z = -260

$$z = -\frac{260}{40}$$
$$z = -\frac{13}{2}$$

z = -6.5

3. Solution

Area of
$$\triangle ABC = \sqrt{\Delta a^2 + \Delta b^2 + \Delta c^2}$$

Where, $\Delta a = \frac{1}{2} \begin{vmatrix} b_1 & c_1 & 1 \\ b_2 & c_2 & 1 \\ b_3 & c_3 & 1 \end{vmatrix}$,

$$\Delta b = \frac{1}{2} \begin{vmatrix} a_1 & c_1 & 1 \\ a_2 & c_2 & 1 \\ a_3 & c_3 & 1 \end{vmatrix},$$

And $\Delta c = \frac{1}{2} \begin{vmatrix} a_1 & b_1 & 1 \\ a_2 & b_2 & 1 \\ a_3 & b_3 & 1 \end{vmatrix}$

Hence it is False.

4. Solution

To solve rewrite the above as

$$2(x-y)=2\pi$$

 $x - y = \pi$

$$y = x - \pi$$

 $\frac{dy}{dx} = 1$ (since differentiating the constant function takes the value)

5. Solution

Option: a

 $x = 2t^2 + 4t; y = 4t$

$$\frac{dx}{dt} = 4t + 4 \text{ and } \frac{dy}{dt} = 4$$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dy}{dt}} = \frac{4}{4t+4} = \frac{1}{t+1}$$

$$dy = 1$$

 $\frac{dy}{dx} = \frac{1}{t+1}$

6. Solution

Option: a

 $\int_{0}^{2} \frac{1}{3x+2}$

$$=\frac{1}{3x+2}dx$$

Substituting u = 3x + 2; dx =
$$\frac{1}{3}$$
 du
= $\frac{1}{3}\int_{0}^{2}\frac{1}{u}du = \frac{1}{3}[\log(u)]_{0}^{2}$
u = 3x + 2
= $\frac{1}{3}[\log(3x+2)]_{0}^{2}$
 $\frac{\log(3x)}{3} + \frac{\log(2)}{3}$
 $\frac{\log 6}{3} + \frac{\log 2}{3} - \frac{\log 2}{3}$
= $\frac{\log 6}{3}$

Answer: true

Explanation: (according to the property of indefinite integral)

$$\int \frac{d}{dx} \left[\int f(x) dx - \int g(x) dx \right] = 0$$

hence $\int f(x)dx - \int g(x)dx = c$, where C is any real number

so, the families of curve $\{\int f(x) dx + C_1, C_1 \in R\}$

and $\{\int g(x)dx + C_2, C_2 \in R\}$ are identical

hence in this sense, $\int f(x) dx$ and $\int g(x) dx$ are equivalent.

8. Solution

Option: a

$$= e^{2-x}dx$$

Substituting $u = 2 - x \rightarrow dx = -du$

 $= -\int e^{u} du$

$$= -\int e^{2-x}$$

$$= -e^{2-x}dx + C$$

Putting value 2 and 0

$$= e^2 - 1$$

= 6.3 is the approximate answer

9. Solution

Option: c

Slope of the tangent at x=3 is

$$\frac{dy}{dx} = (9x^2 - 6)_{x=3}$$

= 9 × 9 - 6 = 81 - 6 = 75
= 75

10. Solution

Option: a

Note that at x-axis; y = 0. So the equation of the curve at y = 0 is given by

$$\frac{dy}{dx} = \frac{x}{(x-9)(x-1)}$$

$$\frac{\frac{d}{dx}(x). [(x-9). (x-1)] - x. \frac{d}{dx}[(x-9)(x-1)]}{(x-9)(x-1)^2}$$

$$= \frac{(x-9)(x-1) - x(2x-10)}{((x-1)(x-9))^2}$$

Therefore the slope at x = 0 is

$$\frac{(-9)(-1) - 0(-10)}{-1^2 \times -9^2} = \frac{9}{81} = \frac{1}{9}$$

Hence equation of the tangent at (0, 0)

$$y-0=\frac{1}{9}(x-0)$$

$$y = \frac{1}{9}x \text{ or } 9y = x$$

SECTION B

11. Solution

Let the original speed be $x \frac{km}{hr}$, therefore $\frac{16}{x} + \frac{9}{x+3} = 5$

$$\frac{16(x+3)+9x}{x(x+3)} = 5$$

 $16x + 48 + 9x = 5x^2 + 15$

 $= 5x^2 - 25x - 33$

$$x = 2.1, -3.1$$

Since x is the speed which cannot be negative hence 2.1 $\frac{km}{h}$ is the original speed

12. Solution

a = 200 first term, b = 400 second term, l =1200 last term.

According to formula when number of terms is not given the SUM of A.P is

Sum =
$$\frac{(b+1-2a)(1+a)}{2(b-a)}$$
:
Sum = $\frac{(400+1200-400)(1200+200)}{2(400-200)}$

$$Sum = \frac{1200 \times 1400}{2 \times 200}; sum = 4200$$

Now as we know Sum of A. P. = $\frac{n}{2}(a+1)$; n=number of terms

Hence sum
$$= \frac{n}{2}(a + 1);$$

 $\frac{n}{2}(a + 1) = 4200$
 $\frac{n}{2}(200 + 1200) = 4200$
 $\frac{n}{2} = \frac{4200}{1400}; \frac{n}{2} = 3; n = 6$

Hence the number of terms is 6

 $\frac{3}{4} = \frac{a+(x-1)d}{a+(x+y-1)d}$

3a + 3(x + y - 1)d = 4a + 4(x - 1)d

3xd + 3a + 3yd - 3d = 4pd - 4d + 4a

3yd + d = a + xd

3y = a

$$y = \frac{a}{3}; y = 1$$

Hence the y^{th} term is 1



The distance between two lines is given by

Distance =
$$\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

= $\sqrt{(2 - 1)^2 + (8 - 3)^2}$
= $\sqrt{1^2 + 5^2}$

Distance = $\sqrt{26}$

Midpoint of a line is given by $\left(\frac{(x_{2+}x_1)}{2}\frac{(y_{2+}y_1)}{2}\right)$

$$= \left(\frac{1+2}{2}, \frac{3+8}{2}\right)$$
$$= \frac{3}{2}, \frac{11}{2}$$

Hence the midpoint of line P (1, 3) to Q (2, 8) is $\frac{3}{2}$, $\frac{11}{2}$

OR

b) Solution

The distance from A to B is

$$=\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$



$$= \sqrt{(3-1)^2 + (-5-5)^2}$$

= $\sqrt{(2)^2 + (-10)^2}$
= $\sqrt{104}$
Diameter = $\sqrt{104}$
Radius = $\frac{\text{Diameter}}{2} = \frac{\sqrt{104}}{2} = \frac{10.19}{2} = 5.099 = 5.1$
Hence the area of the circle is $\pi r^2 = \pi \times (5.1)^2$

Area of the circle is 81.71 unit²

15. a) Solution

$$= \frac{d}{dx} \left(\frac{yx^{3}}{3} + yx - 7 \right)$$

= $\frac{y}{3} \frac{d}{dx} (x^{3}) + \frac{yd}{dx} (x) - \frac{d}{dx} 7.$
= $\frac{3x^{2}y}{3} + y - 0$
= $yx^{2} + y;$

Answer:
$$y(x^2 + 1)$$

OR

b) Solution

$$\frac{d}{dx} = (x^{sinx})$$

Using the generalized formula we use

$$= \frac{d}{dx} [u(x)^{v(x)}] = u(x)^{v(x)} \cdot \frac{d}{dx} = (\log(v(x)) \cdot (v(x)))$$
$$= \frac{d}{dx} (\log(x)) \cdot \sin(x) + (\log(x) \cdot \frac{d}{dx} (\sin(x)) \cdot x^{\sin x}$$
$$= \left(\left(\frac{1}{x}\right) \cdot \sin(x) + \log(x) \cdot \cos(x) \right) \cdot x^{\sin x}$$

$$= x^{sinx} \cdot \left(\frac{sinx}{x} + cosx.logx\right)$$

16. a) Solution

$$\int \frac{6x}{(x-10)(2x-1)} dx$$
$$6 \int \frac{x}{(x-10)(2x-1)} dx$$

Now solving using partial fraction

$$\int \left(\frac{10}{19(x-10)} - \frac{1}{19(2x-1)}\right) dx$$
$$\frac{10}{19} \int \frac{1}{x-10} dx = \frac{1}{19} \int \frac{1}{2x-1} dx$$

Now solving

$$\frac{10}{19} \int \frac{1}{x - 10} dx = \frac{10}{19} \cdot \log(x - 10)$$
$$\frac{1}{19} \int \frac{1}{2x - 1} dx = \frac{1}{19} \cdot \frac{1}{2} \log(2x - 1)$$
$$= \frac{10\log(x - 10)}{19} - \frac{\log(2x - 1)}{38}$$
$$= \frac{60\ln(x - 10)}{19} - \frac{3\log(2x - 1)}{19}$$
$$- \frac{3\ln(|2x - 1|) - 60\log(|x - 10|)}{19} + C$$

OR

b) Solution

$$= 3xe^{3-x}dx$$
$$= 3e^{3-x}\int xdx$$

Solving

$$= \int x e^{-x} dx$$
$$= -x e^{-x} - \int -e^{-x} dx$$



Putting respective values in x = 2, 0 we have

$$= 3(e^3 - 3e)$$

= 35.79 (Approximate value)

17. a) Solution

Separating variable we get

$$(y-3)dy = (2x+1)dx$$

Integrating both sides

$$\int (y-3)dy = \int (2x+1)dx$$

$$\int ydy - 3 \int dy = 2 \int xdx + 1 \int dx$$

$$\frac{y^2}{2} - 3y = 2 \cdot \frac{x^2}{2} + x$$

$$\frac{y(y-6)}{2} = x^2 + x + C_1$$

$$y^2 - 6y = 2x^2 + 2x + 2C_1$$

$$y^2 - 6y - 2x^2 - 2x - 2C_1 = 0; \ 2C_1 = C$$

$$y(y-6) - 2(x^2 - x) - C = 0$$

OR

b) Solution

The slope of tangent to a curve is given $by \frac{dy}{dx}$

$$\frac{dy}{dx} = \frac{2x+3}{3y^2}$$
$$\int 3y^2 \, dy = \int (2x+3) \, dx$$

Solving both sides at the same time.

$$3\int y^2 dy = 2\int x dx + 3\int dx$$

According to formula $y^n = \frac{y^{n+1}}{n+1}$

$$\frac{3y^3}{3} = \frac{x^2}{2} + 3x$$

$$y^3 = x^2 + 6x + C; (x,y) = (1,-2)$$

$$-8 = 1 + 12 + C$$

$$C = -21$$

The equation of the required curve $y^3 = x^2 + 3x - 21$,

18. Solution

Volume=V, surface Area=S.A. and h = x = 4cm

$$V = \frac{1}{3} \pi r^2 h \text{ or } V = \frac{1}{3} \pi r^2 x$$

S. A. = $\pi r \sqrt{r^2 + x^2}$
$$\frac{dv}{dt} = 6 \text{ cm}^3/\text{sec}$$
$$\frac{dv}{dt} = 6 = \frac{d}{dt} \left(\frac{1}{3} \pi r^2 x\right) = \frac{d}{dx} \left(\frac{1}{3} \pi r^2 x\right) \cdot \frac{dx}{dt}$$

$$6 = \left(\frac{1}{3}\pi r^{2}\right)\frac{dx}{dt}$$

$$18 = \pi r^{2} \cdot \frac{dx}{dt} \quad \text{with } r = 3\text{ cm}$$

$$\frac{2}{\pi} = \frac{dx}{dt}$$

$$\frac{ds}{dt} = \pi r \sqrt{r^{2} + x^{2}} = \frac{d}{dx} \left[\pi r^{2} \left[\sqrt{r^{2} + x^{2}}\right] \cdot \frac{dx}{dt}\right]$$

$$\frac{dx}{dt} = \frac{2}{\pi}$$

$$\frac{3\pi}{2} \cdot \left[(x^{2} + 9)^{-\frac{2}{3}}\right] \cdot \frac{d}{dx} (x^{2} + 9) \cdot \frac{2}{\pi}$$

$$\frac{3\pi x}{\sqrt{(x^{2} + 9)}} \cdot \frac{2}{\pi} = \frac{6x}{\sqrt{x^{2} + 9}} \quad \text{When } x = 4$$

$$\frac{6 \times 4}{\sqrt{4^{2} + 9}} = \frac{24}{\sqrt{25}} = 4.8 \text{ cm}^{2}/\text{sec}$$

We have $f(x) = x^2 - 8x + 16$

f'(x) = 2x - 8

=Therefore if f'(x) = 0; then

$$0 = 2x - 8; 2x = 8; x = 4.$$

Now point x=2 cuts the line into two disjoint intervals as $(-\infty, 4)$ and $(4, \infty)$.

In the interval $(-\infty, 4)$; f'(x) = 2x - 8 < 0

Therefore F is strictly decreasing in this interval

In the interval $(4, \infty)$; f'(x) > 0 and therefore function is strictly increasing.

20. Solution

Let x = 5 and $\Delta x = 0.09$

Then $f(5.09) = f(x + \Delta x)$

 $= 4(x + \Delta x)^{2} + 3(x + \Delta x) + 9$

Since
$$\Delta y = f(x + \Delta x) - f(x)$$

 $f(x + \Delta x) = f(x) + \Delta y$
 $f(5.09) = (4x^2 + 3x - 9) + (8x + 3)\Delta x$
 $f(5.09) = (4(5)^2 + 3(5) - 9) + (8(5) + 3) \ 0.09$
 $f(5.09) = 108 + 3.87 = 111.87$

N be the point on CD

ND=(30-x) m and CD=30m

 $MN^2 = MC^2 + CN^2$

 $NO^2 = ND^2 + DO^2$

 $NM^2 + NO^2 = MC^2 + CN^2 + ND^2 + DO^2$

 $196 + x^2 + (x^2 - 60x + 900) + 576$

 $A(x) = 2x^2 - 60x + 1672$

A'(x) = 4x - 60

now at A'(x) = 0; given x = 15

Also
$$A^{\prime\prime}(x) = 4 > 0$$

Therefore by second derivative test; x = 15 is the point of local minima of A

Thus, distance from C to N is 15cm

22. Solution

For every single value of x the deer's position is at point(x, y)

i.e. $(x, y) = (x, x^2 + 4)$, there the closest distance between deer and

Hunter at (1, 1) is.

$$f(x) = \sqrt{(x-1)^2 + (x^2+3)^2}$$
$$f'(x) = 6x^2(x^3+3) + 2(x-1)^2$$

thus f'(x) = 0 gives x = 1.68

Hence as there is no real value of the equation if found, but found only for 1 point i.e. at x=1.68, $f(1.68) = (1.68 - 1)^2 + (1.68^2 + 3)^2$

Thus the distance between the deer and the hunter is $\sqrt{f(x)} = \sqrt{34.46}$

It follows the minimum value of is $\sqrt{f(x)} = \sqrt{34.46} = 5.8$ meters is the shortest distance that the hunter can shoot the deer.

SECTION- C

23. Solution

 $a_{n} = a + (n - 1)d$ $\frac{a_{n}}{a} = \frac{a}{a} + \frac{(n - 1)d}{a}$ $\frac{5}{2} = 1 + \frac{n - 1}{4}$ $\frac{5}{2} - 1 = n - \frac{1}{4}$ 6 = n - 1; n = 7Series No.2 a = 20, d = 2 $S_{7} = \frac{n}{2}[2a + (n - 1)d]$ $S_{7} = \frac{7}{2}[2 \times 20 + 6 \times 2]$ $S_{7} = \frac{7}{2}[52] = 182$

24. Solution

As we know

 $a_{n-1} = a + (n - 1 - 1)d$ $36 = a + (n - 2)d \rightarrow 1$ $a_2 = a + (n - 1)d$

$$20 = a + d \rightarrow 2$$

$$a = 20 - d$$

$$36 = (20 - d) + nd - 2d$$

$$16 = -d + nd - 2d$$

$$16 = d(-1 + n - 2)$$

$$16 = (n - 3)d$$

As the third term from the last of the series is 32

$$a_{n-3} = a + (n-3)d \rightarrow 3$$

 $32 = a + 16$
 $16 = a$
Hence if a=16 and if we put it ina₂, then

20 = 16 + d

Therefore if a=16 and d=4 then the number of terms in the series is

Taking in equation number .1

$$36 = 16 + (n - 2)4$$

$$5 = (n-2)$$

Hence the sum of the series is:

$$S_7 = \frac{n}{2} [2a + (n-1)d]$$

$$S_7 = \frac{7}{2} [2 \times 16 + (7-1) \times 4]$$

$$S_7 = 3.5 \times [32 + 24]$$

$$S_7 = 196.$$

Hence the sum of the series is 196

Area of the rectangular Park is

Area = Length \times breadth

Breadth = x

Area = (x+7)(x)

144 = (x + 7)(x)

 $144 = x^2 + 7x$

 $x^2 + 7x - 144 = 0$

 $x^2 + 16x - 9x - 144 = 0$

x(x+16) - 9(x+16)

(x-9)(x+16)

The negative value for a breadth is not taken hence the value for consideration is -16.

Hence to find out the cost of laying boundary wall around the park is

- = perimeter of the park × Cost per meter.
- $= 2(L + B) \times 10$
- $= 2(9+17) \times 10$
- $= 2(26) \times 10$
- $= 52 \times 10$
- = Rs.520

Therefore the cost of the laying boundary wall around the park is Rs. 520

26. Solution

If equation $2x^2 + 4xy + 2y^2 + 8x + 8y - 10$ gives 2 parallel lines then it should follow the following conditions

 $h^2 - ab = 0$

$$hg = af$$

4 = 2h or h = 2

$$a = 2; b = 2; g = 4; f = 4.$$

Hence after substitution we find that

 $h^2 - ab = 0$

4-4=0

And

hg = af

 $2 \times 4 = 4 \times 2$

Hence the eqn. $2x^2 + 4xy + 2y^2 + 8x + 8y - 10$ does produces parallel lines

Under the following condition $ax + hy + g = \pm \sqrt{g^2 - ca}$

Gives two parallel lines

i)
$$ax + hy + g = +\sqrt{g^2 - ca}$$

ii)
$$ax + hy + g = -\sqrt{g^2 - ca}$$

if both equation i and ii gives lines that are parallel in nature

i)
$$2x + 2y + 4 = +\sqrt{4^2 - (-10)^2}$$

= 2x + 2y + 4 - 6 = 0

line 1

ii)
$$2x + 2y + 4 = +\sqrt{4^2 - (-10)^2}$$

$$= 2x + 2y + 4 + 6 = 0$$
 line 2

Therefore the 2 parallel lines are:

2x + 2y + 4 - 6 = 0 and 2x + 2y + 4 + 6 = 0

27. Solution

The given eqn. $8x^2 + 2xy - 4y^2 - 24x + 6y + 2k = 0$

Here a=8, b=-4, h=1, g=-12, f=3, c=2k

As it represents two straight lines

$$\Delta = 8. -4.2k + 2.8.(12)^2 \cdot 1 - 8(3)^2 - (-4)(12)^2 - k \cdot 1$$

$$\Delta = -64k + 2304 - 216 - (-576) - k$$

$$\Delta = 0, \qquad 65k = 2304 + 576 - 216$$

 $k = 40.98 \text{ or } k \approx 41 \text{ Answer}$

28. Solution



Let AB be straight line on which point f fall perpendicularly the equation is given by

 $a \cos \alpha^{\circ} + b \sin \alpha^{\circ} - t$

 $2\cos 30^\circ + 3\sin 30^\circ - 2$

The lines pass through point F(4,5)

XS =perpendicular distance from point F to AB

XS = DO - SO

$$XS = 2 \times \frac{\sqrt{3}}{2} + 3 \times \frac{1}{2} - 2 = 5 \times 19 + 3 \times 5 - 2$$

Hence the distance between point and the straight line is **6.69**.

29. Solution

$$\frac{(3x^2 + 7x + 4)}{(x-3)^2(x-2)}dx$$

Perform partial fraction decomposition:

$$= \int \left(\frac{30}{(x-2)} - \frac{27}{(x-3)} + \frac{52}{(x-3)^2}\right) dx$$
$$= 30 \int \frac{1}{(x-2)} dx - 27 \int \frac{1}{(x-3)} dx + 52 \int \frac{1}{(x-3)^2} dx$$
$$= 30I_1 - 27I_2 + 52I_3$$

Now solving $I_{\!1\!}$

$$\int \frac{1}{(x-2)} dx$$

Substitute $\mathbf{u} = \mathbf{x} - \mathbf{2} \rightarrow \mathbf{d}\mathbf{x} = \mathbf{d}\mathbf{u}$

$$=\int \frac{1}{u} du$$

 $= \log u$

Substitution $\mathbf{u} = \mathbf{x} - \mathbf{2}$:

$$= \log(x - 2)$$

Now solving I_2

$$\int \frac{1}{x+3} dx$$

Substitute $\mathbf{u} = \mathbf{x} - \mathbf{3} \longrightarrow \mathbf{d}\mathbf{x} = \mathbf{d}\mathbf{u}$

$$=\int \frac{1}{u} du$$

 $= \log u$

Substitution $\mathbf{u} = \mathbf{x} - \mathbf{3}$:

 $= \log(x - 3)$

 $\mathsf{Solving} I_{\mathfrak{B}}$

$$\int \frac{1}{(x+3)^2} dx$$

Substitute $u = x - 3 \rightarrow dx = du$

$$=\int \frac{1}{u^2} \, \mathrm{d} u = -\frac{1}{u}$$

Substitution u = x - 3:

$$=-\frac{1}{(x-3)}$$

Putting all the values of I_1 , I_2 , $I_3 \mbox{we get}$

$$= -\frac{52}{x-3} + 30 \log(x-2) - 27 \log(x-3)$$

Hence the solved integral is

$$\frac{(3x^{2}+7x+4)}{(x-3)^{2}(x-2)}dx = -\frac{52}{x-3} + 30\log(x-2) - 27\log(x-3)$$

30. Solution

$$= \int \frac{x}{(x+3)^2 + (x-3)^2} dx$$

Substitute u = (x + 3)² + (x - 3)² \rightarrow dx = $\frac{1}{(2(x+3)+2(x-3))} du$
= $\int \frac{1}{4u} du$

Simplify:¹/₄∫¹/_udu

Now solving $\int \frac{1}{u} du$

Apply in the solved integral
$$\frac{1}{4}\int \frac{1}{u} du$$

$$\frac{\log(u)}{4}$$

Substitution $u = (x + 3)^2 + (x - 3)^2$

$$=\frac{\log((x+3)^2+(x-3)^2)}{4}$$

Hence the solution for $\int \frac{x}{(x+3)^2+(x-3)^2} dx$ is

$$=\frac{\log((x+3)^2 + (x-3)^2)}{4} + C$$
$$=\frac{\log(x^2 + 9)}{4} + C$$

Hence the solution is $\frac{\log(x^2+9)}{4} + C$

31. Solution

 $3\int x\sqrt{x^4+1}dx + 4\int x^3dx$

Now solving: $\int x\sqrt{x^4 + 1} dx$

Substitute
$$u = x^2 \rightarrow dx = \frac{1}{2x} du$$
 use: $x^4 = u^2$

$$=\frac{1}{2}\int\sqrt{u^2+1}du$$

Now solving $\int \sqrt{u^2 + 1} du$

Applying trigonometric substitution:

Substitute
$$u = tan(v) \rightarrow v = arctan(u)$$
, $du = sec^{2}(v)dv$

$$= \int \sec^2(\mathbf{v})\sqrt{\tan^2(\mathbf{v}) + 1}d\mathbf{v}$$

Simplify $\tan^2(v) + 1 = \sec^2(v)$

$$=\int \sec^{3}(v)dv$$

Apply reduction formula:

$$\int \sec^{n}(v) dv = \frac{(n-2)}{(n-1)} \int \sec^{n-2}(v) dv + \frac{\left(\sec^{(n-2)}(v)\tan(v)\right)}{n-1}$$

With n = 3

$$=\frac{(\sec(v)\tan(v))}{2}+\frac{1}{2}\int\sec(v)dv$$

Now solving ∫ sec(v)dv

$$= \log(\tan(v) + \sec(v))$$

Apply in solved integrals:

$$\frac{(\sec(v)\tan(v))}{2} + \frac{1}{2}\int \sec(v)dv$$
$$= \frac{\log(\tan(v) + \sec(v))}{2} + \frac{(\sec(v)\tan(v))}{2}$$

Substitution v=arc tan(u), use:

$$=\frac{\log(\sqrt{u^{2}+1}+u)}{2}+\frac{(u\sqrt{u^{2}+1})}{2}$$

Applying in solved integrals:

$$\frac{1}{2} \int \sqrt{u^2 + 1} du$$
$$= \frac{\log(\sqrt{u^2 + 1} + u)}{4} + \frac{(u\sqrt{u^2 + 1})}{4}$$

Substitution $\mathbf{u} = \mathbf{x}^2$

$$=\frac{\log(\sqrt{x^4+1}+x^2)}{4}+\frac{(x^2\sqrt{x^4+1})}{4}$$

Now solving $\int x^3 dx = \frac{x^4}{4}$

Again apply in the solved integrals:

$$3\int x\sqrt{x^4 + 1} \, dx + 4\int x^3 \, dx$$
$$= \frac{\left(3\log(\sqrt{x^4 + 1} + x^2)\right)}{4} + \frac{\left(3x^2\sqrt{x^4 + 1}\right)}{4} + x^4 + 6$$

Putting value -3 and 1 we get

$$=\frac{\left(3\log(\sqrt{-3^{4}+1}+-3^{2})\right)}{4}+\frac{\left(3(-3)^{2}\sqrt{(-3)^{4}+1}\right)}{4}+(-3)^{4}-\frac{\left(3\log(\sqrt{1^{4}+1}+1^{2})\right)}{4}+\frac{\left(3(1)^{2}\sqrt{(1)^{4}+1}\right)}{4}+(1)^{4}+C=141.57$$

We get an approximate value of $141.57 \approx 142$ Answer.

32. Solution

 $\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{(2x+y)}{(x+3y)}$

 $F(x,y)=\frac{\scriptscriptstyle (2x+y)}{\scriptscriptstyle (x+3y)}$

Therefore F(x, y) is a homogeneous function of degree zero. So the given differential eqn is a homogeneous differential eqn.

$$\frac{dy}{dx} = \frac{\left(2 + \frac{y}{x}\right)}{\left(1 + \frac{y}{y}\right)} = f\left(\frac{y}{x}\right)$$

To solve it we make substitution y = vx

$$\frac{dy}{dx} = v + \frac{dy}{dx}$$

$$v + x \frac{dv}{dx} = \frac{(2+v)}{(1+3v)}$$

$$x \frac{dv}{dx} = \frac{(2+v)}{(1+3v)} - v$$

$$x \frac{dv}{dx} = \frac{2+v-v(1+3v)}{(1+3v)}$$

$$x \frac{dv}{dx} = \frac{(-3v^3+2)}{(1+3v)}$$

$$\frac{(1+3v)}{(-3v^3+2)} dv = \frac{1}{x} dx$$

$$\int \frac{(1+3v)}{(-3v^3+2)} dv$$

$$= \int \frac{(3v)}{(-3v^3+2)} - \frac{(1)}{(-3v^3+2)} dv$$

$$= 3 \int \frac{(v)}{(-3v^3+2)} dv - \int \frac{(1)}{(-3v^3+2)} dv$$

$$= \int \frac{(v)}{(-3v^3+2)} dv$$

Substituting $u = 3v^2$; $dv = \frac{1}{6v}du$

+2

$$= 3 \times \frac{1}{6} \int \frac{1}{u} du$$
$$= 3 \times \frac{1}{6} \log(u)$$
$$= 3 \times \frac{\log(3v^{2}-2)}{6} = \frac{\log(3v^{2}-2)}{2}$$
$$= now \text{ we solve for } \frac{1}{3v^{2}-2} dv$$
$$= \frac{1}{3v^{2}-2} dv$$

$$=\int \frac{3}{(3v-\sqrt{6})(3v+\sqrt{6})}dv$$

After partial integration we get:

$$=\frac{\log(3v-\sqrt{6})}{2\sqrt{6}}-\frac{\log(3v+\sqrt{6})}{2\sqrt{6}}$$

2√6

Hence after putting all the solved values we get

$$= \frac{\log(3v + \sqrt{6}) - \log(3v - \sqrt{6})}{2\sqrt{6}} - \frac{\log(3v^2 - 2)}{2} + C$$
$$= \frac{\log(3v + \sqrt{6}) - \log(3v - \sqrt{6})}{2} - \frac{\log(3v^2 - 2)}{2} + C$$

2

Putting y=vx

33. Solution

$$\frac{dy}{dx} = \frac{x\left(2 + 3\left(\frac{y}{x}\right)^2\right)}{y}$$
$$\frac{dy}{dx} = v + \frac{xdv}{dx}$$

$$y = vx$$

$$\frac{x\left(2+3\left(\frac{y}{x}\right)^2\right)}{y} = v + \frac{xdv}{dx}$$

$$\frac{x(2+3(v)^2)}{vx} = v + \frac{xdv}{dx}$$

$$\frac{(2+3(v)^2)}{v} - v = \frac{xdv}{dx}$$

$$\frac{(2+3(v)^2) - v^2}{v} = \frac{xdv}{dx}$$

$$\frac{(2+2(v)^2)}{v} = \frac{xdv}{dx}$$

$$\frac{(2+2(v)^2)}{v} = \frac{xdv}{dx}$$

After solving integration on both sides

$$\log x = \frac{1}{4} \cdot \log\left(\left(\frac{y}{x}\right)^2 + 1\right) + C_1$$

$$4\log x - 4C_1 = 1 \cdot \log\left(\left(\frac{y}{x}\right)^2 + 1\right)$$

$$-4C_1 = 1 \cdot \log\left(\left(\frac{y}{x}\right)^2 + 1\right) - 4\log x$$

Dividing each with 4logx

We get

$$-\frac{C_{1}}{\log x} = \frac{1}{4} \cdot \log\left(\frac{y^{2} + x^{2}}{x^{2}} - x\right) - 1$$

$$\frac{y^{2} + x^{2}}{x^{2}} - x = t$$

$$-\frac{C_{1}}{\log x} = \frac{1}{4} \cdot \log(t) - 1$$

$$-\frac{C_{1}}{1} = \frac{1}{4} \cdot (\log x \cdot \log(t)) - \log x$$

$$C_{1} = -\frac{1}{4} \cdot (\log x \cdot \log(t)) + \log x$$

$$C_1 = -\frac{1}{4} \cdot \left(\left(\log(x+t) + \log x \right) \right)$$

$$C_1 = -\frac{1}{4} \cdot \left(\left(\log(x + \frac{y^2 + x^2}{x^2} - x) + \log x \right) \right)$$

$$C_1 = -\frac{1}{4} \cdot \left(\log\left(\frac{y^2 + x^2}{x^2}\right) \cdot x \right)$$

$$P = 2x^2; Q = 3x + 2$$

 $I.\,F.=e^{\int 2x^2}$

 $I.F.=e^{4x}$

According to the formula

$$y.(I.F.) = Q(I.F.)dx + C$$

$$y.(e^{4x}) = Q(e^{4x})dx + C$$

Putting (3x+2)=Q

$$y.(e^{4x}) = (3x + 2)(e^{4x})dx + C$$

After integrating with x we get

$$= \int (3x+2)(e^{4x})dx$$
$$= \frac{(3x+2)e^{4x}}{4} - \int \frac{3e^{4x}}{4}dx$$
$$= \int \frac{3e^{4x}}{4}dx$$
$$= u = 4x, \quad dx = \frac{1}{4}du$$
$$= \frac{3}{16}\int e^{u}du$$
$$= \frac{3}{16}e^{u}$$
$$= u = 4x$$

$$=\frac{3}{16}e^{4x}$$

Apply the solved into the $\frac{(3x+2)e^{4x}}{4} - \int \frac{3e^{4x}}{4} dx$ we get

$$\frac{(3x+2)e^{4x}}{4} - \frac{3}{16}e^{4x} + C$$
$$= \frac{(12x+5)e^{4x}}{16} + C$$

Now put the solution in the eqn $y.(e^{4x}) = Q(e^{4x})dx + C$

С

$$Q(e^{4x})dx + C = \frac{(12x + 5)e^{4x}}{16} + q^{2x}$$

$$y.(e^{4x}) = \frac{(12x + 5)e^{4x}}{16} + C$$

$$y = \frac{(12x + 5)e^{4x}}{16.e^{4x}} + \frac{C}{e^{4x}}$$
Answer $y = \frac{(12x + 5)}{16} + \frac{C}{e^{4x}}$

35. Solution

 $\frac{dv}{dt} = 20 \text{ m}^3/_{\text{sec}}$ $\frac{dv}{dt} = \frac{d}{dt} \left(\frac{4}{3} \cdot \pi r^3\right)$

Since r is the variable hence r=x

$$20 = \frac{4}{3}\pi \int x^{3}$$
$$20 = \frac{4}{3}\pi \frac{d}{dx}(x^{3}) \cdot \frac{dx}{dt}$$
$$20 = \frac{4}{3}\pi 3x^{2} \cdot \frac{dx}{dt}$$
$$\frac{5}{\pi x^{2}} = \frac{dx}{dt}$$

Hence to calculate how fast the surface area of the explosion increases,

We calculate.

$$\frac{ds}{dt} = \frac{d}{dt} \left(\frac{5}{\pi x^2} \right)$$

The surface area of the explosion is $4\pi r^2$, since r is variable r=x.

$$\frac{ds}{st} = \frac{d}{dx} (4\pi x^2) \frac{dx}{dt}$$
$$\frac{ds}{dt} = (4\pi 2x) \cdot \frac{5}{\pi x^2} \frac{dx}{dt}$$
$$\frac{ds}{dt} = \frac{4 \times 2 \times 5}{x} = \frac{40}{x}$$

if x=10m;

$$\frac{ds}{dt} = \frac{40}{10} = 4 \,\mathrm{m}^2/\mathrm{sec}.$$

Answer: The surface area of the explosion increases with $4m^2/sec.$

36. Solution

Let r be the radius of the hemisphere and Δr be the error in measuring the radius, then r=10, Δr =0.04, Now

Volume of the hemisphere $=\frac{2}{3}\pi r^2$

$$\frac{\mathrm{d}v}{\mathrm{d}r} = \frac{2}{3} \pi r^3$$

$$\frac{dv}{dr} = 2 \pi r$$

 $\frac{dv}{dr}\Delta r$

$$= (2\pi r^2)\Delta r = 2\pi (10)^2 \times 0.04$$

$$= 200\pi(0.04)$$

Thus the approximate error value for calculating volume is 8π cm³.

Hence the total error of the 3 hemisphere is $3 \times 8\pi \text{ cm}^3 = 24\pi \text{ cm}^3$.

37. Solution

$$f(x) = x^3 + 6x^2 - 63x + 14$$

$$f'(x) = 3x^{2} + 12x^{1} - 63$$

$$f'(x) = 0; \quad 3(x^{2} + 4x - 21)$$

$$= 3((x - 3)(x + 7))$$

thus $f'(x) = 0$; gives $x = 3, -7$

Hence x=-7 and 3 are the only points which gives us local maxima or local minima resp.

$$f(x) = (3)^{3} + 6(3)^{2} - 63(3) + 14$$

$$f(x) = 27 + 54 - 189 + 14$$

$$f(x) = -94$$

$$f(x) = x^{3} + 6x^{2} - 63x + 14$$

$$f(x) = (-7)^{3} + 6(-7)^{2} - 63(-7) + 14$$

$$f(x) = 406$$

Hence the minima for the eqn is $x^3 + 5x^2 + 6x + 1 - 94$ and the maxima is 406

38. Solution

 $x^{3} + 3x^{2} - 9x + 10 = f(x)$ $f'(x) = 3x^{2} + 6x - 9$ $= 3(x^{2} + 2x - 3)$ = 3((x-1)(x+3))if f'(x) = 0; then for x=1 $f(x) = x^{3} + 3x^{2} - 9x + 10$ $f(-1) = -1^{3} + 3(-1)^{2} - 9(-1) + 10$ f(-1)=5Hence the minima is (x,y) = (1,5) $f(x) = x^{3} + 3x^{2} - 9x + 10$ $f(-3) = (-3)^{3} + 3(-3)^{2} - 9(-3) + 10$

Hence the maxima is (x,y)=(-3,37)



let's take A= (1,5) and B= (-3,37) the distance between (1,5) to (-3,37).

$$D = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$D = \sqrt{(-3 - 1)^2 + (37 - 5)^2}$$

$$D = \sqrt{(4)^2 + (32)^2}$$

$$D = \sqrt{16 + 1024}$$

$$D = 32.24$$

$$r = \frac{D}{2} = 16.12 \text{cm}$$

Area = $\pi r^2 = \pi (16.12)^2 = 260\pi$.

39. Solution

The volume of the cylinder is $V{=}\,\pi r^2h$

$$V = \pi \left(\frac{h}{2}\right)^2 h$$
$$\frac{dV}{dt} = \frac{\pi h^2}{4} h = \frac{h^3 \pi}{4}$$
$$\frac{dv}{dt} = \frac{d}{dh} \left(\frac{\pi h^3}{4}\right) \cdot \frac{dh}{dt}$$

$$= \frac{\pi}{4} \times \frac{\mathrm{h}^3 \mathrm{dh}}{\mathrm{dt}} = \frac{3}{4} \pi \mathrm{h}^2$$

Now the rate of change in vol. of water per hour is $10 \, \frac{\mathrm{cm}^{\mathrm{s}}}{\mathrm{h}}$ and height is 6m

$$10 = \frac{3}{4}\pi \times 6 \times 6 \times \frac{dh}{dt}$$
$$10 = \frac{3}{4} \times \pi \times 36 \times \frac{dh}{dt}$$
$$\frac{10}{3\pi9} = \frac{dh}{dt}; \quad \frac{10}{27\pi} = \frac{dh}{dt}$$

Hence the rate at which the water level rising is $\frac{10}{27\pi}~cm/h$

40. Solution

$$f(x) = \sin^{-1}(\cos 2x + \cos x)$$

$$f'(x) = \frac{-2\sin(2x) - \sin x}{\sqrt{1 - (\cos(2x) + \cos(x))^2}}$$

Hence the denominator for the func. Should always be>0 for the function to be increasing.

$$\sqrt{1 - \left(\cos(2x) + \cos(x)\right)^2} > 0$$

 $\cos 2x + \cos x > 1$

 $\frac{\cos x}{\cos x} > 2sinx.\frac{\sin x}{\cos x}$

1 > 2sinxtanx

Now if we put $\left(\frac{3\pi}{2},\pi\right)$

For $\frac{3\pi}{2}$ We get-3; which is less than 1; -3<1

For π we get 0; which is less than 1 again.

Now for the f'(x)

hence for f'(x)i.e.
$$\frac{-2\sin(2x) - \sin x}{\sqrt{1 - (\cos(2x) + \cos(x))^2}}; \left(\frac{3\pi}{2}, \pi\right)$$

$$f'(x) > 0$$
 in $\left(\frac{3\pi}{2}, \pi\right)$.

Hence f(x) is strictly increasing. Answer

41. Solution

S. P. (x) =
$$\left(100 - \frac{x}{500}\right) \cdot \frac{x}{3}$$

C. P. (x) = $\left(\frac{x}{500} + 200\right)$
profit = S. P. (x) - C. P. (x)
profit(x) = $\left(100 - \frac{x}{500}\right) \times \frac{x}{3} - \left(\frac{x}{500} + 200\right)$

To find the maximum items sold to gain in this transaction we keep profit'(x)= 0

$$\operatorname{profit}'(\mathbf{x}) = \frac{100\mathbf{x}}{3} - \frac{\mathbf{x}^2}{1500} - \left[\frac{\mathbf{x}}{500} + 200\right]$$
$$\mathbf{0} = \frac{100}{3}\mathbf{x} + \frac{\mathbf{x}^2}{1500} - \frac{1}{500}$$

profit " (x) = -2/1500 < 0

x=24998, hence the maximum items he can sell to gain from this transaction is24998 items

Profit=S.P. –C.P.
=
$$\left(100 - \frac{x}{500}\right) \cdot \frac{x}{3} - \left(\frac{x}{500} + 200\right); x = 24998$$

= $\left(100 - \frac{24998}{500}\right) \cdot \frac{24998}{3} - \left(\frac{24998}{500} + 200\right)$

= 416633-249.996

Answer: Rs. 416383 is the profit in this transaction.

SECTION – D

42. a) Solution

The given system of equations can be written as AX = B

Where A = $\begin{bmatrix} 6 & 4 & 2 \\ 8 & 2 & 6 \\ 3 & 3 & 3 \end{bmatrix}$, X = $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$, B = $\begin{bmatrix} 20 \\ 30 \\ 18 \end{bmatrix}$ Now, $|A| = \begin{bmatrix} 6 & 4 & 2 \\ 8 & 2 & 6 \\ 3 & 3 & 3 \end{bmatrix}$

Expanding with respect to R_1

 $= 6 \begin{vmatrix} 2 & 6 \\ 3 & 3 \end{vmatrix} - 4 \begin{vmatrix} 8 & 6 \\ 3 & 3 \end{vmatrix} + 2 \begin{vmatrix} 8 & 2 \\ 3 & 3 \end{vmatrix}$ = 6(6 - 18) - 4(24 - 18) + 2(24 - 6)= -72 - 24 + 36 $= -60 \neq 0$

 $\Rightarrow A^{-1}$ exists, so the given system of equations has a unique solution $X = A^{-1}B$

$$\therefore \operatorname{Adj} A = \begin{bmatrix} \begin{vmatrix} 2 & 6 \\ 3 & 3 \end{vmatrix} & -\begin{vmatrix} 8 & 6 \\ 3 & 3 \end{vmatrix} & \begin{vmatrix} 8 & 2 \\ 3 & 3 \end{vmatrix} & \begin{vmatrix} 6 & 2 \\ 3 & 3 \end{vmatrix} & -\begin{vmatrix} 6 & 4 \\ 3 & 3 \end{vmatrix}$$
$$\begin{vmatrix} 4 & 2 \\ 2 & 6 \end{vmatrix} & -\begin{vmatrix} 6 & 2 \\ 8 & 6 \end{vmatrix} & \begin{vmatrix} 6 & 4 \\ 8 & 2 \end{vmatrix}$$
$$\operatorname{Adj} A = \begin{bmatrix} -12 & -6 & 18 \\ -6 & 12 & -6 \\ 20 & -20 & -20 \end{bmatrix}$$
$$\therefore A^{-1} = \frac{1}{|A|} \operatorname{adj} A$$
$$= \frac{1}{-60} \begin{bmatrix} -12 & -6 & 20 \\ -6 & 12 & -20 \\ 18 & -6 & -20 \end{bmatrix}$$
$$\therefore X = A^{-1}B$$
$$= \frac{1}{-60} \begin{bmatrix} -12 & -6 & 20 \\ -6 & 12 & -20 \\ 18 & -6 & -20 \end{bmatrix} \times \begin{bmatrix} 20 \\ 30 \\ 18 \end{bmatrix}$$
$$= \frac{1}{-60} \begin{bmatrix} -12 \times 20 - 6 \times 30 + 20 \times 18 \\ -6 \times 20 + 12 \times 30 - 20 \times 18 \\ 18 \times 20 - 6 \times 30 - 20 \times 18 \end{bmatrix}$$
$$= \frac{1}{-60} \begin{bmatrix} -60 \\ -120 \\ -180 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

Hence, the solution of the given system of equation is

x = 1, y = 2, z = 3.

OR

b) Solution

 $\text{Given A} = \begin{bmatrix} 5 & 15 & -10 \\ -15 & 0 & -5 \\ 10 & 5 & 0 \end{bmatrix}$ Then A = AI_3 $\Rightarrow \begin{bmatrix} 5 & 15 & -10 \\ -15 & 0 & -5 \\ 10 & 5 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A$ $\Rightarrow 5 \begin{bmatrix} 1 & 3 & -2 \\ -3 & 0 & -1 \\ 2 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A$ $R_2 \rightarrow R_2 + 3R_1; R_3 \rightarrow R_3 - 2R_1$ $\Rightarrow 5 \begin{bmatrix} 1 & 3 & -2 \\ 0 & 9 & -7 \\ 0 & -5 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix} A$ $R_2 \rightarrow R_2 + 2R_3$ $\Rightarrow 5 \begin{bmatrix} 1 & 3 & -2 \\ 0 & -1 & 1 \\ 0 & -5 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 2 \\ -2 & 0 & 1 \end{bmatrix} A$ $R_2 \rightarrow -R_2$ $\Rightarrow 5 \begin{bmatrix} 1 & 3 & -2 \\ 0 & 1 & -1 \\ 0 & -5 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & -2 \\ -2 & 0 & 1 \end{bmatrix} A$ $R_3 \rightarrow R_3 + 5R_2$ $\Rightarrow 5 \begin{bmatrix} 1 & 3 & -2 \\ 0 & 1 & -1 \\ 0 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & -2 \\ 3 & -5 & -9 \end{bmatrix} A$ $R_3 \rightarrow -R_3$

$$\Rightarrow 5 \begin{bmatrix} 1 & 3 & -2 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & -2 \\ -3 & 5 & 9 \end{bmatrix} A$$

$$R_1 \rightarrow R_1 - 3R_2$$

$$\Rightarrow 5 \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -2 & 3 & 6 \\ 1 & -1 & -2 \\ -3 & 5 & 9 \end{bmatrix} A$$

$$R_1 \rightarrow R_1 - R_3 ; R_2 \rightarrow R_2 + R_3$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 1 & -2 & -3 \\ -2 & 4 & 7 \\ -3 & 5 & 9 \end{bmatrix} A$$

$$\Rightarrow I_3 = BA, \text{ where } B = \frac{1}{5} \begin{bmatrix} 1 & -2 & -3 \\ -2 & 4 & 7 \\ -3 & 5 & 9 \end{bmatrix} A$$

$$Hence, A^{-1} = B = \frac{1}{5} \begin{bmatrix} 1 & -2 & -3 \\ -2 & 4 & 7 \\ -3 & 5 & 9 \end{bmatrix}$$

43. a) Solution

Let
$$y = \frac{5x^3}{1-9x^3}$$
 and $z = 15x^3 + 81$, so that $\frac{dy}{dz}$ is wanted.
 $y = \frac{5x^3}{1-9x^3}$

Differentiating both w.r.t. 'x', we get

$$\frac{dy}{dz} = \frac{(1-9x^8)(45x^3) - (15x^8 + 81)(-27x^3)}{(1-9x^8)^2}$$
$$= \frac{45x^3 - 405x^5 + 405x^5 + 1701x^3}{(1-9x^8)^2}$$
$$= \frac{-1746x^3}{(1-9x^8)^2}$$

 $z = 15x^3 + 81$

Differentiating both w.r.t. 'x', we get

$$\frac{dz}{dx} = 45x^2$$

$$\frac{dy}{dx} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}}$$
Provided $\frac{dz}{dx} \neq 0$ i.e., $x \neq 0$

$$= \frac{-1746x^2}{(1-9x^3)^2} \times \frac{1}{45x^2}$$
$$= \frac{-38.8}{(1-9x^3)^2}$$

OR

b) Solution

Given $(u - \sqrt{uv})dv = v du$

$$\Rightarrow \frac{dv}{du} = \frac{v}{u - \sqrt{uv}} \tag{1}$$

Dividing numerator and denominator of R.H.S. of (1) by 'x', we get

$$\frac{dv}{du} = \frac{\frac{v}{u}}{1 - \sqrt{\frac{v}{u}}}, \text{ which is of the form } \frac{dv}{du} = f\left(\frac{v}{u}\right)$$

Therefore, (1) is a homogeneous differential equation

Put
$$v = au \Rightarrow \frac{dv}{du} = a.1 + v.\frac{da}{du}$$

Substituting these values of v and $\frac{dv}{du}$ in (1), we get

$$a + u \frac{aa}{du} = \frac{a}{1 - \sqrt{a}}$$

$$\Rightarrow u \frac{da}{du} = \frac{a}{1 - \sqrt{a}} - a$$

$$= \frac{a^{3/2}}{1 - \sqrt{a}}$$

$$\Rightarrow \frac{1 - \sqrt{a}}{a^{3/2}} da = \frac{1}{u} du$$

$$\Rightarrow \left(a^{-\frac{3}{2}} - \frac{1}{a}\right) da = \frac{1}{u} du$$
Integrating both the side

Integrating both the sides, we get

$$\Rightarrow \frac{a^{-\frac{1}{2}}}{-\frac{1}{2}} - \log|a| = \log|u| + C$$
$$\Rightarrow \frac{-2}{\sqrt{a}} - \log|au| = -C$$
$$\Rightarrow 2\sqrt{\frac{u}{v}} + \log|v| = -C = A \qquad (Say)$$

Hence,

$$2\sqrt{\frac{u}{v}} + \log|v| = A$$
, A is the arbitrary constant.

44. Solution

Let
$$I = \int_{0}^{2} \frac{2x+5}{3x^{2}+16} dx + 5 \int_{0}^{2} \frac{dx}{3x^{2}+16} dx$$

 $= \int_{0}^{2} \frac{2x}{3x^{2}+16} dx + 5 \int_{0}^{2} \frac{dx}{3x^{2}+16} dx$
For $I_{1} = \int_{0}^{2} \frac{2x}{3x^{2}+16} dx$
Put $3x^{2} + 16 = t$
 $6 \times dx = dt$
 $2 \times dx = \frac{1}{3} dt$
When $x = 2$, $t = 3(2)^{2} + 16 = 28$
When $x = 0$, $t = 3(0)^{2} + 16 = 16$
 $\therefore I_{1} = \frac{1}{3} \int_{16}^{28} \frac{dt}{t}$
 $= \frac{1}{3} [\log|t|]_{16}^{28}$
 $= \frac{1}{3} [\log|t|]_{16}^{28}$
 $= \frac{1}{3} \log|\frac{28}{16}|$
 $= \frac{1}{3} \log|\frac{28}{16}|$
 $= \int_{0}^{2} \frac{dx}{(\sqrt{3x})^{2} - (4)^{2}}$
 $= \left[\frac{1}{4} \frac{tan^{-2}(\frac{\sqrt{3x}}{\sqrt{3}})}{\sqrt{3}}\right]_{0}^{2} \left[\int_{x^{2} + a^{2}} dx = \frac{1}{a} tan^{-1}(\frac{x}{a}) + C\right]$

$$= \left[\frac{1}{4\sqrt{3}} \tan^{-1}\left(\frac{\sqrt{3}x}{4}\right)\right]_{0}^{2}$$

$$= \frac{1}{4\sqrt{3}} \tan^{-1}\left(\frac{2\sqrt{3}}{4}\right) - \frac{1}{4\sqrt{3}} \tan^{-1}0$$

$$= \frac{1}{4\sqrt{3}} \tan^{-1}\frac{\sqrt{3}}{2}$$

$$\therefore I = I_{1} + 3I_{2}$$

$$= \frac{1}{3} \log\left|\frac{7}{4}\right| + \frac{3}{4\sqrt{3}} \tan^{-1}\frac{\sqrt{3}}{2}$$

$$= \frac{1}{3} \log\left|\frac{7}{4}\right| + \frac{\sqrt{3}}{4} \tan^{-1}\frac{\sqrt{3}}{2}$$

2

Let I =
$$\int \frac{dx}{x^4 + 81}$$

= $\int \frac{1}{18} \cdot \frac{(x^2 + 9) - (x^2 - 9)}{x^4 + 81} dx$
= $\frac{1}{18} \left[\int \frac{x^2 + 9}{x^4 + 81} dx - \int \frac{x^2 - 9}{x^4 + 81} dx \right]$

(Dividing numerator and denominator by x^2)

$$\begin{split} &= \frac{1}{18} \int \frac{1+\frac{9}{x^2}}{x^2 + \frac{81}{x^2}} dx - \frac{1}{18} \int \frac{1-\frac{9}{x^2}}{x^2 + \frac{81}{x^2}} dx \\ &= \frac{1}{18} I_1 - \frac{1}{18} I_2 \\ &I_1 = \int \frac{1+\frac{9}{x^2}}{x^2 + \frac{81}{x^2}} dx \\ &\left(put \ x - \frac{9}{x} = t \Rightarrow \left(1 + \frac{9}{x^2}\right) dx = dt \ and \ x^2 + \frac{81}{x^2} = t^2 + 18\right) \\ &= \int \frac{dt}{t^2 + 18} = \int \frac{dt}{t^2 + (\sqrt{18})^2} \\ &= \frac{1}{\sqrt{18}} \tan^{-1} \frac{t}{\sqrt{18}} + c_1 \left[\int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a}\right) + C \right] \\ &= \frac{1}{\sqrt{18}} \tan^{-1} \left(\frac{x-\frac{9}{x}}{\sqrt{18}}\right) + C_1 \\ &= \frac{1}{\sqrt{18}} \tan^{-1} \left(\frac{x^2 - 9}{\sqrt{18}}\right) + C_1 \end{split}$$

$$\begin{split} I_2 &= \int \frac{1 - \frac{9}{x^2}}{x^2 + \frac{91}{x^2}} dx \\ \left(Put \ x + \frac{9}{x} = u \Rightarrow \left(1 - \frac{9}{x^2}\right) dx = du \ and \ x^2 + \frac{81}{x^2} = u^2 - 18 \right) \\ &= \int \frac{du}{u^2 - 18} = \int \frac{du}{u^2 - (\sqrt{18})^2} \\ &= \frac{1}{2\sqrt{18}} \log \left| \frac{u - \sqrt{18}}{u + \sqrt{18}} \right| + C_2 \left[\int \frac{dx}{x^2 - \alpha^2} = \frac{1}{2\alpha} \log \left| \frac{x - \alpha}{x + \alpha} \right| + C \right] \\ &= \frac{1}{2\sqrt{18}} \log \left| \frac{x + \frac{9}{x} - \sqrt{18}}{x + \frac{9}{x} + \sqrt{18}} \right| + C_2 \\ &= \frac{1}{2\sqrt{18}} \log \left| \frac{x^2 + 9 - \sqrt{18}x}{x^2 + 9 + \sqrt{18}x} \right| + C_2 \\ &\therefore I = \frac{1}{18} \left[I_1 - I_2 \right] \\ &= \frac{1}{18} \left[\frac{1}{\sqrt{18}} tan^{-1} \left(\frac{x^2 - 9}{\sqrt{18}x} \right) - \frac{1}{2\sqrt{18}} \log \left| \frac{x^2 + 9 - \sqrt{18}x}{x^2 + 9 + \sqrt{18}x} \right| \right] + C \\ &= \frac{1}{18\sqrt{18}} \left[tan^{-1} \left(\frac{x^2 - 9}{\sqrt{18}x} \right) - \frac{1}{2} \log \left| \frac{x^2 + 9 - \sqrt{18}x}{x^2 + 9 + \sqrt{18}x} \right| \right] + C \end{split}$$

Let
$$I = \int \frac{ax}{(4-x)(x^2+6)}$$

Let $\frac{1}{(4-x)(x^2+6)} = \frac{A}{4-x} + \frac{Bx+C}{(x^2+6)}$
 $\Rightarrow 1 = A(x^2+6) + (Bx+C)(4-x)$
 $\Rightarrow 1 = A(x^2+6) + B(4x-x^2) + C(4-x)$

Equating coefficients of x^2 , x and constant terms, we get

$$\begin{split} A &= \frac{1}{22}, B = \frac{1}{22}, C = \frac{2}{5} \\ \therefore I &= \int \frac{1}{22} \cdot \frac{1}{4-x} + \frac{\frac{1}{22}x + \frac{2}{5}}{(x^2+6)} dx \\ &= -\frac{1}{22} \log|4-x| + \frac{1}{22} \int \frac{x}{x^2+6} dx + \frac{2}{5} \int \frac{1}{(x^2+6)} dx \left[\int \frac{f'(x)}{f(x)} dx \text{ form} \right] \\ &= -\frac{1}{22} \log|4-x| + \frac{1}{44} \int \frac{2x}{x^2+6} dx + \frac{2}{5} \cdot \frac{1}{\sqrt{6}} \tan^{-1} \left(\frac{x}{\sqrt{6}} \right) \left[\int \frac{1}{x^2+a^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C \right] \end{split}$$

$$= -\frac{1}{22}\log|4 - x| + \frac{1}{44}\log|x^2 + 6| + \frac{2}{5\sqrt{6}}tan^{-1}\left(\frac{x}{\sqrt{6}}\right) + C$$

Let
$$I = \int \frac{dx}{6+4\sin x + \cos x}$$

$$= \int \frac{dx}{6+4} \frac{dx}{1+\tan^{2}x} + \frac{1-\tan^{2}x}{1+\tan^{2}x} + \frac{1-\tan^{2}x}{1+\tan^{2}x}}$$

$$= \int \frac{1+\tan^{2}x}{6+6\tan^{2}x} + 8\tan^{2}x + 1-\tan^{2}x} dx$$

$$= \int \frac{\sec^{2}x}{5\tan^{2}x} + 8\tan^{2}x + 1-\tan^{2}x} dx$$
Let $\tan^{2}x = 1 \Rightarrow \sec^{2}x \cdot \frac{1}{2} dx = dt \Rightarrow \sec^{2}x dx = 2dt$

$$\therefore I = \int \frac{2dt}{5t^{2}+8t+7}$$

$$= \int \frac{2dt}{5t(t+\frac{4}{5})^{2} - (\frac{\sqrt{19}}{5})^{2}}$$

$$= \frac{2}{5} \cdot \frac{1}{2x \sqrt{\frac{19}{5}}} \log \left| \frac{t+\frac{4}{5} + \frac{\sqrt{19}}{1+\frac{4}{5} + \frac{\sqrt{19}}{5}} \right| + C \left[\int \frac{dx}{x^{2} - a^{2}} = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + C \right]$$

$$= \frac{1}{\sqrt{19}} \log \left| \frac{5t+4-\sqrt{19}}{5t+4+\sqrt{19}} \right| + C$$