




CBSE Class 10 Science Solution PDF


SET 31 / 5 / 3

Q.No.	Value point/ Expected answer	Value	Total marks
1.	Section-A		
	$R = \frac{V}{I}$ $= \frac{5}{0.2}$ $= 2.5 \Omega$	1/2	1
2.	<ul style="list-style-type: none"> Storage and disposal of spent or used fuel. 	1/2	1
3.	Section-B		
	 <ul style="list-style-type: none"> Correct ray diagram labelling of $\angle i$ and $\angle r$ 	1 1/2 x 2 = 1	2
4.	<p>Principle : When current carrying coil is placed in a magnetic field ,it experiences force as per Fleming's left hand rule.</p> <p>Fleming's left hand rule: According to Fleming's left hand rule, stretch the thumb, forefinger and middle finger so that they are mutually perpendicular. If the first finger points in the direction of magnetic field and the second finger in the direction of current, then the thumb will point the direction of force acting on the conductor.</p>  <p style="text-align: center;">atement or for the diagram)</p>	1 1	2
5.	<ul style="list-style-type: none"> Formic acid / Methanoic acid HCOOH Putting a solution of weak /mild base like baking soda. 	1/2 1/2 1	




	Or		
	<p>(a)</p> <ul style="list-style-type: none"> • Acid + Phenolphthalein----->Colourless Base+ Phenolphthalein-----> Pink colour <p>(b)</p> <ul style="list-style-type: none"> • Acid + Methyl Orange -----> Red Base + Methyl Orange -----> Yellow 	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>	2
6.	<p style="text-align: center;">Section-C</p> <p>The series of living organisms taking part at various biotic level forms a food chain.</p> <p>(i) An average of 10% of the food eaten is turned into its own body and made available for the next level of consumers</p> <p>(ii) The energy that is captured by the autotrophs does not revert back to the solar input.</p> <p>(iii) The energy which is passed to the herbivores does not come back to the autotrophs.</p> <p>(iv) As it moves progressively through the various trophic levels it is no longer available to the previous level</p> <p style="text-align: right;">[Any two]</p> <p style="text-align: center;">or</p> <p>(a) Since interference will create disturbances in the protected area (National Park) / To maintain the selfsustainability in the protected area</p> <p>(v) Reuse of materials is better than recycling because the process of recycling use some energy, in the reuse strategy things are used of again and again</p>	<p>1</p> <p>1+1</p> <p>1</p> <p>1+1</p>	3
7.	<p>Dam is a barrier built across a river or a stream for storage of water.</p> <p>Advantage:</p> <ol style="list-style-type: none"> i. Storage of adequate water for irrigation. ii. Generation of electricity <p>Ill effects :</p> <ol style="list-style-type: none"> i. Social problems ii. Economic problems iii. Environmental problems <p>(any two ill effects)</p>	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2} \times 2 = 1$</p>	3
8.	<ul style="list-style-type: none"> • Xylem vessels and Xylem tracheids • At the roots, cells in contact with the soil actively take up ions. } • Creates a difference in concentration of ions • So water moves up. } <p>(b) Since plants do not move and have large proportion of dead cells in many tissues Thus plants have low energy needs.</p>	<p>1</p> <p>1</p> <p>1</p>	3

	$\text{ZnO} + \text{C} \longrightarrow \text{Zn} + \text{CO}$	$\frac{1}{2}$	3
14.	$\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$ $\frac{1}{v} = \frac{1}{f} - \frac{1}{u}$ $\frac{1}{u} = \frac{1}{20} + \frac{1}{30}$ <p>$u = -60\text{cm}$ (without unit (cm) deduct $\frac{1}{2}$ mark)</p> $m = \frac{-v}{u}$ $m = -\left[\frac{-30}{-60}\right]$ $m = -\frac{1}{2}$ $\frac{h_2}{h_1} = m$ $h_2 = h_1 \times m$ $h_2 = 4 \times \left[\frac{-1}{2}\right]$ $h_2 = -2 \text{ cm}$ <p style="text-align: center;">or</p> $m = \frac{v}{u}$ $m = -\frac{2}{3}$ $-\frac{2}{3} = \frac{v}{-12}$ $v = 8 \text{ cm}$ $\frac{1}{f} = \frac{1}{v} - \frac{1}{u}$ $\frac{1}{f} = \frac{1}{8} - \frac{1}{-12}$ $\frac{1}{f} = \frac{1}{8} + \frac{1}{12}$ $\frac{1}{f} = \frac{5}{24}$	$\frac{1}{2}$	
		1	
		$\frac{1}{2}$	
		1	
		$\frac{1}{2}$	
		1	
		$\frac{1}{2}$	
		$\frac{1}{2}$	

	$f = \frac{24}{5}$ $= 4.8 \text{ cm}$		1/2	3								
15.	<ul style="list-style-type: none"> Closeness (crowding) of magnetic field lines is directly related to the strength of the magnetic field. Strength of magnetic field at point 'A' (Pole) is more than at point 'B'. If the student redraws the diagram and mark the arrows correctly (N to S). 		1 1 1	3								
16.	<p style="text-align: center;">Section –D</p> <table border="1" style="width: 100%;"> <thead> <tr> <th style="width: 5%;">(a)</th> <th style="width: 40%;">Cross Pollination</th> <th style="width: 5%;"></th> <th style="width: 50%;">Self Pollination</th> </tr> </thead> <tbody> <tr> <td>1.</td> <td>Pollen is transferred from anther/stamen of one flower to another flower.</td> <td>1.</td> <td>Transfer of pollen from anther/stamen to the stigma of the same flower.</td> </tr> </tbody> </table> <ul style="list-style-type: none"> Site of fertilization – Ovary Product of fertilization – Zygote <p>(b)</p>  <p style="text-align: right;">Correct diagram 1 Correct labelling 1/2 x 4</p>	(a)	Cross Pollination		Self Pollination	1.	Pollen is transferred from anther/stamen of one flower to another flower.	1.	Transfer of pollen from anther/stamen to the stigma of the same flower.		1 1/2 1/2	
(a)	Cross Pollination		Self Pollination									
1.	Pollen is transferred from anther/stamen of one flower to another flower.	1.	Transfer of pollen from anther/stamen to the stigma of the same flower.									

	<p style="text-align: center;">OR</p> <p>(a) </p> <p style="text-align: right;">Correct diagram</p> <p style="text-align: right;">i. Ovary</p> <p style="text-align: right;">ii. Oviduct or fallopian tube</p> <p>(b) Syphilis and Gonorrhoea</p> <p>(c) Chemicals or materials required to avoid pregnancy</p> <p>(i) Controlling human population</p> <p>(ii) To maintain good reproductive health</p> <p>(iii) Maintain gaps between successive birth</p> <p style="text-align: right;">} any two</p>	<p style="text-align: center;">1</p> <p style="text-align: center;">$\frac{1}{2} + \frac{1}{2}$</p> <p style="text-align: center;">1</p> <p style="text-align: center;">1</p> <p style="text-align: center;">$\frac{1}{2} \times 2$</p> <p style="text-align: center;">5</p>	
17.	<p>(a) When the gametes from male and female parent combine during sexual reproduction to form zygote, they contribute equal amount of DNA (half each). The normal body cells of human contain 46 chromosomes each. Human sperm cells and egg cells both have 23 chromosomes. So the combination of these 23 chromosomes from male and female each during sexual reproduction ensures equal genetic contribution to progeny ($23+23 = 46$).</p> <p>(b)</p> <ul style="list-style-type: none"> • Animals have a vast diversity in structures, they probably do not have a common ancestry because common ancestry may greatly limit the extent of diversity. • Many of these diverse animals are inhabiting the same habitat, the evolution by geographical isolation and speciation is not likely to happen. 	<p style="text-align: center;">3</p> <p style="text-align: center;">1</p> <p style="text-align: center;">1</p> <p style="text-align: center;">5</p>	
18.	<p>$f = 20 \text{ cm}, u = -30 \text{ cm}$</p> <p>(a)</p>		

	OR		
	<p>(i) $P = 40 \text{ W}$ $V = 220 \text{ V}$ $P = VI$ $I = \frac{P}{V} = \frac{40 \text{ W}}{220 \text{ V}}$ $= 0.18 \text{ A}$</p> <p>(ii) $R = \frac{V^2}{P}$ $= \frac{220 \times 220}{40}$ $= 1210 \Omega$</p> <p>(iii) $P = 25 \text{ W}$ $V = 220 \text{ V}$ $P = VI$ $I = \frac{P}{V}$ $= \frac{25}{220} = 0.113 \text{ A}$</p> <p>(iv) $R = V^2/P$ $= \frac{220 \times 220}{25}$ $= 1936 \Omega$</p> <p>(v) Yes there is a change in current and resistance</p>	1 1 1 1 1	5
20.	<p>(a) Element with smallest atomic radius ----- Fluorine /F</p> <p>(b) Element with maximum valency----- Carbon /C</p> <p>(c) Element which is metalloid ----- Boron /B</p> <p>(d) Element which is most electropositive ----- Lithium /Li</p> <p>(e) CO and CO₂</p>	1 1 1 1 1	5
21.	<p>(a)</p> <ul style="list-style-type: none"> • Carbon cannot form C⁴⁺ ions as very high energy is required to remove 4 electrons • Carbon cannot gain 4 electrons to form C⁴⁻ ions as 6 protons cannot hold 10 electrons <p>(i) Co-valent compounds are bad conductor of electricity as they do not have free electrons.</p> <p>(ii) Due to weak forces of attraction between the molecules, thus less energy is required for breaking the bonds</p>	1 1 1 1	

	<p>(b)</p>  <p style="text-align: center;">Or</p> <p>(a) Isomers are those compounds which have the same molecular formula but different structural formula.</p> <p>(b)</p> <ul style="list-style-type: none"> • Propanal ----- $\text{CH}_3\text{CH}_2\text{CHO}$ • Propanone ----- CH_3COCH_3 <p>(c) (i) $\text{CH}_3\text{CH}_2\text{OH} \xrightarrow[\text{Conc. H}_2\text{SO}_4]{443 \text{ K}} \text{H}_2\text{C}=\text{CH}_2 + \text{H}_2\text{O}$</p> <p>(ii) $\text{CH}_3\text{CH}_2\text{OH} \xrightarrow[\text{Heat}]{\text{Alkaline KMnO}_4} \text{CH}_3\text{CH}_2\text{COOH} + \text{H}_2\text{O}$</p>	1	
22.	<p style="text-align: center;">Section –E</p> <p>(c) (20 cm, 20 cm) and (inverted and inverted)</p> <p>Reason: Only real and inverted image can be obtained on the screen and in both cases image is formed at the principal focus.</p>	1	
23.	<p>38 mA, 3.2 V</p> <p style="text-align: center;">Or</p> <p>(i) $V \propto I$</p> <p>(ii) at 2.5 V , Current will be 0.25 A</p>	1+1	
24.	 <p style="text-align: center;">OR</p>  <p style="text-align: right;">Correct Diagram and Labelling</p>	$\frac{1}{2} + 1\frac{1}{2}$	
	Diagram Process – Budding	1 1	2
25.	<ul style="list-style-type: none"> • Safranin is used to stain/colour the material for better view. 	1	

	<ul style="list-style-type: none"> Glycerine prevents the leaf peel from getting it dried. 	1	2
26.	<p>The solution turns</p> <ol style="list-style-type: none"> green to colourless black coating is formed on Zinc. <p>Reason: Zinc is more reactive than iron so it displaces the iron from its salt solution.</p>	$\frac{1}{2} + \frac{1}{2}$	
27.	<ul style="list-style-type: none"> No change/As acid turns blue litmus to red, so there is a need of blue litmus paper. To get the blue litmus dip the red litmus paper into a basic solution and get blue colour. <p style="text-align: center;">OR</p> <p>(i) Sodium hydrogen carbonate (NaHCO_3) or Sodium Carbonate (Na_2CO_3)</p> <p>(ii) $2\text{CH}_3\text{COOH} + \text{Na}_2\text{CO}_3 \rightarrow 2\text{CH}_3\text{COONa} + \text{H}_2\text{O} + \text{CO}_2$</p> <p style="text-align: center;">or</p> <p>$\text{CH}_3\text{COOH} + \text{NaHCO}_3 \rightarrow \text{CH}_3\text{COONa} + \text{H}_2\text{O} + \text{CO}_2$</p> <p>(iii) Liberated CO_2 is passed through lime water, which is turned to milky.</p>	2	
		$\frac{1}{2}$	
		1	
		$\frac{1}{2}$	2