

QUESTION PAPER CODE 65/1

EXPECTED ANSWER/VALUE POINTS

SECTION A

$$1. \quad \frac{\pi}{3} - \left(\pi - \frac{\pi}{6} \right) = -\frac{\pi}{2} \quad \frac{1}{2} + \frac{1}{2}$$

Note: $\frac{1}{2}$ m. for any one of the two correct values and $\frac{1}{2}$ m. for final answer

$$2. \quad a = -2, b = 3 \quad \frac{1}{2} + \frac{1}{2}$$

$$3. \quad |\vec{a}| = |\vec{b}| = 3 \quad \frac{1}{2} + \frac{1}{2}$$

$$4. \quad 5 \circ 10 = (5 * 10) + 3 = 10 + 3 = 13 \quad \text{For } 5 * 10 = 10 \quad \frac{1}{2}$$

$$\text{For Final Answer} = 13 \quad \frac{1}{2}$$

SECTION B

$$5. \quad \text{In RHS, put } x = \sin \theta \quad \frac{1}{2}$$

$$\begin{aligned} \text{RHS} &= \sin^{-1} (3 \sin \theta - 4 \sin^3 \theta) \\ &= \sin^{-1} (\sin 3\theta) \end{aligned} \quad 1$$

$$= 3\theta = 3 \sin^{-1} x = \text{LHS.} \quad \frac{1}{2}$$

$$6. \quad |A| = 2, \quad \therefore A^{-1} = \frac{1}{2} \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix} \quad 1$$

$$\text{LHS} = 2A^{-1} = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}, \quad \text{RHS} = 9 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix} = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix} \quad 1$$

$$\therefore \text{LHS} = \text{RHS}$$

$$7. \quad f(x) = \tan^{-1}\left(\frac{1+\cos x}{\sin x}\right) = \tan^{-1}\left(\frac{2\cos^2 x/2}{2\sin x/2 \cos x/2}\right) \quad 1$$

$$= \tan^{-1}\left(\cot x/2\right) = \frac{\pi}{2} - \frac{x}{2} \quad \frac{1}{2}$$

$$\therefore f'(x) = -\frac{1}{2} \quad \frac{1}{2}$$

$$8. \quad \text{Marginal cost} = C'(x) = 0.015x^2 - 0.04x + 30 \quad 1$$

$$\text{At } x = 3, C'(3) = 30.015 \quad 1$$

$$9. \quad I = \int \frac{1 - 2\sin^2 x + 2\sin^2 x}{\cos^2 x} dx \quad \frac{1}{2}$$

$$= \int \sec^2 x dx \quad 1$$

$$= \tan x + C \quad \frac{1}{2}$$

$$10. \quad \frac{dy}{dx} = bae^{bx+5} \Rightarrow \frac{dy}{dx} = by \quad 1$$

$$\Rightarrow \frac{d^2 y}{dx^2} = b \frac{dy}{dx} \quad \frac{1}{2}$$

$$\therefore \text{The differential equation is: } y \frac{d^2 y}{dx^2} = \left(\frac{dy}{dx}\right)^2 \quad \frac{1}{2}$$

$$11. \quad \sin \theta = \frac{|(\hat{i} - 2\hat{j} + 3\hat{k}) \times (3\hat{i} - 2\hat{j} + \hat{k})|}{|\hat{i} - 2\hat{j} + 3\hat{k}| |3\hat{i} - 2\hat{j} + \hat{k}|} \quad \frac{1}{2}$$

$$|(\hat{i} - 2\hat{j} + 3\hat{k}) \times (3\hat{i} - 2\hat{j} + \hat{k})| = |4\hat{i} + 8\hat{j} + 4\hat{k}| = 4\sqrt{6} \quad 1$$

$$\sin \theta = \frac{4\sqrt{6}}{14} = \frac{2\sqrt{6}}{7} \quad \frac{1}{2}$$

$$12. \quad A: \text{Getting a sum of 8, } B: \text{Red die resulted in no. } < 4$$

$$P(A/B) = \frac{P(A \cap B)}{P(B)} \quad 1$$

$$= \frac{2/36}{18/36} = \frac{1}{9} \quad 1$$

SECTION C

13.
$$\text{LHS} = \begin{vmatrix} 1 & 1 & 1+3x \\ 1+3y & 1 & 1 \\ 1 & 1+3z & 1 \end{vmatrix}$$

$$= \begin{vmatrix} 1 & 0 & 3x \\ 1+3y & -3y & -3y \\ 1 & 3z & 0 \end{vmatrix} \quad (\text{Using } C_2 \rightarrow C_2 - C_1 \text{ \& } C_3 \rightarrow C_3 - C_1)$$

1+1
(Any two relevant operations)

$$= 1 \times (9yz) + 3x(3z + 9yz + 3y) \quad (\text{Expanding along } R_1)$$

1

$$= 9(3xyz + xy + yz + zx) = \text{RHS}$$

1

14. Differentiating with respect to 'x'

$$2(x^2 + y^2) \left(2x + 2y \frac{dy}{dx} \right) = x \frac{dy}{dx} + y$$

2

$$\Rightarrow \frac{dy}{dx} = \frac{y - 4x^3 - 4xy^2}{4x^2y + 4y^3 - x}$$

2

OR

$$\frac{dx}{d\theta} = a(2 - 2\cos 2\theta) = 4a \sin^2 \theta$$

1

$$\frac{dy}{d\theta} = 2a \sin 2\theta = 4a \sin \theta \cdot \cos \theta$$

1

$$\therefore \frac{dy}{dx} = \frac{4a \sin \theta \cos \theta}{4a \sin^2 \theta} = \cot \theta$$

1

$$\left. \frac{dy}{dx} \right|_{\theta = \frac{\pi}{3}} = \frac{1}{\sqrt{3}}$$

1

15. $y = \sin(\sin x) \Rightarrow \frac{dy}{dx} = \cos(\sin x) \cdot \cos x$

1

and $\frac{d^2y}{dx^2} = -\sin(\sin x) \cdot \cos^2 x - \sin x \cos(\sin x)$

1+1

$$\text{LHS} = -\sin(\sin x) \cos^2 x - \sin x \cos(\sin x) + \frac{\sin x}{\cos x} \cos(\sin x) \cos x + \sin(\sin x) \cos^2 x$$

1

$$= 0 = \text{RHS}$$

16. $x_1 = 2 \Rightarrow y_1 = 3 \quad (\because y_1 > 0)$

 $\frac{1}{2}$

Differentiating the given equation, we get, $\frac{dy}{dx} = \frac{-16x}{9y}$

 $\frac{1}{2}$

Slope of tangent at $(2, 3) = \left. \frac{dy}{dx} \right|_{(2,3)} = -\frac{32}{27}$

 $\frac{1}{2}$

Slope of Normal at $(2, 3) = \frac{27}{32}$

 $\frac{1}{2}$

Equation of tangent: $32x + 27y = 145$

1

Equation of Normal: $27x - 32y = -42$

1

OR

$f'(x) = x^3 - 3x^2 - 10x + 24$

 $\frac{1}{2}$


$= (x - 2)(x - 4)(x + 3)$

1

$f'(x) = 0 \Rightarrow x = -3, 2, 4.$

 $\frac{1}{2}$

sign of $f'(x)$:



$\therefore f(x)$ is strictly increasing on $(-3, 2) \cup (4, \infty)$

1

and $f(x)$ is strictly decreasing on $(-\infty, -3) \cup (2, 4)$

1

17. Let side of base = x and depth of tank = y

$V = x^2y \Rightarrow y = \frac{V}{x^2}, (V = \text{Quantity of water} = \text{constant})$

Cost of material is least when area of sheet used is minimum.

$A(\text{Surface area of tank}) = x^2 + 4xy = x^2 + \frac{4V}{x}$

 $\frac{1}{2} + \frac{1}{2}$

$\frac{dA}{dx} = 2x - \frac{4V}{x^2}, \frac{dA}{dx} = 0 \Rightarrow x^3 = 2V, y = \frac{x^3}{2x^2} = \frac{x}{2}$

 $\frac{1}{2} + \frac{1}{2}$

$\frac{d^2A}{dx^2} = 2 + \frac{8V}{x^3} > 0, \therefore \text{Area is minimum, thus cost is minimum when } y = \frac{x}{2}$

 $\frac{1}{2} + \frac{1}{2}$

Value: Any relevant value.

1

18. Put $\sin x = t \Rightarrow \cos x \, dx = dt$

 $\frac{1}{2}$

Let $I = \int \frac{2 \cos x}{(1 - \sin x)(1 + \sin^2 x)} dx = \int \frac{2}{(1-t)(1+t^2)} dt$

Let $\frac{2}{(1-t)(1+t^2)} = \frac{A}{1-t} + \frac{Bt+C}{1+t^2}$, solving we get

$A = 1, B = 1, C = 1$

 $1 \frac{1}{2}$

$\therefore I = \int \frac{1}{1-t} dt + \frac{1}{2} \int \frac{2t}{1+t^2} + \int \frac{1}{1+t^2} dt$

$= -\log |1-t| + \frac{1}{2} \log |1+t^2| + \tan^{-1} t + C$

 $1 \frac{1}{2}$

$= -\log (1 - \sin x) + \frac{1}{2} \log (1 + \sin^2 x) + \tan^{-1}(\sin x) + C$

 $\frac{1}{2}$

19. Separating the variables, we get:

$\int \frac{\sec^2 y}{\tan y} dy = \int \frac{e^x}{e^x - 2} dx$

 $1 \frac{1}{2}$

$\Rightarrow \log |\tan y| = \log |e^x - 2| + \log C$

 $1 \frac{1}{2}$

$\Rightarrow \tan y = C(e^x - 2), \text{ for } x = 0, y = \pi/4, C = -1$

 $\frac{1}{2}$

$\therefore$ Particular solution is: $\tan y = 2 - e^x$.

 $\frac{1}{2}$

OR

Integrating factor $= e^{\int 2 \tan x dx} = \sec^2 x$

1

$\therefore$ Solution is: $y \cdot \sec^2 x = \int \sin x \cdot \sec^2 x \, dx = \int \sec x \cdot \tan x \, dx$

1

$\Rightarrow y \cdot \sec^2 x = \sec x + C, \text{ for } x = \frac{\pi}{3}, y = 0, \therefore C = -2$

 $1 + \frac{1}{2}$

$\therefore$ Particular solution is: $y \cdot \sec^2 x = \sec x - 2$

 $\frac{1}{2}$

or $y = \cos x - 2 \cos^2 x$

$$20. \quad \vec{d} = \lambda(\vec{c} \times \vec{b}) = \lambda \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & -1 \\ 1 & -4 & 5 \end{vmatrix} \quad 1$$

$$\therefore \vec{d} = \lambda \hat{i} - 16\lambda \hat{j} - 13\lambda \hat{k} \quad 1$$

$$\vec{d} \cdot \vec{a} = 21 \Rightarrow 4\lambda - 80\lambda + 13\lambda = 21 \Rightarrow \lambda = -\frac{1}{3} \quad 1$$

$$\therefore \vec{d} = -\frac{1}{3}\hat{i} + \frac{16}{3}\hat{j} + \frac{13}{3}\hat{k} \quad 1$$

$$21. \quad \text{Here } \vec{a}_1 = 4\hat{i} - \hat{j}, \vec{a}_2 = \hat{i} - \hat{j} + 2\hat{k}, \vec{a}_2 - \vec{a}_1 = -3\hat{i} + 2\hat{k} \quad 1$$

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -3 \\ 2 & 4 & -5 \end{vmatrix} = 2\hat{i} - \hat{j} \quad 1$$

$$\text{Shortest distance} = \frac{|(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|} \quad 1$$

$$= \left| \frac{-6}{\sqrt{5}} \right| = \frac{6}{\sqrt{5}} \quad \text{or} \quad \frac{6\sqrt{5}}{5} \quad 1$$

$$22. \quad \left. \begin{array}{l} E_1 : \text{She gets 1 or 2 on die.} \\ E_2 : \text{She gets 3, 4, 5 or 6 on die.} \\ A : \text{She obtained exactly 1 tail} \end{array} \right\} \quad 1$$

$$\left. \begin{array}{l} P(E_1) = \frac{1}{3}, P(E_2) = \frac{2}{3} \\ P(A/E_1) = \frac{3}{8}, P(A/E_2) = \frac{1}{2} \end{array} \right\} \quad 1$$

$$P(E_2/A) = \frac{P(E_2) \cdot P(A/E_2)}{P(E_1) \cdot P(A/E_1) + P(E_2) \cdot P(A/E_2)} \quad 1$$

$$= \frac{\frac{2}{3} \times \frac{1}{2}}{\frac{1}{3} \times \frac{3}{8} + \frac{2}{3} \times \frac{1}{2}} = \frac{8}{11} \quad 1$$

23. Let X denote the larger of two numbers

X	2	3	4	5
$P(X)$	1/10	2/10	3/10	4/10
$X \cdot P(X)$	2/10	6/10	12/10	20/10
$X^2 \cdot P(X)$	4/10	18/10	48/10	100/10

$$\text{Mean} = \sum X \cdot P(X) = \frac{40}{10} = 4$$

$$\text{Variance} = \sum X^2 \cdot P(X) - [\sum X \cdot P(X)]^2 = \frac{170}{10} - 4^2 = 1$$

SECTION D

24. Reflexive: $|a - a| = 0$, which is divisible by 4, $\forall a \in A$

$\therefore (a, a) \in R, \forall a \in A \therefore R$ is reflexive

Symmetric: let $(a, b) \in R$

$\Rightarrow |a - b|$ is divisible by 4

$\Rightarrow |b - a|$ is divisible by 4 ($\because |a - b| = |b - a|$)

$\Rightarrow (b, a) \in R \therefore R$ is symmetric.

Transitive: let $(a, b), (b, c) \in R$

$\Rightarrow |a - b|$ & $|b - c|$ are divisible by 4

$\Rightarrow a - b = \pm 4m, b - c = \pm 4n, m, n \in Z$

Adding we get, $a - c = 4(\pm m \pm n)$

$\Rightarrow (a - c)$ is divisible by 4

$\Rightarrow |a - c|$ is divisible by 4 $\therefore (a, c) \in R$

$\therefore R$ is transitive

Hence R is an equivalence relation in A

set of elements related to 1 is $\{1, 5, 9\}$

and $[2] = \{2, 6, 10\}$.

OR

Here $f(2) = f\left(\frac{1}{2}\right) = \frac{2}{5}$ but $2 \neq \frac{1}{2}$

$\therefore f$ is not 1-1

2

for $y = \frac{1}{\sqrt{2}}$ let $f(x) = \frac{1}{\sqrt{2}} \Rightarrow x^2 - \sqrt{2}x + 1 = 0$

As $D = (-\sqrt{2})^2 - 4(1)(1) < 0$, $\therefore$ No real solution

$\therefore f(x) \neq \frac{1}{\sqrt{2}}$, for any $x \in R(D_f)$ $\therefore f$ is not onto

2

$$\text{fog}(x) = f(2x-1) = \frac{2x-1}{(2x-1)^2+1} = \frac{2x-1}{4x^2-4x+2}$$

2

25. $|A| = -1 \neq 0 \therefore A^{-1}$ exists

1

Co-factors of A are:

$A_{11} = 0$;	$A_{12} = 2$;	$A_{13} = 1$	$\left. \begin{array}{l} \text{1 m for any} \\ \text{4 correct} \\ \text{cofactors} \end{array} \right\}$
$A_{21} = -1$;	$A_{22} = -9$;	$A_{23} = -5$	
$A_{31} = 2$;	$A_{32} = 23$;	$A_{33} = 13$	

2

$$\text{adj}(A) = \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix} \Rightarrow A^{-1} = \frac{1}{|A|} \cdot \text{adj}(A) = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix}$$

 $\frac{1}{2}$

For : $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $B = \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix}$, the system of equation is $A \cdot X = B$

 $\frac{1}{2}$

$$\therefore X = A^{-1} \cdot B = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix} \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

1

$\therefore x = 1, y = 2, z = 3$

1

Using elementary Row operations:

let: $A = IA$

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ -2 & -4 & -5 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A$$

1

$$\Rightarrow \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} A \quad \left\{ \text{Using, } R_2 \rightarrow R_2 - 2R_1; R_3 \rightarrow R_3 + 2R_1 \right.$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 5 & -2 & 0 \\ -2 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} A \quad \left\{ \text{Using, } R_1 \rightarrow R_1 - 2R_2 \right.$$

4

$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -2 & -1 \\ -4 & 1 & -1 \\ 2 & 0 & 1 \end{bmatrix} A \quad \left\{ \text{Using, } R_1 \rightarrow R_1 - R_3; R_2 \rightarrow R_2 - R_3 \right.$$

$$\therefore A^{-1} = \begin{bmatrix} 3 & -2 & -1 \\ -4 & 1 & -1 \\ 2 & 0 & 1 \end{bmatrix}$$

1


26.

Correct figure:

1

Pt. of intersection, $x = 4$

1



$$\text{Area of shaded region} = \int_0^4 x \, dx + \int_4^{4\sqrt{2}} \sqrt{32 - x^2} \, dx$$

1

$$= \left[\frac{x^2}{2} \right]_0^4 + \left\{ \frac{x}{2} \sqrt{32 - x^2} + 16 \sin^{-1} \frac{x}{4\sqrt{2}} \right\} \Bigg|_4^{4\sqrt{2}}$$

2

$$= 8 + 16 \frac{\pi}{2} - 8 - 4\pi = 4\pi$$

1

27. Put $\sin x - \cos x = t$, $(\cos x + \sin x) dx = dt$, $1 - \sin 2x = t^2$ 1

when $x = 0, t = -1$ }
and $x = \pi/4, t = 0$ } $\frac{1}{2}$

$$\therefore I = \int_0^{\pi/4} \frac{\sin x + \cos x}{16 + 9 \sin 2x} dx = \int_{-1}^0 \frac{1}{16 + 9(1-t^2)} dt = \int_{-1}^0 \frac{1}{25 - 9t^2} dt \quad 2$$

$$\Rightarrow I = \frac{1}{30} \log \left| \frac{5+3t}{5-3t} \right| \Big|_{-1}^0 \quad 1 \frac{1}{2}$$

$$= \frac{1}{30} \left[0 - \log \frac{1}{4} \right] = -\frac{1}{30} \log \frac{1}{4} \text{ or } \frac{1}{15} \log 2 \quad 1$$

OR

Here $f(x) = x^2 + 3x + e^x$, $a = 1$, $b = 3$, $nh = 2$ 1

$$\therefore \int_1^3 (x^2 + 3x + e^x) dx = \lim_{h \rightarrow 0} [f(1) + f(1+h) + \dots + f(1+\overline{n-1}h)] \quad 1$$

$$= \lim_{h \rightarrow 0} \left[4(nh) + \frac{(nh-h)(nh)(2nh-h)}{6} + \frac{5(nh-h)(nh)}{2} + \frac{h}{e^h-1} \times e \times (e^{nh}-1) \right] \quad 3$$

$$= 8 + \frac{8}{3} + 10 + e(e^2-1) = \frac{62}{3} + e^3 - e \quad 1$$

28. General point on the line is: $(2 + 3\lambda, -1 + 4\lambda, 2 + 2\lambda)$ $1 \frac{1}{2}$


As the point lies on the plane

$$\therefore 2 + 3\lambda + 1 - 4\lambda + 2 + 2\lambda = 5 \Rightarrow \lambda = 0 \quad 1 \frac{1}{2}$$

$$\therefore \text{Point is } (2, -1, 2) \quad 1$$

$$\text{Distance} = \sqrt{(2-(-1))^2 + (-1-(-5))^2 + (2-(-10))^2} = 13 \quad 2$$

29.

Let number of packets of type A = x and number of packets of type B = y $\therefore$ L.P.P. is: Maximize, $Z = 0.7x + y$

1

subject to constraints:

$$\left. \begin{array}{l} 4x + 6y \leq 240 \quad \text{or} \quad 2x + 3y \leq 120 \\ 6x + 3y \leq 240 \quad \text{or} \quad 2x + y \leq 80 \end{array} \right\}$$

2

$$x \geq 0, y \geq 0$$

Correct graph

2

$$Z(0, 0) = 0, Z(0, 40) = 40$$

$$Z(40, 0) = 28, Z(30, 20) = 41 \text{ (Max.)}$$

 $\therefore$ Max. profit is ₹ 41 at $x = 30, y = 20$.

1