

# CBSE Class 9 Maths Solution of Sample Paper

## Answers & Explanations

### Section A

#### 1. Solution:

We have,  $\frac{81}{36}x^2 - \frac{y^2}{25}$

$$= \left(\frac{9}{6}x\right)^2 - \left(\frac{y}{5}\right)^2$$

$$= \left(\frac{9}{6}x + \frac{y}{5}\right)\left(\frac{9}{6}x - \frac{y}{5}\right) \quad [\because a^2 - b^2 = (a + b)(a - b)]$$

Hence,  $\frac{81}{36}x^2 - \frac{y^2}{25} = \left(\frac{9}{6}x + \frac{y}{5}\right)\left(\frac{9}{6}x - \frac{y}{5}\right)$

#### 2. Solution:

Let  $P(x) = x^2 + 9x - 5 + k$

$$\because x - 1 = 0 \Rightarrow x = 1$$

$$\therefore P(1) = 0$$

$$P(1) = 1^2 + 9 \cdot 1 - 5 + k$$

$$\Rightarrow 1 + 9 - 5 + k = 0$$

$$\Rightarrow k + 5 = 0$$

$$\Rightarrow k = -5$$

Hence,  $k = -5$ .

#### 3. Solution:

We have,  $\frac{9^{\frac{2}{3}}}{9^{\frac{1}{5}}}$

$$= \frac{9^{\frac{2}{3}}}{9^{\frac{1}{5}}} = 9^{\frac{2}{3} - \frac{1}{5}} \quad [\because \frac{a^m}{a^n} = a^{m-n}]$$

$$= 9^{\frac{10-3}{15}}$$

$$= 9^{\frac{7}{15}}$$

$$\text{Hence, } \frac{9^{\frac{2}{3}}}{9^{\frac{1}{5}}} = 9^{\frac{7}{15}}.$$

OR

**Solution**

$$(343)^m = \frac{49}{7^m}$$

$$(343)^m = \frac{7^2}{7^m}$$

$$(7^3)^m = 7^{2-m}$$

As bases are equal, we can equate the powers

$$3m = 2 - m$$

$$4m = 2$$

$$m = \frac{1}{2}$$

**4. Solution:**

Given,  $P(4,6)$  and  $Q(-5,-7)$

$\therefore$  Abscissa of  $P = 4$  and abscissa of  $Q = -5$

$\therefore$  (Abscissa of  $P$ ) - (abscissa of  $Q$ ) =  $4 - (-5) = 4 + 5 = 9$

Hence, (abscissa of  $P$ ) - (abscissa of  $Q$ ) = 9.

**5. Solution:**

Given  $a = 25 \text{ cm}$ ,  $b = 20 \text{ cm}$ ,  $c = 15 \text{ cm}$

$\therefore$  Area of the triangle =  $\sqrt{s(s-a)(s-b)(s-c)}$

To find  $s$ :

$$s = \frac{1}{2}(a + b + c)$$

$$s = \frac{25+20+15}{2} = \frac{60}{2} = 30$$

$\therefore$  Area of the triangle =  $\sqrt{30(30-25)(30-20)(30-15)}$

$$= \sqrt{30 \times 5 \times 10 \times 15} = \sqrt{22500} = 150 \text{ cm}^2$$

Hence, the area of the triangle =  $150 \text{ cm}^2$ .

**OR**

**Solution:**

Given, interval = 100 – 110.

Here, lower limit = 100 and upper limit = 110

$$\therefore \text{Class mark} = \frac{\text{upper limit} + \text{lower limit}}{2}$$

$$= \frac{110 + 100}{2} = \frac{210}{2} = 105$$

Hence, the class mark of the interval 100 – 110 is 105.

**6. Solution:**

Here, length ( $l$ ) = 20 m, breadth ( $b$ ) = 20 m and height ( $h$ ) = 10 m.

$\therefore$  Length of the diagonal in the cuboid = Length of the longest rod

$$= \sqrt{l^2 + b^2 + h^2}$$

$$= \sqrt{20^2 + 20^2 + 10^2}$$

$$= \sqrt{400 + 400 + 100}$$

$$= \sqrt{900}$$

$$= 30 \text{ m}$$

Hence, the length of the longest rod = 30 m

## **Section B**

**7. Solution:**

$$\text{Let } x = 0.\overline{78}$$

$$\text{Then, } x = 0.7878 \dots\dots (1)$$

Multiplying 100 on both sides in equation (1), we get

$$100x = 78.7878 \dots\dots (2)$$

On subtracting equation (1) from (2), we get

$$100x - x = 78.7878 - 0.7878$$

$$\Rightarrow 99x = 78$$

$$\Rightarrow x = \frac{78}{99} = \frac{26}{33}$$

$$\text{Hence, } 0.\overline{78} = \frac{26}{33}.$$

OR

**Solution:**

$$\text{We have, } x = 9 - 4\sqrt{5}$$

$$\frac{1}{x} = \frac{1}{9 - 4\sqrt{5}} = \frac{1}{9 - 4\sqrt{5}} \times \frac{9 + 4\sqrt{5}}{9 + 4\sqrt{5}}$$

$$= \frac{9 + 4\sqrt{5}}{81 - 80} = 9 + 4\sqrt{5}$$

$$\therefore x + \frac{1}{x} = (9 - 4\sqrt{5}) + (9 + 4\sqrt{5}) = 18$$

$$\text{Hence, } x + \frac{1}{x} = 18.$$

**8. Solution:**

Let  $a, b$  be the equal and unequal sides of the isosceles triangle.

$$\text{Here, } a = 3\sqrt{2} \text{ cm, } b = 8 \text{ cm}$$

$$\therefore \text{Area of an isosceles triangle} = \frac{1}{4} \times b \cdot \sqrt{4a^2 - b^2}$$

$$= \frac{1}{4} \times 8 \cdot \sqrt{4(3\sqrt{2})^2 - 8^2}$$

$$= 2 \cdot \sqrt{72 - 64} = 2 \cdot \sqrt{8}$$

$$= 2 \times 2\sqrt{2} \text{ cm}^2$$

$$= 4\sqrt{2} \text{ cm}^2$$

Hence, area of an isosceles triangle is  $4\sqrt{2} \text{ cm}^2$ .

**9. Solution:**

The first nine prime numbers are 2, 3, 5, 7, 11, 13, 17, 19 and 23.

$$\begin{aligned}\therefore \text{Mean} &= \frac{\text{Sum of observations}}{\text{Number of observations}} \\ &= \frac{2+3+5+7+11+13+17+19+23}{9} \\ &= \frac{100}{9} = 11.11\end{aligned}$$

Hence, the mean of the first nine prime numbers is 11.11.

**10. Solution:**

Let  $P(x)$  be the polynomial  $x^{997} + x^{886} + x^{775} + x^{654} + x^{113} + 1$ . If  $(x+1)$  is a factor of  $P(x)$ , then  $P(x)$  should be divisible by  $(x + 1)$ .

By remainder theorem,

If  $P(x)$  is divisible by  $(x-a)$ , then  $P(a) = 0$

So, in this case we need to prove that  $P(-1) = 0$  to show that  $x+1$  is a factor of  $P(x)$

$$P(-1) = (-1)^{997} + (-1)^{886} + (-1)^{775} + (-1)^{654} + (-1)^{113} + 1$$

A negative number raised to an odd number will result in a negative number

A negative number raised to an even number will result in a positive number

$$P(-1) = -1 + 1 + (-1) + 1 + (-1) + 1 = 0$$

As we have proved that  $P(-1) = 0$ , so

Yes,  $(x+1)$  is a factor of  $x^{997} + x^{886} + x^{775} + x^{654} + x^{113} + 1$

**11. Solution:**

Let  $(x - 20^\circ)$  and  $x^\circ$  be the first and second angle respectively.

We know that sum of supplementary angle is equal to  $180^\circ$ .

$$\therefore (x - 20^\circ) + x = 180^\circ$$

$$\Rightarrow 2x - 20^\circ = 180^\circ$$

$$\Rightarrow 2x = 200^\circ$$

$$\Rightarrow x = \frac{200^\circ}{2} = 100^\circ$$

$$x - 20^\circ = 100^\circ - 20^\circ = 80^\circ$$

$\therefore$  Larger angle =  $100^\circ$  and smaller angle =  $80^\circ$

Hence, the larger angle is  $100^\circ$ .

**OR**

**Solution:**

Let  $3x, 5x, 6x$  and  $10x$  be the angles of a quadrilateral.

$$\therefore 3x + 5x + 6x + 10x = 360^\circ$$

$$\Rightarrow 24x = 360^\circ$$

$$\Rightarrow x = \frac{360^\circ}{24} = 15^\circ$$

$$\therefore \text{Smallest angle} = 3 \times 15^\circ = 45^\circ$$

**12. Solution:**

$$\text{Area of the circle} = 841\pi \text{ cm}^2$$

Let  $r$  be the radius of the circle.

$$\therefore \text{Area of the circle} = \pi r^2$$

$$\Rightarrow \pi r^2 = 841\pi$$

$$\Rightarrow r^2 = 841 \Rightarrow r^2 = 29^2$$

$$\Rightarrow r = 29 \text{ cm}$$

We know that, the length of the longest chord of the circle is diameter.

$$\therefore \text{The length of the longest chord of the circle} = 2r = 2 \times 29 = 58 \text{ cm}$$

## Section C

### 13. Solution:

$$\begin{aligned} \text{We have, } & \frac{\sqrt{11}-1}{\sqrt{11}+1} \\ &= \frac{\sqrt{11}-1}{\sqrt{11}+1} \times \frac{\sqrt{11}-1}{\sqrt{11}-1} \\ &= \frac{(\sqrt{11}-1)^2}{\sqrt{11}^2-1^2} \\ &= \frac{(\sqrt{11})^2+1^2-2\cdot\sqrt{11}\cdot 1}{11-1} \quad [\because (a-b)^2 = a^2 + b^2 - 2ab \text{ and } a^2 - b^2 = (a+b)(a-b)] \\ &= \frac{11+1-2\cdot\sqrt{11}}{10} \\ &= \frac{12-2\cdot\sqrt{11}}{10} \\ &= \frac{12}{10} - \frac{2\cdot\sqrt{11}}{10} \\ &= \frac{6}{5} - \frac{\sqrt{11}}{5} \\ \therefore \frac{6}{5} - \frac{\sqrt{11}}{5} &= a - b\sqrt{11} \\ \Rightarrow a = \frac{6}{5} \text{ and } b\sqrt{11} &= \frac{\sqrt{11}}{5} \\ \text{Hence, } a = \frac{6}{5} \text{ and } b &= \frac{1}{5} \end{aligned}$$

### 14. Solution:

$$\text{We have, } x^4 + \frac{1}{x^4} = 34$$

Adding 2 on both sides, we get

$$\begin{aligned} x^4 + \frac{1}{x^4} + 2 &= 34 + 2 = 36 \\ \Rightarrow (x^2)^2 + \left(\frac{1}{x^2}\right)^2 + 2 \cdot x^2 \cdot \frac{1}{x^2} &= 36 \\ \Rightarrow \left(x^2 + \frac{1}{x^2}\right)^2 &= 6^2 \\ \therefore x^2 + \frac{1}{x^2} &= 6 \dots\dots (1) \end{aligned}$$

Again adding 2 on both sides in (1), we get

$$x^2 + \frac{1}{x^2} + 2 = 6 + 2 = 8$$

$$\Rightarrow \left(x + \frac{1}{x}\right)^2 = \sqrt{8}^2$$

$$\therefore x + \frac{1}{x} = \sqrt{8}$$

$$\text{Hence, } x + \frac{1}{x} = \sqrt{8}$$

**15. Solution:**

Let  $\alpha$  and  $\beta$  be the two complementary angles.

We know that sum of complementary angle is equal to  $90^\circ$ .

$$\therefore \alpha + \beta = 90^\circ$$

$$\therefore \alpha = 90^\circ - \beta \dots (1)$$

According to question,

$$\beta = \frac{1}{3}\alpha \dots (2)$$

Using (1), we get

$$\beta = \frac{1}{3}(90^\circ - \beta)$$

$$\Rightarrow 3\beta = 90^\circ - \beta \Rightarrow 4\beta = 90^\circ$$

$$\Rightarrow \beta = \frac{90^\circ}{4} = 22.5^\circ$$

Putting  $\beta = 22.5^\circ$  in (1), we get

$$\alpha = 90^\circ - 22.5^\circ = 67.5^\circ$$

Hence, larger angle =  $67.5^\circ$

**OR**

**Solution:**

$\angle POR + \angle ROQ = 180^\circ$  (Linear pairs of lines)

Given,  $\angle POR : \angle ROQ = 5 : 15$

Therefore,  $\angle POR = \frac{5}{20} \times 180^\circ = 45^\circ$



Similarly,  $\angle ROQ = \frac{15}{20} \times 180^\circ = 135^\circ$


Now,  $\angle POS = \angle ROQ^\circ = 135^\circ$  (Vertically opposite)

and  $\angle SOQ = \angle POR = 45^\circ$  (Vertically opposite)

**16. Solution:**

Given, area of the  $\triangle BGC$  is 28 square units

G is the centroid.



We know that,

$$\text{Area of the } \triangle BGC = \frac{1}{3} \times \text{area of the } \triangle ABC$$

$$\therefore \text{Area of the } \triangle ABC = 3 \times \text{area of the } \triangle BGC$$

$$= 3 \times 28 \text{ square units}$$

$$= 84 \text{ square units}$$

Hence, area of the  $\triangle ABC = 84$  square units.

**17. Solution:**

Let  $a$  be the side of the equilateral triangle.

$$\therefore \text{Altitude of an equilateral triangle} = \frac{\sqrt{3}}{2} \times a$$

$$= \frac{\sqrt{3}}{2} \times 6\sqrt{3} \text{ cm}$$

$$= \sqrt{3} \times 3\sqrt{3} \text{ cm}$$

$$= 3 \times 3 \text{ cm}$$

$$= 9 \text{ cm}$$

Hence, the altitude of an equilateral triangle = 9 cm.

### 18. Solution:

Volume of sphere = Volume of wire

Let  $r_s, r_w$  be the radius of sphere and wire respectively.

Let  $h$  be the length of the wire.

Given, radius of the sphere ( $r_s$ ) = 7 cm and radius of the wire ( $r_w$ ) = 0.3 cm

$$\therefore \text{Volume of sphere} = \frac{4}{3}\pi r_s^3 = \frac{4}{3}\pi 7^3 \text{ cm}^3$$

$$\text{Volume of wire} = \pi r_w^2 h = \pi (0.3)^2 h$$

According to question,

Volume of sphere = Volume of wire

$$\Rightarrow \frac{4}{3}\pi 7^3 = \pi (0.3)^2 h$$

$$\Rightarrow h = \frac{\frac{4}{3} \times 343}{0.09} = 5081.48 \text{ m}$$

Hence, the length of the wire is 5081.48 m.

### 19. Solution:

Let the length of the first diagonal be  $x$  cm and the second diagonal is  $5x$  cm respectively.

According to question,

$$x + 5x = 180$$

$$\Rightarrow 6x = 180 \Rightarrow x = 30 \text{ cm}$$

$$\therefore 5x = 5 \times 30 = 150 \text{ cm}$$

$$\text{Area of rhombus} = \frac{1}{2} \times \text{first diagonal} \times \text{second diagonal}$$

$$= \frac{1}{2} \times x \times 5x \text{ square units}$$

$$= \frac{1}{2} \times 30 \times 150 \text{ square units}$$

$$= 2250 \text{ cm}^2$$

Hence, area of rhombus =  $2250 \text{ cm}^2$

**OR**

**Solution:**

Let each base angle of isosceles triangle =  $x$

$$\therefore \text{Angle at vertex} = (x + 15^\circ)$$

We know that

$$\therefore (x + 15^\circ) + x + x = 180^\circ$$

$$\Rightarrow 3x = 180^\circ - 15^\circ = 165^\circ$$


$$\Rightarrow x = 55^\circ$$

Hence, angle at each base is  $55^\circ$ .

**20. Solution:**

According to question,

OBA is a right-angled triangle



Given,  $AC = 18.0 \text{ cm}$

$$OB = \frac{AC}{2} = 9.0 \text{ cm}$$

∴  $OA$  is a hypotenuse.

We know that, hypotenuse is always greater than other two sides.

Hence, the radius of this circle is always greater than  $9.0\text{ cm}$ .

### 21. Solution:

As we know that, in a cyclic quadrilateral, the internal opposite angle is equal to the external angle.

Given, internal opposite angle =  $51^\circ$

∴ External angle = internal opposite angle =  $51^\circ$

Hence, the external angle of a cyclic quadrilateral is equal to  $51^\circ$ .

OR

### Solution:

$$\sqrt{15 + 10\sqrt{2}} + \sqrt{15 - 10\sqrt{2}}$$

Square the given expression to remove the outer square root

Using the formula  $(a + b)^2 = a^2 + 2ab + b^2$

$$\begin{aligned} & \left( \sqrt{15 + 10\sqrt{2}} + \sqrt{15 - 10\sqrt{2}} \right)^2 \\ &= \left( \sqrt{15 + 10\sqrt{2}} \right)^2 + 2\sqrt{15 + 10\sqrt{2}}\sqrt{15 - 10\sqrt{2}} + \left( \sqrt{15 - 10\sqrt{2}} \right)^2 \\ &= 15 + 10\sqrt{2} + 2\sqrt{15 + 10\sqrt{2}}\sqrt{15 - 10\sqrt{2}} + 15 - 10\sqrt{2} \end{aligned}$$

Sum of the first and the third terms simplifies to

$$2\sqrt{15 + 10\sqrt{2}}\sqrt{15 - 10\sqrt{2}} + 30$$

By Law of exponents in multiplication

$$2\sqrt{(15 + 10\sqrt{2})(15 - 10\sqrt{2})} + 30$$

Using the formula  $a^2 - b^2 = (a + b)(a - b)$

$$2\sqrt{15^2 - (10\sqrt{2})^2} + 30 = 2\sqrt{225 - 200} + 30 = 2\sqrt{25} + 30 = 2(5) + 30 = 40$$

We need to take square root of 40 as we have squared in the first step

So, answer is  $\sqrt{40} = 2\sqrt{10}$

## 22. Solution:

The cumulative frequency, is given below:

Class	Frequency( $f_i$ )	Cumulative Frequency
3	6	6
6	12	18
9	9	27
12	14	41
15	24	64
18	11	76
	$\sum f_i = 76$	

$\therefore N = 76$ , which is even.

$$\Rightarrow \frac{N}{2} = \frac{76}{2} = 38$$

$$\Rightarrow \frac{N}{2} + 1 = \frac{76}{2} + 1 = 38 + 1 = 39$$

$$\text{Median} = \frac{1}{2} \{(\text{Value of } 38^{\text{th}} \text{ term}) + (\text{Value of } 39^{\text{th}} \text{ term})\}$$

$$= \frac{1}{2} \{12 + 12\}$$

$$= \frac{1}{2} \{24\}$$

$$= 12$$

Hence, median = 12.

OR

**Solution:**

All possible outcomes are 6, 7, 8, ..., 50.

Total number of all possible outcomes = 45

Favourable outcomes are 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47.

Number of all favourable outcomes = 12

Let E be the event of getting a prime number.

$$\therefore P(E) = \frac{\text{Number of all favourable outcomes}}{\text{Total number of all possible outcomes}} = \frac{12}{45} = \frac{4}{15}$$

Hence, the probability of getting a prime number =  $\frac{4}{15}$

### Section D

**23. Solution:**

$$\text{Given, } x = \frac{\sqrt{7}+\sqrt{6}}{\sqrt{7}-\sqrt{6}} \text{ and } y = \frac{\sqrt{7}-\sqrt{6}}{\sqrt{7}+\sqrt{6}}$$

$$x = \frac{\sqrt{7}+\sqrt{6}}{\sqrt{7}-\sqrt{6}} \times \frac{\sqrt{7}+\sqrt{6}}{\sqrt{7}+\sqrt{6}}$$

$$x = \frac{(\sqrt{7}+\sqrt{6})^2}{\sqrt{7}^2-\sqrt{6}^2} = \frac{7+6+2\cdot\sqrt{7}\cdot\sqrt{6}}{1} = 13 + 2\sqrt{42}$$

$$y = \frac{\sqrt{7}-\sqrt{6}}{\sqrt{7}+\sqrt{6}} \times \frac{\sqrt{7}-\sqrt{6}}{\sqrt{7}-\sqrt{6}}$$

$$y = \frac{(\sqrt{7}-\sqrt{6})^2}{\sqrt{7}^2-\sqrt{6}^2} = \frac{7+6-2\cdot\sqrt{7}\cdot\sqrt{6}}{1} = 13 - 2\sqrt{42}$$

$$\therefore x + y = 13 + 2\sqrt{42} + 13 - 2\sqrt{42} = 26$$

$$\therefore x + y = 26$$

**24. Solution:**

$$\text{Given } P(x) = 2x^3 - 5x^2 + 3x + 7$$

$$2x + 1 = 0 \Rightarrow x = -\frac{1}{2}$$

By remainder theorem, we know that when  $P(x)$  is divided by  $(2x + 1)$ , the remainder is  $P\left(-\frac{1}{2}\right)$ .

$$\text{Now, } P\left(-\frac{1}{2}\right) = 2\left(-\frac{1}{2}\right)^3 - 5\left(-\frac{1}{2}\right)^2 + 3\left(-\frac{1}{2}\right) + 7$$

$$= 2\left(-\frac{1}{8}\right) - 5\left(\frac{1}{4}\right) - \frac{3}{2} + 7$$

$$= -\frac{1}{4} - \frac{5}{4} - \frac{3}{2} + 7$$

$$= \frac{-1-5-6+28}{4}$$

$$= \frac{16}{4}$$

$$= 4$$

Hence, the required remainder is 4.

**25. Solution:**

$$\text{Given, } x + y + z = 6 \text{ and } x^2 + y^2 + z^2 = 18$$

Squaring both sides, we get

$$\Rightarrow (x + y + z)^2 = 6^2$$

$$\Rightarrow x^2 + y^2 + z^2 + 2(xy + yz + zx) = 36$$

$$\Rightarrow 18 + 2(xy + yz + zx) = 36$$

$$\Rightarrow 2(xy + yz + zx) = 36 - 18 = 18$$

$$\Rightarrow xy + yz + zx = \frac{18}{2} = 9$$

$$\therefore x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - (xy + yz + zx))$$

$$= 6(18 - 9) = 54$$

$$\text{Hence, } x^3 + y^3 + z^3 - 3xyz = 54$$

OR

### Solution

Since  $t^2 - 1$  exactly divides the polynomial  $P(t) = a_1t^4 + a_2t^3 + a_3t^2 + a_4t + a_5$ , it means  $t^2 - 1$  is a factor of  $P(t)$

So,  $(t + 1)(t - 1)$  is a factor of  $P(t)$

Therefore  $P(1) = 0$  and  $P(-1) = 0$

Substituting the values in the polynomial we get

$$P(1) = a_1(1)^4 + a_2(1)^3 + a_3(1)^2 + a_4(1) + a_5 = a_1 + a_2 + a_3 + a_4 + a_5 = 0$$

$$P(-1) = a_1(-1)^4 + a_2(-1)^3 + a_3(-1)^2 + a_4(-1) + a_5 = a_1 - a_2 + a_3 - a_4 + a_5 = 0$$

Adding the above two equations, we get

$$2(a_1 + a_3 + a_5) = 0$$

$$a_1 + a_3 + a_5 = 0$$

Subtracting  $P(-1)$  from  $P(1)$

$$2(a_2 + a_4) = 0$$

$$a_2 + a_4 = 0$$


Therefore  $a_1 + a_3 + a_5 = a_2 + a_4 = 0$

### 26. Solution:

Let  $a$  be the side of a rhombus.

Given, perimeter of the rhombus is  $48 \text{ cm}$

$$\therefore \text{Side of the rhombus } (a) = \frac{48}{4} = 12 \text{ cm}$$



$$\therefore \text{Area of rhombus } ABCD = ar(\Delta ABC) + ar(\Delta ABC)$$



$$\text{Area of equilateral triangle } ABC = \frac{\sqrt{3}}{4} a^2$$

$$= \frac{\sqrt{3}}{4} \times 12 \times 12$$

$$= 36\sqrt{3} \text{ cm}^2$$

$$\text{Area of equilateral triangle } ADC = 36\sqrt{3} \text{ cm}^2$$

$$\therefore \text{Area of rhombus } ABCD = ar(\Delta ABC) + ar(\Delta ADC)$$

$$= (36\sqrt{3} + 36\sqrt{3}) \text{ cm}^2$$

$$= 72\sqrt{3} \text{ cm}^2$$

Hence, the area of the rhombus =  $72\sqrt{3} \text{ cm}^2$ .

### 27. Solution:

Let  $r$  be the radius of cylinder.

Given, height ( $h$ ) =  $30 \text{ cm}$  and volume of the cylinder =  $750\pi \text{ cm}^3$

$$\therefore \text{Volume of the cylinder} = \pi r^2 h$$

$$\Rightarrow \pi r^2 \times 30 = 750\pi$$

$$\Rightarrow r^2 = \frac{750\pi}{30\pi} = 25$$

$$\Rightarrow r^2 = 5^2$$

$$\Rightarrow r = 5 \text{ cm}$$

$$\therefore \text{Total surface area} = 2\pi r(r + h)$$

$$= 2 \times \frac{22}{7} \times 5 \times (5 + 30) \text{ cm}^2$$

$$= 10 \times \frac{22}{7} \times 35 \text{ cm}^2$$

$$= 10 \times 22 \times 5 \text{ cm}^2$$

$$= 10 \times 110 \text{ cm}^2$$

$$= 1100 \text{ cm}^2$$

Hence, radius ( $r$ ) = 5 cm and total surface area = 110 cm<sup>2</sup>.

OR

**Solution:**

Let  $h$  be the height of the trapezium.

Given, area of the trapezium = 350 cm<sup>2</sup>,

Sum of the parallel sides of the trapezium = 70 cm

Area of the trapezium =  $\frac{1}{2} \times (\text{sum of parallel sides}) \times \text{height}$

$$\Rightarrow \frac{1}{2} \times 70 \times h = 350$$


$$\Rightarrow 35h = 350$$

$$\Rightarrow h = 10 \text{ cm}$$

Hence, the height of the trapezium = 10 cm.

**28. Solution:**

Let O be the centre of the given circle and P be a point such that  $OP = 40 \text{ cm}$



Let PT be tangent such that  $PT = 32 \text{ cm}$

Join OT.

Now, PT is a tangent at T and OT is the radius through T.

$$\therefore OT \perp PT$$

In the right  $\triangle OTP$ , we have

By Pythagoras's Theorem,

$$OP^2 = OT^2 + PT^2$$

$$OT = \sqrt{OP^2 - PT^2}$$

$$= \sqrt{40^2 - 32^2} = \sqrt{1600 - 1024} = \sqrt{576} = 24 \text{ cm}$$

Hence, radius of the circle is 24 cm.

**29. Solution:**

Let  $b$  cm be the unequal side of an isosceles triangle.

Here, perimeter of an isosceles triangle = 64 cm and equal sides ( $a$ ) = 20 cm.

$$\therefore \text{Perimeter of an isosceles triangle} = (2a + b) \text{ cm}$$

$$\Rightarrow (2 \times 20) + b = 64$$

$$\Rightarrow b = 64 - 40 = 24 \text{ cm}$$

$$\therefore \text{Area of an isosceles triangle} = \frac{1}{4}b\sqrt{4a^2 - b^2} \text{ square units}$$

$$= \frac{1}{4} \times 24\sqrt{4 \times 20^2 - 24^2} \text{ cm}^2$$

$$= \frac{1}{4} \times 24\sqrt{4 \times 400 - 576} \text{ cm}^2$$

$$= 6 \times \sqrt{1600 - 576} \text{ cm}^2$$

$$= 6 \times \sqrt{1024} \text{ cm}^2$$

$$= 6 \times 32 \text{ cm}^2 = 192 \text{ cm}^2$$

Hence, Area of the isosceles triangle = 192 cm<sup>2</sup>.

**OR**

**Solution:**

All possible outcomes are 14, 15, 16, ....., 77.

Total number of all possible outcomes = 64

Favourable outcomes are 14, 21, 28, 35, 42, 49, 56, 63, 70, 77.

Number of all favourable outcomes = 10

Let E be the event of getting a number is divisible by 7.

$$\therefore P(E) = \frac{\text{Number of all favourable outcomes}}{\text{Total number of all possible outcomes}} = \frac{10}{64} = \frac{5}{32}$$

Hence, the probability of the number is divisible by 7 =  $\frac{5}{32}$ .

### 30. Solution:

We prepare the cumulative frequency, as given below:

Class	Frequency ( $f_i$ )	$f_i \times x_i$
3	6	18
5	8	40
7	15	105
9	p	9p
11	8	88
13	4	52
	$\sum f_i = 41 + p$	$\sum f_i \times x_i = 303 + 9p$

$$\therefore \text{Mean} = \frac{\sum f_i \times x_i}{\sum f_i}$$

$$= \frac{303+9p}{41+p}$$

$$\Rightarrow \frac{303+9p}{41+p} = 8$$

$$\Rightarrow 303 + 9p = 8p + 328$$

$$\Rightarrow p = 25$$

Hence,  $p = 25$