

## Exercise 6.1

Page No: 6.2

**Question 1:** Which of the following expressions are polynomials in one variable and which are not?

State reasons for your answer:

(i)  $3x^2 - 4x + 15$

(ii)  $y^2 + 2\sqrt{3}$

(iii)  $3\sqrt{x} + \sqrt{2}x$

(iv)  $x - 4/x$

(v)  $x^{12} + y^3 + t^{50}$

**Solution:**

(i)  $3x^2 - 4x + 15$

It is a polynomial of  $x$ .

(ii)  $y^2 + 2\sqrt{3}$

It is a polynomial of  $y$ .

(iii)  $3\sqrt{x} + \sqrt{2}x$

It is not a polynomial since the exponent of  $3\sqrt{x}$  is a rational term.

(iv)  $x - 4/x$

It is not a polynomial since the exponent of  $-4/x$  is not a positive term.

(v)  $x^{12} + y^3 + t^{50}$

It is a three variable polynomial,  $x$ ,  $y$  and  $t$ .

**Question 2:** Write the coefficient of  $x^2$  in each of the following:

(i)  $17 - 2x + 7x^2$

(ii)  $9 - 12x + x^3$

(iii)  $\pi/6 x^2 - 3x + 4$

(iv)  $\sqrt{3}x - 7$

**Solution:**

(i)  $17 - 2x + 7x^2$

Coefficient of  $x^2 = 7$

(ii)  $9 - 12x + x^3$

Coefficient of  $x^2 = 0$

(iii)  $\frac{1}{6}x^2 - 3x + 4$   
Coefficient of  $x^2 = \frac{1}{6}$

(iv)  $\sqrt{3}x - 7$   
Coefficient of  $x^2 = 0$

**Question 3: Write the degrees of each of the following polynomials:**

(i)  $7x^3 + 4x^2 - 3x + 12$

(ii)  $12 - x + 2x^3$

(iii)  $5y - \sqrt{2}$

(iv) 7

(v) 0

**Solution:**

As we know, degree is the highest power in the polynomial

(i) Degree of the polynomial  $7x^3 + 4x^2 - 3x + 12$  is 3

(ii) Degree of the polynomial  $12 - x + 2x^3$  is 3

(iii) Degree of the polynomial  $5y - \sqrt{2}$  is 1

(iv) Degree of the polynomial 7 is 0

(v) Degree of the polynomial 0 is undefined.

**Question 4: Classify the following polynomials as linear, quadratic, cubic and biquadratic polynomials:**

(i)  $x + x^2 + 4$

(ii)  $3x - 2$

(iii)  $2x + x^2$

(iv)  $3y$

(v)  $t^2 + 1$

(vi)  $7t^4 + 4t^3 + 3t - 2$

**Solution:**

(i)  $x + x^2 + 4$ : It is a quadratic polynomial as its degree is 2.

(ii)  $3x - 2$ : It is a linear polynomial as its degree is 1.

(iii)  $2x + x^2$ : It is a quadratic polynomial as its degree is 2.

(iv)  $3y$ : It is a linear polynomial as its degree is 1.

(v)  $t^2 + 1$ : It is a quadratic polynomial as its degree is 2.

(vi)  $7t^4 + 4t^3 + 3t - 2$ : It is a biquadratic polynomial as its degree is 4.

## Exercise 6.2

Page No: 6.8

**Question 1:** If  $f(x) = 2x^3 - 13x^2 + 17x + 12$ , find

(i)  $f(2)$

(ii)  $f(-3)$

(iii)  $f(0)$

**Solution:**

$$f(x) = 2x^3 - 13x^2 + 17x + 12$$

$$\begin{aligned} \text{(i) } f(2) &= 2(2)^3 - 13(2)^2 + 17(2) + 12 \\ &= 2 \times 8 - 13 \times 4 + 17 \times 2 + 12 \\ &= 16 - 52 + 34 + 12 \\ &= 62 - 52 \\ &= 10 \end{aligned}$$

$$\begin{aligned} \text{(ii) } f(-3) &= 2(-3)^3 - 13(-3)^2 + 17 \times (-3) + 12 \\ &= 2 \times (-27) - 13 \times 9 + 17 \times (-3) + 12 \\ &= -54 - 117 - 51 + 12 \\ &= -222 + 12 \\ &= -210 \end{aligned}$$

$$\begin{aligned} \text{(iii) } f(0) &= 2 \times (0)^3 - 13(0)^2 + 17 \times 0 + 12 \\ &= 0 - 0 + 0 + 12 \\ &= 12 \end{aligned}$$

**Question 2:** Verify whether the indicated numbers are zeros of the polynomials corresponding to them in the following cases:

(i)  $f(x) = 3x + 1, x = -1/3$

(ii)  $f(x) = x^2 - 1, x = 1, -1$

(iii)  $g(x) = 3x^2 - 2, x = 2/\sqrt{3}, -2/\sqrt{3}$

(iv)  $p(x) = x^3 - 6x^2 + 11x - 6, x = 1, 2, 3$

(v)  $f(x) = 5x - \pi, x = 4/5$

(vi)  $f(x) = x^2, x = 0$

(vii)  $f(x) = lx + m, x = -m/l$

(viii)  $f(x) = 2x + 1, x = 1/2$

**Solution:**

(i)  $f(x) = 3x + 1, x = -1/3$

$$f(x) = 3x + 1$$

Substitute  $x = -1/3$  in  $f(x)$

$$f(-1/3) = 3(-1/3) + 1$$

$$= -1 + 1$$

$$= 0$$

Since, the result is 0, so  $x = -1/3$  is the root of  $3x + 1$

**(ii)**  $f(x) = x^2 - 1$ ,  $x = 1, -1$

$$f(x) = x^2 - 1$$

Given that  $x = (1, -1)$

Substitute  $x = 1$  in  $f(x)$

$$f(1) = 1^2 - 1$$

$$= 1 - 1$$

$$= 0$$

Now, substitute  $x = (-1)$  in  $f(x)$

$$f(-1) = (-1)^2 - 1$$

$$= 1 - 1$$

$$= 0$$

Since, the results when  $x = 1$  and  $x = -1$  are 0, so  $(1, -1)$  are the roots of the polynomial  $f(x) = x^2 - 1$

**(iii)**  $g(x) = 3x^2 - 2$ ,  $x = 2/\sqrt{3}, -2/\sqrt{3}$

$$g(x) = 3x^2 - 2$$

Substitute  $x = 2/\sqrt{3}$  in  $g(x)$

$$g(2/\sqrt{3}) = 3(2/\sqrt{3})^2 - 2$$

$$= 3(4/3) - 2$$

$$= 4 - 2$$

$$= 2 \neq 0$$

Now, Substitute  $x = -2/\sqrt{3}$  in  $g(x)$

$$g(2/\sqrt{3}) = 3(-2/\sqrt{3})^2 - 2$$

$$= 3(4/3) - 2$$

$$= 4 - 2$$

$$= 2 \neq 0$$

Since, the results when  $x = 2/\sqrt{3}$  and  $x = -2/\sqrt{3}$  are not 0. Therefore  $(2/\sqrt{3}, -2/\sqrt{3})$  are not zeros of  $3x^2 - 2$ .

**(iv)**  $p(x) = x^3 - 6x^2 + 11x - 6$ ,  $x = 1, 2, 3$

$$p(1) = 1^3 - 6(1)^2 + 11 \times 1 - 6 = 1 - 6 + 11 - 6 = 0$$

$$p(2) = 2^3 - 6(2)^2 + 11 \times 2 - 6 = 8 - 24 + 22 - 6 = 0$$

$$p(3) = 3^3 - 6(3)^2 + 11 \times 3 - 6 = 27 - 54 + 33 - 6 = 0$$

Therefore,  $x = 1, 2, 3$  are zeros of  $p(x)$ .

**(v)**  $f(x) = 5x - \pi$ ,  $x = 4/5$

$$f(4/5) = 5 \times 4/5 - \pi = 4 - \pi \neq 0$$

Therefore,  $x = 4/5$  is not a zeros of  $f(x)$ .

**(vi)**  $f(x) = x^2$ ,  $x = 0$

$$f(0) = 0^2 = 0$$

Therefore,  $x = 0$  is a zero of  $f(x)$ .

(vii)  $f(x) = lx + m$ ,  $x = -m/l$


$$f(-m/l) = l \times -m/l + m = -m + m = 0$$

Therefore,  $x = -m/l$  is a zero of  $f(x)$ .

(viii)  $f(x) = 2x + 1$ ,  $x = \frac{1}{2}$

$$f(1/2) = 2 \times 1/2 + 1 = 1 + 1 = 2 \neq 0$$

Therefore,  $x = \frac{1}{2}$  is not a zero of  $f(x)$ .



## Exercise 6.3

Page No: 6.14

In each of the following, using the remainder theorem, find the remainder when  $f(x)$  is divided by  $g(x)$  and verify the by actual division : (1 – 8)

**Question 1:**  $f(x) = x^3 + 4x^2 - 3x + 10$ ,  $g(x) = x + 4$

**Solution:**

$$f(x) = x^3 + 4x^2 - 3x + 10, g(x) = x + 4$$

Put  $g(x) = 0$

$$\Rightarrow x + 4 = 0 \text{ or } x = -4$$

Remainder =  $f(-4)$

Now,

$$f(-4) = (-4)^3 + 4(-4)^2 - 3(-4) + 10 = -64 + 64 + 12 + 10 = 22$$

**Actual Division:**

$$\begin{array}{r}
 x^2 - 3 \\
 x + 4 \overline{) x^3 + 4x^2 - 3x + 10} \\
 \underline{-(x^3 + 4x^2)} \phantom{+ 10} \\
 0 - 3x + 10 \\
 \underline{-(0 - 3x - 12)} \\
 22
 \end{array}$$

**Question 2:**  $f(x) = 4x^4 - 3x^3 - 2x^2 + x - 7$ ,  $g(x) = x - 1$

**Solution:**

$$f(x) = 4x^4 - 3x^3 - 2x^2 + x - 7$$

Put  $g(x) = 0$

$$\Rightarrow x - 1 = 0 \text{ or } x = 1$$

Remainder =  $f(1)$

Now,

$$f(1) = 4(1)^4 - 3(1)^3 - 2(1)^2 + (1) - 7 = 4 - 3 - 2 + 1 - 7 = -7$$

**Actual Division:**

$$\begin{array}{r}
 4x^3 \quad +x^2 \quad -x \\
 x-1 \overline{) 4x^4 - 3x^3 - 2x^2 + x - 7} \\
 \underline{4x^4 - 4x^3} \phantom{- 2x^2 + x - 7} \\
 x^3 - 2x^2 + x - 7 \\
 \underline{x^3 - x^2} \phantom{+ x - 7} \\
 -x^2 + x - 7 \\
 \underline{-x^2 + x} \phantom{- 7} \\
 0 - 7
 \end{array}$$

**Question 3:**  $f(x) = 2x^4 - 6x^3 + 2x^2 - x + 2$ ,  $g(x) = x + 2$

**Solution:**

$$f(x) = 2x^4 - 6x^3 + 2x^2 - x + 2, g(x) = x + 2$$

$$\text{Put } g(x) = 0$$

$$\Rightarrow x + 2 = 0 \text{ or } x = -2$$

$$\text{Remainder} = f(-2)$$

Now,

$$f(-2) = 2(-2)^4 - 6(-2)^3 + 2(-2)^2 - (-2) + 2 = 32 + 48 + 8 + 2 + 2 = 92$$

**Actual Division:**

$$\begin{array}{r}
 \begin{array}{r} 2x^3 - 10x^2 + 22x - 45 \end{array} \\
 x + 2 \overline{) \begin{array}{r} 2x^4 - 6x^3 + 2x^2 - x + 2 \end{array}} \\
 \underline{\begin{array}{r} 2x^4 + 4x^3 \end{array}} \\
 \begin{array}{r} -10x^3 + 2x^2 - x + 2 \end{array} \\
 \underline{\begin{array}{r} -10x^3 - 20x^2 \end{array}} \\
 \begin{array}{r} 22x^2 - x + 2 \end{array} \\
 \underline{\begin{array}{r} 22x^2 + 44x \end{array}} \\
 \begin{array}{r} -45x + 2 \end{array} \\
 \underline{\begin{array}{r} -45x - 90 \end{array}} \\
 \begin{array}{r} 92 \end{array}
 \end{array}$$

**Question 4:**  $f(x) = 4x^3 - 12x^2 + 14x - 3$ ,  $g(x) = 2x - 1$

**Solution:**

$$f(x) = 4x^3 - 12x^2 + 14x - 3, g(x) = 2x - 1$$

Put  $g(x) = 0$

$$\Rightarrow 2x - 1 = 0 \text{ or } x = 1/2$$

Remainder =  $f(1/2)$

Now,

$$f(1/2) = 4(1/2)^3 - 12(1/2)^2 + 14(1/2) - 3 = \frac{1}{2} - 3 + 7 - 3 = \frac{3}{2}$$

**Actual Division:**

$$\begin{array}{r}
 2x^2 - 5x + \frac{9}{2} \\
 2x - 1 \overline{) 4x^3 - 12x^2 + 14x - 3} \\
 \underline{4x^3 - 2x^2} \phantom{+ 14x - 3} \\
 -10x^2 + 14x - 3 \\
 \underline{-10x^2 + 5x} \phantom{- 3} \\
 9x - 3 \\
 \underline{9x - \frac{9}{2}} \\
 \frac{3}{2}
 \end{array}$$

**Question 5:**  $f(x) = x^3 - 6x^2 + 2x - 4$ ,  $g(x) = 1 - 2x$

**Solution:**

$$f(x) = x^3 - 6x^2 + 2x - 4, g(x) = 1 - 2x$$

$$\text{Put } g(x) = 0$$

$$\Rightarrow 1 - 2x = 0 \text{ or } x = \frac{1}{2}$$

$$\text{Remainder} = f\left(\frac{1}{2}\right)$$

Now,

$$f\left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)^3 - 6\left(\frac{1}{2}\right)^2 + 2\left(\frac{1}{2}\right) - 4 = 1 + \frac{1}{8} - 4 - \frac{3}{2} = -\frac{35}{8}$$

**Actual Division:**

$$\begin{array}{r}
 -\frac{x^2}{2} + \frac{11x}{4} + \frac{3}{8} \\
 -2x + 1 \overline{) x^3 - 6x^2 + 2x - 4} \\
 \underline{x^3 \phantom{- 6x^2} - \frac{x^2}{2}} \\
 -\frac{11x^2}{2} + 2x - 4 \\
 \underline{-\frac{11x^2}{2} + \frac{11x}{4}} \\
 -\frac{3x}{4} - 4 \\
 \underline{-\frac{3x}{4} + \frac{3}{8}} \\
 -\frac{35}{8}
 \end{array}$$

**Question 6:**  $f(x) = x^4 - 3x^2 + 4$ ,  $g(x) = x - 2$

**Solution:**

$$f(x) = x^4 - 3x^2 + 4, g(x) = x - 2$$

$$\text{Put } g(x) = 0$$

$$\Rightarrow x - 2 = 0 \text{ or } x = 2$$

$$\text{Remainder} = f(2)$$

Now,

$$f(2) = (2)^4 - 3(2)^2 + 4 = 16 - 12 + 4 = 8$$

**Actual Division:**

$$\begin{array}{r}
 x^3 + 2x^2 + x + 2 \\
 x - 2 \overline{) x^4 + 0x^3 - 3x^2 + 0x + 4} \\
 \underline{x^4 - 2x^3} \phantom{+ 0x^2 + 0x + 4} \\
 2x^3 - 3x^2 + 0x + 4 \\
 \underline{2x^3 - 4x^2} \phantom{+ 0x + 4} \\
 x^2 + 0x + 4 \\
 \underline{x^2 - 2x} \phantom{+ 4} \\
 2x + 4 \\
 \underline{2x - 4} \\
 8
 \end{array}$$

**Question 7:**  $f(x) = 9x^3 - 3x^2 + x - 5$ ,  $g(x) = x - 2/3$

**Solution:**

$$f(x) = 9x^3 - 3x^2 + x - 5, g(x) = x - 2/3$$

$$\text{Put } g(x) = 0$$

$$\Rightarrow x - 2/3 = 0 \text{ or } x = 2/3$$

$$\text{Remainder} = f(2/3)$$

Now,

$$f(2/3) = 9(2/3)^3 - 3(2/3)^2 + (2/3) - 5 = 8/3 - 4/3 + 2/3 - 5/1 = -3$$

**Actual Division:**

$$\begin{array}{r}
 \phantom{x - \frac{2}{3}} \quad \quad \quad 9x^2 \quad +3x \quad +3 \\
 x - \frac{2}{3} \quad \overline{) \quad 9x^3 \quad -3x^2 \quad +x \quad -5} \\
 \phantom{x - \frac{2}{3}} \quad \quad \quad - \\
 \phantom{x - \frac{2}{3}} \quad 9x^3 \quad -6x^2 \phantom{+x} \phantom{-5} \\
 \phantom{x - \frac{2}{3}} \quad \quad \quad \phantom{9x^3} \quad 3x^2 \quad +x \quad -5 \\
 \phantom{x - \frac{2}{3}} \quad \quad \quad \phantom{9x^3} \quad \quad \quad - \\
 \phantom{x - \frac{2}{3}} \quad \quad \quad \phantom{9x^3} \quad 3x^2 \quad -2x \phantom{-5} \\
 \phantom{x - \frac{2}{3}} \quad \quad \quad \phantom{9x^3} \quad \quad \quad \phantom{3x^2} \quad 3x \quad -5 \\
 \phantom{x - \frac{2}{3}} \quad \quad \quad \phantom{9x^3} \quad \quad \quad \phantom{3x^2} \quad \quad \quad - \\
 \phantom{x - \frac{2}{3}} \quad \quad \quad \phantom{9x^3} \quad \quad \quad \phantom{3x^2} \quad 3x \quad -2 \\
 \phantom{x - \frac{2}{3}} \quad \quad \quad \phantom{9x^3} \quad \quad \quad \phantom{3x^2} \quad \quad \quad \phantom{3x} \quad -3
 \end{array}$$

Exercise 6.4

Page No: 6.24

In each of the following, use factor theorem to find whether polynomial  $g(x)$  is a factor of polynomial  $f(x)$  or, not: (1-7)

**Question 1:**  $f(x) = x^3 - 6x^2 + 11x - 6$ ;  $g(x) = x - 3$

**Solution:**

If  $g(x)$  is a factor of  $f(x)$ , then the remainder will be zero that is  $g(x) = 0$ .

$$g(x) = x - 3 = 0$$

$$\text{or } x = 3$$

$$\text{Remainder} = f(3)$$

Now,

$$f(3) = (3)^3 - 6(3)^2 + 11 \times 3 - 6$$

$$= 27 - 54 + 33 - 6$$

$$= 60 - 60$$

$$= 0$$

Therefore,  $g(x)$  is a factor of  $f(x)$

**Question 2:**  $f(x) = 3x^4 + 17x^3 + 9x^2 - 7x - 10$ ;  $g(x) = x + 5$

**Solution:**

If  $g(x)$  is a factor of  $f(x)$ , then the remainder will be zero that is  $g(x) = 0$ .

$$g(x) = x + 5 = 0, \text{ then } x = -5$$

$$\text{Remainder} = f(-5)$$

Now,

$$f(-5) = 3(-5)^4 + 17(-5)^3 + 9(-5)^2 - 7(-5) - 10$$

$$= 3 \times 625 + 17 \times (-125) + 9 \times (25) - 7 \times (-5) - 10$$

$$= 1875 - 2125 + 225 + 35 - 10$$

$$= 0$$

Therefore,  $g(x)$  is a factor of  $f(x)$ .

**Question 3:**  $f(x) = x^5 + 3x^4 - x^3 - 3x^2 + 5x + 15$ ,  $g(x) = x + 3$

**Solution:**

If  $g(x)$  is a factor of  $f(x)$ , then the remainder will be zero that is  $g(x) = 0$ .

$$g(x) = x + 3 = 0, \text{ then } x = -3$$

$$\text{Remainder} = f(-3)$$

Now,

$$\begin{aligned} f(-3) &= (-3)^5 + 3(-3)^4 - (-3)^3 - 3(-3)^2 + 5(-3) + 15 \\ &= -243 + 3 \times 81 - (-27) - 3 \times 9 + 5(-3) + 15 \\ &= -243 + 243 + 27 - 27 - 15 + 15 \\ &= 0 \end{aligned}$$

Therefore,  $g(x)$  is a factor of  $f(x)$ .

**Question 4:**  $f(x) = x^3 - 6x^2 - 19x + 84$ ,  $g(x) = x - 7$

**Solution:**

If  $g(x)$  is a factor of  $f(x)$ , then the remainder will be zero that is  $g(x) = 0$ .

$$g(x) = x - 7 = 0, \text{ then } x = 7$$

$$\text{Remainder} = f(7)$$

Now,

$$\begin{aligned} f(7) &= (7)^3 - 6(7)^2 - 19 \times 7 + 84 \\ &= 343 - 294 - 133 + 84 \\ &= 343 + 84 - 294 - 133 \\ &= 0 \end{aligned}$$

Therefore,  $g(x)$  is a factor of  $f(x)$ .

**Question 5:**  $f(x) = 3x^3 + x^2 - 20x + 12$ ,  $g(x) = 3x - 2$

**Solution:**

If  $g(x)$  is a factor of  $f(x)$ , then the remainder will be zero that is  $g(x) = 0$ .

$$g(x) = 3x - 2 = 0, \text{ then } x = 2/3$$

$$\text{Remainder} = f(2/3)$$

Now,

$$\begin{aligned} f(2/3) &= 3(2/3)^3 + (2/3)^2 - 20(2/3) + 12 \\ &= 3 \times 8/27 + 4/9 - 40/3 + 12 \\ &= 8/9 + 4/9 - 40/3 + 12 \\ &= 0/9 \\ &= 0 \end{aligned}$$

Therefore,  $g(x)$  is a factor of  $f(x)$ .

**Question 6:**  $f(x) = 2x^3 - 9x^2 + x + 12$ ,  $g(x) = 3 - 2x$

**Solution:**

If  $g(x)$  is a factor of  $f(x)$ , then the remainder will be zero that is  $g(x) = 0$ .

$$g(x) = 3 - 2x = 0, \text{ then } x = 3/2$$

$$\text{Remainder} = f(3/2)$$

Now,

$$f(3/2) = 2(3/2)^3 - 9(3/2)^2 + (3/2) + 12$$

$$= 2 \times 27/8 - 9 \times 9/4 + 3/2 + 12$$

$$= 27/4 - 81/4 + 3/2 + 12$$

$$= 0/4$$

$$= 0$$

Therefore,  $g(x)$  is a factor of  $f(x)$ .

**Question 7:**  $f(x) = x^3 - 6x^2 + 11x - 6$ ,  $g(x) = x^2 - 3x + 2$

**Solution:**

If  $g(x)$  is a factor of  $f(x)$ , then the remainder will be zero that is  $g(x) = 0$ .

$$g(x) = 0$$

$$\text{or } x^2 - 3x + 2 = 0$$

$$x^2 - x - 2x + 2 = 0$$

$$x(x - 1) - 2(x - 1) = 0$$

$$(x - 1)(x - 2) = 0$$

Therefore  $x = 1$  or  $x = 2$

Now,

$$f(1) = (1)^3 - 6(1)^2 + 11(1) - 6 = 1 - 6 + 11 - 6 = 12 - 12 = 0$$

$$f(2) = (2)^3 - 6(2)^2 + 11(2) - 6 = 8 - 24 + 22 - 6 = 30 - 30 = 0$$

$$\Rightarrow f(1) = 0 \text{ and } f(2) = 0$$

Which implies  $g(x)$  is factor of  $f(x)$ .

**Question 8:** Show that  $(x - 2)$ ,  $(x + 3)$  and  $(x - 4)$  are factors of  $x^3 - 3x^2 - 10x + 24$ .

**Solution:**

$$\text{Let } f(x) = x^3 - 3x^2 - 10x + 24$$

$$\text{If } x - 2 = 0, \text{ then } x = 2,$$

$$\text{If } x + 3 = 0 \text{ then } x = -3,$$

$$\text{and If } x - 4 = 0 \text{ then } x = 4$$

Now,

$$f(2) = (2)^3 - 3(2)^2 - 10 \times 2 + 24 = 8 - 12 - 20 + 24 = 32 - 32 = 0$$

$$f(-3) = (-3)^3 - 3(-3)^2 - 10(-3) + 24 = -27 - 27 + 30 + 24 = -54 + 54 = 0$$

$$f(4) = (4)^3 - 3(4)^2 - 10 \times 4 + 24 = 64 - 48 - 40 + 24 = 88 - 88 = 0$$

$$f(2) = 0$$

$$f(-3) = 0$$

$$f(4) = 0$$

Hence  $(x - 2)$ ,  $(x + 3)$  and  $(x - 4)$  are the factors of  $f(x)$

**Question 9: Show that  $(x + 4)$ ,  $(x - 3)$  and  $(x - 7)$  are factors of  $x^3 - 6x^2 - 19x + 84$ .**

**Solution:**

$$\text{Let } f(x) = x^3 - 6x^2 - 19x + 84$$

$$\text{If } x + 4 = 0, \text{ then } x = -4$$

$$\text{If } x - 3 = 0, \text{ then } x = 3$$

$$\text{and if } x - 7 = 0, \text{ then } x = 7$$

Now,

$$f(-4) = (-4)^3 - 6(-4)^2 - 19(-4) + 84 = -64 - 96 + 76 + 84 = 160 - 160 = 0$$

$$f(-4) = 0$$

$$f(3) = (3)^3 - 6(3)^2 - 19 \times 3 + 84 = 27 - 54 - 57 + 84 = 111 - 111 = 0$$

$$f(3) = 0$$

$$f(7) = (7)^3 - 6(7)^2 - 19 \times 7 + 84 = 343 - 294 - 133 + 84 = 427 - 427 = 0$$

$$f(7) = 0$$

Hence  $(x + 4)$ ,  $(x - 3)$ ,  $(x - 7)$  are the factors of  $f(x)$ .

### Exercise 6.5

Page No: 6.32

Using factor theorem, factorize each of the following polynomials:

**Question 1:**  $x^3 + 6x^2 + 11x + 6$

**Solution:**

$$\text{Let } f(x) = x^3 + 6x^2 + 11x + 6$$

Step 1: Find the factors of constant term

Here constant term = 6

Factors of 6 are  $\pm 1, \pm 2, \pm 3, \pm 6$

Step 2: Find the factors of  $f(x)$

$$\text{Let } x + 1 = 0$$

$$\Rightarrow x = -1$$

Put the value of  $x$  in  $f(x)$

$$f(-1) = (-1)^3 + 6(-1)^2 + 11(-1) + 6$$

$$= -1 + 6 - 11 + 6$$

$$= 12 - 12$$

$$= 0$$

So,  $(x + 1)$  is the factor of  $f(x)$

$$\text{Let } x + 2 = 0$$

$$\Rightarrow x = -2$$

Put the value of  $x$  in  $f(x)$

$$f(-2) = (-2)^3 + 6(-2)^2 + 11(-2) + 6 = -8 + 24 - 22 + 6 = 0$$

So,  $(x + 2)$  is the factor of  $f(x)$

$$\text{Let } x + 3 = 0$$

$$\Rightarrow x = -3$$

Put the value of  $x$  in  $f(x)$

$$f(-3) = (-3)^3 + 6(-3)^2 + 11(-3) + 6 = -27 + 54 - 33 + 6 = 0$$

So,  $(x + 3)$  is the factor of  $f(x)$

$$\text{Hence, } f(x) = (x + 1)(x + 2)(x + 3)$$

**Question 2:  $x^3 + 2x^2 - x - 2$**

**Solution:**

$$\text{Let } f(x) = x^3 + 2x^2 - x - 2$$

$$\text{Constant term} = -2$$

Factors of  $-2$  are  $\pm 1, \pm 2$

$$\text{Let } x - 1 = 0$$

$$\Rightarrow x = 1$$

Put the value of  $x$  in  $f(x)$

$$f(1) = (1)^3 + 2(1)^2 - 1 - 2 = 1 + 2 - 1 - 2 = 0$$

So,  $(x - 1)$  is factor of  $f(x)$

$$\text{Let } x + 1 = 0$$

$$\Rightarrow x = -1$$

Put the value of  $x$  in  $f(x)$

$$f(-1) = (-1)^3 + 2(-1)^2 - 1 - 2 = -1 + 2 + 1 - 2 = 0$$

$(x + 1)$  is a factor of  $f(x)$

$$\text{Let } x + 2 = 0$$

$$\Rightarrow x = -2$$

Put the value of  $x$  in  $f(x)$

$$f(-2) = (-2)^3 + 2(-2)^2 - (-2) - 2 = -8 + 8 + 2 - 2 = 0$$

$(x + 2)$  is a factor of  $f(x)$

$$\text{Let } x - 2 = 0$$

$$\Rightarrow x = 2$$

Put the value of  $x$  in  $f(x)$

$$f(2) = (2)^3 + 2(2)^2 - 2 - 2 = 8 + 8 - 2 - 2 = 12 \neq 0$$

$(x - 2)$  is not a factor of  $f(x)$

$$\text{Hence } f(x) = (x + 1)(x - 1)(x + 2)$$

**Question 3:  $x^3 - 6x^2 + 3x + 10$**

**Solution:**

$$\text{Let } f(x) = x^3 - 6x^2 + 3x + 10$$

Constant term = 10

Factors of 10 are  $\pm 1, \pm 2, \pm 5, \pm 10$

$$\text{Let } x + 1 = 0 \text{ or } x = -1$$

$$f(-1) = (-1)^3 - 6(-1)^2 + 3(-1) + 10 = 10 - 10 = 0$$

$$f(-1) = 0$$

$$\text{Let } x + 2 = 0 \text{ or } x = -2$$

$$f(-2) = (-2)^3 - 6(-2)^2 + 3(-2) + 10 = -8 - 24 - 6 + 10 = -28$$

$$f(-2) \neq 0$$

$$\text{Let } x - 2 = 0 \text{ or } x = 2$$

$$f(2) = (2)^3 - 6(2)^2 + 3(2) + 10 = 8 - 24 + 6 + 10 = 0$$

$$f(2) = 0$$

$$\text{Let } x - 5 = 0 \text{ or } x = 5$$

$$f(5) = (5)^3 - 6(5)^2 + 3(5) + 10 = 125 - 150 + 15 + 10 = 0$$

$$f(5) = 0$$

Therefore,  $(x + 1)$ ,  $(x - 2)$  and  $(x - 5)$  are factors of  $f(x)$

$$\text{Hence } f(x) = (x + 1)(x - 2)(x - 5)$$

**Question 4:  $x^4 - 7x^3 + 9x^2 + 7x - 10$** **Solution:**

$$\text{Let } f(x) = x^4 - 7x^3 + 9x^2 + 7x - 10$$

Constant term = -10

Factors of -10 are  $\pm 1, \pm 2, \pm 5, \pm 10$ Let  $x - 1 = 0$  or  $x = 1$ 

$$f(1) = (1)^4 - 7(1)^3 + 9(1)^2 + 7(1) - 10 = 1 - 7 + 9 + 7 - 10 = 0$$

$$f(1) = 0$$

Let  $x + 1 = 0$  or  $x = -1$ 

$$f(-1) = (-1)^4 - 7(-1)^3 + 9(-1)^2 + 7(-1) - 10 = 1 + 7 + 9 - 7 - 10 = 0$$

$$f(-1) = 0$$

Let  $x - 2 = 0$  or  $x = 2$ 

$$f(2) = (2)^4 - 7(2)^3 + 9(2)^2 + 7(2) - 10 = 16 - 56 + 36 + 14 - 10 = 0$$

$$f(2) = 0$$

Let  $x - 5 = 0$  or  $x = 5$ 

$$f(5) = (5)^4 - 7(5)^3 + 9(5)^2 + 7(5) - 10 = 625 - 875 + 225 + 35 - 10 = 0$$

$$f(5) = 0$$

Therefore,  $(x - 1)$ ,  $(x + 1)$ ,  $(x - 2)$  and  $(x - 5)$  are factors of  $f(x)$ 

$$\text{Hence } f(x) = (x - 1)(x + 1)(x - 2)(x - 5)$$

**Question 5:  $x^4 - 2x^3 - 7x^2 + 8x + 12$** **Solution:**

$$f(x) = x^4 - 2x^3 - 7x^2 + 8x + 12$$

Constant term = 12

Factors of 12 are  $\pm 1, \pm 2, \pm 3, \pm 4, \pm 6, \pm 12$ Let  $x - 1 = 0$  or  $x = 1$ 

$$f(1) = (1)^4 - 2(1)^3 - 7(1)^2 + 8(1) + 12 = 1 - 2 - 7 + 8 + 12 = 12$$

$$f(1) \neq 0$$

Let  $x + 1 = 0$  or  $x = -1$ 

$$f(-1) = (-1)^4 - 2(-1)^3 - 7(-1)^2 + 8(-1) + 12 = 1 + 2 - 7 - 8 + 12 = 0$$

$$f(-1) = 0$$

Let  $x + 2 = 0$  or  $x = -2$

$$f(-2) = (-2)^4 - 2(-2)^3 - 7(-2)^2 + 8(-2) + 12 = 16 + 16 - 28 - 16 + 12 = 0$$

$$f(-2) = 0$$

Let  $x - 2 = 0$  or  $x = 2$

$$f(2) = (2)^4 - 2(2)^3 - 7(2)^2 + 8(2) + 12 = 16 - 16 - 28 + 16 + 12 = 0$$

$$f(2) = 0$$

Let  $x - 3 = 0$  or  $x = 3$

$$f(3) = (3)^4 - 2(3)^3 - 7(3)^2 + 8(3) + 12 = 0$$

$$f(3) = 0$$

Therefore,  $(x - 1)$ ,  $(x + 2)$ ,  $(x - 2)$  and  $(x - 3)$  are factors of  $f(x)$

$$\text{Hence } f(x) = (x - 1)(x + 2)(x - 2)(x - 3)$$

**Question 6:  $x^4 + 10x^3 + 35x^2 + 50x + 24$**

**Solution:**

$$\text{Let } f(x) = x^4 + 10x^3 + 35x^2 + 50x + 24$$

Constant term = 24

Factors of 24 are  $\pm 1, \pm 2, \pm 3, \pm 4, \pm 6, \pm 8, \pm 12, \pm 24$

Let  $x + 1 = 0$  or  $x = -1$

$$f(-1) = (-1)^4 + 10(-1)^3 + 35(-1)^2 + 50(-1) + 24 = 1 - 10 + 35 - 50 + 24 = 0$$

$$f(-1) = 0$$

$(x + 1)$  is a factor of  $f(x)$

Likewise,  $(x + 2), (x + 3), (x + 4)$  are also the factors of  $f(x)$

$$\text{Hence } f(x) = (x + 1)(x + 2)(x + 3)(x + 4)$$

**Question 7:  $2x^4 - 7x^3 - 13x^2 + 63x - 45$**

**Solution:**

$$\text{Let } f(x) = 2x^4 - 7x^3 - 13x^2 + 63x - 45$$

Constant term = -45

Factors of -45 are  $\pm 1, \pm 3, \pm 5, \pm 9, \pm 15, \pm 45$

Here coefficient of  $x^4$  is 2. So possible rational roots of  $f(x)$  are

$$\pm 1, \pm 3, \pm 5, \pm 9, \pm 15, \pm 45, \pm 1/2, \pm 3/2, \pm 5/2, \pm 9/2, \pm 15/2, \pm 45/2$$

Let  $x - 1 = 0$  or  $x = 1$

$$f(1) = 2(1)^4 - 7(1)^3 - 13(1)^2 + 63(1) - 45 = 2 - 7 - 13 + 63 - 45 = 0$$

$$f(1) = 0$$

$f(x)$  can be written as,

$$f(x) = (x-1)(2x^3 - 5x^2 - 18x + 45)$$

$$\text{or } f(x) = (x-1)g(x) \dots (1)$$

Let  $x - 3 = 0$  or  $x = 3$

$$f(3) = 2(3)^4 - 7(3)^3 - 13(3)^2 + 63(3) - 45 = 162 - 189 - 117 + 189 - 45 = 0$$

$$f(3) = 0$$

Now, we are available with 2 factors of  $f(x)$ ,  $(x - 1)$  and  $(x - 3)$

$$\text{Here } g(x) = 2x^2(x-3) + x(x-3) - 15(x-3)$$

Taking  $(x-3)$  as common

$$= (x-3)(2x^2 + x - 15)$$

$$= (x-3)(2x^2 + 6x - 5x - 15)$$

$$= (x-3)(2x-5)(x+3)$$

$$= (x-3)(x+3)(2x-5) \dots (2)$$

From (1) and (2)

$$f(x) = (x-1)(x-3)(x+3)(2x-5)$$

## Exercise VSAQs

Page No: 6.33

**Question 1: Define zero or root of a polynomial**

**Solution:**

zero or root, is a solution to the polynomial equation,  $f(y) = 0$ .

It is that value of  $y$  that makes the polynomial equal to zero.

**Question 2: If  $x = 1/2$  is a zero of the polynomial  $f(x) = 8x^3 + ax^2 - 4x + 2$ , find the value of  $a$ .**

**Solution:**

If  $x = 1/2$  is a zero of the polynomial  $f(x)$ , then  $f(1/2) = 0$

$$8(1/2)^3 + a(1/2)^2 - 4(1/2) + 2 = 0$$

$$8 \times 1/8 + a/4 - 2 + 2 = 0$$

$$1 + a/4 = 0$$

$$a = -4$$

**Question 3: Write the remainder when the polynomial  $f(x) = x^3 + x^2 - 3x + 2$  is divided by  $x + 1$ .**

**Solution:**

Using factor theorem,

Put  $x + 1 = 0$  or  $x = -1$

$f(-1)$  is the remainder.

Now,

$$f(-1) = (-1)^3 + (-1)^2 - 3(-1) + 2$$

$$= -1 + 1 + 3 + 2$$

$$= 5$$

Therefore 5 is the remainder.

**Question 4:** Find the remainder when  $x^3 + 4x^2 + 4x - 3$  if divided by  $x$

**Solution:**

Using factor theorem,

Put  $x = 0$

$f(0)$  is the remainder.

Now,

$$f(0) = 0^3 + 4(0)^2 + 4 \times 0 - 3 = -3$$

Therefore -3 is the remainder.

**Question 5:** If  $x+1$  is a factor of  $x^3 + a$ , then write the value of  $a$ .

**Solution:**

$$\text{Let } f(x) = x^3 + a$$

If  $x+1$  is a factor of  $x^3 + a$  then  $f(-1) = 0$

$$(-1)^3 + a = 0$$

$$-1 + a = 0$$

$$\text{or } a = 1$$

**Question 6:** If  $f(x) = x^4 - 2x^3 + 3x^2 - ax - b$  when divided by  $x - 1$ , the remainder is 6, then find the value of  $a+b$ .

**Solution:**

From the statement, we have  $f(1) = 6$

$$(1)^4 - 2(1)^3 + 3(1)^2 - a(1) - b = 6$$

$$1 - 2 + 3 - a - b = 6$$

$$2 - a - b = 6$$

$$a + b = -4$$