




CBSE Class 12 Chemistry Question Paper Solution 2016


MARKING SCHEME -CHEMISTRY 2016


SET -56/1/N

Q.N.	Value Points	Marks
1.	(i) , Inversion of configuration	$\frac{1}{2} + \frac{1}{2}$
2.	NO ₂	1
3.	Due to presence of free electrons at interstitial sites, / metal excess defect	1
4.	N-methyl-2-methylpropanamine / 2-methyl-N-methylpropanamine	1
5.	Like Charged particles cause repulsion/ Brownian motion/ solvation	1
6.	(i) Osmotic pressure (ii) Positive deviation from Raouls' law/ Positive deviation	1 1
7	<div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> <p>(i)</p>  </div> <div style="text-align: center;"> <p>(ii)</p>  </div> </div>	1 +1
8.	(i) [Ni(H ₂ O) ₆] Cl ₂ (ii) Hexaaquanickel(II) chloride	1 1
9.	(i) zero order , bimolecular/ unimolecular (II) mol L ⁻¹ s ⁻¹	$\frac{1}{2} + \frac{1}{2}$ 1

	$t_{1/2} = 0.693 / k$ $k = 0.693 / 200$ $= 0.0034 \text{ min}^{-1} / 3.4 \times 10^{-3} \text{ min}^{-1}$	$\frac{1}{2}$ 1
13	(i) Oil as dispersed phase and water as dispersion medium (ii) The potential difference between fixed layer and diffused / double layer of opposite charges. (iii) Large number of atoms or smaller molecules of a substance aggregate together to form species having size in colloidal range.	1 1 1
14	1. Chromatography 2. To Separate two sulphide ores 3. It decomposes to CaO which removes impurity (silica) as slag/ Acts as flux.	1 1 1

15	$\Delta T_b = i \frac{K_b w_b \times 1000}{M_b \times w_a}$ $\Delta T_b = \frac{3 \times 0.52 \times 2 \times 1000}{142 \times 50}$ $= 0.439 \text{ K}$ $\Delta T_b = T_b - T_b^0$ $T_b = 0.439 + 373 = 373.439 \text{ K} \quad (\text{OR } 373.589 \text{ K})$	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1
16.	(i) Due to presence of two P-H bonds in H_3PO_2 / In H_3PO_2 O.S of P = +1 which can increase but in H_3PO_4 O.S of P = +5 (max.) (ii) Due to stronger S-S bond than O-O bond. (iii) Size of halogen increases / bond length increases / bond dissociation enthalpy decreases (any one)	1 1 1
17.		1 1 1
18.	(a) . In phenols lone pair of electron on oxygen are delocalized over benzene ring due to resonance but in alcohol lone pair of electron on oxygen are localized & hence available for protonation / + R- effect in phenol but not in	1

	(b)	
		1
23.	(i) Aware, concerned or any other two correct values (ii) Side effects/ health problems (iii) Neurologically active drugs/ stress relievers/drugs used to treat mental diseases example- valium, equanil (or any other two correct example)	½ + ½ 1 1 ½ + ½
24.	(a) $E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.059}{n} \log \frac{[\text{Cr}^{3+}]^2}{[\text{Fe}^{2+}]^3}$ $0.261 \text{ V} = E_{\text{cell}}^0 - \frac{0.059}{6} \log \frac{[0.01]^2}{[0.01]^3}$ $0.261 \text{ V} = E_{\text{cell}}^0 - \frac{0.059}{6} \log 100$ $E_{\text{cell}}^0 = 0.261 + 0.0197$ $= 0.2807 \text{ V}$ (b) A, due to its more negative E^0 value.	½ 1 ½ 1 1+1
	OR	
24	(a). $\Lambda_m^c = \kappa \times 1000 / C$ $= 3.905 \times 10^{-5} \times 1000 / 0.001$ $= 39.05 \text{ S cm}^2/\text{mole}$ $\text{CH}_3\text{COOH} \rightarrow \text{CH}_3\text{COO}^- + \text{H}^+$ $\Lambda^0 \text{CH}_3\text{COOH} = \Lambda^0 \text{CH}_3\text{COO}^- + \Lambda^0 \text{H}^+$ $= 40.9 + 349.6$ $\Lambda^0 \text{CH}_3\text{COOH} = 390.5 \text{ S cm}^2/\text{mol}$ $\alpha = \frac{\Lambda_m}{\Lambda_m^0}$ $= 39.05 / 390.5$ $= 0.1$ (b). Device used for the production of electricity from energy released during spontaneous chemical reaction and the use of electrical energy to bring about a chemical change. The reaction gets reversed / It starts acting as an electrolytic cell & vice – versa.	½ 1 ½ 1 1 1

25.	<p>(a) (i) Ability of oxygen to form multiple bond . (ii) Due to lanthanoid contraction. (iii) Due to variable oxidation state/unpaired electrons</p> <p>(b) (i) $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \rightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$ (ii) $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{I}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 3\text{I}_2$</p>	1 1 1 1 1
	OR	
25	<p>(i) Zn , because of not having partially filled d-orbital in its ground state or ionic state. (ii) Cr (iii) Cu (iv) Mn , because Mn^{+2} has extra stability due to half filled d-orbital</p>	$\frac{1}{2}+1$ 1 1 $\frac{1}{2}+1$
26.	<p>a). A : CH_3CHO B : $\begin{array}{c} \text{OH} \\ \\ \text{CH}_3-\text{CH}-\text{CH}_2\text{CHO} \end{array}$</p> <p>C : $\text{CH}_3-\text{CH}=\text{CH}-\text{CHO}$ D : $\begin{array}{c} \text{OH} \\ \\ \text{CH}_3-\text{CH}-\text{CN} \end{array}$</p> <p>b) i) Heat both the compounds with NaOH and I_2, $\text{C}_6\text{H}_5-\text{CH}=\text{CH}-\text{COCH}_3$ gives yellow ppt of iodoform while $\text{C}_6\text{H}_5-\text{CH}=\text{CH}-\text{COCH}_2\text{CH}_3$ does not. ii) Add ammoniacal silver nitrate solution (Tollens' reagent) , HCOOH gives silver mirror while $\text{CH}_3\text{CH}_2\text{COOH}$ does not. c) $\text{CH}_3\text{COCH}_3 < \text{CH}_3\text{CH}_2\text{OH} < \text{CH}_3\text{COOH}$</p>	$\frac{1}{2}$ $\times 4 = 2$ 1 1 1
	OR	
26	<p>a.)</p> <p></p> <p>b) $\text{C}_6\text{H}_5\text{COCH}_3 < \text{CH}_3\text{CHO} < \text{HCHO}$ c) stronger -I effect of Cl , stronger acid less pK_a / strong electron withdrawing power of Cl. d) $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}-\text{CH}_2\text{CHO}$ e) A: CH_3COCH_3 B: $\text{CH}_3\text{CH}_2\text{CHO}$</p>	1 1 1 1 $\frac{1}{2}$ $\frac{1}{2}$