


CBSE Class 12 Chemistry Question Paper Solution 2016



MARKING SCHEME 2016- CHEMISTRY(043)





Set 56 / 1/S




Q. No.	Value Points	Marks
1	$\text{CH}_2=\text{CH}-\text{CH}_2\text{Cl}$	1
2	NO_2	1
3	Anti-ferromagnetism	1
4	2,4-dibromoaniline / 2,4-dibromobenzenamine	1
5	Like Charged particles cause repulsion/ Brownian motion/ solvation	1
6	(i) Mercury cell (ii) Fuel cell (iii) Lead storage cell (iv) Dry cell	$4 \times \frac{1}{2} = 2$
7	(i) A : $\text{K}_2\text{MnO}_4 / \text{MnO}_4^{2-}$, B : $\text{KMnO}_4 / \text{MnO}_4^-$, (ii) On heating it decomposes forming K_2MnO_4 and oxygen gas OR $2\text{KMnO}_4 \longrightarrow \text{K}_2\text{MnO}_4 + \text{MnO}_2 + \text{O}_2$	$\frac{1}{2} + \frac{1}{2}$ 1
8	(i) $[\text{Pd}(\text{NH}_3)_4]\text{Cl}_2$ (ii) Tetraamminepalladium(II) chloride	1 1
9	(i) Order is zero, and molecularity is two / one . (ii) $\text{mol L}^{-1} \text{ s}^{-1}$	$\frac{1}{2} + \frac{1}{2}$ 1
10	(i) $\text{CH}_3\text{CHO} \xrightarrow{\text{Zn-Hg/HCl}} \text{CH}_3\text{CH}_3 + \text{H}_2\text{O}$ (ii) $\text{R}-\text{CH}_2-\text{COOH} \xrightarrow[\text{(ii) H}_2\text{O}]{\text{(i) X}_2 / \text{Red P.}} \text{R}-\underset{\substack{ \\ \text{X (X = Cl or Br)}}}{\text{CH}}-\text{COOH}$ (any other correct examples)	1 1
OR		
10	(i) $\text{C}_6\text{H}_5-\text{CH}_3 + \text{CrO}_2\text{Cl}_2 \xrightarrow{\text{CS}_2} \text{C}_6\text{H}_5-\text{CH}(\text{OCrOHCl}_2)_2 \xrightarrow{\text{H}_3\text{O}^+} \text{C}_6\text{H}_5-\text{CHO}$ (ii) $\text{CH}_3-\text{COCl} \xrightarrow{\text{H}_2 / \text{Pd}, \text{BaSO}_4} \text{CH}_3-\text{CHO} + \text{HCl}$ (any other correct method)	1 1
11	M x z	

	$d = \frac{\quad}{a^3 \times N_A}$ $N_A = (M \times z) / a^3 \times d = (280\text{g} \times 4) / (400 \times 10^{-10}\text{cm})^3 \times 7\text{gcm}^{-3}$ $= 2.5 \times 10^{24} \text{ atoms}$ <p style="text-align: right;">(or any other correct method)</p>	<p>1</p> <p>1</p> <p>1</p>
12	$\log k = \log A - E_a / 2.303RT ; \quad \log k = 14.2 - (1.0 \times 10^4 \text{K}) / T$ $\frac{E_a}{2.303RT} = \frac{1.0 \times 10^4 \text{K}}{T}$ $E_a = 2.303 \times 8.314 \text{ J mol}^{-1} \text{K}^{-1} \times 1.0 \times 10^4 \text{K}$ $E_a = 19.15 \times 10^4 \text{ J mol}^{-1} = 191.5 \text{ kJ mol}^{-1}.$ $\text{Rate constant, } k = 0.693 / t_{1/2} = 0.693 / 200 \text{ min}$ $= 0.0034 \text{ min}^{-1} / 3.4 \times 10^{-3} \text{ min}^{-1}$	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p>
13	<p>(i) Silica gel</p> <p>(ii) H_3PO_4 is more effective in causing coagulation because of greater negative charge / Hardy Schulze Rule .</p> <p>(iii) Proteins</p>	<p>1</p> <p>1/2 + 1/2</p> <p>1</p>
14	<p>(i) van Arkel method</p> <p>(ii) Leaching / Bayer's Process</p> <p>(iii) Limestone decomposes to CaO (flux) which removes silica impurity as slag.</p> $\begin{array}{ccc} \text{CaCO}_3 & \xrightarrow[\text{CaO} + \text{SiO}_2]{\text{Or } \Delta} & \text{CaO} + \text{CO}_2 \\ & \longrightarrow & \text{CaSiO}_3 \\ & & \text{Slag} \end{array}$	<p>1</p> <p>1</p> <p>1</p>
15	$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.0591}{6} \log \frac{[\text{Cr}^{3+}]^2}{[\text{Fe}^{2+}]^3}$ $0.261 \text{ V} = E_{\text{cell}}^{\circ} - \frac{0.0591}{6} \log \frac{(0.01)^2}{\quad}$	<p>1</p> <p>1</p>

	$0.261 \text{ V} = E_{\text{cell}}^{\circ} - \frac{0.0591}{6} \log (10^2)$ $0.261 \text{ V} = E_{\text{cell}}^{\circ} - (0.0591 / 6) \times 2$ $E_{\text{cell}}^{\circ} = E_{\text{cell}} + 0.0197 \text{ V} = 0.2807 \text{ V}$	1
16	<p>(i) Due to multiple bonding ability of Oxygen with transition Metals / $p\pi$- $d\pi$ bonding.</p> <p>(ii) Due to absence of unpaired electrons in zinc atom and the presence of unpaired electrons in Chromium atom.</p> <p>(iii) Eu^{2+} gets oxidized to more stable +3 state.</p>	1 1 1
17	<p>(i) </p> <p>(ii) $\text{CH}_3\text{-}\underset{\text{Cl}}{\text{CH}}\text{-CH}_2\text{-CH}_3 \xrightarrow{\text{alc. KOH}} \text{CH}_3\text{-CH=CH-CH}_3$</p> <p>(iii) $\text{CH}_3\text{-CH}_2\text{-Cl} + \text{Na} \xrightarrow{\text{dry ether}} \text{CH}_3\text{-CH}_2\text{-CH}_2\text{-CH}_3 + \text{NaCl}$</p>	1 1 1
18	<p>(a) In ketones presence of two electron releasing alkyl groups reduce the electrophilicity of the carbonyl group more effectively than in aldehydes wherein only one alkyl group occurs / Presence of two alkyl groups in ketones provide more steric hinderance to incoming nucleophile than in aldehydes where only one alkyl group occurs.</p> <p>(b) Due to the absence of alpha hydrogen.</p> <p>(c) Because the carboxyl group is deactivating and the Lewis acid AlCl_3 gets bonded to the the carboxyl group.</p>	1 1 1
19	<p>(i) A : $\text{C}_6\text{H}_5\text{CONH}_2$; B : $\text{C}_6\text{H}_5\text{NH}_2$; C : $\text{C}_6\text{H}_5\text{NHCOCH}_3$</p> <p>(ii) A : $\text{C}_6\text{H}_5\text{NO}_2$; B : $\text{C}_6\text{H}_5\text{NH}_2$; C : $\text{C}_6\text{H}_5\text{NC}$.</p>	$\frac{1}{2} + \frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2} + \frac{1}{2}$

20	(i) It acts as initiator of free radical / catalyst. (ii) $\text{CH}_2\text{OH}-\text{CH}_2\text{OH}$ and  or Ethylene glycol and phthalic acid / IUPAC name. (iii) Buna-N < PVC < Nylon-6	1 1 1
	OR	
20	<p><i>Chain initiation steps</i></p> $\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}\text{C}-\text{O}-\overset{\text{O}}{\parallel}\text{C}-\text{C}_6\text{H}_5 \longrightarrow 2\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}\text{C}-\dot{\text{O}} \longrightarrow 2\dot{\text{C}}_6\text{H}_5$ <p style="text-align: center;">Benzoyl peroxide Phenyl radical</p> $\dot{\text{C}}_6\text{H}_5 + \text{CH}_2=\text{CH}_2 \longrightarrow \text{C}_6\text{H}_5-\text{CH}_2-\dot{\text{C}}\text{H}_2$ <p><i>Chain propagating step</i></p> $\text{C}_6\text{H}_5-\text{CH}_2-\dot{\text{C}}\text{H}_2 + \text{CH}_2=\text{CH}_2 \longrightarrow \text{C}_6\text{H}_5-\text{CH}_2-\text{CH}_2-\text{CH}_2-\dot{\text{C}}\text{H}_2$ \downarrow $\text{C}_6\text{H}_5-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\dot{\text{C}}\text{H}_2$ <p><i>Chain terminating step</i></p> <p>For termination of the long chain, these free radicals can combine in different ways to form polythene. One mode of termination of chain is shown as under:</p> $\text{C}_6\text{H}_5-\text{CH}_2-\text{CH}_2-\text{CH}_2-\dot{\text{C}}\text{H}_2 + \text{C}_6\text{H}_5-\text{CH}_2-\dot{\text{C}}\text{H}_2-\text{CH}_2-\text{CH}_2-\dot{\text{C}}\text{H}_2 \longrightarrow \text{C}_6\text{H}_5-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{C}_6\text{H}_5$ <p style="text-align: center;">Polythene</p>	1 1 1
21	(i) α -D-Glucose and α -D-Glucose / Glucose and Glucose. (ii) Vitamin-B ₆ / Pyridoxine. (iii) Fibrous protien : Keratin / Myosin / Kephalin Globular protien : Insulin / Albumin / Haemoglobin (or any other one)	1 1 1/2 1/2
22	(a) sp^3d^2 hybridisation ; Paramagnetic ; High spin complex. (b) 	1 + 1/2 + 1/2 1
23	i) Aware, concerned or any other correct two values. (ii) Side effects, unknown health problems (iii) Neurologically active drugs/ stress relievers example- valium, equanil (or any other correct two examples)	1/2 + 1/2 1 1 1/2 + 1/2
24	(a) (i) Due to decrease in bond enthalpy from H_2S to H_2Te / Larger H-Te bond than H-S bond allowing more dissociation of H_2Te . (ii) +5 oxidation state of P in PCl_5 makes it more covalent/ high charge to size ratio.	1 1

	<p>(iii) Interhalogen compounds are slightly polar having dipole-dipole forces but pure halogens non-polar, have weak van der Waals forces.</p> <p>(b) (i)  (ii) </p>	<p>1</p> <p>1+1</p>
	OR	
24	<p>(i) $\text{H}_3\text{PO}_4 < \text{H}_3\text{PO}_3 < \text{H}_3\text{PO}_2$</p> <p>(ii) Xe ; Lower ionization enthalpy of Xe than He.</p> <p>(iii) High pressure , optimum temperature , Use of catalyst</p> <p>(iv) For bleaching woodpulp / cotton / textiles/ Extraction of gold / Platinum/ Manufacture of dyes/ drugs/ CHCl_3/ CCl_4/ DDT/ Sterilising water, etc (or any other two uses)</p> <p>(v) SO_2 decolourises acidified dilute solution of KMnO_4 / changes orange color of acidified $\text{K}_2\text{Cr}_2\text{O}_7$ to green.</p>	<p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>1</p>
25	<p>(a) (i) $\text{CH}_2 = \text{CH} - \text{CHO}$</p> <p>(ii) C_6H_6 / </p> <p>(iii) </p> <p>(b) _____</p> <p>(i) $\text{CH}_3 - \text{CH}_2 - \ddot{\text{O}} - \text{H} + \text{H}^+ \rightarrow \text{CH}_3 - \text{CH}_2 - \overset{\text{H}}{\underset{\cdot\cdot}{\text{O}}}^+ - \text{H}$</p> <p>(ii) $\text{CH}_3\text{CH}_2 - \ddot{\text{O}}: + \text{CH}_3 - \text{CH}_2 - \overset{\text{H}}{\underset{\cdot\cdot}{\text{O}}}^+ - \text{H} \rightarrow \text{CH}_3\text{CH}_2 - \overset{\text{H}}{\underset{\cdot\cdot}{\text{O}}}^+ - \text{CH}_2\text{CH}_3 + \text{H}_2\text{O}$</p> <p>(iii) $\text{CH}_3\text{CH}_2 - \overset{\text{H}}{\underset{\cdot\cdot}{\text{O}}}^+ - \text{CH}_2\text{CH}_3 \rightarrow \text{CH}_3\text{CH}_2 - \text{O} - \text{CH}_2\text{CH}_3 + \text{H}^+$</p>	<p>1</p> <p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$</p>
	OR	
25	(a) (i)	

	 <p>(ii)</p>  <p>(iii)</p> 	1
	<p>(b) (i) On treatment with acetic acid or acetic anhydride in presence of drops of H_2SO_4, ethanol gives pleasant smell but Diethyl ether does not.</p> <p>(ii) On treatment with anhy. ZnCl_2 and HCl, ter-butyl alcohol gives immediate turbidity but Propanol does not. (or any other correct test)</p>	1 1
26	<p>(a)</p> $\Delta T_b = i K_b m$ $\Delta T_b = i K_b \frac{w_b \times 1000}{M_b \times w_a}$ $T_b - T_b^0 = \frac{3 \times 0.52 \text{ kg/mol} \times 2 \times 1000 \text{ g kg}^{-1}}{142 \text{ g/mol} \times 50 \text{ g}}$ $T_b - 373 \text{ K} = 0.44 \text{ K} ; T_b = 373.44 \text{ K} / 100.44^\circ\text{C}$ <p>(b) (i) Properties of dilute solutions that depend on the number of particles of solute but not on nature of the solute particles are called colligative properties.</p> <p>(ii) The solutions which obey Raoult's law over the entire range of concentration are known as ideal solutions.</p>	1 1 1 1
	OR	
26	<p>(a)</p> $\Delta T_f = K_f m; \text{ or } M_B = K_f (w_B \times 1000) / (\Delta T_f \times w_A)$ $M_B = (3.83 \times 2.56 \times 1000) / (0.383 \times 100)$ $= 256 \text{ g mol}^{-1}$ <p>Atomicity = $256 / 32 = 8$</p> <p>Formula of Sulphur = S_8.</p>	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1

	(b) (i) Shrinks (ii) Swells	1 1
--	--------------------------------	--------

