





CBSE Class 12 Chemistry Question Paper Solution 2019


MARKING SCHEME- 2019




CHEMISTRY (043) CLASS XII

56/5/1




Q.No.	Expected answers/ Value points	Marks
SECTION A		
1	A_3B_2	1
2	$As_2O_3 + 3H_2S \rightarrow As_2S_3(sol) + 3H_2O$ / By Double decomposition of Arsenic oxide with Hydrogen sulphide.	1
OR		
2	$N_2(g) + 3H_2(g) \xrightarrow{Fe(s)} 2NH_3(g)$ or any other suitable example.	1
3	Steric reason / Electronic reason/ Inductive effect	1
4	Mixture of amines including quaternary ammonium salts / RNH_2 , R_2NH , R_3N , $R_4N^+X^-$	1
5	Ionization isomerism	1
OR		
5	$[Cr(Cl)_6]^{3-} < [Cr(NH_3)_6]^{3+} < [Cr(CN)_6]^{3-}$	1
SECTION B		
6	At cathode- $H_2(g)$ is produced due to greater E° value of H^+ ion At anode- $Cl_2(g)$ is produced due to over voltage /over potential of oxygen	1 1
7	$t = (2.303/k) \log [R]_0 / [R]$ Let $[R]_0 = 100$ For 99% completion reaction- $t_{99\%} = (2.303/k) \log (100/1)$ $k = (2.303 \times 2) / t_{99\%}$ For 90% completion $t_{90\%} = (2.303/k) \log (100/10)$ Putting the value of k $t_{90\%} = (2.303 \times t_{99\%} \times \log 10) / 2.303 \times 2$ $2 \times t_{90\%} = t_{99\%}$ (Or any other suitable method)	½ ½ ½ ½
8	$H_2S < H_2Se < H_2Te < H_2O$ From H_2S to H_2Te boiling point increases as magnitude of van der Waal force increases. The boiling point of water is maximum due to H-bonding H_2O / water	1 ½ ½

9	<p>1. When pyrolusite is fused with KOH in presence of air or oxidizing agent , potassium manganate is produced . $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \rightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$</p> <p>2. Potassium manganate upon further oxidation or disproportionation in a neutral or acidic medium gives potassium permanganate.</p> <p style="text-align: center;">$3\text{MnO}_4^{2-} + 4\text{H}^+ \rightarrow 2\text{MnO}_4^- + \text{MnO}_2 + 2\text{H}_2\text{O} / \text{MnO}_4^{2-} \xrightarrow[\text{alkaline solution}]{\text{Electrolytic oxidation in}} \text{MnO}_4^-$</p> <p style="text-align: center;">(Award full marks if only correct reactions are given)</p>	1 1						
10	<p style="text-align: center;">  </p> <p style="text-align: center;">(Full marks may be awarded if only correct names are given)</p>	$\frac{1}{2} \times 4$						
OR								
10	<p>a)</p> <p style="text-align: center;">  </p> <p>b)</p> <p style="text-align: center;"> $\text{CH}_3\text{CH}_2\text{OH} \xrightarrow[\Delta]{\text{PCC}} \text{CH}_3\text{CHO}$ </p> <p style="text-align: center;">(Or any other suitable method)</p>	1 1						
11	<p>a) n-hexane is formed</p> <p style="text-align: center;">  </p> <p>b) Saccharic acid is formed</p> <p style="text-align: center;">  </p> <p style="text-align: center;">(Award full marks if correct reactions are given)</p>	1 1						
OR								
11	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%; text-align: center;">Fibrous</th> <th style="width: 50%; text-align: center;">Globular</th> </tr> </thead> <tbody> <tr> <td>i)The polypeptide chains run parallel and are held together by hydrogen and disulphide bonds.</td> <td>The polypeptide chains coil around to give a spherical shape.</td> </tr> <tr> <td>ii)Insoluble in water</td> <td>Soluble in water</td> </tr> </tbody> </table> <p style="text-align: center;">(Or any other correct difference)</p>	Fibrous	Globular	i)The polypeptide chains run parallel and are held together by hydrogen and disulphide bonds.	The polypeptide chains coil around to give a spherical shape.	ii)Insoluble in water	Soluble in water	1+1
Fibrous	Globular							
i)The polypeptide chains run parallel and are held together by hydrogen and disulphide bonds.	The polypeptide chains coil around to give a spherical shape.							
ii)Insoluble in water	Soluble in water							
12	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%; text-align: center;">DNA</th> <th style="width: 50%; text-align: center;">RNA</th> </tr> </thead> <tbody> <tr> <td>i)Deoxyribose sugar</td> <td>Ribose sugar</td> </tr> <tr> <td>ii)contain bases ATGC</td> <td>Contain bases AUGC</td> </tr> </tbody> </table>	DNA	RNA	i)Deoxyribose sugar	Ribose sugar	ii)contain bases ATGC	Contain bases AUGC	1+1
DNA	RNA							
i)Deoxyribose sugar	Ribose sugar							
ii)contain bases ATGC	Contain bases AUGC							

	(Or any other correct difference)	
SECTION C		
13	a) Ferromagnetism b) let the number of O^{2-} ions be 100 then total number of $Ni^{2+} = 98$ Let the number of $Ni^{2+} = x$ Then number of $Ni^{3+} = 0.98 - x$ Total charge of cation = Total charge of anion $2x + 3(0.98 - x) = 2$ $x = 0.94$ Fraction of $Ni^{2+} = 0.94/0.98 = 0.96$ Fraction of $Ni^{3+} = 1 - 0.96 = 0.04$	1 1 $\frac{1}{2}$ $\frac{1}{2}$
14	$n = 6$ $E^{\circ} = E^{\circ}_{\text{cathode}} - E^{\circ}_{\text{anode}}$ $= (-0.25) - (-1.66)$ $= 1.41 \text{ V}$ $E_{\text{cell}} = E^{\circ}_{\text{cell}} - (0.059/n) \log [Al^{+3}]^2 / [Ni^{+2}]^3$ $E_{\text{cell}} = 1.41 - (0.059/6) \log (10^{-3})^2 / (0.1)^3$ $E_{\text{cell}} = 1.4395 \text{ V}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1 $\frac{1}{2}$
15	For experiment II $[A] = 0.2 \text{ mol L}^{-1}$ For Experiment III Rate = $0.08 \text{ mol L}^{-1} \text{ min}^{-1}$ For Experiment IV $[A] = 0.1 \text{ mol L}^{-1}$	1 1 1
16	a) Negative sol is formed Due to adsorption of I^{-} from dispersion medium b)  As extent of adsorption decreases with increase in temperature c) 'A'; With higher critical temperature, it will liquify more easily	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
17	a) The ore particles are wetted with oil, while gangue particles are wetted by water / Preferential wettability b) The impurities are more soluble in the melt than in the solid state of metal c) The metal forms a volatile compound with a suitable reagent. The volatile compound is easily decomposed on heating.	1 1 1
OR		
17	a) Cast iron contain 3% carbon while pig iron contain 4% carbon b) Hydraulic washing is the method of concentration of an ore while liquation is the method of refining of metal (or any other suitable difference) c) Leaching is the method of concentration of an ore while roasting is the method used to convert a sulphide ore to oxide	1 1 1

	(or any other suitable difference)	
18	a) At + 3, Stable d^0 is obtained b) Absence of unpaired electron / no d-d transition occurs c) MnO has Mn in +2 Oxidation State Mn ₂ O ₇ has Mn in +7 Oxidation State . Higher the Oxidation State , Higher is the acidic character.	1 1 1
	OR	
18	a) $4f^{1-14} 5d^{0-1} 6s^2$ b) +3 and +4 c) 5f 6d 7s orbitals/levels are of comparable energies	1 1 1
19	a) d^2sp^3 , diamagnetic b) sp^3 , diamagnetic	$1 + \frac{1}{2}$ $1 + \frac{1}{2}$
20	a) The stereoisomers related to each other as non superimposable mirror images are called enantiomers. b) Equimolar mixture of d- and l- form is known as racemic mixture c) Resonance effect / difference in hybridisation of carbon atom in C – X bond / instability of phenyl cation /partial double bond character	1 1 1
21	a) Add neutral FeCl ₃ to both the compounds. Phenol gives violet colouration while, 1 – propanol does not. b) Add I ₂ / NaOH(aq) to both the compounds, Ethanol gives yellow precipitate while ether does not c) Add HCl and ZnCl ₂ to both the compounds, 2-methyl-2-propanol gives turbidity immediately while 1-propanol does not (Or any other suitable chemical test)	1 1 1
	OR	
21	a) CH ₃ -CH ₂ OH + CH ₃ I b)  c) 	1 1 1
22	a) Due to resonance stabilisation of arene diazonium ion /  b) Methyl amine being basic, gains a proton from water and releases hydroxyl ions which precipitate hydrated ferric oxide.	$1 + \frac{1}{2}$ $1 + \frac{1}{2}$

	$\text{CH}_3\text{NH}_2 \xrightarrow{\text{H}_2\text{O}} \text{CH}_3\text{NH}_3^+ + \text{OH}^-$ $\text{FeCl}_3 \xrightarrow{3\text{OH}^-} \text{Fe}(\text{OH})_3 \downarrow + 3\text{Cl}^-$	
	OR	
22	a) p-nitroaniline > aniline > p-toluidine pK_b decreases, basicity increases / due to inductive effect / EWG $-\text{NO}_2$ and EDG $-\text{CH}_3$ group b) $\text{C}_2\text{H}_5\text{NH}_2 > (\text{C}_2\text{H}_5)_2\text{NH} > (\text{C}_2\text{H}_5)_3\text{N}$ Due to increase in number of ethyl groups, inductive effect increases	1 $\frac{1}{2}$ 1 $\frac{1}{2}$
23	a) Homopolymer as it is formed only from one monomer. b) Addition polymer as it is formed by addition of monomeric units. c) Bakelite is a thermosetting plastic due to extensive crosslinking.	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
24	a) 2-3% solution of iodine in alcohol – water mixture It is an antiseptic / is applied on wounds b) Chloroxylenol and terpineol c) $\text{CH}_3(\text{CH}_2)_6$ - Hydrophobic ; $-\text{COO}(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_2\text{CH}_2\text{OH}$ - Hydrophilic	$\frac{1}{2}$ $\frac{1}{2}$ 1 $\frac{1}{2} + \frac{1}{2}$
	SECTION D	
25	a) <div style="text-align: center;"> </div> <p>When a solute is added to a solvent, the vapour pressure of the solvent decreases and it becomes equal to atmospheric pressure at a higher temperature.</p> b) $i = 3$ $\pi = i CRT$ $\pi = 3 \times 0.025 \times 0.0821 \times 298/174 \times 2$ $\pi = 5.27 \times 10^{-3} \text{ atm.}$ (Deduct half mark if unit is not given or wrong)	1+1 $\frac{1}{2}$ $\frac{1}{2}$ 1 1
	OR	
25	a) Solution does not obey Raoult's law over the entire range of concentration, $\Delta H_{\text{mix}} \neq 0$ (or any two suitable characteristics) b) $n = 2$ (dimer) $\Delta T_f = i K_f m$ $i = 1.62 \times 122 \times 25 / 4.9 \times 2 \times 1000$ $i = 0.504$ $\alpha = 2(1-i)$ $\alpha = 2(1 - 0.504)$	1 + 1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

	<p>= 0.992 = 99.2%</p> <p>(Or any other suitable method)</p>	½
26	<p>a) Due to the presence of an unpaired / unbonded / odd electron on nitrogen atom</p> <p>b) Bleaching effect of chlorine is due to oxidation / oxidation is caused by nascent oxygen released by reaction of Cl₂ and H₂O / $\text{Cl}_2 + \text{H}_2\text{O} \rightarrow 2\text{HCl} + [\text{O}]$ Coloured substance + [O] → colourless substance</p> <p>c) Due to small size of oxygen, the added electron suffers inter electronic repulsion.</p> <p>d) Unavailability of d-orbital in Fluorine / Due to high electronegativity and small size.</p> <p>e) Due to weak dispersion forces.</p>	1 1 1 1 1
OR		
26	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>a) i) See – saw ii) Bent T</p> </div> <div style="text-align: center;">  <p>ii) Bent T</p> </div> </div> <p>b) i) It forms chloride and chlorate. $6 \text{NaOH} + 3 \text{Cl}_2 \rightarrow 5 \text{NaCl} + \text{NaClO}_3 + 3 \text{H}_2\text{O}$</p> <p>ii) On complete hydrolysis XeO₃ is formed. $\text{XeF}_6 + 3 \text{H}_2\text{O} \rightarrow \text{XeO}_3 + 6 \text{HF}$</p> <p>iii) Charring of cane sugar takes place. $\text{C}_{12}\text{H}_{22}\text{O}_{11} + \text{H}_2\text{SO}_4 \rightarrow 12 \text{C} + 11 \text{H}_2\text{O}$</p> <p style="text-align: right;">(Balancing may be ignored)</p>	½ + ½ ½ + ½ ½ + ½ ½ + ½ ½ + ½
27	<p>a) But-2-enal</p> <p>b) To both the compounds add Tollen's reagent, ethanal gives silver mirror while ethanol does not (Or any other correct chemical test)</p> <p>c) i)</p> <div style="text-align: center;">  </div> <p>ii)</p> $\text{CH}_3\text{CH}_2\text{OH} \xrightarrow{\text{PCC, Heat}} \text{CH}_3\text{CHO} \xrightarrow[\text{ii) H}_3\text{O}^+]{\text{i) CH}_3\text{MgBr}} \text{CH}_3\text{CH}(\text{OH})\text{CH}_3$ <p>iii)</p> $\text{CH}_3\text{-CH}_2\text{-CHO} \xrightarrow[\text{H}^+]{\text{KMnO}_4} \text{CH}_3\text{-CH}_2\text{-COOH} \xrightarrow[\text{(2) NaOH (aq)}]{\text{(1) Cl}_2 / \text{P}_4} \text{CH}_3\text{-CH}(\text{OH})\text{-COOH}$	1 1 1 1 1
OR		
27	a) 2-Hydroxy benzoic acid	1

b) Due to Inductive effect of Cl^- / presence of EWG(Cl^-)	1
c) i) $(\text{CH}_3)_3\text{C}-\text{CH}_3$	1
ii) $(\text{CH}_3)_3\text{C}-\text{CH}_2\text{OH}$ and $(\text{CH}_3)_3\text{C}-\text{COONa}$	1
iii) $(\text{CH}_3)_3\text{C}-\text{CH}=\text{NNHCONH}_2$	1

