

# CBSE Class 12 Maths Question Paper Solution 2016

65/1/N

QUESTION PAPER CODE 65/1/N

## EXPECTED ANSWER/VALUE POINTS

### SECTION A

1. Finding  $A^T = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$   $\frac{1}{2}$   
 Getting  $\alpha = \frac{\pi}{4}$  or  $45^\circ$   $\frac{1}{2}$
  
2.  $k = 27$  1
  
3. For a unique solution  $\frac{1}{2}$   

$$\begin{vmatrix} 1 & 1 & 1 \\ 2 & 1 & -1 \\ 3 & 2 & k \end{vmatrix} \neq 0$$
  
 $\Rightarrow k \neq 0$   $\frac{1}{2}$
  
4. Getting equation as  $\frac{x}{\frac{5}{2}} + \frac{y}{5} + \frac{z}{-5} = 1$   $\frac{1}{2}$   
 Sum of intercepts  $\frac{5}{2} + 5 - 5 = \frac{5}{2}$   $\frac{1}{2}$
  
5. Getting  $\lambda = -9$  and  $\mu = 27$   $\frac{1}{2}$  each
  
6.  $\vec{a} + \vec{b} = 6\hat{i} - 3\hat{j} + 2\hat{k}$   $\frac{1}{2}$   
 Unit vector parallel to  $\vec{a} + \vec{b}$  is  $\frac{1}{7}(6\hat{i} - 3\hat{j} + 2\hat{k})$   $\frac{1}{2}$

### SECTION B

7.  $\tan^{-1}(x-1) + \tan^{-1}(x+1) = \tan^{-1} 3x - \tan^{-1} x$   $\frac{1}{2}$   
 $\Rightarrow \tan^{-1}\left(\frac{2x}{2-x^2}\right) = \tan^{-1}\left(\frac{2x}{1+3x^2}\right)$   $\frac{1}{2}$   

$$\frac{2x}{2-x^2} = \frac{2x}{1+3x^2}$$
  $\frac{1}{2}$   
 $2x(1+3x^2-2+x^2) = 0$   $\frac{1}{2}$   
 $x = 0, \frac{1}{2}, -\frac{1}{2}$  1

OR

$$\text{Let } 2x = \tan \theta \quad 1$$

$$\text{L. H. S} = \tan^{-1} \left( \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \right) - \tan^{-1} \left( \frac{2 \tan \theta}{1 - \tan^2 \theta} \right) \quad 1$$

$$= \tan^{-1}(\tan 3\theta) - \tan^{-1}(\tan 2\theta) \quad 1$$

$$= 3\theta - 2\theta$$

$$= \theta \text{ or } \tan^{-1} 2x$$

$$\therefore \text{L. H. S} = \text{R. H. S} \quad 1$$

$$8. \text{ Getting matrix equation as } \begin{pmatrix} 10 & 3 \\ 3 & 10 \end{pmatrix} \begin{pmatrix} E \\ H \end{pmatrix} = \begin{pmatrix} 145 \\ 180 \end{pmatrix} \quad 1 \frac{1}{2}$$

$$\Rightarrow \begin{pmatrix} E \\ H \end{pmatrix} = \begin{pmatrix} 10 & 3 \\ 3 & 10 \end{pmatrix}^{-1} \begin{pmatrix} 145 \\ 180 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} E \\ H \end{pmatrix} = \begin{pmatrix} 10 \\ 15 \end{pmatrix}$$

$$\Rightarrow E = 10, H = 15 \quad 1 \frac{1}{2}$$

The poor boy was charged ₹ 65 less

Value: Helping the poor

$$9. \text{ L.H.L} = a + 3 \quad 1 \frac{1}{2}$$

$$\text{R.H.L} = b/2 \quad 1 \frac{1}{2}$$

$$f(x) \text{ is continuous at } x = 0. \text{ So, } a + 3 = 2 = b/2 \quad 1 \frac{1}{2}$$

$$\Rightarrow a = -1 \text{ and } b = 4 \quad 1 \frac{1}{2}$$

$$10. \frac{dx}{dy} = \frac{\sin a}{\cos^2(a+y)} \quad 1 \frac{1}{2}$$

$$\Rightarrow \frac{dy}{dx} = \frac{\cos^2(a+y)}{\sin a} \quad 1 \frac{1}{2}$$

$$\frac{d^2y}{dx^2} = \frac{-2\cos(a+y) \sin(a+y)}{\sin a} \frac{dy}{dx}$$

$$= \frac{-\sin 2(a+y)}{\sin a} \frac{dy}{dx} \quad 1 \frac{1}{2}$$

$$\Rightarrow \sin a \frac{d^2y}{dx^2} + \sin 2(a+y) \frac{dy}{dx} = 0 \quad 1 \frac{1}{2}$$

**OR**

Let  $2x = \sin \theta$

1

$$\therefore y = \sin^{-1} \left( \frac{6x - \sqrt{1 - 4x^2}}{5} \right)$$

$$= \sin^{-1} \left( \frac{3}{5} \sin \theta - \frac{4}{5} \cos \theta \right)$$

$$= \sin^{-1} (\cos \alpha \sin \theta - \sin \alpha \cos \theta)$$

$$[\cos \alpha = \frac{3}{5}; \sin \alpha = \frac{4}{5}]$$

1

$$= \sin^{-1} (\sin(\theta - \alpha))$$

$$= \theta - \alpha$$

1

$$= \sin^{-1} (2x) - \alpha$$

$$\Rightarrow \frac{dy}{dx} = \frac{2}{\sqrt{1 - 4x^2}}$$

1

11. Slope of the tangent =  $3x^2 + 2 = 14$

1

Points of contact (2, 8) and (-2, -16)

1

Equations of tangent

$14x - y - 20 = 0$

1

and  $14x - y + 12 = 0$

1

12. Let  $2x = t$

$$I = \frac{1}{2} \int \frac{(t-5)}{(t-3)^3} e^t dt$$

1

$$= \frac{1}{2} \int \left[ \frac{1}{(t-3)^2} - \frac{2}{(t-3)^3} \right] e^t dt$$

2

$$= \frac{1}{2} \frac{1}{(t-3)^2} e^t + C = \frac{1}{2} \frac{1}{(2x-3)^2} e^{2x} + C$$

1

**OR**

Writing  $\frac{x^2 + x + 1}{(x^2 + 1)(x + 2)} = \frac{Ax + B}{x^2 + 1} + \frac{C}{x + 2}$

$$\Rightarrow A = \frac{2}{5}, B = \frac{1}{5}, C = \frac{3}{5}$$

1

$$\therefore I = \frac{1}{5} \int \frac{2x}{x^2 + 1} dx + \frac{1}{5} \int \frac{dx}{x^2 + 1} + \frac{3}{5} \int \frac{dx}{x + 2}$$

2

$$\Rightarrow I = \frac{1}{5} \log|x^2 + 1| + \frac{1}{5} \tan^{-1} x + \frac{3}{5} \log|x + 2| + C$$

1

13. Using property:  $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$  1

$$I = \int_{-2}^2 \left( \frac{x^2}{1+5^x} \right) dx = \int_{-2}^2 \left( \frac{x^2}{1+5^{-x}} \right) dx$$
 1

$$2I = \int_{-2}^2 x^2 dx$$
 1

$$2I = \frac{16}{3} \text{ or } I = \frac{8}{3}$$
 1

14. Writing  $x + 3 = A(-4 - 2x) + B$

$$\Rightarrow A = -\frac{1}{2}, B = 1$$
 1

$$\therefore I = -\frac{1}{2} \int (-4 - 2x) \sqrt{3 - 4x - x^2} dx + \int \sqrt{(\sqrt{7})^2 - (x+2)^2} dx$$
 1

$$I = -\frac{1}{3} (3 - 4 - x^2)^{3/2} + \frac{x+2}{2} \sqrt{3 - 4x - x^2} + \frac{7}{2} \sin^{-1} \frac{x+2}{\sqrt{7}} + C$$
 2

15. Writing linear equation  $\frac{dy}{dx} + \frac{\cos x}{1 + \sin x} y = -\frac{x}{1 + \sin x}$  1

$$I.F = e^{\int \frac{\cos x}{1 + \sin x} dx} = 1 + \sin x$$
 1

General solution is:  $y(1 + \sin x) = -\frac{x^2}{2} + C$  1

Particular solution is:  $y(1 + \sin x) = 1 - \frac{x^2}{2}$  1

16.  $\frac{dx}{dy} = \frac{2xe^v - y}{2ye^v}$

$$\frac{x}{y} = v, \text{ then } \frac{dx}{dy} = v + y \frac{dv}{dy}$$
 1

$$v + y \frac{dv}{dy} = \frac{2vye^v - y}{2ye^v}$$

$$2 \int e^v dv = - \int \frac{dy}{y}$$
 1

General solution is:  $2e^v = -\log |y| + C$  or  $2e^{x/y} = -\log |y| + C$  1

Particular solution is:  $2e^{x/y} + \log |y| = 2$  1

17.  $\overline{AB} = -4\hat{i} - 6\hat{j} - 2\hat{k}, \overline{AC} = -\hat{i} + 4\hat{j} + 3\hat{k}, \overline{AD} = -8\hat{i} - \hat{j} + 3\hat{k}$   $\frac{1}{2}$

For 4 points to be coplanar,  $[\overline{AB} \overline{AC} \overline{AD}] = 0$


$$\text{i.e., } \begin{vmatrix} -4 & -6 & -2 \\ -1 & 4 & 3 \\ -8 & -1 & 3 \end{vmatrix} = 0 \quad 1 \frac{1}{2}$$

$$= -4(12 + 3) + 6(-3 + 24) - 2(1 + 32)$$

$$= -60 + 126 - 66 = 0 \text{ which is true}$$

Hence, points are coplanar. 1

18.



$$\text{Equation of line BC: } \frac{x}{2} = \frac{y+1}{-2} = \frac{z-3}{-4} = r \quad 1$$

$$\text{General point on BC: } (2r, -2r-1, -4r+3)$$

$$\Rightarrow \text{d.r.'s of AP: } (2r+1, -2r-9, -4r-1) \quad 1$$

$$\text{As } AP \perp BC \Rightarrow r = -1$$

$$\Rightarrow \text{Co-ordinates of P: } (-2, 1, 7) \quad 1$$

$$\text{Hence, coordinates of Image of A: } (-3, -6, 10) \quad 1$$

19. Let  $E_1$  and  $E_2$  be the events of drawing bag X and bag Y respectively.

$$\text{Then, } P(E_1) = P(E_2) = \frac{1}{2} \quad 1 \frac{1}{2}$$

Let A be the event of drawing one white and one black ball from any one of the bag without replacement.

Then,

$$\Rightarrow P(A/E_1) = \frac{4}{6} \times \frac{2}{5} + \frac{2}{6} \times \frac{4}{5} = \frac{16}{30}$$

$$P(A/E_2) = \frac{3}{6} \times \frac{3}{5} + \frac{3}{6} \times \frac{3}{5} = \frac{18}{30} \quad 1 \frac{1}{2}$$

Using Bayes' Theorem, we have

$$P(E_2/A) = \frac{P(E_2) P(A/E_2)}{P(E_1) P(A/E_1) + P(E_2) P(A/E_2)} \quad 1$$

$$= \frac{\frac{1}{2} \times \frac{18}{30}}{\frac{1}{2} \times \frac{16}{30} + \frac{1}{2} \times \frac{18}{30}} = \frac{9}{17} \quad 1$$

**OR**

Let  $A_i$  and  $B_i$  be the events of throwing 10 by A and B in the respective  $i$ th turn. Then,

$$P(A_i) = P(B_i) = \frac{1}{12} \text{ and } P(\bar{A}_i) = P(\bar{B}_i) = \frac{11}{12} \quad 1 + \frac{1}{2}$$

Probability of winning A, when A starts first

$$= \frac{1}{12} + \left(\frac{11}{12}\right)^2 \frac{1}{12} + \left(\frac{11}{12}\right)^4 \frac{1}{12} + \dots \quad 1$$

$$= \frac{1/12}{1 - (11/12)^2}$$

$$= \frac{12}{23} \quad 1$$

$$\text{Probability of winning of B} = 1 - P(A) = 1 - \frac{12}{23} = \frac{11}{23} \quad 1 \frac{1}{2}$$

## SECTION C

20. The variate  $X$  takes values 3, 4, 5, and 6 2

$$P(X=3) = \frac{1}{20}; P(X=4) = \frac{3}{20}; P(X=5) = \frac{6}{20}; P(X=6) = \frac{10}{20}; \quad 1$$

Probability distribution is:

X	3	4	5	6
P(X)	$\frac{1}{20}$	$\frac{3}{20}$	$\frac{6}{20}$	$\frac{10}{20}$

1

$$\text{Mean} = \sum XP(X) = \frac{105}{20} = \frac{21}{4} \quad 1$$

$$\text{Variance} = \sum X^2P(X) - (\sum X P(X))^2 = \frac{63}{80} \quad 1$$

21. Proving \* is commutative  $1\frac{1}{2}$

Proving \* is associative  $1\frac{1}{2}$

Getting identity element as (0, 0)  $1\frac{1}{2}$

Getting inverse of (a, b) as (-a, -b)  $1\frac{1}{2}$

22. Getting  $\frac{dy}{d\theta} = \frac{\cos\theta(4 - \cos\theta)}{(2 + \cos\theta)^2}$  2

Equating  $\frac{dy}{d\theta}$  to 0 and getting critical point as  $\cos\theta = 0$  i.e.,  $\theta = \frac{\pi}{2}$  1

For all  $\theta, 0 \leq \theta \leq \frac{\pi}{2}, \frac{dy}{d\theta} \geq 0$  2

Hence,  $y$  is an increasing function of  $\theta$  on  $\left[0, \frac{\pi}{2}\right]$  1

OR



Correct Figure 1

$$\text{Writing } V = \frac{1}{3}\pi r^2 h = \frac{\pi}{2} l^3 \sin^2 \theta \cos \theta \quad 1$$

$$\text{Getting } \frac{dV}{d\theta} = \frac{\pi}{2} l^3 [2 \sin \theta \cos^2 \theta - \sin^3 \theta] \quad 1$$

$$\text{For maxima and minima, } \frac{dV}{d\theta} = 0 \quad 1\frac{1}{2}$$

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{3}} \text{ or } \theta = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right) \quad 1\frac{1}{2}$$


$$\text{Getting } \frac{d^2V}{d\theta^2} \text{ negative} \quad 1$$

Hence, volume of the cone is maximum when semi-vertical angle is  $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$

23.

Correct Figure

1



Writing equations of three sides in terms of y as

$$x_{AB} = \frac{2}{5}(y+2) + 2; \quad x_{BC} = 3(y-3) + 4; \quad x_{AC} = -\frac{1}{4}(y+2) + 2$$

$$\text{Area} = \int_{-2}^3 \left( \frac{2}{5}(y+2) + 2 \right) dy - \int_{-2}^2 \left( -\frac{1}{4}(y+2) + 2 \right) dy - \int_2^3 (3(y-3) + 4) dy$$

$$= \left[ \frac{2}{10}(y+2)^2 + 2y \right]_{-2}^3 - \left[ -\frac{1}{8}(y+2)^2 + 2y \right]_{-2}^2 - \left[ \frac{3}{2}(y-3)^2 + 4y \right]_2^3$$

$$= 15 - 6 - \frac{5}{2} \text{ or } \frac{13}{2}$$

24. Required equation of the plane is

$$\vec{r} \cdot [(1-2\lambda)\hat{i} + (-2+\lambda)\hat{j} + (3+\lambda)\hat{k}] = 4-5\lambda$$

Intercept of the plane on x-axis = Intercept of the plane on y-axis

$$\Rightarrow \frac{4-5\lambda}{1-2\lambda} = \frac{4-5\lambda}{\lambda-2} \text{ i.e., } \lambda = 1, \frac{4}{5} \left( \text{rejecting } \lambda = \frac{4}{5} \right)$$

Required equation of the plane is  $\vec{r} \cdot (-\hat{i} - \hat{j} + 4\hat{k}) + 1 = 0$ 

25.

Let the investment in bond A be ₹ x and in bond B be ₹ y


$$\text{Objective function is: } Z = \frac{x}{10} + \frac{9}{100}y$$

Subject to constraints

$$x + y \geq 50000; \quad x \geq 20,000; \quad y \geq 10,000; \quad x \geq y (*)$$

Correct Figure

Vertices of feasible region are A, B, C, and D



Point	$Z = \frac{x}{10} + \frac{9}{100}y$	Value
A(25,000, 25000)	$2500 + 2250$	4750
B(20,000, 20,000)	$2000 + 1800$	3800
C(20,000, 10,000)	$2000 + 900$	2900
D(40,000, 10,000)	$4000 + 900$	4900

Return is maximum when ₹ 40000 are invested in Bond A and ₹ 10000 in Bond B

Maximum return is ₹ 4900

Since there are more than 3 constraints, student may be given full 6 marks even if reaches upto (\*).

$$26. \Delta = \begin{vmatrix} (x+y)^2 & zx & zy \\ zx & (z+y)^2 & xy \\ zy & xy & (z+x)^2 \end{vmatrix}$$

$$R_1 \rightarrow zR_1, R_2 \rightarrow xR_2, R_3 \rightarrow yR_3$$

$$\Delta = \frac{1}{xyz} \begin{vmatrix} z(x+y)^2 & z^2x & z^2y \\ x^2z & x(z+y)^2 & x^2y \\ y^2z & y^2x & y(z+x)^2 \end{vmatrix} \quad 1$$

$$= \frac{xyz}{xyz} \begin{vmatrix} (x+y)^2 & z^2 & z^2 \\ x^2 & (z+y)^2 & x^2 \\ y^2 & y^2 & (z+x)^2 \end{vmatrix} \quad 1$$

$$C_1 \rightarrow C_1 - C_3 \text{ and } C_2 \rightarrow C_2 - C_3$$

$$\Delta = \begin{vmatrix} (x+y)^2 - z^2 & 0 & z^2 \\ 0 & (z+y)^2 - x^2 & x^2 \\ y^2 - (z+x)^2 & y^2 - (z+x)^2 & (z+x)^2 \end{vmatrix}$$

$$= (x+y+z)^2 \begin{vmatrix} x+y-z & 0 & z^2 \\ 0 & z+y-x & x^2 \\ y-z-x & y-z-x & (z+x)^2 \end{vmatrix} \quad 1$$

$$R_3 \rightarrow R_3 - R_1 - R_2 \text{ we get}$$

$$= (x+y+z)^2 \begin{vmatrix} x+y-z & 0 & z^2 \\ 0 & z+y-x & x^2 \\ -2x & -2z & 2xz \end{vmatrix} \quad 1$$

$$C_1 \rightarrow C_1 + \frac{C_3}{z}, C_2 \rightarrow C_2 + \frac{C_3}{x} \text{ we get}$$

$$\Delta = (x+y+z)^2 \begin{vmatrix} x+y & \frac{z^2}{x} & z^2 \\ \frac{x^2}{z} & z+y & x^2 \\ 0 & 0 & 2xz \end{vmatrix} \quad 1$$

$$= 2xyz (x+y+z)^3 \quad 1$$

**OR**

$$\text{For getting } A^2 = \begin{pmatrix} 5 & 0 & 8 \\ 2 & 4 & 5 \\ 8 & 0 & 13 \end{pmatrix} \quad 1\frac{1}{2}$$

$$\text{For getting } A^3 = \begin{pmatrix} 21 & 0 & 34 \\ 12 & 8 & 23 \\ 34 & 0 & 55 \end{pmatrix} \quad 1\frac{1}{2}$$

Simplifying  $A^3 - 6A^2 + 7A + kI_3$  as  $\begin{pmatrix} k-2 & 0 & 0 \\ 0 & k-2 & 0 \\ 0 & 0 & k-2 \end{pmatrix}$  2

Equating  $\begin{pmatrix} k-2 & 0 & 0 \\ 0 & k-2 & 0 \\ 0 & 0 & k-2 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

$$\Rightarrow k - 2 = 0$$

$$k = 2$$

1

