



# GEOMETRY

Inspiration is needed in geometry just as much as in poetry. - Alexander Pushkin

Thales (Pronounced THAYLEES) was born in the Greek city of Miletus. He was known for theoretical and practical understanding of geometry, especially



triangles. He used geometry to solve many problems such as calculating the height of pyramids and the distance of ships from the sea shore. He was one of the so-called Seven Sagas or Seven Wise Men of Greece and many regarded him as the first philosopher in the western tradition.

# Learning Outcomes

- **To understand theorems on linear pairs and vertically opposite angles.**
- **To understand the angle sum property of triangle.**
- To understand the properties of quadrilaterals and use them in problem solving.
- To understand, interpret and apply theorems on the chords and the angles subtended by arcs of a circle
- **To understand, interpret and apply theorems on the cyclic quadrilaterals.**
- To construct and locate centroid, orthocentre, circumcentre and incentre of a triangle.

#### 4.1 Introduction

In geometry, we study **shapes**. But what is there to *study* in shapes, you may ask. Think first, what are all the things we do with shapes? We draw shapes, we compare shapes, we *measure* shapes. *What* do we measure in shapes?

138 9<sup>th</sup> Standard Mathematics

 $( \bullet )$ 

Take some shapes like this:



In both of them, there is a *curve* forming the shape: one is a closed curve, enclosing a region, and the other is an open curve. We can use a rope (or a thick string) to measure the **length** of the open curve and the length of the boundary of the region in the case of the closed curve.

Curves are tricky, aren't they? It is so much

easier to measure length of straight lines using the scale, isn't it? Consider the two shapes below.

۲



Now we are going to focus our attention only on shapes made up of straight lines and on closed figures. As you will see, there is plenty of interesting things to do. Fig.4.2 shows an open figure.

We not only want to draw such shapes, we

want to compare them, measure them and do much more. For doing so, we want to **describe** them. How would you describe these closed shapes? (See Fig 4.3) They are all made up of straight lines and are closed.

#### 4.2 Types of Angles-Recall

Plumbers measure the angle between connecting pipes to make a good fitting. Wood workers adjust their saw blades to cut wood at the correct angle. Air Traffic Controllers (ATC) use angles to direct planes. Carom and billiards players must know their angles to plan their shots. An angle is formed by two rays that share a common end point provided that the two rays are non-collinear.



Fig. 4.3



Geometry 139

4-Geometry\_Term1.indd 139

 $( \bullet )$ 

#### **Complementary Angles**

Two angles are Complementary if their sum is 90°. For example, if  $\angle ABC = 64^{\circ}$  and  $\angle DEF = 26^{\circ}$ , then angles  $\angle ABC$  and  $\angle DEF$  are complementary to each other because  $\angle ABC + \angle DEF = 90^{\circ}$ 



#### Supplementary Angles



Two angles are Supplementary if their sum is 180°. For example if  $\angle ABC=110^{\circ}$  and  $\angle XYZ=70^{\circ}$ 

Here  $\angle ABC + \angle XYZ = 180^{\circ}$ 

۲

 $\therefore \angle ABC$  and  $\angle XYZ$  are supplementary to each other

#### **Adjacent Angles**

Two angles are called adjacent angles if

- (i) They have a common vertex.
- (ii) They have a common arm.
- (iii) The common arm lies between the two non-common arms.



#### **Linear Pair of Angles**

If a ray stands on a straight line then the sum of two adjacent angle is 180°. We then say that the angles so formed is a linear pair.



 $\angle AOC + \angle BOC = 180^{\circ}$ 

 $\therefore \angle AOC$  and  $\angle BOC$  form a linear pair

$$\angle XOZ + \angle YOZ = 180^{\circ}$$

 $\angle XOZ$  and  $\angle YOZ$  form a linear pair

#### **Vertically Opposite Angles**

If two lines intersect each other, then vertically opposite angles are equal.

In this figure  $\angle POQ = \angle SOR$ 

 $\angle POS = \angle QOR$ 



Fig. 4.10

#### 4.2.1 Transversal

A line which intersects two or more lines at distinct points is called a transversal of those lines.

۲

Case (i) When a transversal intersect two lines, we get eight angles.

In the figure the line l is the transversal for the lines m and n

- (i) Corresponding Angles:  $\angle 1$  and  $\angle 5$ ,  $\angle 2$  and  $\angle 6$ ,  $\angle 3$  and  $\angle 7$ ,  $\angle 4$  and  $\angle 8$
- (ii) Alternate Interior Angles:  $\angle 4$  and  $\angle 6$ ,  $\angle 3$  and  $\angle 5$
- (iii) Alternate Exterior Angles:  $\angle 1$  and  $\angle 7$ ,  $\angle 2$  and  $\angle 8$
- (iv)  $\angle 4$  and  $\angle 5, \angle 3$  and  $\angle 6$  are interior angles on the same side of the transversal.
- (v)  $\angle 1$  and  $\angle 8$ ,  $\angle 2$  and  $\angle 7$  are exterior angles on the same side of the transversal.
- Case (ii) If a transversal intersects two parallel lines. The transversal forms different pairs of angles.



Geometry 141



#### 4.2.2 Triangles



Activity 1

Take three different colour sheets; place one over the other and draw a triangle on the top sheet. Cut the sheets to get triangles of different colour which are identical. Mark the vertices and the angles as shown. Place the interior angles  $\angle 1$ ,  $\angle 2$  and  $\angle 3$  on a straight line, adjacent to each other, without leaving any gap. What can you say about the total measure of the three angles  $\angle 1$ ,  $\angle 2$  and  $\angle 3$ ?





to explain the "Exterior angle property" of a triangle?

- If a side of a triangle is stretched, the exterior angle so formed is equal to the sum of the two interior opposite angles. That is d=a+b (see Fig 4.14)

#### **4.2.3 Congruent Triangles**

Two triangles are congruent if the sides and angles of one triangle are equal to the corresponding sides and angles of another triangle.

Can you use

the same figure

۲

Rule	Diagrams	Reason
SSS	A O R	AB = PQ
		BC = QR
		AC = PR
	В С Р	$\Delta ABC \cong \Delta PQR$
SAS	AX	AB = XY
		$\angle BAC = \angle YXZ$
		AC = XZ
	B C Y Z	$\Delta ABC \cong \Delta XYZ$





1. In the figure, *AB* is parallel to *CD*, find x



- 2. The angles of a triangle are in the ratio 1: 2 : 3, find the measure of each angle of the triangle.
- 3. Consider the given pairs of triangles and say whether each pair is that of congruent triangles. If the triangles are congruent, say 'how'; if they are not congruent say 'why' and also say if a small modification would make them congruent:



Geometry 143

( )

۲



- 4. ΔABC and ΔDEF are two triangles in which
  AB=DF, ∠ACB=70°, ∠ABC=60°; ∠DEF=70° and ∠EDF=60°.
  Prove that the triangles are congruent.
- 5. Find all the three angles of the  $\triangle ABC$



# 4.3 Quadrilaterals

۲



Four Tamil Nadu State Transport buses take the following routes. The first is a one-way journey, and the rest are round trips. Find the places on the map, put points on them and connect them by lines to draw the routes. The places connecting four different routes are given as follows.

- (i) Nagercoil, Tirunelveli, Virudhunagar, Madurai
- (ii) Sivagangai, Puthukottai, Thanjavur, Dindigul, Sivagangai
- (iii) Erode, Coimbatore, Dharmapuri, Karur, Erode
- (iv) Chennai, Cuddalore, Krishnagiri, Vellore, Chennai



۲



Label the vertices with city names, draw the shapes exactly as they are shown on the map without rotations.

We observe that the first is a single line, the four points are collinear. The other three are closed shapes made of straight lines, of the kind we have seen before. We need names to call such closed shapes, we will call them **polygons** from now on.





How do polygons look? They have sides, with points at either end. We call these points as **vertices** of the polygon. The sides are line segments joining the vertices. The word *poly* stands for many, and a polygon is a many-sided figure.



**Concave polygon:** Polygon having any one of the interior angle greater than 180° **Convex Polygon:** Polygon having each interior angle less than 180° (Diagonals should be inside the polygon)

How many sides can a polygon have? One? But that is just a line segment. Two ? But how can you get a closed shape with two sides? Three? Yes, and this is what we know as a triangle. Four sides?

Squares and rectangles are examples of polygons with 4 sides but they are not the only ones. Here (Fig. 4.20) are some examples of 4-sided polygons. We call them *quadrilaterals*.

#### 4.3.1 Special Names for Some Quadrilaterals

- 1. A **parallelogram** is a quadrilateral in which opposite sides are parallel and equal.
- 2. A **rhombus** is a quadrilateral in which opposite sides are parallel and all sides are equal.
- 3. A **trapezium** is a quadrilateral in which *one pair of* opposite sides are parallel.

Draw a few parallellograms, a few rhombuses (correctly called rhombii, like cactus and cactii) and a few trapeziums (correctly written trapezia).

Geometry 145

4-Geometry\_Term1.indd 145

۲



The great advantage of knowing properties of quadrilateral is that we can see the relationships among them immediately.

- Every parallelogram is a trapezium, but not necessarily the other way.
- Every rhombus is a parallelogram, but not necessarily the other way.
- Every rectangle is a parallelogram, but not necessarily the other way.
- Every square is a rhombus and hence every square is a parallelogram as well.

For "not necessarily the other way" mathematicians usually say "the converse is not true". A smart question then is: just *when* is the other way also true? For instance, when is a parallelogram also a rectangle? Any parallelogram in which all angles are also equal is a rectangle. (Do you see why?) Now we can observe many more interesting properties. For instance, we see that a rhombus is a parallelogram in which all **sides** are also equal.



You know *bi*-cycles and *tri*-cycles? When we attach *bi or tri* to the front of any word, they stand for 2 (bi) or 3 (tri) of them. Similarly *quadri* stands for 4 of them. We should really speak of quadri-cycles also, but we don't. *Lateral* stands for sideways, thus quadrilateral means a 4-sided figure. You know *trilaterals*; they are also called triangles !

After 4 ? We have: 5 – *penta*, 6 – *hexa*, 7 – *hepta*, 8 – *octa*, 9 – *nano*, 10 – *deca*. Conventions are made by history. Trigons are called triangles, quadrigons are called quadrilaterals. Continuing in th same way we get pentagons, hexagons, heptagons, octagons, nanogons and decagons. Beyond these, we have 11-gons, 12-gons etc. Perhaps you can draw a 23-gon !

#### 4.3.2 More Special Names

When all sides of a quadrilateral are equal, we call it **equilateral**. When all angles of a quadrilateral are equal, we call it **equiangular**. In triangles, we talked of equilateral triangles as those with all sides equal. Now we can call them equiangular triangles as well!

#### We thus have:

A rhombus is an equilateral parallelogram.

- A rectangle is an equiangular parallelogram.
- A square is an equilateral and equiangular parallelogram.

146 9<sup>th</sup> Standard Mathematics

( )



Α

В

A

( )

В

Geometry 147

۲

В

Fig. 4.24

Α

В

Α



Step – 3: Fold the quadrilaterals along both of their diagonals. Press to make creases.



Fig. 4.26

We observe that two imposed triangles are congruent to each other. Measure the lengths of portions of diagonals and angles between the diagonals.

Also do the same for the quadrilaterals such as Trapezium, Isosceles Trapezium and Kite.

From the above activity, measure the lengths of diagonals and angles between the diagonals and record them in the table below:

S.	Name of the	Length along diagonals				Measure of angles					
No.	quadrilateral	AC	BD	OA	OB	<b>0</b> C	OD	∠AOB	∠BOC	∠ <i>COD</i>	∠ <b>DO</b> A
1	Trapezium										
2	Isosceles										
	Trapezium										
3	Parallelogram										
4	Rectangle										
5	Rhombus										
6	Square										
7	Kite										

148 9<sup>th</sup> Standard Mathematics

( )

۲



# Activity 4

#### Angle sum for a polygon

Draw any quadrilateral ABCD.

Mark a point *P* in its interior.

Join the segments PA, PB, PC and PD.

You have 4 triangles now.

How much is the sum of all the angles of the 4 triangles?

How much is the sum of the angles at P?

Can you now find the 'angle sum' of the quadrilateral ABCD?

Can you extend this idea to any polygon?

# Thinking Corner

1. If there is a polygon of *n* sides  $(n \ge 3)$ , then the sum of all interior angles is  $(n-2) \times 180^{\circ}$ 

۲

D

Р

R

Fig. 4.27

- 2. For the regular polygon (All the sides of a polygon are equal in size)
  - **c** Each interior angle is  $\frac{(n-2)}{n} \times 180^{\circ}$
  - **Characteristics** Each exterior angle is  $\frac{360^{\circ}}{n}$
  - The sum of all the exterior angles formed by producing the sides of a convex polygon in the order is 360°.
  - The polygon has 'n' sides , then the number of diagonals of the polygon is  $\frac{n(n-3)}{2}$

Geometry 149

# 4.3.4 Properties of Quadrilaterals

Name	Diagram	Sides	Angles	Diagonals	
Parallelogram	E F L I	Opposite sides are parallel and equal	Opposite angles are equal and sum of any two adjacent angles is 180°	Diagonals bisect each other.	
Rhombus		All sides are equal and opposite sides are parallel	Opposite angles are equal and sum of any two adjacent angles is 180°	Diagonals bisect each other at right angle.	
Trapezium		One pair of opposite sides are parallel	The angles at the ends of each non-parallel sides are supplementary	Diagonals need not be equal	
Isosceles Trapezium	B C	One pair of opposite sides are parallel and non-parallel sides are equal in length.	The angles at the ends of each parallel sides are equal.	Diagonals are of equal length.	
Kite $F$ $G$		Two pairs of adjacent sides are equal	One pair of opposite angles are equal	<ol> <li>Diagonals intersect at right angle.</li> <li>Shorter diagonal bisected by longer diagonal</li> <li>Longer diagonal divides the kite into two congruent triangles</li> </ol>	

150 9<sup>th</sup> Standard Mathematics

۲

۲



Note

- (i) A rectangle is an equiangular parallelogram.
- (ii) A rhombus is an equilateral parallelogram.
- (iii) A square is an equilateral and equiangular parallelogram.
- (iv) A square is a rectangle , a rhombus and a parallelogram.



- 1. State the reasons for the following.
  - (i) A square is a special kind of a rectangle.
  - (ii) A rhombus is a special kind of a parallelogram.
  - (iii) A rhombus and a kite have one common property.
  - (iv) A square and a rhombus have one common property.
- 2. What type of quadrilateral is formed when the following pairs of congruent triangles are joined together?
  - (i) Equilateral triangle.
  - (ii) Right angled triangle.
  - (iii) Isosceles triangle.
- 3. Identify which ones are parallelograms and which are not.



5. Identify which ones are trapeziums and which are not.



Geometry 151

4.

( )

۲

#### 4.3.5 Properties of Parallelogram

We can now embark on an interesting journey. We can tour among lots of quadrilaterals, noting down interesting properties. What properties do we look for, and how do we know they are true?

 $( \mathbf{0} )$ 

For instance, opposite sides of a parallelogram are parallel, but are they also **equal**? We could draw any number of parallelograms and verify whether this is true or not. In fact, we see that opposite sides are equal in **all of** them. Can we then conclude that opposite sides are equal in *all* parallelograms? No, because we might later find a parallelogram, one which we had not thought of until then, in which opposite sides are unequal. So, we need an argument, a **proof**.

Consider the parallelogram ABCD in the given Fig. 4.28. We believe that AB = CD and AD = BC, but how can we be sure? We know triangles and their properties. So we can try and see if we can use that knowledge. But we don't have any triangles in the parallelogram ABCD.





This is easily taken care of by joining *AC*. (We could equally well have joined *BD*, but let it be *AC* for now.) We now have 2 triangles *ADC* and *ABC* with a common side *AC*. If we could somehow prove that these two triangles are congruent, we would get AB = CD and AD = BC, which is what we want!

Is there any hope of proving that  $\triangle ADC$  and  $\triangle ABC$  are congruent? There are many criteria for congruence, it is not clear which one is relevant here.

So far we have not used the fact that ABCD is a parallelogram at all. So we need to use the facts that AB||DC and AD||BC to show that  $\triangle ADC$  and  $\triangle ABC$  are congruent. From sides being parallel we have to get into some angles being equal. Do we know any such properties? we do, and that is all about **transversals**!

Now we can see it clearly.  $AD \| BC$  and AC is a transversal, hence  $\angle DAC = \angle BCA$ . Similarly,  $AB \| DC$ , AC is a transversal, hence  $\angle BAC = \angle DCA$ . With AC as common side, the ASA criterion tells us that  $\triangle ADC$  and  $\triangle ABC$  are congruent, just what we needed. From this we can conclude that AB = CD and AD = BC.

Thus opposite sides are indeed equal in a parallelogram.

The argument we now constructed is written down as a *formal proof* in the following manner.

152 9<sup>th</sup> Standard Mathematics

#### Theorem 1

In a parallelogram, opposite sides are equal

GivenABCD is a parallelogramTo ProveAB=CD and DA=BCConstructionJoin AC

Proof

Since *ABCD* is a parallelogram

 $AD \parallel BC$  and AC is the transversal

 $\angle DAC = \angle BCA$ 

 $AB \parallel DC$  and AC is the transversal

$$\angle BAC = \angle DCA$$

In  $\triangle ADC$  and  $\triangle CBA$ 

 $\angle DAC = \angle BCA$  from (1)

AC is common

	$\angle DCA = \angle BAC$	from (2)
	$\Delta ADC \cong \Delta CBA$	(By ASA)
Hence	AD = CB and $DC = BA$	(Corresponding sides are equal)

۲

Along the way in the proof above, we have proved another property that is worth recording as a theorem.

#### Theorem 2

A diagonal of a parallelogram divides it into two congruent triangles.

Notice that the proof above established that  $\angle DAC = \angle BCA$  and  $\angle BAC = \angle DCA$ . Hence we also have, in the figure above,

 $\angle BCA + \angle BAC = \angle DCA + \angle DAC$ 

But we know that:

 $\angle B + \angle BCA + \angle BAC = 180$ 

and  $\angle D + \angle DCA + \angle DAC = 180$ 

Therefore we must have that  $\angle B = \angle D$ .

With a little bit of work, proceeding similarly, we could have shown that  $\angle A = \angle C$  as well. Thus we have managed to prove the following theorem:

Geometry 153

В

C



 $\rightarrow$ (1) (alternate angles are equal)

 $\Rightarrow$ (2) (alternate angles are equal)

D

Fig. 4.31

A

 $( \bullet )$ 

#### Theorem 3

#### The opposite angles of a parallelogram are equal.

Now that we see congruence of triangles as a good "strategy", we can look for more triangles. Consider both diagonals AC and DB. We already know that  $\triangle ADC$  and  $\triangle CBA$  are congruent. By a similar argument we can show that  $\triangle DAB$  and  $\triangle BCD$  are congruent as well. Are there more congruent triangles to be found in this figure ?

۲

Yes. The two diagonals intersect at point *O*. We now see 4 new  $\triangle AOB$ ,  $\triangle BOC$ ,  $\triangle COD$  and  $\triangle DOA$ . Can you see any congruent pairs among them?

Since *AB* and *CD* are parallel and equal, one good guess is that  $\triangle AOB$  and  $\triangle COD$  are congruent. We could again try <sup>A</sup> the ASA crierion, in which case we want  $\angle OAB = \angle OCD$  and



 $\angle ABO = \angle CDO$ . But the first of these follows from the fact that  $\angle CAB = \angle ACD$  (which we already established) and observing that  $\angle CAB$  and  $\angle OAB$  are the same (and so also  $\angle OCD$  and  $\angle ACD$ ). We now use the fact that *BD* is a transversal to get that  $\angle ABD = \angle CDB$ , but then  $\angle ABD$  is the same as  $\angle ABO$ ,  $\angle CDB$  is the same as  $\angle CDO$ , and we are done.

۲

Again, we need to write down the formal proof, and we have another theorem.

#### **Theorem 4**

#### The diagonals of a parallelogram bisect each other.

It is time now to reinforce our concepts on parallelograms. Consider each of the given statements, in the adjacent box, one by one. Identify the type of parallelogram which satisfies each of the statements. Support your answer with reason.

Now we begin with lots of interesting properties of parallelograms. Can we try and prove some property relating to two or more parallelograms ? A simple case to try is when two parallelograms



- Each pair of opposite sides is equal.
- All of its angles are right angles.
- Its diagonals bisect each other.
- The diagonals are equal.
- The diagonals are perpendicular and equal.
- The diagonals are perpendicular bisectors of each other.
- Each pair of its consecutive angles is supplementary.



154 9<sup>th</sup> Standard Mathematics

We see parallelograms *ABCD* and *ABEF* are on the common base *AB*. At once we can see a pair of triangles for being congruent  $\triangle ADF$  and  $\triangle BCE$ . We already have that AD = BC and AF = BE. But then since  $AD \parallel BC$  and  $AF \parallel BE$ , the angle formed by *AD* and *AF* must be the same

( )

as the angle formed by BC tand BE. Therefore  $\angle DAF = \angle CBE$ . Thus  $\triangle ADF$  and  $\triangle BCE$  are congruent.

6

That is an interesting observation; can we infer anything more from this ? Yes, we know that congruent triangles have the same area. This makes us think about the areas of the parallelograms ABCD and ABEF.

> Area of *ABCD* = area of quadrilateral *ABED* + area of  $\triangle BCE$ = area of quadrilateral ABED + area of  $\triangle ADF$ = area of *ABEF*

Thus we have proved another interesting theorem:

#### **Theorem 5:**

Parallelograms on the same base and between the same parallels are equal in area.

In this process, we have also proved other interesting statements. These are called *Corollaries*, which do not need separate detailed proofs.

**Corollary 1:** Triangles on the same base and between the same parallels are equal in area.

Corollary 2: A rectangle and a parallelogram on the same base and between the same parallels are equal in area.

These statements that we called Theorems and Corollaries, hold for all parallelograms, however large or small, with whatever be the lengths of sides and angles at vertices.

#### **Example 4.1**



In a parallelogram *ABCD*, the bisectors of the consecutive angles  $\angle A$  and  $\angle B$  insersect at *P*. Show that  $\angle APB = 90^{\circ}$ 

#### **Solution**

ABCD is a parallelogram AP and BP are bisectors of consecutive angles  $\angle A$  and  $\angle B$ .

Since the consecutive angles of a parallelogram are supplementary

$$\angle A + \angle B = 180^{\circ}$$
$$\frac{1}{2}\angle A + \frac{1}{2}\angle B = \frac{180^{\circ}}{2}$$

$$\implies \angle PAB + \angle PBA = 90^{\circ}$$

In  $\triangle APB$ ,

 $\angle PAB + \angle APB + \angle PBA = 180^{\circ}$  (angle sum property of triangle)  $\angle APB = 180^{\circ} - [\angle PAB + \angle PBA]$  $= 180^{\circ} - 90^{\circ} = 90^{\circ}$ 

Hence Proved.





Geometry 155

Example 4.2

In the Fig.4.35 *ABCD* is a parallelogram, *P* and *Q* are the midpoints of sides *AB* and *DC* respectively. Show that *APCQ* is a parallelogram.

#### Solution

Since *P* and *Q* are the mid points of



۲

Thus , in quadrilateral *APCQ* we have AP = QC and  $AP \parallel QC$  [from (2) and (3)] Hence , quadrilateral *APCQ* is a parallelogram.

#### Example 4.3

*ABCD* is a parallelogram Fig.4.36 such that  $\angle BAD = 120^{\circ}$  and *AC* bisects  $\angle BAD$  show that *ABCD* is a rhombus.

#### **Solution**

Given

$$\angle BAD = 120^{\circ} \text{ and } AC \text{ bisects } \angle BAD$$
  
 $\angle BAC = \frac{1}{2} \times 120^{\circ} = 60^{\circ}$   
 $\angle 1 = \angle 2 = 60^{\circ}$ 

 $AD \parallel BC$  and AC is the traversal

 $\angle 2 = \angle 4 = 60^{\circ}$ 

 $\triangle ABC$  is isosceles triangle  $[\because \angle 1 = \angle 4 = 60^{\circ}]$ 

 $\Rightarrow \qquad AB = BC$ 

Parallelogram *ABCD* is a rhombus.

156 9th Standard Mathematics



۲



**Example 4.4** In a parallelogram ABCD, P and Q are the points on line DB such that PD = BQ show that APCQ is a parallelogram

#### Solution

ABCD is a parallelogram.

now

OB = OD (:: Diagonals bisect each other)

 $W \qquad OB + BQ = OD + DP$ 

OQ = OP and OA = OC

OA = OC and

APCQ is a parallelogram.



- 1. The angles of a quadrilateral are in the ratio 2 : 4 : 5 : 7. Find all the angles.
- 2. In a quadrilateral *ABCD*,  $\angle A = 72^{\circ}$  and  $\angle C$  is the supplementary of  $\angle A$ . The other two angles are 2x-10 and x + 4. Find the value of *x* and the measure of all the angles.
- 3. ABCD is a rectangle whose diagonals AC and BD intersect at O. If  $\angle OAB = 46^{\circ}$ , find  $\angle OBC$
- 4. The lengths of the diagonals of a Rhombus are 12 cm and 16 cm . Find the side of the rhombus.
- 5. Show that the bisectors of angles of a parallelogram form a rectangle .
- 6. If a triangle and a parallelogram lie on the same base and between the same parallels, then prove that the area of the triangle is equal to half of the area of parallelogram.
- 7. Iron rods a, b, c, d, e, and f are making a design in a bridge as shown in the figure. If a || b, c || d, e || f, find the marked angles between
  - (i) b and c
  - (ii) d and e
  - (iii d and f
  - (iv) c and f



Fig. 4.37

Fig. 4.38

Geometry 157

23-02-2019 14:18:01

9. In the given Fig. 4.40, if AB = 2, BC = 6, AE = 6, BF = 8, CE = 7, and CF = 7, compute the ratio of the area of quadrilateral *ABDE* to the area of  $\triangle CDF$ .

- 10. In the Fig. 4.41 , *ABCD* is a rectangle and *EFGH* is a parallelogram. Using the measurements given in the figure, what is the length d of the segment that is perpendicular to  $\overline{HE}$  and  $\overline{FG}$  ?
- 11. In parallelogram *ABCD* of the accompanying diagram, line *DP* is drawn bisecting *BC* at *N* and meeting *AB* (extended) at *P*. From vertex *C*, line *CQ* is drawn bisecting side *AD* at *M* and meeting *AB* (extended) at *Q*. Lines *DP* and *CQ* meet at *O*. Show that the area of triangle *QPO* is  $\frac{9}{8}$  of the area of the parallelogram *ABCD*.



۲

158 9<sup>th</sup> Standard Mathematics



#### 4.4 Parts of a Circle

Circles are geometric shapes you can see all around you. The significance of the concept of a circle can be well understood from the fact that the wheel is one of the ground-breaking inventions in the history of mankind.



A circle, you can describe, is the set of all points in a plane at a constant distance from a fixed point. The fixed point is the centre of the circle; the constant distance corresponds to a radius of the circle.

A line that cuts the circle in two points is called a secant of the circle.

A line segment whose end points lie on the circle is called a chord of the circle.



A circle notably differs from a polygon. A polygon (for example, a quadrilateral) has edges and corners while, a circle is a 'smooth' curve.

A chord of a circle that has the centre is called a diameter of the circle. The circumference of a circle is its boundary. (We use the term perimeter in the case of polygons).



In Fig.4.44, we see that all the line segments meet at two points on the circle. These line segments are called the chords of the circle. So, a line segment joining any two points on the circle is called a chord of the circle. In this figure AB, PQ and RS are the chords of the circle.

Now place four points P, R, Q and S on the same

Fig. 4.44 circle (Fig.4.45), then PRQ and QSP are the continuous parts (sections) of the circle. These parts (sections) are to be denoted by PRQ and QSP or simply by PQ and QP. This continuous part of a circle is called an arc of the circle. Usually the arcs are denoted in anti-clockwise direction.



R

Fig. 4.45

4-Geometry\_Term1.indd 159



Now in (Fig.4.46), consider the region which is surrounded by the chord PQ and major arc  $\overrightarrow{QP}$ . This is called the major segment of the circle. In the same way, the segment containing the minor

arc and the same chord is called the minor segment.



In (Fig.4.47), if two arcs  $\overrightarrow{AB}$  and  $\overrightarrow{CD}$  of a circle subtend the same angle at the centre, they are said to be congruent arcs and we write,

$$AB \equiv CD$$
 implies  $mAB = mCD$ 

implies 
$$\angle AOB = \angle COD$$

۲

A diameter of a circle is:

- the line segment which bisects the circle.
- the largest chord of a circle.
- a line of symmetry for the circle.
- twice in length of a radius in a circle.



 $( \bullet )$ 

Now, let us observe (Fig.4.48). Is there any special name for the region surrounded by two radii and arc? Yes, its name is sector. Like segment, we find that the minor arc corresponds to the minor sector and the major arc corresponds to the major sector.

#### **Concentric Circles**

Circles with the same centre but different radii are said to be concentric.

Here are some real-life examples:



An Archery target



A carom board coin Fig. 4.49



Water ripples

#### **Congruent Circles**

Two circles are congruent if they are copies of one another or identical. That is, they have the same size. Here are some real life examples:





#### Position of a Point with respect to a Circle

Consider a circle in a plane (Fig.4.51). Consider any point *P* on the circle. If the distance from the centre *O* to the point *P* is *OP*, then

- (i) OP =radius (If the point *P* lies on the circle)
- (ii) *OP* < radius (If the point *P* Point lies inside the circle)
- (iii) *OP* > radius(If the point P lies outside the circle)

So, a circle divides the plane on which it lies into three parts.





### Progress Check

#### Say True or False

- 1. Every chord of a circle contains exactly two points of the circle.
- 2. All radii of a circle are of same length.
- 3. Every radius of a circle is a chord.
- 4. Every chord of a circle is a diameter.
- 5. Every diameter of a circle is a chord.
- 6. There can be any number of diameters for a circle.
- 7. Two diameters cannot have the same end-point.
- 8. A circle divides the plane into three disjoint parts.
- 9. A circle can be partitioned into a major arc and a minor arc.
- 10. The distance from the centre of a circle to the circumference is that of a diameter



- How many sides does a circle have ?
- 2. Is circle, a polygon?

Geometry 161

 $( \bullet )$ 

#### **4.4.1 Circle Through Three Points**

We have already learnt that there is one and only one line passing through two points. In the same way, we are going to see how many circles can be drawn through a given point, and through two given points. We see that in both cases there can be infinite number of circles passing through a given point P (Fig.4.52), and through two given points A and B (Fig.4.53).

۲



Now consider three collinear points A, B and C (Fig.4.14). Can we draw a circle passing through these three points? Think over it. If the points are collinear, we can't?

If the three points are non collinear, they form a triangle (Fig.4.55). Recall the construction of the circumcentre. The intersecting point of the perpendicular bisector of the sides is the circumcentre and the circle is circumcircle.



Therefore from this we know that, there is a unique circle which passes through A, B and C. Now, the above statement leads to a result as follows.

**Theorem 6** There is one and only one circle passing through three non-collinear points.

#### 4.5 **Properties of Chords of a Circle**

In this chapter, already we come across lines, angles, triangles and quadrilaterals. Recently we have seen a new member circle. Using all the properties of these, we get some standard results one by one. Now, we are going to discuss some properties based on chords of the circle.

Considering a chord and a perpendicular line from the centre to a chord, we are going to see an interesting property.

#### 4.5.1 Perpendicular from the Centre to a Chord

Consider a chord AB of the circle with centre O. Draw  $OC \perp AB$  and join the points OA, OB. Here, easily we get two triangles  $\triangle AOC$  and  $\triangle BOC$  (Fig.4.56).

162 9<sup>th</sup> Standard Mathematics

Can we prove these triangles are congruent? Now we try to prove this using the congruence of triangle rule which we have already learnt.  $\angle OCA = \angle OCB = 90^{\circ}(OC \perp AB)$  and OA = OB is the radius of the circle. The side OC is common. RHS criterion tells us that  $\triangle AOC$  and  $\Delta BOC$  are congruent. From this we can conclude that AC = BC. This argument leads to the result as follows.



**Theorem 7** The perpendicular from the centre of a circle to a chord bisects the chord.

Converse of Theorem 7 The line joining the centre of the circle and the midpoint of a chord is perpendicular to the chord.

#### Example 4.5

Find the length of a chord which is at a distance of  $2\sqrt{11}$  cm from the centre of a circle of radius 12cm.

#### **Solution**

Let AB be the chord and C be the mid point of AB

Therefore,  $OC \perp AB$ 

Join OA and OC.

*OA* is the radius

Given  $OC = 2\sqrt{11}$ cm and OA = 12cm

In a right  $\triangle OAC$ ,

using Pythagoras Theorem, we get,

 $AC^2 = OA^2 - OC^2$  $= 12^2 - (2\sqrt{11})^2$ = 144 - 44= 100 cm $AC^{2} = 100 \text{cm}$ AC = 10cm Therefore, length of the chord AB = 2AC $= 2 \times 10$ cm = 20cm





Α

#### Pythagoras theorem

One of the most important and well known results in geometry is Pythagoras Theorem. "In a right angled triangle, the <sup>*B*</sup> square of the hypotenuse

is equal to the sum of the squares of the other two sides".

In right  $\triangle ABC, BC^2 = AB^2 + AC^2$ . Application of this theorem is most useful in this unit.

**Example 4.6** 

In the concentric circles, chord AB of the outer circle cuts the inner circle at C and D as shown in the diagram. Prove that, AB - CD = 2AC



Geometry 163

#### Solution

Given : Chord AB of the outer circle cuts the inner circle at C and D.

To prove : AB - CD = 2ACConstruction : Draw  $OM \perp AB$ Proof : Since,  $OM \perp AB$  (By construction) Also,  $OM \perp CD$ Therefore, AM = MB ... (1) (Perpendicular drawn from centre to chord bisect it) CM = MD ... (2)

Now, AB - CD = 2AM - 2CM

= 2(AM - CM) from (1) and (2)

AB - CD = 2AC



- 1. The radius of the circle is 25 cm and the length of one of its chord is 40cm. Find the distance of the chord from the centre.
- 2. Draw three circles passing through the points P and Q, where PQ = 4cm.

#### 4.5.2 Angle Subtended by Chord at the Centre

Instead of a single chord we consider two equal chords. Now we are going to discuss another property.

Let us consider two equal chords in the circle with centre O. Join the end points of the chords with the centre to get the triangles  $\triangle AOB$  and  $\triangle OCD$ , chord AB = chord CD (because the given chords are equal). The other sides are radii, therefore OA=OC and OB=OD. By SSS rule, the triangles are congruent, that is  $\triangle OAB \equiv \triangle OCD$ . This gives  $m \angle AOB = m \angle COD$ . Now this leads to the following result.





Theorem 8 Equal chords of a circle subtend equal angles at the centre.

164 9<sup>th</sup> Standard Mathematics



# Activity - 5

Procedure

- 1. Draw a circle with centre *O* and with suitable radius.
- 2. Make it a semi-circle through folding. Consider the point *A*, *B* on it.

6

- 3. Make crease along AB in the semi circles and open it.
- 4. We get one more crease line on the another part of semi circle,  $_C$  name it as CD (observe AB = CD)
- 5. Join the radius to get the  $\triangle OAB$  and  $\triangle OCD$ .
- 6. Using trace paper, take the replicas of triangle  $\triangle OAB$  and  $\triangle OCD$ .
- 7. Place these triangles  $\triangle OAB$  and  $\triangle OCD$  one on the other.

#### Observation

- 1. What do you observe? Is  $\triangle OAB \equiv \triangle OCD$ ?
- 2. Construct perpendicular line to the chords *AB* and *CD* passing through the centre *O*. Measure the distance from *O* to the chords.

Now we are going to find out the length of the chords AB and CD, given the angles subtended by two chords at the centre of the circle are equal. That is,  $\angle AOB = \angle COD$  and the two sides which include these angles of the D and  $\Delta COD$  are radii and are equal.

By SAS rule,  $\Delta AOB \equiv \Delta COD$ . This gives chord AB = chord CD. Now let us write the converse result as follows:

#### **Converse of theorem 8**

If the angles subtended by two chords at the centre of a circle are equal, then the chords are equal.

In the same way we are going to discuss about the distance from the centre, when the equal chords are given. Draw the perpendicular  $OL \perp AB$  and  $OM \perp CD$ . From theorem 7, these perpendicular divides the chords equally. So AL = CM. By comparing the  $\triangle OAL$  and  $\triangle OCM$ , the angles  $\angle OLA = \angle OMC = 90^{\circ}$  and OA = OC are radii. By RHS rule, the  $\triangle OAL \equiv \triangle OCM$ . It gives the distance from the centre OL = OM and write the conclusion as follows.

**Theorem 9** Equal chords of a circle are equidistant from the centre.

Let us know the converse of theorem 9, which is very useful in solving problems.





Geometry 165

#### **Converse of theorem 9**

The chords of a circle which are equidistant from the centre are equal.

۲

#### 4.5.3 Angle Subtended by an Arc of a Circle

#### Activity - 6 **Procedure :** Draw three circles of any radius with centre *O* on a chart paper. 1. 2. From these circles, cut a semi-circle, a minor segment and a major segment. Consider three points on these segment and name them as A, B and C. 3. Semi circle •0 В •0 0 C<sup>B</sup> В Fig. 4.63 Major segment (iv) Cut the triangles and paste it on the graph sheet 4. •0 so that the point A coincides with the origin as shown in the figure. В A Minor **Observation :** segment Angle in a Semi-Circle is \_\_\_\_\_ angle. (i) В (ii) Angle in a major segment is \_\_\_\_\_ angle. Fig. 4.64 (iii) Angle in a minor segment is \_\_\_\_\_ angle.

Now we are going to verify the relationship between the angle subtended by an arc at the centre and the angle subtended on the circumference.

#### 4.5.4 Angle at the Centre and the Circumference

Let us consider any circle with centre O. Now place the points A, B and C on the circumference.



( )

Here AB is a minor arc in Fig.4.65, a semi circle in Fig.4.66 and a major arc in Fig.4.67. The point C makes different types of angles in different positions (Fig. 4.65 to 4.67). In all these circles,  $\widehat{AXB}$  subtends  $\angle AOB$  at the centre and  $\angle ACB$  at a point on the circumference of the circle.

۲

We want to prove  $\angle AOB = 2 \angle ACB$ . For this purpose extend *CO* to *D* and join *CD*.

 $\angle OCA = \angle OAC$  since OA = OC (radii)

Exterior angle = sum of two interior opposite angles.

 $\angle AOD = \angle OAC + \angle OCA$ =  $2\angle OCA$  .... (1)

Similarly,

$$\angle BOD = \angle OBC + \angle OCB$$
$$= 2\angle OCB \qquad \dots (2)$$

From (1) and (2),  $\angle AOD + \angle BOD = 2(\angle OCA + \angle OCB)$ 

Finally we reach our result  $\angle AOB = 2 \angle ACB$ .

From this we get the result as follows :

#### Theorem 10

The angle subtended by an arc of the circle at the centre is double the angle subtended by it at any point on the remaining part of the circle.





Progress Check

- i. Draw the outline of different size of bangles and try to find out the centre of each using set square.
- ii. Trace the given cresent and complete as full moon using ruler and compass.



#### Solution

Using the theorem the angle subtended by an arc of a circle at the centre is double the angle subtended by it at any point on the remaining part of a circle.

۲



**Example 4.8** If *O* is the centre of the circle and  $\angle ABC = 30^{\circ}$  then find  $\angle AOC$ . (see Fig. 4.73)

#### Solution

Given  $\angle ABC = 30^{\circ}$ 

$$\angle AOC = 2 \angle ABC$$

(The angle subtended by an arc at the centre is double the angle at any point on the circle)



$$= 2 \times 30^{\circ}$$
$$= 60^{\circ}$$

Now we shall see, another interesting theorem. We have learnt that minor arc subtends obtuse angle, major arc subtends acute angle and semi circle subtends right angle on the circumference. If a chord AB is given and C and D are two different points on the circumference of the circle, then find  $\angle ACB$  and  $\angle ADB$ . Is there any difference in these angles?

168 9<sup>th</sup> Standard Mathematics

۲

#### 4.5.5 Angles in the same segment of a circle

Consider the circle with centre O and chord AB. C and D are the points on the circumference of the circle in the same segment. Join the radius OA and OB.

$$\frac{1}{2} \angle AOB = \angle ACB \text{ (by theorem 10)}$$
  
and 
$$\frac{1}{2} \angle AOB = \angle ADB \text{ (by theorem 10)}$$
$$\angle ACB = \angle ADB$$

This conclusion leads to the new result.

Theorem 11 Angles in the same segment of a circle are equal.

**Example 4.9** In the given figure, *O* is the center of the circle. If the measure of  $\angle OQR = 48^\circ$ , what is the measure of  $\angle P$ ?

#### **Solution**

Given  $\angle OQR = 48^{\circ}$ .

Therefore,  $\angle ORQ$  also is 48°. (Why?\_\_\_\_\_

 $\angle QOR = 180^{\circ} - (2 \times 48^{\circ}) = 84^{\circ}.$ 

The central angle made by chord QR is twice the inscribed angle at P.

Thus, measure of  $\angle QPR = \frac{1}{2} \times 84^\circ = 42^\circ$ .

1. The diameter of the circle is 52cm and the length of one of its chord is 20cm. Find the distance of the chord from the centre.

Exercise 4.3

2. The chord of length 30 cm is drawn at the distance of 8cm from the centre of the circle. Find the radius of the circle

- 3. Find the length of the chord AC where AB and CD are the two diameters perpendicular to each other of a circle with radius  $4\sqrt{2}$  cm and also find  $\angle OAC$  and  $\angle OCA$ .
- 4. A chord is 12cm away from the centre of the circle of radius 15cm. Find the length of the chord.

Geometry 169







 $igodoldsymbol{\Theta}$ 

5. In a circle, AB and CD are two parallel chords with centre O and radius 10 cm such that AB = 16 cm and CD = 12 cm determine the distance between the two chords?

۲

- 6. Two circles of radii 5 cm and 3 cm intersect at two points and the distance between their centres is 4 cm. Find the length of the common chord.
- 7. Find the value of  $x^{\circ}$  in the following figures:



#### 4.6 Cyclic Quadrilaterals

Now, let us see a special quadrilateral with its properties called "Cyclic Quadrilateral". A quadrilateral is called cyclic quadrilateral if all its four vertices lie on the circumference of the circle. Now we are going to learn the special property of cyclic quadrilateral.

Consider the quadrilateral ABCD whose vertices lie on a circle. We want to show that its opposite angles are supplementary. Connect the centre O of the circle with each vertex. You now see four radii OA, OB, OC and OD giving rise to four isosceles triangles OAB, OBC, OCD and ODA. The sum of the angles around the centre of the circle is  $360^{\circ}$ . The angle sum of each isosceles triangle is  $180^{\circ}$ 

Thus, we get from the figure,

 $2 \times (\angle 1 + \angle 2 + \angle 3 + \angle 4)$  + Angle at centre  $O = 4 \times 180^{\circ}$ 

 $2 \times (\angle 1 + \angle 2 + \angle 3 + \angle 4) + 360^{\circ} = 720^{\circ}$ 

Simplifying this,

 $(\angle 1 + \angle 2 + \angle 3 + \angle 4) = 180^{\circ}.$ 

You now interpret this as

170 9th Standard Mathematics



D



(i)  $(\angle 1 + \angle 2) + (\angle 3 + \angle 4) = 180^{\circ}$  (Sum of opposite angles *B* and *D*)

(ii)  $(\angle 1 + \angle 4) + (\angle 2 + \angle 3) = 180^{\circ}$  (Sum of opposite angles A and C)

Now the result is given as follows.

Theorem 12 Opposite angles of a cyclic quadrilateral are supplementary.

Let us see the converse of theorem 12, which is very useful in solving problems

 $( \mathbf{0} )$ 

**Converse of Theorem 12** If a pair of opposite angles of a quadrilateral is supplementary, then the quadrilateral is cyclic.



#### Procedure

- 1. Draw a circle of any radius with centre *O*.
- 2. Mark any four points A, B, C and D on the boundary. Make a cyclic quadrilateral ABCD and name the angles as in Fig. 4.78
- 3. Make a replica of the cyclic quadrilateral *ABCD* with the help of tracing paper.
- 4. Make the cutout of the angles A, B, C and D as in Fig. 4.79
- 5. Paste the angle cutout  $\angle 1, \angle 2, \angle 3$  and  $\angle 4$  adjacent to the angles opposite to A, B, C and D as in Fig. 4.80
- 6. Measure the angles  $\angle 1 + \angle 3$ , and  $\angle 2 + \angle 4$ .

#### Observe and complete the following:

- 1. (i)  $\angle A + \angle C = \_\_\_\_$  (ii)  $\angle B + \angle D = \_\_\_\_$
- (iii)  $\angle C + \angle A =$  (iv)  $\angle D + \angle B =$
- 2. Sum of opposite angles of a cyclic quadrilateral is \_
- 3. The opposite angles of a cyclic quadrilateral is \_\_\_\_\_

#### Example 4.10

If PQRS is a cyclic quadrilateral in which  $\angle PSR = 70^{\circ}$  and  $\angle QPR = 40^{\circ}$ , then find  $\angle PRQ$  (see Fig. 4.81).

#### Solution

PQRS is a cyclic quadrilateral Given  $\angle PSR = 70^{\circ}$ 



Fig. 4.80



Fig. 4.78

Fig. 4.79

3

$$\angle PSR + \angle PQR = 180^{\circ} \quad \text{(state reason})$$

$$70^{\circ} + \angle PQR = 180^{\circ}$$

$$\angle PQR = 180^{\circ} - 70^{\circ}$$

$$\angle PQR = 110^{\circ}$$
In  $\triangle PQR$  we have,
$$\angle PQR + \angle PRQ + \angle QPR = 180^{\circ} \quad \text{(state reason})$$

$$110^{\circ} + \angle PRQ + 40^{\circ} = 180^{\circ}$$

$$\angle PRQ = 180^{\circ} - 150^{\circ}$$
$$\angle PRQ = 30^{\circ}$$

#### **Exterior Angle of a Cyclic Quadrilateral**

An exterior angle of a quadrilateral is an angle in its exterior formed by one of its sides and the extension of an adjacent side.



Let the side AB of the cyclic quadrilateral ABCD be extended to E. Here  $\angle ABC$  and  $\angle CBE$  are linear pair, their sum is 180° and the angles  $\angle ABC$  and  $\angle ADC$  are the opposite angles of a cyclic

quadrilateral, and their sum is also 180°. From this,  $\angle ABC + \angle CBE = \angle ABC + \angle ADC$ and finally we get  $\angle CBE = \angle ADC$ . Similarly it can be proved for other angles.

**Theorem 13** If one side of a cyclic quadrilateral is produced then the exterior angle is equal to the interior opposite angle.

# Progress Check

- 1. If a pair of opposite angles of a quadrilateral is supplementary, then the quadrilateral is
- 2. As the length of the chord decreases, the distance from the centre \_\_\_\_\_
- 3. If one side of a cyclic quadrilateral is produced then the exterior angle is \_\_\_\_\_\_ to the interior opposite angle.
- 4. Opposite angles of a cyclic quadrilateral are

```
Example 4.11
```

In the figure given, find the value of  $x^{\circ}$  and  $y^{\circ}$ .

#### Solution

By the exterior angle property of a cyclic quadrilateral,

we get,  $y^{\circ} = 100^{\circ}$  and

 $x^{\circ} + 30^{\circ} = 60^{\circ}$  and so  $x^{\circ} = 30^{\circ}$ 

172 9<sup>th</sup> Standard Mathematics

100°

30°

Fig. 4.83



1. Find the value of x in the given figure.

In the given figure, AC is the diameter of the circle with centre
O. If ∠ADE = 30°; ∠DAC = 35° and ∠CAB = 40°.
Find (i) ∠ACD (ii) ∠ACB (iii) ∠DAE

- 3. Find all the angles of the given cyclic quadrilateral *ABCD* in the figure.
- 4. In the given figure, ABCD is a cyclic quadrilateral where diagonals intersect at P such that ∠DBC = 40° and ∠BAC = 60° find
  (i) ∠CAD
  (ii) ∠BCD
- 5. In the given figure, AB and CD are the parallel chords of a circle with centre O. Such that AB = 8cm and CD = 6cm. If  $OM \perp AB$  and  $OL \perp CD$  distance between LM is 7cm. Find the radius of the circle?
- 6. The arch of a bridge has dimensions as shown, where the arch measure 2m at its highest point and its width is 6m. What is the radius of the circle that contains the arch?
- 7. In figure,  $\angle ABC = 120^{\circ}$ , where *A*, *B* and *C* are points on the circle with centre *O*. Find  $\angle OAC$ ?
- 8. A school wants to conduct tree plantation programme. For this a teacher allotted a circle of radius 6m ground to nineth



D

120°

Ř

4-Geometry\_Term1.indd 173



standard students for planting sapplings. Four students plant trees at the points A, B, C and D as shown in figure. Here AB = 8m, CD = 10m and  $AB \perp CD$ . If another student places a flower pot at the point P, the intersection of AB and CD, then find the distance from the centre to P.

9. In the given figure,  $\angle POQ = 100^{\circ}$  and  $\angle PQR = 30^{\circ}$ , then find  $^{R}$   $\angle RPO$ .

#### 4.7 Practical Geometry

Practical geometry is the method of applying the rules of geometry dealt with the properties of points, lines and other figures to construct geometrical figures. "Construction" in Geometry means to draw shapes, angles or lines accurately. The geometric constructions have been discussed in detail in Euclid's book 'Elements'. Hence these constructions are also known as Euclidean constructions. These constructions use only compass and straightedge (i.e. ruler). The compass establishes equidistance and the straightedge establishes collinearity. All geometric constructions are based on those two concepts.

 $( \mathbf{0} )$ 

It is possible to construct rational and irrational numbers using straightedge and a compass as seen in chapter II. In 1913 the Indian mathematical Genius, Ramanujam gave a geometrical construction for  $355/113 = \pi$ . Today with all our accumulated skill in exact measurements. it is a noteworthy feature that lines driven through a mountain meet and make a tunnel.In the earlier classes, we have learnt the construction of angles and triangles with the given measurements.

In this chapter we are going to learn to construct Centroid, Orthocentre, Circumcentre and Incentre of a triangle by using concurrent lines.

#### 4.7.1 Construction of the Centroid of a Triangle

#### Centroid

The point of concurrency of the medians of a triangle is called the centroid of the triangle and is usually denoted by *G*.





**Objective** To find the mid-point of a line segment using paper folding

**Procedure** Make a line segment on a paper by folding it and name it PQ. Fold the line segment PQ in such a way that P falls on Q and mark the point of intersection of the line segment and the crease formed by folding the paper as M. M is the midpoint of PQ.

174 9<sup>th</sup> Standard Mathematics

 $( \bullet )$ 

Example 4.12

Construct the centroid of

۲

 $\Delta PQR$  whose sides are PQ = 8cm; QR = 6cm; RP = 7cm.





Step 1 : Draw  $\Delta PQR$  using the given measurements PQ = 8cm QR = 6cm and RP = 7cm and construct the perpendicular bisector of any two sides (PQand QR) to find the mid-points M and N of PQ and QRrespectively.

G

M 8 cm Ν

Fig. 4.86

6 cm

Ο

7 cm

Step 2 : Draw the medians PN and<br/>RM and let them meet at G.<br/>The point G is the centroid<br/>of the given  $\Delta PQR$ .

# Note

( )

- Three medians can be drawn in a triangle
- The centroid divides each median in the ratio 2:1 from the vertex.
- The centroid of any triangle always lie inside the triangle.
- Centroid is often described as the triangle's centre of gravity (where the triangle balances evenly) and also as the barycentre.

# 4.7.2 Construction of Orthocentre of a Triangle

#### Orthocentre

The orthocentre is the point of concurrency of the altitudes of a triangle. Usually it is denoted by H.



۲

۲

Р



**Objective** To construct a perpendicular to a line segment from an external point using paper folding.

۲

**Procedure** Draw a line segment *AB* and mark an external point *P*. Move *B* along *BA* till the fold passes through *P* and crease it along that line. The crease thus formed is the perpendicular to *AB* through the external point *P*.

Activity 10

**Objective** To locate the Orthocentre of a triangle using paper folding.

**Procedure** Using the above Activity with any two vertices of the triangle as external points, construct the perpendiculars to opposite sides. The point of intersection of the perpendiculars is the Orthocentre of the given triangle.



#### Step 2:

Construct altitudes from any two vertices (say) R and P, to their opposite sides PQ and QR respectively.

The point of intersection of the altitude H is the Orthocentre of the given  $\Delta PQR$ .

176 9<sup>th</sup> Standard Mathematics

( )

 $( \bullet )$ 

Note								
Where do the Orthocentre lie in the given triangles.								
	Acute Triangle	Obtuse Triangle	<b>Right Triangle</b>					
Orthocentre	Inside of Triangle	Outside of Triangle	Vertex at Right Angle					



- 1. Construct the  $\Delta LMN$  such that LM=7.5cm, MN=5cm and LN=8cm. Locate its centroid.
- 2. Draw and locate the centroid of the triangle ABC where right angle at A, AB = 4cm and AC = 3cm.
- 3. Draw the  $\triangle ABC$ , where AB = 6cm,  $\angle B = 110^{\circ}$  and AC = 9cm and construct the centroid.
- 4. Construct the  $\triangle PQR$  such that PQ = 5cm, PR = 6cm and  $\angle QPR = 60^{\circ}$  and locate its centroid.
- 5. Draw  $\triangle PQR$  with sides PQ = 7 cm, QR = 8 cm and PR = 5 cm and construct its Orthocentre.
- 6. Draw an equilateral triangle of sides 6.5 cm and locate its Orthocentre.
- 7. Draw  $\triangle ABC$ , where AB = 6 cm,  $\angle B = 110^{\circ}$  and BC = 5 cm and construct its Orthocentre.
- 8. Draw and locate the Orthocentre of a right triangle *PQR* where PQ = 4.5 cm, QR = 6 cm and PR = 7.5 cm.

#### 4.7.3 Construction of the Circumcentre of a Triangle

#### Circumcentre

The Circumcentre is the point of concurrency of the Perpendicular bisectors of the sides of a triangle. It is usually denoted by S.

#### Circumcircle

The circle passing through all the three vertices of the triangle with circumcentre (*S*) as centre is called circumcircle.



Geometry 177

4-Geometry\_Term1.indd 177



**Procedure** Make a line segment on a paper by folding it and name it as *PQ*. Fold *PQ* in such a way that *P* falls on *Q* and thereby creating a crease RS. This line *RS* is the perpendicular bisector of *PQ*.



**Objective** To locate the circumcentre of a triangle using paper folding.

**Procedure** Using Activity 12, find the perpendicular bisectors for any two sides of the given triangle. The meeting point of these is the circumcentre of the given triangle.

# Example 4.14

Construct the circumcentre of the  $\triangle ABC$  with AB = 5 cm,  $\angle A = 60^{\circ}$  and  $\angle B = 80^{\circ}$ . Also draw the circumcircle and find the circumradius of the  $\triangle ABC$ .



 $( \bullet )$ 

#### Solution

**Step 1** Draw the  $\triangle ABC$  with the given measurements

#### Step 2

Construct the perpendicular bisector of any two sides (*AC* and *BC*) and let them meet at *S* which is the circumcentre.

#### Step 3

S as centre and SA = SB = SC as radius,

draw the Circumcircle to passes through *A*,*B* and *C*.

Circumradius = 3.9 cm.









# 4.7.4 Construction of the Incircle of a Triangle

#### Incentre

۲

The incentre is (one of the triangle's points of concurrency formed by) the intersection of the triangle's three angle bisectors.

The incentre is the centre of the incircle ; It is usually denoted by I; it <sub>A</sub> is the one point in the triangle whose distances to the sides are equal.



Geometry 179

В

🖌 Fig. 4.94

۲



- 1 Draw a triangle *ABC*, where AB = 8 cm, BC = 6 cm and  $\angle B = 70^{\circ}$  and locate its circumcentre and draw the circumcircle.
- 2 Construct the right triangle *PQR* whose perpendicular sides are 4.5 cm and 6 cm. Also locate its circumcentre and draw the circumcircle.
- 3. Construct  $\triangle ABC$  with AB = 5 cm  $\angle B = 100^{\circ}$  and BC = 6 cm. Also locate its circumcentre draw circumcircle.
- 4. Construct an isosceles triangle *PQR* where *PQ*= *PR* and  $\angle Q = 50^{\circ}$ , *QR* = 7cm. Also draw its circumcircle.
- 5. Draw an equilateral triangle of side 6.5 cm and locate its incentre. Also draw the incircle.

180 9<sup>th</sup> Standard Mathematics

۲

6. Draw a right triangle whose hypotenuse is 10 cm and one of the legs is 8 cm. Locate its incentre and also draw the incircle.

۲

- 7. Draw  $\triangle ABC$  given AB = 9 cm,  $\angle CAB = 115^{\circ}$  and  $\triangle ABC = 40^{\circ}$ . Locatge its incentre and also draw the incircle. (Note: You can check from the above examples that the incentre of any triangle is always in its interior).
- 8. Construct  $\triangle ABC$  in which AB = BC = 6cm and  $\angle B = 80^{\circ}$ . Locate its incentre and draw the incircle.



Geometry 181

۲

(4) Parallelogram but not a square

7. If bisectors of  $\angle A$  and  $\angle B$  of a quadrilateral *ABCD* meet at *O*, then  $\angle AOB$  is (2)  $\frac{1}{2}(\angle C + \angle D)$ (1)  $\angle C + \angle D$ (4)  $\frac{1}{3} \angle C + \frac{1}{2} \angle D$ (3)  $\frac{1}{2} \angle C + \frac{1}{3} \angle D$ The interior angle made by the side in a parallelogram is 90° then the parallelogram is a 8. (2) rectangle (1) rhombus (3) trapezium (4) kite 9. Which of the following statement is correct? (1) Opposite angles of a parallelogram are not equal. (2) Adjacent angles of a parallelogram are complementary. (3) Diagonals of a parallelogram are always equal. (4) Both pairs of opposite sides of a parallelogram are always equal. The angles of the triangle are 3x-40, x+20 and 2x-10 then the value of x is 10. (4) 45° (1) 40° (2) 35° (3) 50° PQ and RS are two equal chords of a circle with centre  $\mathit{O}$  such that  $\angle POQ = 70^\circ$  , 11. then  $\angle ORS =$  $(1) 60^{\circ}$ (2)  $70^{\circ}$ (3) 55°  $(4) 80^{\circ}$ A chord is at a distance of 15cm from the centre of the circle of radius 25cm. The 12. length of the chord is (1) 25cm (4) 18cm (2) 20cm (3) 40cm In the figure, O is the centre of the circle and 4013. 0  $\angle ACB = 40^{\circ}$  then  $\angle AOB =$ (4) 65°  $(3) 70^{\circ}$ (1)  $80^{\circ}$ (2)  $85^{\circ}$ In a cyclic quadrilaterals *ABCD*,  $\angle A = 4x$ ,  $\angle C = 2x$  the value of x is 14. (2) 20°  $(3) 15^{\circ}$  $(1) 30^{\circ}$  $(4) 25^{\circ}$ 0 In the figure, O is the centre of a circle and diameter AB bisects A 15. В F the chord *CD* at a point *E* such that CE=ED=8 cm and EB=4cm. The radius of the circle is (1) 8cm (2) 4cm (3) 6cm (4)10cm S R In the figure, PQRS and PTVS are two cyclic 16. 100° quadrilaterals, If  $\angle QRS = 100^{\circ}$ , then  $\angle TVS =$ (1)  $80^{\circ}$ (2) 100°  $(3) 70^{\circ}$  $(4) 90^{\circ}$ Т 17. If one angle of a cyclic quadrilateral is  $75^{\circ}$ , then the Q opposite angle is Р  $(1) 100^{\circ}$  $(3) 85^{\circ}$ (2)  $105^{\circ}$  $(4) 90^{\circ}$ 

۲

182 9th Standard Mathematics

18. In the figure, ABCD is a cyclic quadrilateral in which DC produced to E and CF is drawn parallel to ABsuch that  $\angle ADC = 80^{\circ}$  and  $\angle ECF = 20^{\circ}$ , then  $\angle BAD = ?$ (1) 100° (2) 20° (3) 120° (4) 110°

۲



S

(4) 6cm.

- 19. AD is a diameter of a circle and AB is a chord If AD = 30 cm and AB = 24 cm then the distance of AB from the centre of the circle is
  - (1) 10cm (2) 9cm (3) 8cm
- 20. In the given figure, If OP = 17cm, PQ = 30 cm and OS is perpendicular to PQ, then RS is
  - (1) 10cm (2) 6cm
  - (3) 7cm (4) 9cm.

#### Points to Remember

- In a parallelogram the opposite sides and opposite angles are equal
- The diagonals of a parallelogram bisect each other.
- The diagonals of a parallelogram divides it into two congruent triangles
- A quadrilateral is a parallelogram if its opposite sides are equal.
- Parallelogram on the same base and between same parallel are equal in area.
- Triangles on the same base and between same parallel are equal in area.
- Parallelogram is a rhombus if its diagonals are perpendicular.
- There is one and only one circle passing through three non-collinear points.
- Equal chords of a circle subtend equal angles at the centre.
- Perpendicular from the centre of a circle to a chord bisects the chord.
- Equal chords of a circle are equidistant from the centre.
- The angle substended by an arc of a circle at the centre is double the angle subtended by it at any point on the remaining part of the circle.
- The angle in a semi circle is a right angle.
- Angles in the same segment of a circle are equal.
- The sum of either pair of opposite angle of a cyclic quadrilateral is 180°.
- If one side of a cyclic quadrilateral is produced then the exterior angle is equal to the interior opposite angle.

Geometry 183

# ICT Corner-1

Expected Result is shown in this picture

# Step – 1

Open the Browser and copy and paste the Link given below (or) by typing the URL given (or) Scan the QR Code.

# Step - 2

GeoGebra worksheet "Properties: Parallelogram" will appear. There are two sliders named "Rotate" and "Page"

# Step-3

Drag the slider named "Rotate "and see that the triangle is doubled as parallelogram.

# Step-4

( )

Drag the slider named "Page" and you will get three pages in which the Properties are explained.

# BSGG\_9\_MAT\_EM\_T3

# ICT Corner-2

Expected Result is shown in this picture



#### Step – 1

Open the Browser type the URL Link given below (or) Scan the QR Code. GeoGebra work sheet named "Angles in a circle" will open. In the work sheet there are two activities on Circles.

The first activity is the relation between Angle at the circumference and the angle at the centre. You can change the angle by moving the slider. Also, you can drag on the point A, C and D to change the position and the radius. Compare the angles at A and O.

۲

#### Step - 2

The second activity is "Angles in the segment of a circle". Drag the points B and D and check the angles. Also drag "Move" to change the radius and chord length of the circle.



#### Scan the QR Code.

184 9<sup>th</sup> Standard Mathematics

