



GOVERNMENT OF TAMIL NADU

HIGHER SECONDARY SECOND YEAR

MATHEMATICS

VOLUME - I

A publication under Free Textbook Programme of Government of Tamil Nadu

Department of School Education

Untouchability is Inhuman and a Crime

۲

 \bigcirc

Government of Tamil Nadu

First Edition - 2019

(Published under New Syllabus)

NOT FOR SALE

Content Creation



۲

State Council of Educational Research and Training © SCERT 2019

Printing & Publishing



Tamil NaduTextbook and Educational Services Corporation

www.textbooksonline.tn.nic.in

HOW TO USE THE BOOK? Awareness on the scope of higher educational opportunities; courses, **Scope of Mathematics** institutions and required competitive examinations. Possible financial assistance to help students climb academic ladder. • Overview of the unit Learning Objectives: Give clarity on the intended learning outcomes of the unit. Visual representation of concepts with illustrations Videos, animations, and tutorials. To increase the span of attention of concepts To visualize the concepts for strengthening and understanding ICT • To link concepts related to one unit with other units. To utilize the digital skills in classroom learning and providing students experimental learning. Recapitulation of the salient points of each chapter for recalling the Summary concepts learnt. Assessing student's understanding of concepts and get them acquainted **Evaluation** with solving exercise problems. **Books for Reference** · List of relevant books for further reading. • To motivate students aspiring to take up competitive examinations such **Scope for** as JEE, KVPY, Math olympiad, etc., the concepts and questions based on **Higher Order Thinking** Higher Order Thinking are incorporated in the content of this book. · Frequently used Mathematical terms have been given with their Glossary Tamil equivalents.

Mathematics Learning

The correct way to learn is to understand the concepts throughly. Each chapter opens with an Introduction, Learning Objectives, Various Definitions, Theorems, Results and Illustrations. These in turn are followed by solved examples and exercise problems which have been classified in to various types for quick and effective revision. One can develop the skill of solving mathematical problems only by doing them. So the teacher's role is to teach the basic concepts and problems related to it and to scaffold students to try the other problems on their own. Since the second year of Higher Secondary is considered to be the foundation for learning higher mathematics, the students must be given more attention to each and every concept mentioned in this book.

۲

۲

()

CONTENTS

۲

|--|

CHAPTER 1	- Applications of Matrices and Determinants	1
1.1	Introduction	1
1.2	Inverse of a Non-Singular Square Matrix	2
1.3	Elementary Transformations of a Matrix	16
1.4	Applications of Matrices: Solving System of Linear Equations	27
1.5	Applications of Matrices: Consistency of system of linear equations by rank method	38
CHAPTER 2	2 – Complex Numbers	52
2.1	Introduction to Complex Numbers	52
2.2	Complex Numbers	54
2.3	Basic Algebraic Properties of Complex Numbers	58
2.4	Conjugate of a Complex Number	60
2.5	Modulus of a Complex Number	66
2.6	Geometry and Locus of Complex Numbers	73
2.7	Polar and Euler form of a Complex Number	75
2.8	de Moivre's Theorem and its Applications	83
CHAPTER 3	3 – Theory of Equations	97
3.1	Introduction	97
3.2	Basics of Polynomial Equations	99
3.3	Vieta's Formulae and Formation of Polynomial Equations	100
3.4	Nature of Roots and Nature of Coefficients of Polynomial Equations	107
3.5	Applications to Geometrical Problems	111
3.6	Roots of Higher Degree Polynomial Equations	112
3.7	Polynomials with Additional Information	113
3.8	Polynomial Equations with no additional information	118
3.9	Descartes Rule	124
CHAPTER 4	- Inverse Trigonometric Functions	129
4.1	Introduction	129
4.2	Some Fundamental Concepts	130
4.3	Sine Function and Inverse Sine Function	133
4.4	The Cosine Function and Inverse Cosine Function	138
4.5	The Tangent Function and the Inverse Tangent Function	143

۲

۲

4.6	The Cosecant Function and the Inverse Cosecant Function	148
4.7	The Secant Function and Inverse Secant Function	149
4.8	The Cotangent Function and the Inverse Cotangent Function	151
4.9	Principal Value of Inverse Trigonometric Functions	153
4.10	Properties of Inverse Trigonometric Functions	155
CHAPTER 5	- Two Dimensional Analytical Geometry-II	172
5.1	Introduction	172
5.2	Circle	173
5.3	Conics	182
5.4	Conic Sections	197
5.5	Parametric form of Conics	199
5.6	Tangents and Normals to Conics	201
5.7	Real life Applications of Conics	207
CHAPTER 6	- Applications of Vector Algebra	221
6.1	Introduction	221
6.2	Geometric Introduction to Vectors	222
6.3	Scalar Product and Vector Product	224
6.4	Scalar triple product	231
6.5	Vector triple product	238
6.6	Jacobi's Identity and Lagrange's Identity	239
6.7	Different forms of Equation of a Straight line	242
6.8	Different forms of Equation of a plane	255
6.9	Image of a point in a plane	273
6.10	Meeting point of a line and a plane	274
ANSWERS		282
CLOSSARV		292



E-book





DIGI links



Lets use the QR code in the text books ! How ?

- Download the QR code scanner from the Google PlayStore/ Apple App Store into your smartphone
 Open the QR code scanner application

• Once the scanner button in the application is clicked, camera opens and then bring it closer to the QR code in the text book.

۲

Scope for students after completing Higher Secondary

۲

EXAMS

EXAMS	

J	oint Entrance Examination (JEE) Main
Purpose	For Admission in B. E./B. Tech., B. Arch., B. Planning
Eligibility	Class 12 pass (PCM)
Application Mode	Online
Source	http://jeemain.nic.in
	JEE Advanced
Purpose	Admission in UG programmes in IITs and ISM Dhanbad
Eligibility	Class 12 Pass (PCM)
Application Mode	Online
Source	http://jeeadv.iitd.ac.in/
Indian	Maritime University Common Entrance Test
Purpose	Admission in Diploma in Nautical Science (DNS) leading to BSc. (Nautical Science)
Eligibility	Class 12 (PCM)
Application Mode	By post
Source	www.imu.edu.in/index.php
	Indian Navy B.Tech Entry Scheme
Purpose	Admission in Indian Navy B.Tech course
Eligibility	Class 12 passed

	indian Navy Sailors Recruitment
Purpose	Admission in 24 weeks Basic training at INS Chilka followed by Professional training
Eligibility	Class 12 (Maths and Physics/Chem./Bio/Computer Sc.)
Application Mode	Online, By Post
Source	www.nausena-bharti.nic.in/index.php
Defence Ir	idian Army Technical Entry Scheme (TES)
Purpose	Technical Entry to Army
Eligibility	Class 12 PCM
Application Mode	Online
Source	www.joinindianarmy.nic.in/
National Defence Acad	emy and Naval Academy Examination (I)
Purpose	Admission to Army and Air Force wings of NDA and
	4 years B. Tech course for the Indian Naval Academy
	Course (INAC)
Eligibility	Class 12 Passed
Application Mode	Online
Source	www.upsc.gov.in/

www.nausena-bharti.nic.in/index.php

Application Mode Online

Source

۲

۲

Scope for students after completing Higher Secondary

۲



If you have aspiration to become a scientist/teacher, you can quired in XII standard mathematics will definitely elevate you to a do degree course in mathematics in any college of your choice. mathematics with your knowledge which ac-Doing B.Sc., better career.

There are some institutions such as IIT's, IISc., ISI's and Anna University which admit students at XII Standard level for their Integrated Courses leading to the award of M.Sc., degree in mathematics/engineering/Statistics/ Computer Science

- 4-year Integrated B.Sc-B.Ed. In Mathematics, the following degrees programmes are offered:

- 3-year BSc

- 3-year B. Math - 4-year B. Tech

- 5-year Integrated M.Sc /MS - 4-year BS

BITSAT

Purpose- For admission to Integrated First Degree Programmes in BITS Pilani Goa & Hyderabad campuses.

National Aptitude Test in Architecture (NATA)

Purpose- Admission to B.Arch. program

Kishore Vaigyanik Protsahan Yojana (KVPY)

Purpose-Fellowship and admission to IISc Banglore in 4 year BS Degree

Indian Statistical Institute (ISI) Admission

Purpose-Admission in B Stat (Hons), B Math (Hons)

Chennai Mathematical Institute

Purpose- Admission in BSc (Hons) in Mathematics and Computer Science, BSc (Hons) in Mathematics and Physics

Job Opportunities

- You have acquired sufficient analytical still to appear for competitive examinations conducted by various organizations such as TNPSC, UPSC, BSRB, RSB.
- You are placed with better opportunity to get recruitment directly into some of the top Multi National Companies.
- You can plan to take up engineering degree course to acquire BE/B.TECH es. Institutions such as IIT's, IISc, NIT's conduct All India Level Entrance in any of the leading Technical Institutions/Universities/Engineering colleg-Examinations to admit students.

۲

 Scientist Jobs in ISRO, DRDO, CSIR labs 	Staff Selection Commission	Public Sector Bank	Statistical Investigator	Tamil Nadu Public Service Commission	
Indian Forest Services	Union Public Service Commission	Indian Defense Services	Tax Assistant	Combined Graduate Level Exam	Teaching Profession

۲

Scholarship and Research Opportunities

۲

FINANCIAL ASSISTANCE

- Scholarship for Graduate and Post Graduate courses
- NTSE at the end of X (from class XI to Ph.D)
- International Olympiad: for getting stipend for Higher Education in Science and Mathematics
- DST INSPIRE Scholarships (for UG and PG)
- DST INSPIRE Fellowships (for Ph.D)
- UGC National Fellowships (for Ph.D)
- Indira Gandhi Fellowship for Single Girl Child (for UG and PG)
- Moulana Azad Fellowship for minorities (for Ph.D)
- In addition various fellowships for SC / ST / PWD / OBC etc are available. (Visit website of University Grants Commission (UGC) and Department of Science and Technology (DST))
- University Fellowships
- Tamil Nadu Collegiate Education fellowship.

100

ິ

NAME OF THE INSTITUTION	WEBSITE
ndian Institute of Science (IISc) Bangalore	www.iisc.ac.in
Chennai Mathematical Institute (CMI) Chennaie	www.cmi.ac.in
ata Institue of Fundamental Research (TIFR) Mumbai	www.tifr.res.in
ndian Institute of Space Science and Technology (IIST) invandrum	www.iist.ac.in
Vational Institute of Science Education and Research (NIS-ER)	www.niser.ac.in
sirla Institute of Technology and Science, Pilani	www.bits-pilani.ac.in
ndian Institute of Science Education and Research	www.iiseradmission.in
Anna University	https://www.annauniv.edu/
ndian Institute of Technology in various places (IIT's)	www.iitm.ac.in
Vational Institute of Technology (NITs)	www.nitt.edu
Central Universities	www.cucet.ac.in
state Universities	www.ugc.ac.in
amil Nadu agricultural University (tnau.ac.in)	tnau.ac.in
ntemational Institute of Information Technology	www.iiit.ac.in
he Institute of Mathematical Sciences (IMSC) Chennai.	www.imsc.res.in
łyderabad Central university, Hyderabad.	www.uohyd.ac.in
Jelhi University, Delhi	www.du.ac.in
Aumbai University, Mumbai	www.mu.ac.in
savithiri Bai Phule Pune University, Pune	www.unipune.ac.in

۲

۲

VIII

Chapter

Applications of Matrices and Determinants



"The greatest mathematicians, as Archimedes, Newton, and Gauss, always united theory and applications in equal measure." -Felix Klein

1.1 Introduction



Carl Friedrich Gauss

Matrices are very important and indispensable in handling system of linear equations which arise as mathematical models of real-world problems. Mathematicians Gauss, Jordan, Cayley, and Hamilton have developed the theory of matrices which has been used in investigating solutions of systems of linear equations.

In this chapter, we present some applications of matrices in solving system of linear equations. To be specific, we study four methods, namely (i) Matrix inversion method, (ii) Cramer's rule (iii) Gaussian elimination method, and (iv) Rank method. Before knowing these methods, we introduce the following: (i) Inverse of a non-singular

(1777-1855) square matrix, (ii) Rank of a matrix, (iii) Elementary row and column German mathematician and transformations, and (iv) Consistency of system of linear equations. physicist

Solution Learning Objectives

Upon completion of this chapter, students will be able to

- Demonstrate a few fundamental tools for solving systems of linear equations:
 - Adjoint of a square matrix
 - Inverse of a non-singular matrix
 - Elementary row and column operations
 - Row-echelon form
 - Rank of a matrix
- Use row operations to find the inverse of a non-singular matrix
 - Illustrate the following techniques in solving system of linear equations by
 - Matrix inversion method
 - Cramer's rule
 - Gaussian elimination method
- Test the consistency of system of non-homogeneous linear equations
- Test for non-trivial solution of system of homogeneous linear equations

1.2 Inverse of a Non-Singular Square Matrix

We recall that a square matrix is called a **non-singular matrix** if its determinant is not equal to zero and a square matrix is called **singular** if its determinant is zero. We have already learnt about multiplication of a matrix by a scalar, addition of two matrices, and multiplication of two matrices. But a rule could not be formulated to perform division of a matrix by another matrix since a matrix is just an arrangement of numbers and has no numerical value. When we say that, a matrix A is of order n, we mean that A is a square matrix having n rows and n columns.

۲

In the case of a real number $x \neq 0$, there exists a real number $y\left(=\frac{1}{x}\right)$, called the inverse (or reciprocal) of x such that xy = yx = 1. In the same line of thinking, when a matrix A is given, we search for a matrix B such that the products AB and BA can be found and AB = BA = I, where I is a unit matrix.

In this section, we define the inverse of a non-singular square matrix and prove that a non-singular square matrix has a unique inverse. We will also study some of the properties of inverse matrix. For all these activities, we need a matrix called the adjoint of a square matrix.

1.2.1 Adjoint of a Square Matrix

We recall the properties of the cofactors of the elements of a square matrix. Let A be a square matrix of by order n whose determinant is denoted |A| or det(A). Let a_{ij} be the element sitting at the intersection of the i^{th} row and j^{th} column of A. Deleting the i^{th} row and j^{th} column of A, we obtain a sub-matrix of order (n-1). The determinant of this sub-matrix is called **minor** of the element a_{ij} . It is denoted by M_{ij} . The product of M_{ij} and $(-1)^{i+j}$ is called **cofactor** of the element a_{ij} . It is denoted by A_{ij} . Thus **the cofactor of** a_{ij} is $A_{ij} = (-1)^{i+j} M_{ij}$.

An important property connecting the elements of a square matrix and their cofactors is that the sum of the products of the entries (elements) of a row and the corresponding cofactors of the elements of the same row is equal to the determinant of the matrix; and the sum of the products of the entries (elements) of a row and the corresponding cofactors of the elements of any other row is equal to 0. That is,

$$a_{i1}A_{j1} + a_{i2}A_{j2} + \dots + a_{in}A_{jn} = \begin{cases} |A| & \text{if } i = j \\ 0 & \text{if } i \neq j, \end{cases}$$

where |A| denotes the determinant of the square matrix A. Here |A| is read as "determinant of A" and not as "modulus of A". Note that |A| is just a real number and it can also be negative. For

instance, we have $\begin{vmatrix} 2 & 1 & 1 \\ 1 & 1 & 1 \\ 2 & 2 & 1 \end{vmatrix} = 2(1-2) - 1(1-2) + 1(2-2) = -2 + 1 + 0 = -1.$

Definition 1.1

Let *A* be a square matrix of order *n*. Then the **matrix of cofactors** of *A* is defined as the matrix obtained by replacing each element a_{ij} of *A* with the corresponding cofactor A_{ij} . The **adjoint matrix** of *A* is defined as the transpose of the matrix of cofactors of *A*. It is denoted by adj *A*.

()

Note

adj *A* is a square matrix of order *n* and adj $A = \begin{bmatrix} A_{ij} \end{bmatrix}^T = \begin{bmatrix} (-1)^{i+j} M_{ij} \end{bmatrix}^T$.

In particular, adj A of a square matrix of order 3 is given below:

adj
$$A = \begin{bmatrix} +M_{11} & -M_{12} & +M_{13} \\ -M_{21} & +M_{22} & -M_{23} \\ +M_{31} & -M_{32} & +M_{33} \end{bmatrix}^{T} = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{bmatrix}^{T} = \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix}.$$

Theorem 1.1

For every square matrix A of order n, $A(\operatorname{adj} A) = (\operatorname{adj} A)A = |A|I_n$.

Proof

For simplicity, we prove the theorem for n = 3 only.

Consider
$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$
. Then, we get
$$a_{11}A_{11} + a_{12}A_{12} + a_{13}A_{13} = |A|, \quad a_{11}A_{21} + a_{12}A_{22} + a_{13}A_{23} = 0, \qquad a_{11}A_{31} + a_{12}A_{32} + a_{13}A_{33} = 0;$$
$$a_{21}A_{11} + a_{22}A_{12} + a_{23}A_{13} = 0, \qquad a_{21}A_{21} + a_{22}A_{22} + a_{23}A_{23} = |A|, \quad a_{21}A_{31} + a_{22}A_{32} + a_{23}A_{33} = 0;$$
$$a_{31}A_{11} + a_{32}A_{12} + a_{33}A_{13} = 0, \qquad a_{31}A_{21} + a_{32}A_{22} + a_{33}A_{23} = 0, \qquad a_{31}A_{31} + a_{32}A_{32} + a_{33}A_{33} = |A|.$$

By using the above equations, we get

$$A(\operatorname{adj} A) = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix} = \begin{bmatrix} |A| & 0 & 0 \\ 0 & |A| & 0 \\ 0 & 0 & |A| \end{bmatrix} = |A| \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = |A| I_3 \qquad \dots (1)$$
$$(\operatorname{adj} A) A = \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} |A| & 0 & 0 \\ 0 & |A| & 0 \\ 0 & 0 & |A| \end{bmatrix} = |A| \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = |A| I_3, \qquad \dots (2)$$

where I_3 is the identity matrix of order 3.

So, by equations (1) and (2), we get $A(\operatorname{adj} A) = (\operatorname{adj} A)A = |A|I_3$.

Note

۲

If A is a singular matrix of order n, then |A| = 0 and so $A(\operatorname{adj} A) = (\operatorname{adj} A)A = O_n$, where O_n denotes zero matrix of order n.

Example 1.1

If
$$A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$$
, verify that $A(adj A) = (adj A)A = |A||I_3$.
Solution
We find that $|A| = \begin{vmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{vmatrix} = 8(21-16) + 6(-18+8) + 2(24-14) = 40 - 60 + 20 = 0.$

3

By the definition of adjoint, we get

adj
$$A = \begin{bmatrix} (21-16) & -(-18+8) & (24-14) \\ -(-18+8) & (24-4) & -(-32+12) \\ (24-14) & -(-32+12) & (56-36) \end{bmatrix}^{T} = \begin{bmatrix} 5 & 10 & 10 \\ 10 & 20 & 20 \\ 10 & 20 & 20 \end{bmatrix}$$

So, we get

$$A(\operatorname{adj} A) = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix} \begin{bmatrix} 5 & 10 & 10 \\ 10 & 20 & 20 \\ 10 & 20 & 20 \end{bmatrix}$$
$$= \begin{bmatrix} 40 - 60 + 20 & 80 - 120 + 40 \\ -30 + 70 - 40 & -60 + 140 - 80 \\ 10 - 40 + 30 & 20 - 80 + 60 & 20 - 80 + 60 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = 0I_3 = |A|I_3,$$

Similarly, we get

$$(adj A)A = \begin{bmatrix} 5 & 10 & 10 \\ 10 & 20 & 20 \\ 10 & 20 & 20 \end{bmatrix} \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$$
$$= \begin{bmatrix} 40 - 60 + 20 & -30 + 70 - 40 & 10 - 40 + 30 \\ 80 - 120 + 40 & -60 + 140 - 80 & 20 - 80 + 60 \\ 80 - 120 + 40 & -60 + 140 - 80 & 20 - 80 + 60 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = 0I_3 = |A|I_3.$$

Hence, $A(\operatorname{adj} A) = (\operatorname{adj} A)A = |A|I_3$.

1.2.2 Definition of inverse matrix of a square matrix

Now, we define the inverse of a square matrix.

Definition 1.2

Let A be a square matrix of order n. If there exists a square matrix B of order n such that $AB = BA = I_n$, then the matrix B is called an **inverse** of A.

Theorem 1.2

If a square matrix has an inverse, then it is unique.

Proof

Let *A* be a square matrix order *n* such that an inverse of *A* exists. If possible, let there be two inverses *B* and *C* of *A*. Then, by definition, we have $AB = BA = I_n$ and $AC = CA = I_n$.

Using these equations, we get

 $C = CI_n = C(AB) = (CA)B = I_nB = B.$

Hence the uniqueness follows.

Notation The inverse of a matrix A is denoted by A^{-1} .

Note

 $AA^{-1} = A^{-1}A = I_n$.

XII - Mathematics

Theorem 1.3

Let A be square matrix of order n. Then, A^{-1} exists if and only if A is non-singular.

Proof

Suppose that A^{-1} exists. Then $AA^{-1} = A^{-1}A = I_n$.

By the product rule for determinants, we get

 $\det(AA^{-1}) = \det(A)\det(A^{-1}) = \det(A^{-1})\det(A) = \det(I_n) = 1.$ So, $|A| = \det(A) \neq 0.$

Hence A is non-singular.

Conversely, suppose that A is non-singular.

Then $|A| \neq 0$. By Theorem 1.1, we get

 $A(\operatorname{adj} A) = (\operatorname{adj} A)A = |A|I_n.$

So, dividing by |A|, we get $A\left(\frac{1}{|A|} \operatorname{adj} A\right) = \left(\frac{1}{|A|} \operatorname{adj} A\right) A = I_n$.

Thus, we are able to find a matrix $B = \frac{1}{|A|} \operatorname{adj} A$ such that $AB = BA = I_n$.

Hence, the inverse of A exists and it is given by $A^{-1} = \frac{1}{|A|} \operatorname{adj} A$.

Remark

The determinant of a singular matrix is 0 and so a singular matrix has no inverse.

Example 1.2

If
$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$
 is non-singular, find A^{-1} .

Solution

We first find adj A. By definition, we get adj $A = \begin{bmatrix} +M_{11} & -M_{12} \\ -M_{21} & +M_{22} \end{bmatrix}^{T} = \begin{bmatrix} d & -c \\ -b & a \end{bmatrix}^{T} = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$

Since A is non-singular, $|A| = ad - bc \neq 0$.

As
$$A^{-1} = \frac{1}{|A|} \operatorname{adj} A$$
, we get $A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$.

Example 1.3

Find the inverse of the matrix $\begin{bmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{bmatrix}$.

Solution

Let
$$A = \begin{bmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{bmatrix}$$
. Then $|A| = \begin{vmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{vmatrix} = 2(7) + (-12) + 3(-1) = -1 \neq 0$.

Therefore, A^{-1} exists. Now, we get

Applications of Matrices and Determinants

Chapter 1 Matrices.indd 5

$$\operatorname{adj} A = \begin{bmatrix} + \begin{vmatrix} 3 & 1 \\ 2 & 3 \end{vmatrix} - \begin{vmatrix} -5 & 1 \\ -3 & 3 \end{vmatrix} + \begin{vmatrix} -5 & 3 \\ -3 & 2 \end{vmatrix} \\ - \begin{vmatrix} -1 & 3 \\ 2 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 3 \\ -3 & 3 \end{vmatrix} - \begin{vmatrix} 2 & -1 \\ -3 & 2 \end{vmatrix} \\ + \begin{vmatrix} -1 & 3 \\ -3 & 3 \end{vmatrix} - \begin{vmatrix} 2 & 3 \\ -3 & 2 \end{vmatrix} + \begin{vmatrix} 2 & -1 \\ -3 & 2 \end{vmatrix} \\ = \begin{bmatrix} 7 & 12 & -1 \\ 9 & 15 & -1 \\ -10 & -17 & 1 \end{bmatrix}^{T} = \begin{bmatrix} 7 & 9 & -10 \\ 12 & 15 & -17 \\ -1 & -1 & 1 \end{bmatrix}.$$

Hence, $A^{-1} = \frac{1}{|A|} (\operatorname{adj} A) = \frac{1}{(-1)} \begin{bmatrix} 7 & 9 & -10 \\ 12 & 15 & -17 \\ -1 & -1 & 1 \end{bmatrix} = \begin{bmatrix} -7 & -9 & 10 \\ -12 & -15 & 17 \\ 1 & 1 & -1 \end{bmatrix}.$

1.2.3 Properties of inverses of matrices

We state and prove some theorems on non-singular matrices.

If A is non-s

Theorem 1.4

(i)
$$|A^{-1}| = \frac{1}{|A|}$$
 (ii) $(A^{T})^{-1} = (A^{-1})^{T}$ (iii) $(\lambda A)^{-1} = \frac{1}{\lambda} A^{-1}$, where λ is a non-zero scalar.

Proof

Let A be non-singular. Then $|A| \neq 0$ and A^{-1} exists. By definition,

$$AA^{-1} = A^{-1}A = I_n.$$
 ...(1)

(i) By (1), we get
$$|AA^{-1}| = |A^{-1}A| = |I_n|$$
.

Using the product rule for determinants, we get $|A||A^{-1}| = |I_n| = 1$.

Hence, $|A^{-1}| = \frac{1}{|A|}$.

(ii) From (1), we get
$$(AA^{-1})^T = (A^{-1}A)^T = (I_n)^T$$

Using the reversal law of transpose, we get $(A^{-1})^T A^T = A^T (A^{-1})^T = I_n$. Hence

$$\left(A^{T}\right)^{-1} = \left(A^{-1}\right)^{T}.$$

(iii) Since λ is a non-zero scalar, from (1), we get $(\lambda A) \left(\frac{1}{\lambda} A^{-1}\right) = \left(\frac{1}{\lambda} A^{-1}\right) (\lambda A) = I_n$.

So,
$$(\lambda A)^{-1} = \frac{1}{\lambda} A^{-1}$$

Theorem 1.5 (Left Cancellation Law)

Let A, B, and C be square matrices of order n. If A is non-singular and AB = AC, then B = C.

Proof

Since *A* is non-singular, A^{-1} exists and $AA^{-1} = A^{-1}A = I_n$. Taking AB = AC and pre-multiplying both sides by A^{-1} , we get $A^{-1}(AB) = A^{-1}(AC)$. By using the associative property of matrix multiplication and property of inverse matrix, we get B = C.

6

XII - Mathematics

Theorem1.6 (Right Cancellation Law)

Let A, B, and C be square matrices of order n. If A is non-singular and BA = CA, then B = C.

۲

Proof

Since *A* is non-singular, A^{-1} exists and $AA^{-1} = A^{-1}A = I_n$. Taking BA = CA and post-multiplying both sides by A^{-1} , we get $(BA)A^{-1} = (CA)A^{-1}$. By using the associative property of matrix multiplication and property of inverse matrix, we get B = C.

Note

If A is singular and AB = AC or BA = CA, then B and C need not be equal. For instance, consider the following matrices:

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}, B = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \text{ and } C = \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix}.$$

We note that |A| = 0 and AB = AC; but $B \neq C$.

Theorem 1.7 (Reversal Law for Inverses)

If A and B are non-singular matrices of the same order, then the product AB is also non-singular and $(AB)^{-1} = B^{-1}A^{-1}$.

Proof

Assume that *A* and *B* are non-singular matrices of same order *n*. Then, $|A| \neq 0$, $|B| \neq 0$, both A^{-1} and B^{-1} exist and they are of order *n*. The products *AB* and $B^{-1}A^{-1}$ can be found and they are also of order *n*. Using the product rule for determinants, we get $|AB| = |A||B| \neq 0$. So, *AB* is non-singular and

$$(AB)(B^{-1}A^{-1}) = (A(BB^{-1}))A^{-1} = (AI_n)A^{-1} = AA^{-1} = I_n;$$

$$(B^{-1}A^{-1})(AB) = (B^{-1}(A^{-1}A))B = (B^{-1}I_n)B = B^{-1}B = I_n.$$

Hence $(AB)^{-1} = B^{-1}A^{-1}$.

Theorem 1.8 (Law of Double Inverse)

If A is non-singular, then A^{-1} is also non-singular and $(A^{-1})^{-1} = A$.

Proof

Assume that A is non-singular. Then $|A| \neq 0$, and A^{-1} exists. Now $|A^{-1}| = \frac{1}{|A|} \neq 0 \Rightarrow A^{-1}$ is also non-singular, and $AA^{-1} = A^{-1}A = I$. Now, $AA^{-1} = I \Rightarrow (AA^{-1})^{-1} = I \Rightarrow (A^{-1})^{-1} A^{-1} = I$ (1) Post-multiplying by A on both sides of equation (1), we get $(A^{-1})^{-1} = A$.



Proof

۲

Since A is a non-singular square matrix, we have $|A| \neq 0$ and so, we get

(i)
$$A^{-1} = \frac{1}{|A|} (\operatorname{adj} A) \Rightarrow \operatorname{adj} A = |A| A^{-1} \Rightarrow (\operatorname{adj} A)^{-1} = (|A| A^{-1})^{-1} = \frac{1}{|A|} (A^{-1})^{-1} = \frac{1}{|A|} A$$

Replacing A by A^{-1} in $\operatorname{adj} A = |A| A^{-1}$, we get $\operatorname{adj} (A^{-1}) = |A^{-1}| (A^{-1})^{-1} = \frac{1}{|A|} A$.
Hence, we get $(\operatorname{adj} A)^{-1} = \operatorname{adj} (A^{-1}) = \frac{1}{|A|} A$.
(ii) $A(\operatorname{adj} A) = (\operatorname{adj} A)A = |A| I_n \Rightarrow \det(A(\operatorname{adj} A)) = \det((\operatorname{adj} A)A) = \det(|A| I_n)$
 $\Rightarrow |A| |\operatorname{adj} A| = |A|^n \Rightarrow |\operatorname{adj} A| = |A|^{n-1}$.

(iii) For any non-singular matrix B of order n, we have $B(\operatorname{adj} B) = (\operatorname{adj} B)B = |B|I_n$.

Put $B = \operatorname{adj} A$. Then, we get $(\operatorname{adj} A)(\operatorname{adj}(\operatorname{adj} A)) = |\operatorname{adj} A| I_n$. So, since $|\operatorname{adj} A| = |A|^{n-1}$, we get $(\operatorname{adj} A)(\operatorname{adj}(\operatorname{adj} A)) = |A|^{n-1} I_n$. Pre-multiplying both sides by A, we get $A((\operatorname{adj} A)(\operatorname{adj} (\operatorname{adj} A))) = A(|A|^{n-1} I_n)$. Using the associative property of matrix multiplication, we get $(A(\operatorname{adj} A))\operatorname{adj}(\operatorname{adj} A) = A(|A|^{n-1} I_n)$.

Hence, we get $(|A|I_n)(\operatorname{adj}(\operatorname{adj} A)) = |A|^{n-1} A$. That is, $\operatorname{adj}(\operatorname{adj} A) = |A|^{n-2} A$.

(iv) Replacing A by λA in $\operatorname{adj}(A) = |A| A^{-1}$ where λ is a non-zero scalar, we get $\operatorname{adj}(\lambda A) = |\lambda A| (\lambda A)^{-1} = \lambda^n |A| \frac{1}{2} A^{-1} = \lambda^{n-1} |A| A^{-1} = \lambda^{n-1} \operatorname{adj}(A)$.

- (v) By (iii), we have $\operatorname{adj}(\operatorname{adj} A) = |A|^{n-2} A$. So, by taking determinant on both sides, we get $|\operatorname{adj}(\operatorname{adj} A)| = ||A|^{n-2} A| = (|A|^{(n-2)})^n |A| = |A|^{n^2-2n+1} = |A|^{(n-1)^2}$.
- (vi) Replacing A by A^T in $A^{-1} = \frac{1}{|A|} \operatorname{adj} A$, we get $(A^T)^{-1} = \frac{1}{|A^T|} \operatorname{adj}(A^T)$ and hence, we get $\operatorname{adj}(A^T) = |A^T| (A^T)^{-1} = |A| (A^{-1})^T = (|A|A^{-1})^T = (|A|\frac{1}{|A|}\operatorname{adj} A)^T = (\operatorname{adj} A)^T$.

XII - Mathematics

Note

If A is a non-singular matrix of order 3, then, $|A| \neq 0$. By theorem 1.9 (*ii*), we get $|\operatorname{adj} A| = |A|^2$

and so, |adj A| is positive. Then, we get $|A| = \pm \sqrt{|adj A|}$.

So, we get $A^{-1} = \pm \frac{1}{\sqrt{|\operatorname{adj} A|}} \operatorname{adj} A$.

Further, by property (*iii*), we get $A = \frac{1}{|A|} \operatorname{adj}(\operatorname{adj} A)$.

Hence, if A is a non-singular matrix of order 3, then we get $A = \pm \frac{1}{\sqrt{|\operatorname{adj} A|}} \operatorname{adj}(\operatorname{adj} A)$.

Example 1.4

If A is a non-singular matrix of odd order, prove that |adj A| is positive.

Solution

Let *A* be a non-singular matrix of order 2m+1, where $m = 0, 1, 2, \cdots$. Then, we get $|A| \neq 0$ and, by theorem 1.9 (ii), we have $|\operatorname{adj} A| = |A|^{(2m+1)-1} = |A|^{2m}$.

Since $|A|^{2m}$ is always positive, we get that |adj A| is positive.

Example 1.5

Find a matrix A if $adj(A) = \begin{bmatrix} 7 & 7 & -7 \\ -1 & 11 & 7 \\ 11 & 5 & 7 \end{bmatrix}$.

Solution

First, we find $|adj(A)| = \begin{vmatrix} 7 & 7 & -7 \\ -1 & 11 & 7 \\ 11 & 5 & 7 \end{vmatrix} = 7(77 - 35) - 7(-7 - 77) - 7(-5 - 121) = 1764 > 0.$

So, we get

$$A = \pm \frac{1}{\sqrt{|\text{adj } A|}} \operatorname{adj}(\text{adj } A) = \pm \frac{1}{\sqrt{1764}} \begin{bmatrix} +(77-35) & -(-7-77) & +(-5-121) \\ -(49+35) & +(49+77) & -(35-77) \\ +(49+77) & -(49-7) & +(77+7) \end{bmatrix}^{T}$$
$$= \pm \frac{1}{42} \begin{bmatrix} 42 & 84 & -126 \\ -84 & 126 & 42 \\ 126 & -42 & 84 \end{bmatrix}^{T} = \pm \begin{bmatrix} 1 & -2 & 3 \\ 2 & 3 & -1 \\ -3 & 1 & 2 \end{bmatrix}.$$

Example 1.6

If adj $A = \begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, find A^{-1} .

Applications of Matrices and Determinants

۲

Solution

We compute
$$|\operatorname{adj} A| = \begin{vmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{vmatrix} = 9$$
.
So, we get $A^{-1} = \pm \frac{1}{\sqrt{|\operatorname{adj}(A)|}} \operatorname{adj}(A) = \pm \frac{1}{\sqrt{9}} \begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} = \pm \frac{1}{3} \begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$.

Example1.7

If A is symmetric, prove that adj A is also symmetric.

Solution

Suppose A is symmetric. Then, $A^T = A$ and so, by theorem 1.9 (vi), we get $\operatorname{adj}(A^T) = (\operatorname{adj} A)^T \Rightarrow \operatorname{adj} A = (\operatorname{adj} A)^T \Rightarrow \operatorname{adj} A$ is symmetric.

Theorem 1.10

If A and B are any two non-singular square matrices of order n, then

 $\operatorname{adj}(AB) = (\operatorname{adj} B)(\operatorname{adj} A).$

Proof

۲

Replacing A by AB in $adj(A) = |A|A^{-1}$, we get

$$adj(AB) = |AB|(AB)^{-1} = (|B|B^{-1})(|A|A^{-1}) = adj(B)adj(A).$$

Example 1.8

Verify the property $(A^T)^{-1} = (A^{-1})^T$ with $A = \begin{bmatrix} 2 & 9 \\ 1 & 7 \end{bmatrix}$.

Solution

For the given A, we get |A| = (2)(7) - (9)(1) = 14 - 9 = 5. So, $A^{-1} = \frac{1}{5} \begin{bmatrix} 7 & -9 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} \frac{7}{5} & -\frac{9}{5} \\ -\frac{1}{5} & \frac{2}{5} \end{bmatrix}$.

Then,
$$\left(A^{-1}\right)^{T} = \begin{bmatrix} \frac{7}{5} & -\frac{1}{5} \\ -\frac{9}{5} & \frac{2}{5} \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 7 & -1 \\ -9 & 2 \end{bmatrix}$$
. ... (1)

For the given *A*, We get $A^T = \begin{bmatrix} 2 & 1 \\ 9 & 7 \end{bmatrix}$. So $|A^T| = (2)(7) - (1)(9) = 5$.

Then,
$$(A^T)^{-1} = \frac{1}{5} \begin{bmatrix} 7 & -1 \\ -9 & 2 \end{bmatrix}$$
. ... (2)

From (1) and (2), we get $(A^{-1})^T = (A^T)^{-1}$. Thus, we have verified the given property.

XII - Mathematics

۲

Example 1.9

Verify
$$(AB)^{-1} = B^{-1}A^{-1}$$
 with $A = \begin{bmatrix} 0 & -3 \\ 1 & 4 \end{bmatrix}, B = \begin{bmatrix} -2 & -3 \\ 0 & -1 \end{bmatrix}.$
Solution
We get $AB = \begin{bmatrix} 0 & -3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} -2 & -3 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0+0 & 0+3 \\ -2+0 & -3-4 \end{bmatrix} = \begin{bmatrix} 0 & 3 \\ -2 & -7 \end{bmatrix}$
 $(AB)^{-1} = \frac{1}{(0+6)} \begin{bmatrix} -7 & -3 \\ 2 & 0 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} -7 & -3 \\ 2 & 0 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} -7 & -3 \\ 2 & 0 \end{bmatrix}$... (1)
 $A^{-1} = \frac{1}{(0+3)} \begin{bmatrix} 4 & 3 \\ -1 & 0 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 4 & 3 \\ -1 & 0 \end{bmatrix}$

۲

As the matrices in (1) and (2) are same, $(AB)^{-1} = B^{-1}A^{-1}$ is verified.

Example 1.10

If
$$A = \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix}$$
, find x and y such that $A^2 + xA + yI_2 = O_2$. Hence, find A^{-1} .

 $B^{-1} = \frac{1}{(2-0)} \begin{bmatrix} -1 & 3 \\ 0 & -2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} -1 & 3 \\ 0 & -2 \end{bmatrix}$

 $B^{-1}A^{-1} = \frac{1}{2} \begin{bmatrix} -1 & 3 \\ 0 & -2 \end{bmatrix} \frac{1}{3} \begin{bmatrix} 4 & 3 \\ -1 & 0 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} -7 & -3 \\ 2 & 0 \end{bmatrix}.$

Solution

۲

Since
$$A^2 = \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix} = \begin{bmatrix} 22 & 27 \\ 18 & 31 \end{bmatrix}$$
,
 $A^2 + xA + yI_2 = O_2 \Rightarrow \begin{bmatrix} 22 & 27 \\ 18 & 31 \end{bmatrix} + x \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix} + y \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
 $\Rightarrow \begin{bmatrix} 22 + 4x + y & 27 + 3x \\ 18 + 2x & 31 + 5x + y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$.
 $+ y = 0.31 + 5x + y = 0.27 + 3x = 0$ and $18 + 2x = 0$.

So, we get 22 + 4x + y = 0, 31 + 5x + y = 0, 27 + 3x = 0 and 18 + 2x = 0.

Hence x = -9 and y = 14. Then, we get $A^2 - 9A + 14I_2 = O_2$.

Post-multiplying this equation by A^{-1} , we get $A - 9I_2 + 14A^{-1} = O_2$. Hence, we get

$$A^{-1} = \frac{1}{14} (9I_2 - A) = \frac{1}{14} \left(9 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix} \right) = \frac{1}{14} \begin{bmatrix} 5 & -3 \\ -2 & 4 \end{bmatrix}$$

11

Applications of Matrices and Determinants

... (2)

1.2.4 Application of matrices to Geometry

There is a special type of non-singular matrices which are widely used in applications of matrices to geometry. For simplicity, we consider two-dimensional analytical geometry.

۲

Let *O* be the origin, and x'Ox and y'Oy be the *x*-axis and *y*-axis. Let *P* be a point in the plane whose coordinates are (x, y) with respect to the coordinate system. Suppose that we rotate the *x*-axis and *y*-axis about the origin, through an angle θ as shown in the figure. Let *X'OX* and *Y'OY* be the new *X*-axis and new *Y*-axis. Let (X,Y) be the new set of coordinates of *P* with respect to the new coordinate system. Referring to Fig.1.1, we get



$$x = OL = ON - LN = X\cos\theta - QT = X\cos\theta - Y\sin\theta$$

$$y = PL = PT + TL = QN + PT = X\sin\theta + Y\cos\theta$$

These equations provide transformation of one coordinate system into another coordinate system. The above two equations can be written in the matrix form

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} X \\ Y \end{bmatrix}.$$

Let $W = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$. Then $\begin{bmatrix} x \\ y \end{bmatrix} = W \begin{bmatrix} X \\ Y \end{bmatrix}$ and $|W| = \cos^2\theta + \sin^2\theta = 1$.

So, *W* has inverse and $W^{-1} = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$. We note that $W^{-1} = W^T$. Then, we get the

inverse transformation by the equation

$$\begin{bmatrix} X \\ Y \end{bmatrix} = W^{-1} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x\cos\theta - y\sin\theta \\ x\sin\theta + y\cos\theta \end{bmatrix}.$$

Hence, we get the transformation $X = x \cos \theta - y \sin \theta$, $Y = x \sin \theta + y \cos \theta$.

This transformation is used in **Computer Graphics** and determined by the matrix $W = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$ We note that the matrix W satisfies a special property $W^{-1} = W^{T}$; that is, $WW^{T} = W^{T}W = I$

Definition 1.3

A square matrix A is called **orthogonal** if $AA^{T} = A^{T}A = I$.

Note

A is orthogonal if and only if A is non-singular and $A^{-1} = A^{T}$.

XII - Mathematics

Example 1.11

Prove that
$$\begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$
 is orthogonal.

Solution

Let
$$A = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$
. Then, $A^T = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}^T = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$.

So, we get

$$AA^{T} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$$
$$= \begin{bmatrix} \cos^{2}\theta + \sin^{2}\theta & \cos\theta\sin\theta - \sin\theta\cos\theta \\ \sin\theta\cos\theta - \cos\theta\sin\theta & \sin^{2}\theta + \cos^{2}\theta \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_{2}.$$

Similarly, we get $A^T A = I_2$. Hence $AA^T = A^T A = I_2 \Rightarrow A$ is orthogonal.

Example 1.12

If
$$A = \frac{1}{7} \begin{bmatrix} 6 & -3 & a \\ b & -2 & 6 \\ 2 & c & 3 \end{bmatrix}$$
 is orthogonal, find *a*, *b* and *c*, and hence A^{-1} .

Solution

۲

If A is orthogonal, then $AA^{T} = A^{T}A = I_{3}$. So, we have

$$AA^{T} = I_{3} \Rightarrow \frac{1}{7} \begin{bmatrix} 6 & -3 & a \\ b & -2 & 6 \\ 2 & c & 3 \end{bmatrix}^{\frac{1}{7}} \begin{bmatrix} 6 & b & 2 \\ -3 & -2 & c \\ a & 6 & 3 \end{bmatrix}^{\frac{1}{7}} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\Rightarrow \begin{bmatrix} 45 + a^{2} & 6b + 6 + 6a & 12 - 3c + 3a \\ 6b + 6 + 6a & b^{2} + 40 & 2b - 2c + 18 \\ 12 - 3c + 3a & 2b - 2c + 18 & c^{2} + 13 \end{bmatrix}^{\frac{1}{7}} = 49 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\Rightarrow \begin{bmatrix} 45 + a^{2} = 49 \\ b^{2} + 40 = 49 \\ c^{2} + 13 = 49 \\ 6b + 6 + 6a = 0 \\ 12 - 3c + 3a = 0 \\ 2b - 2c + 18 = 0 \end{bmatrix} \Rightarrow \begin{bmatrix} a^{2} = 4, b^{2} = 9, c^{2} = 36, \\ a + b = -1, a - c = -4, b - c = -9 \end{bmatrix} \Rightarrow a = 2, b = -3, c = 6$$
So, we get $A = \frac{1}{7} \begin{bmatrix} 6 & -3 & 2 \\ -3 & -2 & 6 \\ 2 & 6 & 3 \end{bmatrix}$ and hence, $A^{-1} = A^{T} = \frac{1}{7} \begin{bmatrix} 6 & -3 & 2 \\ -3 & -2 & 6 \\ 2 & 6 & 3 \end{bmatrix}$.

13

Applications of Matrices and Determinants

1.2.5 Application of matrices to Cryptography

One of the important applications of inverse of a non-singular square matrix is in cryptography. Cryptography is an art of communication between two people by keeping the information not known to others. It is based upon two factors, namely encryption and decryption. Encryption means the process of transformation of an information (plain form) into an unreadable form (coded form). On the other hand, **Decryption** means the

transformation of the coded message back into original form. Encryption and decryption require a secret technique which is known only to the sender and the receiver.

This secret is called a key. One way of generating a key is by using a non-singular matrix to encrypt a message by the sender. The receiver decodes (decrypts) the message to retrieve the original message by using the inverse of the matrix. The matrix used for encryption is called encryption matrix (encoding matrix) and that used for decoding is called decryption matrix (decoding matrix).

We explain the process of encryption and decryption by means of an example.

Suppose that the sender and receiver consider messages in alphabets A-Z only, both assign the numbers 1-26 to the letters A-Z respectively, and the number 0 to a blank space. For simplicity, the sender employs a key as post-multiplication by a non-singular matrix of order 3 of his own choice. The receiver uses post-multiplication by the inverse of the matrix which has been chosen by the sender.

Let the encoding matrix be

$$A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

Let the message to be sent by the sender be "WELCOME".

Since the key is taken as the operation of post-multiplication by a square matrix of order 3, the message is cut into pieces (WEL), (COM), (E), each of length 3, and converted into a sequence of row matrices of numbers:

[23 5 12],[3 15 13],[5 0 0].

Note that, we have included two zeros in the last row matrix. The reason is to get a row matrix with 5 as the first entry.

Next, we encode the message by post-multiplying each row matrix as given below:

Uncoded Encoding Coded
row matrix matrix Coded
[23 5 12]
$$\begin{bmatrix} 1 & -1 & 1 \\ 2 & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = [45 - 28 \ 23];$$

[3 15 13] $\begin{bmatrix} 1 & -1 & 1 \\ 2 & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = [46 \ -18 \ 3];$
[5 0 0] $\begin{bmatrix} 1 & -1 & 1 \\ 2 & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = [5 \ -5 \ 5].$

XII - Mathematics

A





So the encoded message is [45 - 28 23] [46 - 18 3] [5 - 5 5]

The receiver will decode the message by the reverse key, post-multiplying by the inverse of A. So the decoding matrix is

۲

$$A^{-1} = \frac{1}{|A|} \operatorname{adj} A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & -1 & 2 \\ 1 & -1 & 1 \end{bmatrix}.$$

The receiver decodes the coded message as follows:

Coded
row matrixDecoding
matrixDecoded
row matrix
$$[45 - 28 \ 23]$$
 $\begin{bmatrix} 0 & 0 & 1 \\ 0 & -1 & 2 \\ 1 & -1 & 1 \end{bmatrix}$ = $[23 \ 5 \ 12];$ $[46 - 18 \ 3]$ $\begin{bmatrix} 0 & 0 & 1 \\ 0 & -1 & 2 \\ 1 & -1 & 1 \end{bmatrix}$ = $[3 \ 15 \ 13];$ $[5 \ -5 \ 5]$ $\begin{bmatrix} 0 & 0 & 1 \\ 0 & -1 & 2 \\ 1 & -1 & 1 \end{bmatrix}$ = $[5 \ 0 \ 0].$

So, the sequence of decoded row matrices is [23 5 12],[3 15 13],[5 0 0]. Thus, the receiver reads the message as **"WELCOME"**.

EXERCISE 1.1

	□ 2	47		2	3	1	1	2	2	1
(i)	-3	4	(ii)	3	4	1	(iii) $\frac{1}{2}$	-2	1	2
	[0	2		3	7	2	3	1	-2	2

2. Find the inverse (if it exists) of the following:

(i)
$$\begin{bmatrix} -2 & 4 \\ 1 & -3 \end{bmatrix}$$
 (ii) $\begin{bmatrix} 5 & 1 & 1 \\ 1 & 5 & 1 \\ 1 & 1 & 5 \end{bmatrix}$ (iii) $\begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$

3. If $F(\alpha) = \begin{bmatrix} \cos \alpha & 0 & \sin \alpha \\ 0 & 1 & 0 \\ -\sin \alpha & 0 & \cos \alpha \end{bmatrix}$, show that $[F(\alpha)]^{-1} = F(-\alpha)$.

4. If
$$A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$$
, show that $A^2 - 3A - 7I_2 = O_2$. Hence find A^{-1}
5. If $A = \frac{1}{9} \begin{bmatrix} -8 & 1 & 4 \\ 4 & 4 & 7 \\ 1 & -8 & 4 \end{bmatrix}$, prove that $A^{-1} = A^T$.

15

Applications of Matrices and Determinants

Chapter 1 Matrices.indd 15

۲

6. If $A = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$, verify that $A(\operatorname{adj} A) = (\operatorname{adj} A)A = |A|I_2$. 7. If $A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & -3 \\ 5 & 2 \end{bmatrix}$, verify that $(AB)^{-1} = B^{-1}A^{-1}$. 8. If $adj(A) = \begin{bmatrix} 2 & -4 & 2 \\ -3 & 12 & -7 \\ -2 & 0 & 2 \end{bmatrix}$, find A. 9. If $adj(A) = \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}$, find A^{-1} . 10. Find $\operatorname{adj}(\operatorname{adj}(A))$ if $\operatorname{adj} A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}$. 11. $A = \begin{bmatrix} 1 & \tan x \\ -\tan x & 1 \end{bmatrix}$, show that $A^T A^{-1} = \begin{bmatrix} \cos 2x & -\sin 2x \\ \sin 2x & \cos 2x \end{bmatrix}$. 12. Find the matrix A for which $A\begin{bmatrix} 5 & 3\\ -1 & -2 \end{bmatrix} = \begin{bmatrix} 14 & 7\\ 7 & 7 \end{bmatrix}$. 13. Given $A = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -2 \\ 1 & 1 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$, find a matrix X such that AXB = C. 14. If $A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$, show that $A^{-1} = \frac{1}{2} (A^2 - 3I)$.

۲

15. Decrypt the received encoded message $\begin{bmatrix} 2 & -3 \end{bmatrix} \begin{bmatrix} 20 & 4 \end{bmatrix}$ with the encryption matrix $\begin{bmatrix} -1 & -1 \\ 2 & 1 \end{bmatrix}$

and the decryption matrix as its inverse, where the system of codes are described by the numbers 1-26 to the letters A-Z respectively, and the number 0 to a blank space.

1.3 Elementary Transformations of a Matrix

A matrix can be transformed to another matrix by certain operations called elementary row operations and elementary column operations.

1.3.1 Elementary row and column operations

Elementary row (column) operations on a matrix are as follows:

- (i) The interchanging of any two rows (columns) of the matrix
- (ii) Replacing a row (column) of the matrix by a non-zero scalar multiple of the row (column) by a non-zero scalar.
- (iii) Replacing a row (column) of the matrix by a sum of the row (column) with a non-zero scalar multiple of another row (column) of the matrix.

XII - Mathematics

16

۲

 \bigcirc

Elementary row operations and elementary column operations on a matrix are known as elementary transformations.

We use the following notations for elementary row transformations:

- (i) Interchanging of i^{th} and j^{th} rows is denoted by $R_i \leftrightarrow R_i$.
- (ii) The multiplication of each element of *i*th row by a non-zero constant λ is denoted by $R_i \rightarrow \lambda R_i$.
- (iii) Addition to i^{th} row, a non-zero constant λ multiple of j^{th} row is denoted by $R_i \rightarrow R_i + \lambda R_j$.

Similar notations are used for elementary column transformations.

Definition 1.4

Two matrices A and B of same order are said to be equivalent to one another if one can be obtained from the other by the applications of elementary transformations. Symbolically, we write $A \sim B$ to mean that the matrix A is equivalent to the matrix B.

For instance, let us consider a matrix
$$A = \begin{bmatrix} 1 & -2 & 2 \\ -1 & 1 & 3 \\ 1 & -1 & -4 \end{bmatrix}$$
.

After performing the elementary row operation $R_2 \rightarrow R_2 + R_1$ on A, we get a matrix B in which the second row is the sum of the second row in A and the first row in A.

Thus, we get
$$A \sim B = \begin{bmatrix} 1 & -2 & 2 \\ 0 & -1 & 5 \\ 1 & -1 & -4 \end{bmatrix}$$
.

The above elementary row transformation is also represented as follows:

[1	-2	2		1	-2	2	
-1	1	3	$\xrightarrow{R_2 \to R_2 + R_1} \to$	0	-1	5	•
1	-1	-4		1	-1	-4	

Note

An elementary transformation transforms a given matrix into another matrix which need not be equal to the given matrix.

1.3.2 Row-Echelon form

Using the row elementary operations, we can transform a given non-zero matrix to a simplified form called a **Row-echelon form**. In a row-echelon form, we may have rows all of whose entries are zero. Such rows are called **zero rows**. A non-zero row is one in which at least one of the entries is not

zero. For instance, in the matrix $\begin{bmatrix} 6 & 0 & -1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$, R_1 and R_2 are non-zero rows and R_3 is a zero row.

Definition 1.5

A non-zero matrix *E* is said to be in a **row-echelon form** if:

- (i) All zero rows of E occur below every non-zero row of E.
- (ii) The first non-zero element in any row *i* of *E* occurs in the j^{th} column of *E*, then all other entries in the j^{th} column of *E* below the first non-zero element of row *i* are zeros.
- (iii) The first non-zero entry in the i^{th} row of *E* lies to the left of the first non-zero entry in $(i+1)^{th}$ row of *E*.

Note

A non-zero matrix is in a row-echelon form if all zero rows occur as bottom rows of the matrix, and if the first non-zero element in any lower row occurs to the right of the first non-zero entry in the higher row.

	0	1	1		1	0	-1	2
The following matrices are in row-echelon form:(i)	0	0	3	,(ii)	0	0	2	8
	0	0	0_		0	0	0	6

Consider the matrix in (i). Go up row by row from the last row of the matrix. The third row is a zero row. The first non-zero entry in the second row occurs in the third column and it lies to the right of the first non-zero entry in the first row which occurs in the second column. So the matrix is in row-echelon form.

Consider the matrix in (ii). Go up row by row from the last row of the matrix. All the rows are non-zero rows. The first non-zero entry in the third row occurs in the fourth column and it occurs to the right of the first non-zero entry in the second row which occurs in the third column. The first non-zero entry in the second row occurs in the third column and it occurs to the right of the first non-zero entry in the first column and it occurs to the right of the first non-zero entry in the first non-zero entry in the first column. So the matrix is in row-echelon form.

The following matrices are not in row-echelon form:

	1	-2	0			$\begin{bmatrix} 0 \end{bmatrix}$	3	-2	
(i)	0	0	5	,	(ii)	5	0	0	
	0	1	0			3	2	0	

Consider the matrix in (i). In this matrix, the first non-zero entry in the third row occurs in the second column and it is on the left of the first non-zero entry in the second row which occurs in the third column. So the matrix is not in row-echelon form.

Consider the matrix in (ii). In this matrix, the first non-zero entry in the second row occurs in the first column and it is on the left of the first non-zero entry in the first row which occurs in the second column. So the matrix is not in row-echelon form.

Method to reduce a matrix $[a_{ij}]_{m \times n}$ to a row-echelon form.

Step 1

Inspect the first row. If the first row is a zero row, then the row is interchanged with a non-zero row below the first row. If a_{11} is not equal to 0, then go to step 2. Otherwise, interchange the first row R_1 with any other row below the first row which has a non-zero element in the first column; if no row below the first row has non-zero entry in the first column, then consider a_{12} . If a_{12} is not equal to 0, then go to step 2. Otherwise, interchange the first row which has a non-zero element in the first row which has a non-zero element in the second column; if no row below the first row below the first row which has a non-zero element in the second column; if no row below the first row has non-zero entry in the second column. The second column, then consider a_{13} . Proceed in the same way till we get a non-zero entry in the first row. This is called **pivoting** and the first non-zero element in the first row is called the **pivot** of the first row.

Step 2

Use the first row and elementary row operations to transform all elements under the pivot to become zeros.

Step 3

Consider the next row as first row and perform steps 1 and 2 with the rows below this row only. Repeat the step until all rows are exhausted.

XII - Mathematics

Example 1.13

Reduce the matrix $\begin{bmatrix} 3 & -1 & 2 \\ -6 & 2 & 4 \\ -3 & 1 & 2 \end{bmatrix}$ to a row-echelon form.

Solution

$$\begin{bmatrix} 3 & -1 & 2 \\ -6 & 2 & 4 \\ -3 & 1 & 2 \end{bmatrix} \xrightarrow{R_2 \to R_2 + 2R_1, R_2 \to R_3 + R_1} \begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 8 \\ 0 & 0 & 4 \end{bmatrix} \xrightarrow{R_3 \to R_3 - \frac{1}{2}R_2} \begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 8 \\ 0 & 0 & 0 \end{bmatrix}.$$

Note

$$\begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 8 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_2 \to R_2/8} \begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$
. This is also a row-echelon form of the given matrix.

So, a row-echelon form of a matrix is not necessarily unique.

Example 1.14

Reduce the matrix $\begin{bmatrix} 0 & 3 & 1 & 6 \\ -1 & 0 & 2 & 5 \\ 4 & 2 & 0 & 0 \end{bmatrix}$ to a row-echelon form.

Solution

 \bigcirc

$$\begin{bmatrix} 0 & 3 & 1 & 6 \\ -1 & 0 & 2 & 5 \\ 4 & 2 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} -1 & 0 & 2 & 5 \\ 0 & 3 & 1 & 6 \\ 4 & 2 & 0 & 0 \end{bmatrix} \xrightarrow{R_3 \to R_3 + 4R_1} \begin{bmatrix} -1 & 0 & 2 & 5 \\ 0 & 3 & 1 & 6 \\ 0 & 2 & 8 & 20 \end{bmatrix}$$
$$\xrightarrow{R_3 \to R_3 - \frac{2}{3}R_2} \begin{bmatrix} -1 & 0 & 2 & 5 \\ 0 & 3 & 1 & 6 \\ 0 & 0 & \frac{22}{3} & 16 \end{bmatrix} \xrightarrow{R_3 \to 3R_2} \begin{bmatrix} -1 & 0 & 2 & 5 \\ 0 & 3 & 1 & 6 \\ 0 & 0 & 22 & 48 \end{bmatrix}.$$

1.3.3 Rank of a Matrix

To define the rank of a matrix, we have to know about sub-matrices and minors of a matrix.

Let A be a given matrix. A matrix obtained by deleting some rows and some columns of A is called a **sub-matrix** of A. A matrix is a sub-matrix of itself because it is obtained by leaving zero number of rows and zero number of columns.

Recall that the determinant of a square sub-matrix of a matrix is called a minor of the matrix.

Definition 1.6

The rank of a matrix A is defined as the order of a highest order non-vanishing minor of the matrix A. It is denoted by the symbol $\rho(A)$. The rank of a zero matrix is defined to be 0.

19

()

Note

- (i) If a matrix contains at-least one non-zero element, then $\rho(A) \ge 1$.
- (ii) The rank of the identity matrix I_n is n.
- (iii) If the rank of a matrix A is r, then there exists at-least one minor of A of order r which does not vanish and every minor of A of order r+1 and higher order (if any) vanishes.
- (iv) If A is an $m \times n$ matrix, then $\rho(A) \le \min\{m, n\} = \min \{m, n\}$
- (v) A square matrix A of order n has inverse if and only if $\rho(A) = n$.

Example 1.15

Find the rank of each of the following matrices: (i) $\begin{bmatrix} 3 & 2 & 5 \\ 1 & 1 & 2 \\ 3 & 3 & 6 \end{bmatrix}$ (ii) $\begin{bmatrix} 4 & 3 & 1 & -2 \\ -3 & -1 & -2 & 4 \\ 6 & 7 & -1 & 2 \end{bmatrix}$

Solution

(i) Let $A = \begin{bmatrix} 3 & 2 & 3 \\ 1 & 1 & 2 \\ 3 & 3 & 6 \end{bmatrix}$. Then A is a matrix of order 3×3 . So $\rho(A) \le \min\{3,3\} = 3$. The highest

order of minors of A is 3. There is only one third order minor of A.

It is $\begin{vmatrix} 3 & 2 & 5 \\ 1 & 1 & 2 \\ 3 & 3 & 6 \end{vmatrix} = 3(6-6) - 2(6-6) + 5(3-3) = 0$. So, $\rho(A) < 3$.

Next consider the second-order minors of A.

We find that the second order minor
$$\begin{vmatrix} 3 & 2 \\ 1 & 1 \end{vmatrix} = 3 - 2 = 1 \neq 0$$
. So $\rho(A) = 2$.

(ii) Let $A = \begin{bmatrix} 4 & 3 & 1 & -2 \\ -3 & -1 & -2 & 4 \\ 6 & 7 & -1 & 2 \end{bmatrix}$. Then A is a matrix of order 3×4 . So $\rho(A) \le \min\{3, 4\} = 3$.

The highest order of minors of A is 3. We search for a non-zero third-order minor of A. But we find that all of them vanish. In fact, we have

4	3	1		4	3	-2	4	1	-2		3	1	-2	
-3	-1	-2	= 0;	-3	-1	4	=0; -3	-2	4	=0;	-1	-2	4	=0.
6	7	-1		6	7	2	6	-1	2		7	-1	2	

So, $\rho(A) < 3$. Next, we search for a non-zero second-order minor of A.

We find that
$$\begin{vmatrix} 4 & 3 \\ -3 & -1 \end{vmatrix} = -4 + 9 = 5 \neq 0$$
. So, $\rho(A) = 2$.

Remark

Finding the rank of a matrix by searching a highest order non-vanishing minor is quite tedious when the order of the matrix is quite large. There is another easy method for finding the rank of a matrix even if the order of the matrix is quite high. This method is by computing the rank of an equivalent row-echelon form of the matrix. If a matrix is in row-echelon form, then all entries below the **leading diagonal** (it is the line joining the positions of the diagonal elements $a_{11}, a_{22}, a_{33}, \cdots$ of the matrix) are zeros. So, checking whether a minor is zero or not, is quite simple.

XII - Mathematics

۲

Example 1.16

Find the rank of the following matrices which are in row-echelon form :

	[²	0	_7]		□_2	2	_17		6	0	-9
$\langle \cdot \rangle$		0	_/	<i>(</i>)		2	_1	()	0	2	0
(1)	0	3	1	(11)	0	5	1	(111)	0	0	0
	0	0	1		0	0	0			0	0
										U	U

Solution

(i) Let
$$A = \begin{bmatrix} 2 & 0 & -7 \\ 0 & 3 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$
. Then A is a matrix of order 3×3 and $\rho(A) \le 3$

The third order minor
$$|A| = \begin{vmatrix} 2 & 0 & -7 \\ 0 & 3 & 1 \\ 0 & 0 & 1 \end{vmatrix} = (2)(3)(1) = 6 \neq 0$$
. So, $\rho(A) = 3$.

Note that there are three non-zero rows.

(ii) Let
$$A = \begin{bmatrix} -2 & 2 & -1 \\ 0 & 5 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$
. Then A is a matrix of order 3×3 and $\rho(A) \le 3$.

The only third order minor is $|A| = \begin{vmatrix} -2 & 2 & -1 \\ 0 & 5 & 1 \\ 0 & 0 & 0 \end{vmatrix} = (-2)(5)(0) = 0$. So $\rho(A) \le 2$.

There are several second order minors. We find that there is a second order minor, for example, $\begin{vmatrix} -2 & 2 \\ 0 & 5 \end{vmatrix} = (-2)(5) = -10 \neq 0$. So, $\rho(A) = 2$.

Note that there are two non-zero rows. The third row is a zero row.

(iii) Let $A = \begin{bmatrix} 6 & 0 & -9 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$. Then A is a matrix of order 4×3 and $\rho(A) \le 3$.

The last two rows are zero rows. There are several second order minors. We find that there

is a second order minor, for example, $\begin{vmatrix} 6 & 0 \\ 0 & 2 \end{vmatrix} = (6)(2) = 12 \neq 0$. So, $\rho(A) = 2$.

Note that there are two non-zero rows. The third and fourth rows are zero rows.

We observe from the above example that the rank of a matrix in row echelon form is equal to the number of non-zero rows in it. We state this observation as a theorem without proof.

Theorem 1.11

The rank of a matrix in row echelon form is the number of non-zero rows in it.

The rank of a matrix which is not in a row-echelon form, can be found by applying the following result which is stated without proof.

Theorem 1.12

The rank of a non-zero matrix is equal to the number of non-zero rows in a row-echelon form of the matrix.

۲

Example 1.17

Find the rank of the matrix $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \\ 3 & 0 & 5 \end{bmatrix}$ by reducing it to a row-echelon form.

Solution

Let
$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \\ 3 & 0 & 5 \end{bmatrix}$$
. Applying elementary row operations, we get
$$A \xrightarrow{R_2 \to R_2 - 2R_1} A \xrightarrow{R_3 \to R_3 - 3R_1} \begin{bmatrix} 1 & 2 & 3 \\ 0 & -3 & -2 \\ 0 & -6 & -4 \end{bmatrix} \xrightarrow{R_3 \to R_3 - 2R_2} \begin{bmatrix} 1 & 2 & 3 \\ 0 & -3 & -2 \\ 0 & 0 & 0 \end{bmatrix}$$
.

The last equivalent matrix is in row-echelon form. It has two non-zero rows. So, $\rho(A) = 2$.

Example 1.18

Find the rank of the matrix $\begin{bmatrix} 2 & -2 & 4 & 3 \\ -3 & 4 & -2 & -1 \\ 6 & 2 & -1 & 7 \end{bmatrix}$ by reducing it to an echelon form.

Solution

Let A be the matrix. Performing elementary row operations, we get

$$A = \begin{bmatrix} 2 & -2 & 4 & 3 \\ -3 & 4 & -2 & -1 \\ 6 & 2 & -1 & 7 \end{bmatrix} \xrightarrow{R_2 \to 2R_2} \begin{bmatrix} 2 & -2 & 4 & 3 \\ -6 & 8 & -4 & -2 \\ 6 & 2 & -1 & 7 \end{bmatrix} \xrightarrow{R_2 \to R_2 + 3R_1} \begin{bmatrix} 2 & -2 & 4 & 3 \\ 0 & 2 & 8 & 7 \\ 0 & 8 & -13 & -2 \end{bmatrix} \cdot \frac{R_3 \to R_3 - 4R_2}{R_3 \to R_3 - 4R_2} \xrightarrow{R_2 \to R_2 + 3R_1} \begin{bmatrix} 2 & -2 & 4 & 3 \\ 0 & 2 & 8 & 7 \\ 0 & 0 & -45 & -30 \end{bmatrix} \xrightarrow{R_3 \to R_3 \div (-15)} \begin{bmatrix} 2 & -2 & 4 & 3 \\ R_3 \to R_3 - R_3 - 3R_1 \\ 0 & 2 & 8 & 7 \\ 0 & 0 & 3 & 2 \end{bmatrix} \cdot \frac{R_3 \to R_3 + (-15)}{R_3 \to R_3 \div (-15)} \xrightarrow{R_3 \to R_3 \div (-15)} \begin{bmatrix} 2 & -2 & 4 & 3 \\ 0 & 2 & 8 & 7 \\ 0 & 0 & 3 & 2 \end{bmatrix} \cdot \frac{R_3 \to R_3 + (-15)}{R_3 \to R_3 \div (-15)} \xrightarrow{R_3 \to R_3 \div (-15)} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 - 3R_1} \xrightarrow{R_3 \to R_3 - 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 - 3R_1} \xrightarrow{R_3 \to R_3 - 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 + 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 + 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 + 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 + 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 + 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 + 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 + 3R_1} \frac{R_3 \to R_3 + 3R_1}{R_3 \to R_3 + 3R_1} \xrightarrow{R_3 \to R_3 +$$

The last equivalent matrix is in row-echelon form. It has three non-zero rows. So, $\rho(A) = 3$. Elementary row operations on a matrix can be performed by pre-multiplying the given matrix by a special class of matrices called elementary matrices.

Definition 1.7

An **elementary matrix** is defined as a matrix which is obtained from an identity matrix by applying only one elementary transformation.

Remark

If we are dealing with matrices with three rows, then all elementary matrices are square matrices of order 3 which are obtained by carrying out only one elementary row operations on the unit matrix I_3 . Every elementary row operation that is carried out on a given matrix A can be obtained by pre-multiplying A with elementary matrix. Similarly, every elementary column operation that is carried out on a given matrix A can be obtained by post-multiplying A with an elementary matrix. In the present chapter, we use elementary row operations only.

XII - Mathematics

For instance, let us consider the matrix
$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$
.

Suppose that we do the transformation $R_2 \rightarrow R_2 + \lambda R_3$ on A, where $\lambda \neq 0$ is a constant. Then, we get

$$A \xrightarrow{R_{1} \to R_{2} + \lambda R_{1}} \xrightarrow{a_{11}} a_{12} a_{13} a_{22} + \lambda a_{31} a_{22} + \lambda a_{32} a_{23} + \lambda a_{33} a_{33} = a_{32} a_{33} = a_{33} = a_{32} a_{33} = a_{33} = a_{32} a_{33} = a_{33} =$$

So, the effect of applying the elementary transformation $R_2 \rightarrow R_2 + \lambda R_3$ on A is the same as that of pre-multiplying the matrix A with the elementary matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & \lambda \\ 0 & 0 & 1 \end{bmatrix}$.

Similarly, we can show that

- (i) the effect of applying the elementary transformation $R_2 \leftrightarrow R_3$ on A is the same as that of
 - pre-multiplying the matrix A with the elementary matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$.
- (ii) the effect of applying the elementary transformation $R_2 \rightarrow \lambda R_2$ on A is the same as that of $\begin{bmatrix} 1 & 0 & 0 \end{bmatrix}$

pre-multiplying the matrix A with the elementary matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & 1 \end{bmatrix}.$

We state the following result without proof.

23

Chapter 1 Matrices.indd 23

۲

Theorem 1.13

Every non-singular matrix can be transformed to an identity matrix, by a sequence of elementary row operations.

۲

As an illustration of the above theorem, let us consider the matrix $A = \begin{bmatrix} 2 & -1 \\ 3 & 4 \end{bmatrix}$.

Then, $|A| = 12 + 3 = 15 \neq 0$. So, A is non-singular. Let us transform A into I_2 by a sequence of elementary row operations. First, we search for a row operation to make a_{11} of A as 1. The elementary

row operation needed for this is $R_1 \rightarrow \left(\frac{1}{2}\right) R_1$. The corresponding elementary matrix is $E_1 = \begin{bmatrix} \frac{1}{2} & 0\\ 0 & 1 \end{bmatrix}$.

Then, we get $E_1 A = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & \frac{-1}{2} \\ 3 & 4 \end{bmatrix}.$

Next, let us make all elements below a_{11} of E_1A as 0. There is only one element a_{21} .

The elementary row operation needed for this is $R_2 \rightarrow R_2 + (-3)R_1$.

The corresponding elementary matrix is $E_2 = \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix}$. Then, we get $E_2(E_1A) = \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & -\frac{1}{2} \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & -\frac{1}{2} \\ 0 & \frac{11}{2} \end{bmatrix}$.

Next, let us make a_{22} of $E_2(E_1A)$ as 1. The elementary row operation needed for this is $R_2 \rightarrow \left(\frac{2}{11}\right)R_2.$

The corresponding elementary matrix is $E_3 = \begin{bmatrix} 1 & 0 \\ 0 & \frac{2}{11} \end{bmatrix}$. Then, we get $E_3(E_2(E_1A)) = \begin{bmatrix} 1 & 0 \\ 0 & \frac{2}{11} \end{bmatrix} \begin{bmatrix} 1 & -\frac{1}{2} \\ 0 & \frac{11}{2} \end{bmatrix} = \begin{bmatrix} 1 & -\frac{1}{2} \\ 0 & 1 \end{bmatrix}$.

Finally, let us find an elementary row operation to make a_{12} of $E_3(E_2(E_1A))$ as 0. The elementary row operation needed for this is $R_1 \rightarrow R_1 + (\frac{1}{2})R_2$. The corresponding elementary matrix is

 $E_4 = \begin{bmatrix} 1 & \frac{1}{2} \\ 0 & 1 \end{bmatrix}.$

XII - Mathematics

۲

Then, we get
$$E_4(E_3(E_1A)) = \begin{bmatrix} 1 & \frac{1}{2} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -\frac{1}{2} \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2.$$

We write the above sequence of elementary transformations in the following manner:

$$A = \begin{bmatrix} 2 & -1 \\ 3 & 4 \end{bmatrix} \xrightarrow{R_1 \to \left(\frac{1}{2}\right)R_1} \begin{bmatrix} 1 & \frac{-1}{2} \\ 2 & 3 & 4 \end{bmatrix} \xrightarrow{R_2 \to R_2 + (-3)R_1} \begin{bmatrix} 1 & -\frac{1}{2} \\ 0 & \frac{11}{2} \end{bmatrix} \xrightarrow{R_2 \to \left(\frac{2}{11}\right)R_2} \begin{bmatrix} 1 & -\frac{1}{2} \\ 0 & 1 \end{bmatrix} \xrightarrow{R_1 \to R_1 + \left(\frac{1}{2}\right)R_2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Example 1.19

Show that the matrix $\begin{bmatrix} 3 & 1 & 4 \\ 2 & 0 & -1 \\ 5 & 2 & 1 \end{bmatrix}$ is non-singular and reduce it to the identity matrix by

elementary row transformations.

Solution

Let
$$A = \begin{bmatrix} 3 & 1 & 4 \\ 2 & 0 & -1 \\ 5 & 2 & 1 \end{bmatrix}$$
. Then, $|A| = 3(0+2) - 1(2+5) + 4(4-0) = 6 - 7 + 16 = 15 \neq 0$. So, A is

non-singular. Keeping the identity matrix as our goal, we perform the row operations sequentially on *A* as follows:

$$\begin{bmatrix} 3 & 1 & 4 \\ 2 & 0 & -1 \\ 5 & 2 & 1 \end{bmatrix} \xrightarrow{R_1 \to \frac{1}{3}R_1} \begin{bmatrix} 1 & \frac{1}{3} & \frac{4}{3} \\ 2 & 0 & -1 \\ 5 & 2 & 1 \end{bmatrix} \xrightarrow{R_2 \to R_2 - 2R_1, R_2 \to R_3 - 5R_1} \xrightarrow{R_2 \to R_3 - 5R_1} \begin{bmatrix} 1 & \frac{1}{3} & \frac{4}{3} \\ 0 & -\frac{2}{3} & -\frac{11}{3} \\ 0 & \frac{1}{3} & -\frac{17}{3} \end{bmatrix} \xrightarrow{R_2 \to \left(-\frac{3}{2}\right)R_2} \begin{bmatrix} 1 & \frac{1}{3} & \frac{4}{3} \\ 0 & 1 & \frac{11}{2} \\ 0 & \frac{1}{3} & -\frac{17}{3} \end{bmatrix}$$

1.3.4 Gauss-Jordan Method

Let A be a non-singular square matrix of order n. Let B be the inverse of A. Then, we have $AB = BA = I_n$. By the property of I_n , we have $A = I_n A = AI_n$.

Consider the equation $A = I_n A$

Since A is non-singular, pre-multiplying by a sequence of elementary matrices (row operations) on both sides of (1), A on the left-hand-side of (1) is transformed to the identity matrix I_n and the same sequence of elementary matrices (row operations) transforms I_n of the right-hand-side of (1) to a matrix B. So, equation (1) transforms to $I_n = BA$. Hence, the inverse of A is B. That is, $A^{-1} = B$.

25

...(1)

Note

If E_1, E_2, \dots, E_k are elementary matrices (row operations) such that $(E_k \cdots E_2 E_1) A = I_n$, then $A^{-1} = E_k \cdots E_2 E_1.$

۲

Transforming a non-singular matrix A to the form I_n by applying elementary row operations, is called Gauss-Jordan method. The steps in finding A^{-1} by Gauss-Jordan method are given below:

Step 1

Augment the identity matrix I_n on the right-side of A to get the matrix $[A | I_n]$.

Step 2

Obtain elementary matrices (row operations) E_1, E_2, \dots, E_k such that $(E_k \dots E_2 E_1) A = I_n$.

Apply
$$E_1, E_2, \dots, E_k$$
 on $[A | I_n]$. Then $[(E_k \cdots E_2 E_1)A | (E_k \cdots E_2 E_1)I_n]$. That is, $[I_n | A^{-1}]$.

Example 1.20

Find the inverse of the non-singular matrix $A = \begin{bmatrix} 0 & 5 \\ -1 & 6 \end{bmatrix}$, by Gauss-Jordan method.

Solution

۲

Applying Gauss-Jordan method, we get

$$\begin{bmatrix} A \mid I_2 \end{bmatrix} = \begin{bmatrix} 0 & 5 & | & 1 & 0 \\ -1 & 6 & | & 0 & 1 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} -1 & 6 & | & 0 & 1 \\ 0 & 5 & | & 1 & 0 \end{bmatrix} \xrightarrow{R_1 \to (-1)R_1} \begin{bmatrix} 1 & -6 & | & 0 & -1 \\ 0 & 5 & | & 1 & 0 \end{bmatrix}$$
$$\xrightarrow{R_2 \to \frac{1}{5}R_2} \begin{bmatrix} 1 & -6 & | & 0 & -1 \\ 0 & 1 & | & (1/5) & 0 \end{bmatrix} \xrightarrow{R_1 \to R_1 + 6R_2} \begin{bmatrix} 1 & 0 & | & (6/5) & -1 \\ 0 & 1 & | & (1/5) & 0 \end{bmatrix}.$$
So, we get $A^{-1} = \begin{bmatrix} (6/5) & -1 \\ (1/5) & 0 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 6 & -5 \\ 1 & 0 \end{bmatrix}.$

Example 1.21

Find the inverse of $A = \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix}$ by Gauss-Jordan method.

Solution

Applying Gauss-Jordan method, we get

$$\begin{bmatrix} A \mid I_3 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1 & | & 1 & 0 & 0 \\ 3 & 2 & 1 & | & 0 & 1 & 0 \\ 2 & 1 & 2 & | & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_1 \to \frac{1}{2}R_1} \begin{bmatrix} 1 & (1/2) & (1/2) & | & (1/2) & 0 & 0 \\ 3 & 2 & 1 & | & 0 & 1 & 0 \\ 2 & 1 & 2 & | & 0 & 0 & 1 \end{bmatrix}$$
$$\xrightarrow{R_2 \to R_2 - 3R_1} \begin{bmatrix} 1 & (1/2) & (1/2) & | & (1/2) & 0 & 0 \\ 0 & (1/2) & -(1/2) & | & -(3/2) & 1 & 0 \\ 0 & 0 & 1 & | & -1 & 0 & 1 \end{bmatrix} \xrightarrow{R_2 \to 2R_2} \begin{bmatrix} 1 & (1/2) & (1/2) & | & (1/2) & 0 & 0 \\ 0 & 1 & -1 & | & -3 & 2 & 0 \\ 0 & 0 & 1 & | & -1 & 0 & 1 \end{bmatrix}$$

XII - Mathematics

۲

۲

EXERCISE 1.2

1. Find the rank of the following matrices by minor method:

Га	47		-1	3]		[1	-2	3		$\begin{bmatrix} 0 \end{bmatrix}$	1	2	1
(i) $\begin{bmatrix} 2\\ 1 \end{bmatrix}$	-4	(ii)	4	-7	(iii) $\begin{vmatrix} 1 & -2 & -1 & 0 \\ 2 & (-2 & 1) \end{vmatrix}$ (iv)	2	4	-6	(v)	0	2	4	3
[-1	2]		3	_4		5	1	-1		8	1	0	2_

2. Find the rank of the following matrices by row reduction method:

(i)	$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 2 & -1 & 3 & 4 \\ 5 & -1 & 7 & 11 \end{bmatrix}$	(ii)	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	(ii	i)	3 2 -1	-8 -5 2	5 1 3	2 4 -2
			[1 -1 1]		L		_	-	-1

3. Find the inverse of each of the following by Gauss-Jordan method:

Гр	1]		[1	-1	0		[1	2	3	
(i) $\begin{vmatrix} 2 & -1 \\ 5 & 2 \end{vmatrix}$		(ii)	1	0	-1	(iii)	2	5	3	
[3	-2		6	-2	-3_		1	0	8_	

1.4 Applications of Matrices: Solving System of Linear Equations

One of the important applications of matrices and determinants is solving of system of linear equations. Systems of linear equations arise as mathematical models of several phenomena occurring in biology, chemistry, commerce, economics, physics and engineering. For instance, analysis of circuit theory, analysis of input-output models, and analysis of chemical reactions require solutions of systems of linear equations.

1.4.1 Formation of a System of Linear Equations

The meaning of a system of linear equations can be understood by formulating a mathematical model of a simple practical problem.

Three persons A, B and C go to a supermarket to purchase same brands of rice and sugar. Person A buys 5 Kilograms of rice and 3 Kilograms of sugar and pays ₹ 440. Person B purchases 6 Kilograms of rice and 2 Kilograms of sugar and pays ₹ 400. Person C purchases 8 Kilograms of rice and 5 Kilograms of sugar and pays ₹ 720. Let us formulate a mathematical model to compute the price per Kilogram of rice and the price per Kilogram of sugar. Let *x* be the price in rupees per Kilogram of rice and 9 we the price in rupees per Kilogram of sugar. Person A buys 5 Kilograms of rice and 3 Kilograms sugar and pays ₹ 440. So, 5x + 3y = 440. Similarly, by considering Person B and Person C, we get 6x + 2y = 400 and 8x + 5y = 720. Hence the mathematical model is to obtain *x* and *y* such that

$$5x + 3y = 440$$
, $6x + 2y = 400$, $8x + 5y = 720$.

27

Applications of Matrices and Determinants

۲

()

Note

In the above example, the values of x and y which satisfy one equation should also satisfy all the other equations. In other words, the equations are to be satisfied by the same values of x and y simultaneously. If such values of x and y exist, then they are said to form a **solution** for the system of linear equations. In the three equations, x and y appear in first degree only. Hence they are said to form a system of linear equations in two unknowns x and y. They are also called **simultaneous** linear equations in two unknowns x and y. The system has three linear equations in two unknowns x and y.

The equations represent three straight lines in two-dimensional analytical geometry. In this section, we develop methods using matrices to find solutions of systems of linear equations.

1.4.2 System of Linear Equations in Matrix Form

A system of m linear equations in n unknowns is of the following form:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1,$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2,$$

$$\dots \qquad \dots \qquad \dots \qquad \dots$$

$$a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \dots + a_{mn}x_n = b_m,$$

(1)

where the coefficients a_{ij} , $i = 1, 2, \dots, m$; $j = 1, 2, \dots, n$ and b_k , $k = 1, 2, \dots, m$ are constants. If all the b_k 's are zeros, then the above system is called a **homogeneous system** of linear equations. On the other hand, if at least one of the b_k 's is non-zero, then the above system is called a **non-homogeneous system** of linear equations. If there exist values $\alpha_1, \alpha_2, \dots, \alpha_n$ for x_1, x_2, \dots, x_n respectively which satisfy every equation of (1), then the ordered n – tuple $(\alpha_1, \alpha_2, \dots, \alpha_n)$ is called a **solution** of (1). The above system (1) can be put in a matrix form as follows:

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}$ be the $m \times n$ matrix formed by the coefficients of

 $x_1, x_2, x_3, \dots, x_n$. The first row of *A* is formed by the coefficients of $x_1, x_2, x_3, \dots, x_n$ in the same order in which they occur in the first equation. Likewise, the other rows of *A* are formed. The first column is formed by the coefficients of x_1 in the *m* equations in the same order. The other columns are formed in a similar way.

Let $X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$ be the $n \times 1$ order column matrix formed by the unknowns $x_1, x_2, x_3, \dots, x_n$. Let $B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$ be the $m \times 1$ order column matrix formed by the right-hand side constants $b_1, b_2, b_3, \dots, b_m$.

XII - Mathematics

Then we get

$$AX = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n \\ \cdots & \cdots & \cdots \\ a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \cdots + a_{mn}x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} = B.$$

۲

Then AX = B is a matrix equation involving matrices and it is called the **matrix form** of the system of linear equations (1). The matrix A is called the **coefficient matrix** of the system and the

matrix $\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} & b_2 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} & b_m \end{bmatrix}$ is called the **augmented matrix** of the system. We denote the

augmented matrix by [A | B].

As an example, the matrix form of the system of linear equations

$$2x + 3y - 5z + 7 = 0, 7y + 2z - 3x = 17, 6x - 3y - 8z + 24 = 0 \text{ is} \begin{bmatrix} 2 & 3 & -5 \\ -3 & 7 & 2 \\ 6 & -3 & -8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -7 \\ 17 \\ -24 \end{bmatrix}$$

1.4.3 Solution to a System of Linear equations

The meaning of solution to a system of linear equations can be understood by considering the following cases :

Case (i)

۲

Consider the system of linear equations

$$2x - y = 5,$$
 ... (1)
 $x + 3y = 6.$... (2)

These two equations represent a pair of straight lines in two dimensional analytical geometry (see the Fig. 1.2). Using (1), we get

$$x = \frac{5+y}{2} \dots (3)$$

Substituting (3) in (2) and simplifying, we get y = 1.

Substituting y=1 in (1) and simplifying, we get x=3.

Both equations (1) and (2) are satisfied by x = 3 and y = 1.

That is, a solution of (1) is also a solution of (2).

So, we say that the system is consistent and has unique solution (3,1).

The point (3,1) is the point of intersection of the two lines 2x - y = 5 and x + 3y = 6.

29

Х



$$Fig.1.2$$



Substituting (3) in (2) and simplifying, we get 0 = 0.

3x + 2y = 5,

6x + 4y = 10

This informs us that equation (2) is an elementary transformation of equation (1). In fact, by dividing equation (2) by 2, we get equation (1). It is not possible to find uniquely x and y with just a single equation (1).



So we are forced to assume the value of one unknown, say y = t, where t is any real number. Then, $x = \frac{5-2t}{3}$. The two equations (1) and (2) represent one and only one straight line (coincident lines) in two dimensional analytical geometry (see Fig. 1.3). In other words, the system is consistent (a solution of (1) is also a solution of (2)) and has infinitely many solutions, as t can assume any real value.

(1)

... (2)

... (3)

Case (iii)

 \bigcirc

Case (ii)

Consider the system of linear equations

$$4x + y = 6, \qquad \dots$$

Using equation (1), we get

$$x = \frac{6-y}{4}$$

Substituting (3) in (2) and simplifying, we get 12 = 18.

8x + 2y = 18.



This is a contradicting result, which informs us that equation (2) is inconsistent with equation (1). So, a solution of (1) is not a solution of (2).

In other words, the system is inconsistent and has no solution. We note that the two equations represent two parallel straight lines (non-coincident) in two dimensional analytical geometry (see Fig. 1.4). We know that two non-coincident parallel lines never meet in real points.

Note

- (1) Interchanging any two equations of a system of linear equations does not alter the solution of the system.
- (2) Replacing an equation of a system of linear equations by a non-zero constant multiple of itself does not alter the solution of the system.
- (3) Replacing an equation of a system of linear equations by addition of itself with a non-zero multiple of any other equation of the system does not alter the solution of the system.

XII - Mathematics

30

Definition 1.8

A system of linear equations having at least one solution is said to be **consistent**. A system of linear equations having no solution is said to be **inconsistent**.

Remark

If the number of the equations of a system of linear equations is equal to the number of unknowns of the system, then the coefficient matrix A of the system is a square matrix. Further, if A is a non-singular matrix, then the solution of system of equations can be found by any one of the following methods : (i) matrix inversion method, (ii) Cramer's rule, (iii) Gaussian elimination method.

1.4.3 (i) Matrix Inversion Method

This method can be applied only when the coefficient matrix is a square matrix and non-singular. Consider the matrix equation

AX = B, ... (1) where A is a square matrix and non-singular. Since A is non-singular, A^{-1} exists and $A^{-1}A = AA^{-1} = I$.

Pre-multiplying both sides of (1) by A^{-1} , we get $A^{-1}(AX) = A^{-1}B$. That is, $(A^{-1}A)X = A^{-1}B$.

Hence, we get $X = A^{-1}B$.

Example 1.22

Solve the following system of linear equations, using matrix inversion method: 5x + 2y = 3, 3x + 2y = 5.

Solution

The matrix form of the system is AX = B, where $A = \begin{bmatrix} 5 & 2 \\ 3 & 2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 3 \\ 5 \end{bmatrix}$.

47

We find
$$|A| = \begin{vmatrix} 5 & 2 \\ 3 & 2 \end{vmatrix} = 10 - 6 = 4 \neq 0$$
. So, A^{-1} exists and $A^{-1} = \frac{1}{4} \begin{bmatrix} 2 & -2 \\ -3 & 5 \end{bmatrix}$.

Then, applying the formula $X = A^{-1}B$, we get

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 2 & -2 \\ -3 & 5 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 6-10 \\ -9+25 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} -4 \\ 16 \end{bmatrix} = \begin{bmatrix} \frac{-4}{4} \\ \frac{16}{4} \end{bmatrix} = \begin{bmatrix} -1 \\ 4 \end{bmatrix}.$$

So the solution is (x = -1, y = 4).

Example 1.23

Solve the following system of equations, using matrix inversion method:

 $2x_1 + 3x_2 + 3x_3 = 5$, $x_1 - 2x_2 + x_3 = -4$, $3x_1 - x_2 - 2x_3 = 3$.

Solution

The matrix form of the system is AX = B, where

$$A = \begin{bmatrix} 2 & 3 & 3 \\ 1 & -2 & 1 \\ 3 & -1 & -2 \end{bmatrix}, X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}, B = \begin{bmatrix} 5 \\ -4 \\ 3 \end{bmatrix}.$$

31

()

We find
$$|A| = \begin{vmatrix} 2 & 3 & 3 \\ 1 & -2 & 1 \\ 3 & -1 & -2 \end{vmatrix} = 2(4+1) - 3(-2-3) + 3(-1+6) = 10 + 15 + 15 = 40 \neq 0$$
.

So, A^{-1} exists and

$$A^{-1} = \frac{1}{|A|} (\operatorname{adj} A) = \frac{1}{40} \begin{bmatrix} +(4+1) & -(-2-3) & +(-1+6) \\ -(-6+3) & +(-4-9) & -(-2-9) \\ +(3+6) & -(2-3) & +(-4-3) \end{bmatrix}^{T} = \frac{1}{40} \begin{bmatrix} 5 & 3 & 9 \\ 5 & -13 & 1 \\ 5 & 11 & -7 \end{bmatrix}$$

Then, applying $X = A^{-1}B$, we get

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \frac{1}{40} \begin{bmatrix} 5 & 3 & 9 \\ 5 & -13 & 1 \\ 5 & 11 & -7 \end{bmatrix} \begin{bmatrix} 5 \\ -4 \\ 3 \end{bmatrix} = \frac{1}{40} \begin{bmatrix} 25 - 12 + 27 \\ 25 + 52 + 3 \\ 25 - 44 - 21 \end{bmatrix} = \frac{1}{40} \begin{bmatrix} 40 \\ 80 \\ -40 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}.$$

So, the solution is $(x_1 = 1, x_2 = 2, x_3 = -1)$.

Example 1.24

If
$$A = \begin{bmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{bmatrix}$$
 and $B = \begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{bmatrix}$, find the products *AB* and *BA* and hence solve the

system of equations x - y + z = 4, x - 2y - 2z = 9, 2x + y + 3z = 1.

Solution

We find
$$AB = \begin{bmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{bmatrix} \begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{bmatrix} = \begin{bmatrix} -4+4+8 & 4-8+4 & -4-8+12 \\ -7+1+6 & 7-2+3 & -7-2+9 \\ 5-3-2 & -5+6-1 & 5+6-3 \end{bmatrix} = \begin{bmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 8 \end{bmatrix} = 8I_3$$

and $BA = \begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{bmatrix} \begin{bmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{bmatrix} = \begin{bmatrix} -4+7+5 & 4-1-3 & 4-3-1 \\ -4+14-10 & 4-2+6 & 4-6+2 \\ -8-7+15 & 8+1-9 & 8+3-3 \end{bmatrix} = \begin{bmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 8 \end{bmatrix} = 8I_3.$

So, we get
$$AB = BA = 8I_3$$
. That is, $\left(\frac{1}{8}A\right)B = B\left(\frac{1}{8}A\right) = I_3$. Hence, $B^{-1} = \frac{1}{8}A_3$.
Writing the given system of equations in matrix form, we get

Writing the given system of equations in matrix form, we get

$$\begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 9 \\ 1 \end{bmatrix}.$$
 That is, $B\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 9 \\ 1 \end{bmatrix}.$
So, $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = B^{-1} \begin{bmatrix} 4 \\ 9 \\ 1 \end{bmatrix} = \left(\frac{1}{8}A\right) \begin{bmatrix} 4 \\ 9 \\ 1 \end{bmatrix} = \frac{1}{8} \begin{bmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{bmatrix} \begin{bmatrix} 4 \\ 9 \\ 1 \end{bmatrix} = \frac{1}{8} \begin{bmatrix} -16+36+4 \\ -28+9+3 \\ 20-27-1 \end{bmatrix} = \frac{1}{8} \begin{bmatrix} 24 \\ -16 \\ -8 \end{bmatrix} = \begin{bmatrix} 3 \\ -2 \\ -1 \end{bmatrix}$

Hence, the solution is (x = 3, y = -2, z = -1).

XII - Mathematics

Chapter 1 Matrices.indd 32

EXERCISE 1.3

۲

1. Solve the following system of linear equations by matrix inversion method:

(i) 2x + 5y = -2, x + 2y = -3(ii) 2x - y = 8, 3x + 2y = -2(iii) 2x + 3y - z = 9, x + y + z = 9, 3x - y - z = -1(iv) x + y + z - 2 = 0, 6x - 4y + 5z - 31 = 0, 5x + 2y + 2z = 132. If $A = \begin{bmatrix} -5 & 1 & 3 \\ 7 & 1 & -5 \\ 1 & -1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$, find the products *AB* and *BA* and hence solve the

system of equations x + y + 2z = 1, 3x + 2y + z = 7, 2x + y + 3z = 2.

- 3. A man is appointed in a job with a monthly salary of certain amount and a fixed amount of annual increment. If his salary was ₹ 19,800 per month at the end of the first month after 3 years of service and ₹ 23,400 per month at the end of the first month after 9 years of service, find his starting salary and his annual increment. (Use matrix inversion method to solve the problem.)
- 4. Four men and 4 women can finish a piece of work jointly in 3 days while 2 men and 5 women can finish the same work jointly in 4 days. Find the time taken by one man alone and that of one woman alone to finish the same work by using matrix inversion method.
- 5. The prices of three commodities A, B and C are $\gtrless x, y$ and z per units respectively. A person P purchases 4 units of B and sells two units of A and 5 units of C. Person Q purchases 2 units of C and sells 3 units of A and one unit of B. Person R purchases one unit of A and sells 3 unit of B and one unit of C. In the process, P,Q and R earn $\gtrless 15,000, \gtrless 1,000$ and $\end{Bmatrix} 4,000$ respectively. Find the prices per unit of A, B and C. (Use matrix inversion method to solve the problem.)

1.4.3 (ii) Cramer's Rule

This rule can be applied only when the coefficient matrix is a square matrix and non-singular. It is explained by considering the following system of equations:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1,$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2,$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3,$$

where the coefficient matrix $\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ is non-singular. Then $\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \neq 0.$

Let us put $\Delta = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$. Then, we have

$$x_{1}\Delta = x_{1} \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11}x_{1} & a_{12} & a_{13} \\ a_{21}x_{1} & a_{22} & a_{23} \\ a_{31}x_{1} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11}x_{1} & a_{12} & a_{13} \\ a_{21}x_{1} + a_{22}x_{2} + a_{23}x_{3} & a_{22} & a_{23} \\ a_{31}x_{1} + a_{32}x_{2} + a_{33}x_{3} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} b_{1} & a_{12} & a_{13} \\ b_{2} & a_{22} & a_{23} \\ b_{3} & a_{32} & a_{33} \end{vmatrix} = \Delta_{1}$$

XII - Mathematics

۲

Since
$$\Delta \neq 0$$
, we get $x_1 = \frac{\Delta_1}{\Delta}$.

Similarly, we get
$$x_2 = \frac{\Delta_2}{\Delta}$$
, $x_3 = \frac{\Delta_3}{\Delta}$, where $\Delta_2 = \begin{vmatrix} a_{11} & b_1 & a_{13} \\ a_{21} & b_2 & a_{23} \\ a_{31} & b_3 & a_{33} \end{vmatrix}$, $\Delta_3 = \begin{vmatrix} a_{11} & a_{12} & b_1 \\ a_{21} & a_{22} & b_2 \\ a_{31} & a_{32} & b_3 \end{vmatrix}$.

Thus, we have the Cramer's rule $x_1 = \frac{\Delta_1}{\Lambda}, x_2 = \frac{\Delta_2}{\Lambda}, x_3 = \frac{\Delta_3}{\Lambda},$

where
$$\Delta = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$
, $\Delta_1 = \begin{vmatrix} b_1 & a_{12} & a_{13} \\ b_2 & a_{22} & a_{23} \\ b_3 & a_{32} & a_{33} \end{vmatrix}$, $\Delta_2 = \begin{vmatrix} a_{11} & b_1 & a_{13} \\ a_{21} & b_2 & a_{23} \\ a_{31} & b_3 & a_{33} \end{vmatrix}$, $\Delta_3 = \begin{vmatrix} a_{11} & a_{12} & b_1 \\ a_{21} & a_{22} & b_2 \\ a_{31} & a_{32} & b_3 \end{vmatrix}$

Note

Replacing the first column elements a_{11}, a_{21}, a_{31} of Δ with b_1, b_2, b_3 respectively, we get Δ_1 . Replacing the second column elements a_{12}, a_{22}, a_{32} of Δ with b_1, b_2, b_3 respectively, we get Δ_2 . Replacing the third column elements a_{13}, a_{23}, a_{33} of Δ with b_1, b_2, b_3 respectively, we get Δ_3 . If $\Delta = 0$, Cramer's rule cannot be applied.

Example 1.25

Solve, by Cramer's rule, the system of equations

$$x_1 - x_2 = 3, 2x_1 + 3x_2 + 4x_3 = 17, x_2 + 2x_3 = 7.$$

Solution

First we evaluate the determinants

$$\Delta = \begin{vmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{vmatrix} = 6 \neq 0, \ \Delta_1 = \begin{vmatrix} 3 & -1 & 0 \\ 17 & 3 & 4 \\ 7 & 1 & 2 \end{vmatrix} = 12, \ \Delta_2 = \begin{vmatrix} 1 & 3 & 0 \\ 2 & 17 & 4 \\ 0 & 7 & 2 \end{vmatrix} = -6, \ \Delta_3 = \begin{vmatrix} 1 & -1 & 3 \\ 2 & 3 & 17 \\ 0 & 1 & 7 \end{vmatrix} = 24.$$

By Cramer's rule, we get $x_1 = \frac{\Delta_1}{\Delta} = \frac{12}{6} = 2$, $x_2 = \frac{\Delta_2}{\Delta} = \frac{-6}{6} = -1$, $x_3 = \frac{-24}{6} = 4$. So, the solution is $(x_1 = 2, x_2 = -1, x_3 = 4)$.

Example 1.26

In a T20 match, Chennai Super Kings needed just 6 runs to win with 1 ball left to go in the last over. The last ball was bowled and the batsman at the crease hit it high up. The ball traversed along a path in a vertical plane and the equation of the path is $y = ax^2 + bx + c$ with respect to a xy-coordinate system in the vertical plane and the ball traversed through the points (10,8),(20,16),(30,18), can you conclude that Chennai Super Kings won the match?

Justify your answer. (All distances are measured in metres and the meeting point of the plane of the path with the farthest boundary line is (70,0).)

Solution

The path $y = ax^2 + bx + c$ passes through the points (10,8), (20,16), (40,22). So, we get the system of equations 100a + 10b + c = 8,400a + 20b + c = 16,1600a + 40b + c = 22. To apply Cramer's ind

Chapter 1 Matrices.indd 34



$$\Delta = \begin{vmatrix} 100 & 10 & 1 \\ 400 & 20 & 1 \\ 1600 & 40 & 1 \end{vmatrix} = 1000 \begin{vmatrix} 1 & 1 & 1 \\ 4 & 2 & 1 \\ 16 & 4 & 1 \end{vmatrix} = 1000[-2+12-16] = -6000,$$

$$\Delta_{1} = \begin{vmatrix} 8 & 10 & 1 \\ 16 & 20 & 1 \\ 22 & 40 & 1 \end{vmatrix} = 20 \begin{vmatrix} 4 & 1 & 1 \\ 8 & 2 & 1 \\ 11 & 4 & 1 \end{vmatrix} = 20[-8+3+10] = 100,$$

$$\Delta_{2} = \begin{vmatrix} 100 & 8 & 1 \\ 400 & 16 & 1 \\ 1600 & 22 & 1 \end{vmatrix} = 200 \begin{vmatrix} 1 & 4 & 1 \\ 4 & 8 & 1 \\ 16 & 11 & 1 \end{vmatrix} = 200[-3+48-84] = -7800,$$

$$\Delta_{3} = \begin{vmatrix} 100 & 10 & 8 \\ 400 & 20 & 16 \\ 1600 & 40 & 22 \end{vmatrix} = 2000 \begin{vmatrix} 1 & 1 & 4 \\ 4 & 2 & 8 \\ 16 & 4 & 11 \end{vmatrix} = 2000[-10+84-64] = 20000.$$
By Cramer's rule, we get $a = \frac{\Delta_{1}}{\Delta} = -\frac{1}{60}, b = \frac{\Delta_{2}}{\Delta} = \frac{7800}{6000} = \frac{78}{60} = \frac{13}{10}, c = \frac{\Delta_{3}}{\Delta} = -\frac{20000}{6000} = -\frac{20}{6} = -\frac{10}{3}.$

So, the equation of the path is $y = -\frac{1}{60}x^2 + \frac{13}{10}x - \frac{10}{3}$.

When x = 70, we get y = 6. So, the ball went by 6 metres high over the boundary line and it is impossible for a fielder standing even just before the boundary line to jump and catch the ball. Hence the ball went for a super six and the Chennai Super Kings won the match.

EXERCISE 1.4

- 1. Solve the following systems of linear equations by Cramer's rule:
 - (i) 5x 2y + 16 = 0, x + 3y 7 = 0
 - (ii) $\frac{3}{x} + 2y = 12, \frac{2}{x} + 3y = 13$
 - (iii) 3x + 3y z = 11, 2x y + 2z = 9, 4x + 3y + 2z = 25
 - (iv) $\frac{3}{x} \frac{4}{y} \frac{2}{z} 1 = 0, \frac{1}{x} + \frac{2}{y} + \frac{1}{z} 2 = 0, \frac{2}{x} \frac{5}{y} \frac{4}{z} + 1 = 0$
- 2. In a competitive examination, one mark is awarded for every correct answer while $\frac{1}{4}$ mark is

deducted for every wrong answer. A student answered 100 questions and got 80 marks. How many questions did he answer correctly ? (Use Cramer's rule to solve the problem).

- 3. A chemist has one solution which is 50% acid and another solution which is 25% acid. How much each should be mixed to make 10 litres of a 40% acid solution ? (Use Cramer's rule to solve the problem).
- 4. A fish tank can be filled in 10 minutes using both pumps A and B simultaneously. However, pump B can pump water in or out at the same rate. If pump B is inadvertently run in reverse, then the tank will be filled in 30 minutes. How long would it take each pump to fill the tank by itself? (Use Cramer's rule to solve the problem).

35

Applications of Matrices and Determinants

Chapter 1 Matrices.indd 35

5. A family of 3 people went out for dinner in a restaurant. The cost of two dosai, three idlies and two vadais is ₹ 150. The cost of the two dosai, two idlies and four vadais is ₹ 200. The cost of five dosai, four idlies and two vadais is ₹ 250. The family has ₹ 350 in hand and they ate 3 dosai and six idlies and six vadais. Will they be able to manage to pay the bill within the amount they had ?

۲

1.4.3 (iii) Gaussian Elimination Method

This method can be applied even if the coefficient matrix is singular matrix and rectangular matrix. It is essentially the method of substitution which we have already seen. In this method, we transform the augmented matrix of the system of linear equations into row-echelon form and then by back-substitution, we get the solution.

Example 1.27

Solve the following system of linear equations, by Gaussian elimination method :

$$4x + 3y + 6z = 25$$
, $x + 5y + 7z = 13$, $2x + 9y + z = 1$.

Solution

Transforming the augmented matrix to echelon form, we get

4 3 6	25			1	5	7	13		$a \rightarrow b$	$R_{a} - 4R_{a}$	[1	5	7	13]
1 5 7	13	R_1	$\leftrightarrow R_2 \rightarrow$	4	3	6	25		$R_3 \rightarrow K$	$R_3^2 - 2R_1$	$\rightarrow 0$	-17	-22	-27
2 9 1	1			2	9	1	1				0	-1	-13	-25
$R_{-} \rightarrow R_{-} \div (-1)$	[1	5	7	1	3				1	5	7	13]	
$R_3 \rightarrow R_3 \div (-1)$	$\rightarrow 0$	17	22	2	27	R	$_3 \rightarrow 17R_3 - K_3$	\rightarrow	0	17	22	27		
	0	1	13	2	25				0	0	199	398		

The equivalent system is written by using the echelon form:

3

$$x + 5y + 7z = 13, \qquad \dots (1)$$

$$17y + 22z = 27$$
, ... (2)

$$199z = 398.$$
 ... (3)

From (3), we get $z = \frac{398}{199} = 2$.

Substituting z = 2 in (2), we get $y = \frac{27 - 22 \times 2}{17} = \frac{-17}{17} = -1$. Substituting z = 2, y = -1 in (1), we get $x = 13 - 5 \times (-1) - 7 \times 2 = 4$. So, the solution is (x = 4, y = -1, z = 2).

Note. The above method of going from the last equation to the first equation is called the **method** of back substitution.

Example 1.28

The upward speed v(t) of a rocket at time *t* is approximated by $v(t) = at^2 + bt + c$, $0 \le t \le 100$ where *a*,*b*, and *c* are constants. It has been found that the speed at times t = 3, t = 6, and t = 9 seconds are respectively, 64, 133, and 208 miles per second respectively. Find the speed at time t = 15 seconds. (Use Gaussian elimination method.)



XII - Mathematics

36

۲

Solution

Since v(3) = 64, v(6) = 133, and v(9) = 208, we get the following system of linear equations

$$9a+3b+c = 64$$
,
 $36a+6b+c = 133$,
 $81a+9b+c = 208$.

We solve the above system of linear equations by Gaussian elimination method.

Reducing the augmented matrix to an equivalent row-echelon form by using elementary row operations, we get

$$\begin{bmatrix} A \mid B \end{bmatrix} = \begin{bmatrix} 9 & 3 & 1 & | & 64 \\ 36 & 6 & 1 & | & 133 \\ 81 & 9 & 1 & | & 208 \end{bmatrix} \xrightarrow{R_2 \to R_2 - 4R_1, R_3 \to R_3 - 9R_1} \begin{bmatrix} 9 & 3 & 1 & | & 64 \\ 0 & -6 & -3 & | & -123 \\ 0 & -18 & -8 & | & -368 \end{bmatrix} \xrightarrow{R_2 \to R_2 \div (-3), R_3 \to R_3 \div (-2)} \begin{bmatrix} 9 & 3 & 1 & | & 64 \\ 0 & 2 & 1 & | & 41 \\ 0 & 9 & 4 & | & 184 \end{bmatrix}$$
$$\xrightarrow{R_1 \to 2R_3} \begin{bmatrix} 9 & 3 & 1 & | & 64 \\ 0 & 2 & 1 & | & 41 \\ 0 & 18 & 8 & | & 368 \end{bmatrix} \xrightarrow{R_3 \to R_1 - 9R_2} \begin{bmatrix} 9 & 3 & 1 & | & 64 \\ 0 & 2 & 1 & | & 41 \\ 0 & 0 & -1 & | & -1 \end{bmatrix} \xrightarrow{R_3 \to (-1)R_3} \begin{bmatrix} 9 & 3 & 1 & | & 64 \\ 0 & 2 & 1 & | & 41 \\ 0 & 0 & 1 & | & 1 \end{bmatrix}.$$

Writing the equivalent equations from the row-echelon matrix, we get 9a+3b+c=64, 2b+c=41, c=1.

By back substitution, we get c = 1, $b = \frac{(41-c)}{2} = \frac{(41-1)}{2} = 20$, $a = \frac{64-3b-c}{9} = \frac{64-60-1}{9} = \frac{1}{3}$. So, we get $v(t) = \frac{1}{3}t^2 + 20t + 1$. Hence, $v(15) = \frac{1}{3}(225) + 20(15) + 1 = 75 + 300 + 1 = 376$.

EXERCISE 1.5

- 1. Solve the following systems of linear equations by Gaussian elimination method:
 - (i) 2x 2y + 3z = 2, x + 2y z = 3, 3x y + 2z = 1
 - (ii) 2x + 4y + 6z = 22, 3x + 8y + 5z = 27, -x + y + 2z = 2
- 2. If $ax^2 + bx + c$ is divided by x+3, x-5, and x-1, the remainders are 21,61 and 9 respectively. Find *a*, *b* and *c*. (Use Gaussian elimination method.)
- 3. An amount of ₹ 65,000 is invested in three bonds at the rates of 6%, 8% and 10% per annum respectively. The total annual income is ₹ 4,800. The income from the third bond is ₹ 600 more than that from the second bond. Determine the price of each bond. (Use Gaussian elimination method.)
- 4. A boy is walking along the path $y = ax^2 + bx + c$ through the points (-6,8),(-2,-12), and (3,8). He wants to meet his friend at P(7,60). Will he meet his friend? (Use Gaussian elimination method.)

37



Chapter 1 Matrices.indd 37

1.5 Applications of Matrices: Consistency of System of Linear Equations by Rank Method

In section 1.3.3, we have already defined consistency of a system of linear equation. In this section, we investigate it by using rank method. We state the following theorem without proof:

۲

Theorem 1.14 (Rouché-Capelli Theorem)

A system of linear equations, written in the matrix form as AX = B, is consistent if and only if the rank of the coefficient matrix is equal to the rank of the augmented matrix; that is, $\rho(A) = \rho([A | B])$.

We apply the theorem in the following examples.

1.5.1 Non-homogeneous Linear Equations

Example 1.29

Test for consistency of the following system of linear equations and if possible solve: x+2y-z=3, 3x-y+2z=1, x-2y+3z=3, x-y+z+1=0.

Solution

۲

Here the number of unknowns is 3.

The matrix form of the system is AX = B, where

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 1 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 3 \\ 1 \\ 3 \\ -1 \end{bmatrix}$$
$$\begin{bmatrix} A \mid B \end{bmatrix} = \begin{bmatrix} 1 & 2 & -1 & 3 \\ 3 & -1 & 2 & 1 \\ 1 & -2 & 3 & 3 \\ 1 & -1 & 1 & -1 \end{bmatrix}.$$

The augmented matrix is

Applying Gaussian elimination method on [A | B], we get

$$\begin{bmatrix} A \mid B \end{bmatrix} \xrightarrow{R_{2} \to R_{2} - 3R_{1}, \\ R_{3} \to R_{3} - R_{1}, \\ R_{4} \to R_{4} - R_{1}, \\ R_{4} \to R_{4} - R_{1}, \\ \hline \\ 0 \quad -4 \quad 4 \quad 0 \\ 0 \quad -3 \quad 2 \quad | -4 \end{bmatrix} \xrightarrow{R_{2} \to (-1)R_{3}, \\ R_{4} \to (-1)R_{4}} \begin{bmatrix} 1 & 2 & -1 & | & 3 \\ 0 & 7 & -5 & | & 8 \\ 0 & 4 & -4 & | & 0 \\ 0 & 3 & -2 \quad | & 4 \end{bmatrix} \xrightarrow{R_{3} \to 7R_{3} - 4R_{2}, \\ R_{4} \to 7R_{4} - 3R_{2}} \xrightarrow{R_{4} \to 7R_{4} - 3R_{2}} \xrightarrow{R_{3} \to (-3)} \begin{bmatrix} 1 & 2 & -1 & | & 3 \\ 0 & -4 & 4 & | & 0 \\ 0 & -3 & 2 & | & -4 \end{bmatrix} \xrightarrow{R_{3} \to R_{3} \div (-8)} \xrightarrow{R_{3} \to R_{3} \div (-8)} \xrightarrow{R_{4} \to R_{4} - R_{3}} \xrightarrow{R_{4} \to R_{4} - R_{4}$$

There are three non-zero rows in the row-echelon form of [A | B]. So, $\rho([A | B]) = 3$.

So, the row-echelon form of A is $\begin{bmatrix} 1 & 2 & -1 \\ 0 & 7 & -5 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$. There are three non-zero rows in it. So $\rho(A) = 3$.

XII - Mathematics

۲

Hence, $\rho(A) = \rho([A | B]) = 3$.

From the echelon form, we write the equivalent system of equations x + 2y - z = 3, 7y - 5z = 8, z = 4, 0 = 0.

The last equation 0 = 0 is meaningful. By the method of back substitution, we get

$$z = 4$$

$$7y-20 = 8 \implies y=4,$$

$$x = 3-8+4 \implies x=-1.$$

So, the solution is (x = -1, y = 4, z = 4). (Note that A is not a square matrix.)

Here the given system is consistent and the solution is unique.

Example 1.30

Test for consistency of the following system of linear equations and if possible solve: 4x-2y+6z=8, x+y-3z=-1, 15x-3y+9z=21.

Solution

Here the number of unknowns is 3.

The matrix form of the system is AX = B, where

$$A = \begin{bmatrix} 4 & -2 & 6 \\ 1 & 1 & -3 \\ 15 & -3 & 9 \end{bmatrix}, \quad X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \quad B = \begin{bmatrix} 8 \\ -1 \\ 21 \end{bmatrix}.$$

Applying elementary row operations on the augmented matrix [A | B], we get

$$[A | B] = \begin{bmatrix} 4 & -2 & 6 & | & 8 \\ 1 & 1 & -3 & | & -1 \\ 15 & -3 & 9 & | & 21 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} 1 & 1 & -3 & | & -1 \\ 4 & -2 & 6 & | & 8 \\ 15 & -3 & 9 & | & 21 \end{bmatrix} \xrightarrow{R_2 \rightarrow R_2 - 4R_1,} \begin{bmatrix} 1 & 1 & -3 & | & -1 \\ 0 & -6 & 18 & | & 12 \\ 0 & -18 & 54 & | & 36 \end{bmatrix}$$
$$\xrightarrow{R_2 \rightarrow R_2 \div (-6),} \begin{bmatrix} 1 & 1 & -3 & | & -1 \\ 0 & 1 & -3 & | & -2 \\ 0 & 1 & -3 & | & -2 \end{bmatrix} \xrightarrow{R_3 \rightarrow R_3 - R_2} \begin{bmatrix} 1 & 1 & -3 & | & -1 \\ 0 & 1 & -3 & | & -2 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}$$

So, $\rho(A) = \rho([A | B]) = 2 < 3$. From the echelon form, we get the equivalent equations

$$x + y - 3z = -1, y - 3z = -2, 0 = 0$$

The equivalent system has two non-trivial equations and three unknowns. So, one of the unknowns should be fixed at our choice in order to get two equations for the other two unknowns. We fix z arbitrarily as a real number t, and we get y = 3t-2, x = -1-(3t-2)+3t = 1. So, the solution is (x = 1, y = 3t - 2, z = t), where t is real. The above solution set is a one-parameter family of solutions. Here, the given system is consistent and has infinitely many solutions which form a one parameter family of solutions.

Note

In the above example, the square matrix A is singular and so matrix inversion method cannot be applied to solve the system of equations. However, Gaussian elimination method is applicable and we are able to decide whether the system is consistent or not. The next example also confirms the supremacy of Gaussian elimination method over other methods.

39

Applications of Matrices and Determinants

Example 1.31

Test for consistency of the following system of linear equations and if possible solve: x-y+z=-9, 2x-2y+2z=-18, 3x-3y+3z+27=0.

Solution

Here the number of unknowns is 3.

The matrix form of the system is AX = B, where

$$A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & -3 & 3 \end{bmatrix}, \quad X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \quad B = \begin{bmatrix} -9 \\ -18 \\ -27 \end{bmatrix}.$$

Applying elementary row operations on the augmented matrix [A | B], we get

$$[A | B] = \begin{bmatrix} 1 & -1 & 1 & | & -9 \\ 2 & -2 & 2 & | & -18 \\ 3 & -3 & 3 & | & -27 \end{bmatrix} \xrightarrow{R_2 \to R_2 - 2R_1, \\ R_3 \to R_3 - 3R_1} \begin{bmatrix} 1 & -1 & 1 & | & -9 \\ 0 & 0 & 0 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}$$

So, $\rho(A) = \rho([A | B]) = 1 < 3$.

From the echelon form, we get the equivalent equations x - y + z = -9, 0 = 0, 0 = 0. The equivalent system has one non-trivial equation and three unknowns.

Taking y = s, z = t arbitrarily, we get x - s + t = -9; or x = -9 + s - t.

So, the solution is (x = -9 + s - t, y = s, z = t), where *s* and *t* are parameters. The above solution set is a two-parameter family of solutions.

Here, the given system of equations is consistent and has infinitely many solutions which form a two parameter family of solutions.

Example 1.32

Test the consistency of the following system of linear equations

x - y + z = -9, 2x - y + z = 4, 3x - y + z = 6, 4x - y + 2z = 7.

Solution

Here the number of unknowns is 3.

The matrix form of the system of equations is AX = B, where

$$A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -1 & 1 \\ 3 & -1 & 1 \\ 4 & -1 & 2 \end{bmatrix}, \quad X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \quad B = \begin{bmatrix} -9 \\ 4 \\ 6 \\ 7 \end{bmatrix}.$$

Applying elementary row operations on the augmented matrix [A | B], we get

$$[A|B] = \begin{bmatrix} 1 & -1 & 1 & | & -9 \\ 2 & -1 & 1 & | & 4 \\ 3 & -1 & 1 & | & 6 \\ 4 & -1 & 2 & | & 7 \end{bmatrix} \xrightarrow{R_2 \to R_2 - 2R_1, \\ R_4 \to R_4 - 4R_1 \to R_4 - 2R_2, \\ \hline R_3 \to R_3 - 2R_2, \\ R_4 \to R_4 - 3R_2 \to R_4 - 3R_2 \to R_4 \to R_4 - 2R_4 \to R_4 - 2R_4 \to R_4 \to R_4 - 2R_4 \to R_4 \to R_4 - 2R_4 \to R_4 - 2R_4 \to R_4 \to R_4 - 2R_4 \to R_4 \to$$

So, $\rho(A) = 3$ and $\rho([A | B]) = 4$. Hence $\rho(A) \neq \rho([A | B])$.

XII - Mathematics

If we write the equivalent system of equations using the echelon form, we get

۲

x - y + z = -9, y - z = 22, z = -23, 0 = -11.

The last equation is a contradiction.

So the given system of equations is inconsistent and has no solution.

By Rouché-Capelli theorem, we have the following rule:

- If there are *n* unknowns in the system of equations and $\rho(A) = \rho([A | B]) = n$, then the system AX = B, is consistent and has a unique solution.
- If there are *n* unknowns in the system AX = B, and $\rho(A) = \rho([A | B]) = n k, k \neq 0$ then the system is consistent and has infinitely many solutions and these solutions form a k – parameter family. In particular, if there are 3 unknowns in a system of equations and $\rho(A) = \rho([A | B]) = 2$, then the system has infinitely many solutions and these solutions form a one parameter family. In the same manner, if there are 3 unknowns in a system of equations and $\rho(A) = \rho([A | B]) = 1$, then the system has infinitely many solutions and these solutions form a two parameter family.
- If $\rho(A) \neq \rho([A | B])$, then the system AX = B is inconsistent and has no solution.

Example 1.33

Find the condition on a, b and c so that the following system of linear equations has one parameter family of solutions: x + y + z = a, x + 2y + 3z = b, 3x + 5y + 7z = c.

Solution

Here the number of unknowns is 3.

The matrix form of the system is AX = B, where $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 3 & 5 & 7 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} a \\ b \\ c \end{bmatrix}$.

Applying elementary row operations on the augmented matrix [A | B], we get

$$\begin{bmatrix} A \mid B \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & | & a \\ 1 & 2 & 3 & | & b \\ 3 & 5 & 7 & | & c \end{bmatrix} \xrightarrow{R_2 \to R_3 - 3R_1} \begin{bmatrix} 1 & 1 & 1 & | & a \\ 0 & 1 & 2 & | & b - a \\ 0 & 2 & 4 & | & c - 3a \end{bmatrix}$$
$$\xrightarrow{R_3 \to R_3 - 2R_2} \begin{bmatrix} 1 & 1 & 1 & | & a \\ 0 & 1 & 2 & | & b - a \\ 0 & 1 & 2 & | & b - a \\ 0 & 0 & | & (c - 3a) - 2(b - a) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & | & a \\ 0 & 1 & 2 & | & b - a \\ 0 & 0 & | & (c - 2b - a) \end{bmatrix}.$$

In order that the system should have one parameter family of solutions, we must have

 $\rho(A) = \rho([A, B]) = 2$. So, the third row in the echelon form should be a zero row.

So,
$$c - 2b - a = 0 \implies c = a + 2b$$
.

Example 1.34

Investigate for what values of λ and μ the system of linear equations

x + 2y + z = 7, $x + y + \lambda z = \mu$, x + 3y - 5z = 5

has (i) no solution (ii) a unique solution (iii) an infinite number of solutions.

41

Solution

Here the number of unknowns is 3.

The matrix form of the system is AX = B, where $A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 1 & \lambda \\ 1 & 3 & -5 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 7 \\ \mu \\ 5 \end{bmatrix}$.

Applying elementary row operations on the augmented matrix [A | B], we get

$$\begin{bmatrix} A \mid B \end{bmatrix} = \begin{bmatrix} 1 & 2 & 1 & | & 7 \\ 1 & 1 & \lambda & | & \mu \\ 1 & 3 & -5 & | & 5 \end{bmatrix} \xrightarrow{R_2 \leftrightarrow R_3} \begin{bmatrix} 1 & 2 & 1 & | & 7 \\ 1 & 3 & -5 & | & 5 \\ 1 & 1 & \lambda & | & \mu \end{bmatrix}$$
$$\xrightarrow{R_2 \rightarrow R_2 - R_1, \\ R_3 \rightarrow R_3 - R_1 \longrightarrow \begin{bmatrix} 1 & 2 & 1 & | & 7 \\ 0 & 1 & -6 & | & -2 \\ 0 & -1 & \lambda - 1 & | & \mu - 7 \end{bmatrix} \xrightarrow{R_3 \rightarrow R_3 + R_2} \begin{bmatrix} 1 & 2 & 1 & | & 7 \\ 0 & 1 & -6 & | & -2 \\ 0 & 0 & \lambda - 7 & | & \mu - 9 \end{bmatrix}.$$

- (i) If $\lambda = 7$ and $\mu \neq 9$, then $\rho(A) = 2$ and $\rho([A | B]) = 3$. So $\rho(A) \neq \rho([A | B])$. Hence the given system is inconsistent and has no solution.
- (ii) If λ ≠ 7 and μ is any real number, then ρ(A) = 3 and ρ([A | B]) = 3.
 So ρ(A) = ρ([A | B]) = 3 = Number of unknowns. Hence the given system is consistent and has a unique solution.

(iii) If $\lambda = 7$ and $\mu = 9$, then $\rho(A) = 2$ and $\rho([A | B]) = 2$.

So, $\rho(A) = \rho([A | B]) = 2 <$ Number of unknowns. Hence the given system is consistent and has infinite number of solutions.

EXERCISE 1.6

- 1. Test for consistency and if possible, solve the following systems of equations by rank method.
 - (i) x y + 2z = 2, 2x + y + 4z = 7, 4x y + z = 4
 - (ii) 3x + y + z = 2, x 3y + 2z = 1, 7x y + 4z = 5
 - (iii) 2x + 2y + z = 5, x y + z = 1, 3x + y + 2z = 4
 - (iv) 2x y + z = 2, 6x 3y + 3z = 6, 4x 2y + 2z = 4
- 2. Find the value of k for which the equations kx 2y + z = 1, x 2ky + z = -2, x 2y + kz = 1 have
 - (i) no solution (ii) unique solution (iii) infinitely many solution
- 3. Investigate the values of λ and μ the system of linear equations 2x + 3y + 5z = 9,

7x + 3y - 5z = 8, $2x + 3y + \lambda z = \mu$, have

(i) no solution (ii) a unique solution (iii) an infinite number of solutions.

XII - Mathematics

1.5.2 Homogeneous system of linear equations

We recall that a homogeneous system of linear equations is given by

۲

 $a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \dots + a_{mn}x_n = 0,$

where the coefficients a_{ij} , $i = 1, 2, \dots, m$; $j = 1, 2, \dots, n$ are constants. The above system is always satisfied by $x_1 = 0, x_2 = 0, \dots, x_n = 0$. This solution is called the **trivial solution** of (1). In other words, the system (1) always possesses a solution.

The above system (1) can be put in the matrix form $AX = O_{m \times 1}$, where

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}, X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, O_{m \times 1} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

We will denote $O_{m \times 1}$ simply by the capital letter O.SinceO is the zero column matrix, it is always true that $\rho(A) = \rho([A | O]) \le m$. So, by Rouché-Capelli Theorem, any system of homogeneous linear equations is always consistent.

Suppose that m < n, then there are more number of unknowns than the number of equations. So $\rho(A) = \rho([A | O]) < n$. Hence, system (1) possesses a non-trivial solution.

Suppose that m = n, then there are equal number of equations and unknowns:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = 0,$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = 0,$$

$$\dots \qquad \dots \qquad \dots \qquad \dots$$

$$a_{n1}x_1 + a_{n2}x_2 + a_{n3}x_3 + \dots + a_{nn}x_n = 0,$$

$$\dots \qquad (2)$$

Two cases arise.

Case (i)

If $\rho(A) = \rho([A | O]) = n$, then system (2) has a **unique solution** and it is the **trivial solution**. Since $\rho(A) = n$, $|A| \neq 0$. So for **trivial solution** $|A| \neq 0$.

Case (ii)

If $\rho(A) = \rho([A | O]) < n$, then system (2) has a non-trivial solution. Since $\rho(A) < n$, |A| = 0. In other words, the homogeneous system (2) has a non-trivial solution if and only if the determinant of the coefficient matrix is zero.

Suppose that m > n, then there are more number of equations than the number of unknowns. Reducing the system by elementary transformations, we get $\rho(A) = \rho([A | O]) \le n$.

43

Example 1.35

Solve the following system:

x + 2y + 3z = 0, 3x + 4y + 4z = 0, 7x + 10y + 12z = 0.

Solution

Here the number of equations is equal to the number of unknowns.

Transforming into echelon form (Gaussian elimination method), the augmented matrix becomes

$$\begin{bmatrix} 1 & 2 & 3 & 0 \\ 3 & 4 & 4 & 0 \\ 7 & 10 & 12 & 0 \end{bmatrix} \xrightarrow{R_2 \to R_2 - 3R_1, \\ R_3 \to R_3 - 7R_1} \begin{bmatrix} 1 & 2 & 3 & 0 \\ 0 & -2 & -5 & 0 \\ 0 & -4 & -9 & 0 \end{bmatrix} \xrightarrow{R_2 \to R_2 \div (-1), \\ R_3 \to R_3 \div (-1)} \begin{bmatrix} 1 & 2 & 3 & 0 \\ 0 & 2 & 5 & 0 \\ 0 & 4 & 9 & 0 \end{bmatrix}$$
$$\xrightarrow{R_3 \to R_3 - 2R_2} \begin{bmatrix} 1 & 2 & 3 & 0 \\ 0 & 2 & 5 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix} \xrightarrow{R_3 \to R_3 \div (-1)} \begin{bmatrix} 1 & 2 & 3 & 0 \\ 0 & 2 & 5 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

So, $\rho(A) = \rho([A | O]) = 3$ = Number of unknowns.

Hence, the system has a unique solution. Since x = 0, y = 0, z = 0 is always a solution of the homogeneous system, the only solution is the trivial solution x = 0, y = 0, z = 0.

Note

In the above example, we find that

$$\begin{vmatrix} A \end{vmatrix} = \begin{vmatrix} 1 & 2 & 3 \\ 3 & 4 & 4 \\ 7 & 10 & 12 \end{vmatrix} = 1(48 - 40) - 2(36 - 28) + 3(30 - 28) = 8 - 16 + 6 = -2 \neq 0.$$

Example 1.36

Solve the system: x + 3y - 2z = 0, 2x - y + 4z = 0, x - 11y + 14z = 0.

Solution

Here the number of unknowns is 3.

Transforming into echelon form (Gaussian elimination method), the augmented matrix becomes

$$\begin{bmatrix} 1 & 3 & -2 & | & 0 \\ 2 & -1 & 4 & | & 0 \\ 1 & -11 & 14 & | & 0 \end{bmatrix} \xrightarrow{R_2 \to R_2 - 2R_1, \\ R_3 \to R_1 \to R_1} \begin{bmatrix} 1 & 3 & -2 & | & 0 \\ 0 & -7 & 8 & | & 0 \\ 0 & -14 & 16 & | & 0 \end{bmatrix} \xrightarrow{R_2 \to R_2 \div (-1), \\ R_3 \to R_1 \div (-2) \to R_1 \div (-2) \to R_1 \to R_2 \to$$

So, $\rho(A) = \rho([A | O]) = 2 < 3 =$ Number of unknowns.

Hence, the system has a one parameter family of solutions.

Writing the equations using the echelon form, we get

x + 3y - 2z = 0, 7y - 8z = 0, 0 = 0.

Taking z = t, where t is an arbitrary real number, we get by back substitution,

XII - Mathematics

44

$$z = t,$$

$$7y - 8t = 0 \implies y = \frac{8t}{7},$$

$$x + 3\left(\frac{8t}{7}\right) - 2t = 0 \implies x + \frac{24t - 14t}{7} = 0 \implies x = -\frac{10t}{7}.$$

So, the solution is $\left(x = -\frac{10t}{7}, y = \frac{8t}{7}, z = t\right)$, where t is any real number.

Example 1.37

Solve the system: x + y - 2z = 0, 2x - 3y + z = 0, 3x - 7y + 10z = 0, 6x - 9y + 10z = 0.

Solution

Here the number of equations is 4 and the number of unknowns is 3. Reducing the augmented matrix to echelon-form, we get

$$\begin{bmatrix} A \mid O \end{bmatrix} = \begin{bmatrix} 1 & 1 & -2 \mid 0 \\ 2 & -3 & 1 \mid 0 \\ 3 & -7 & 10 \mid 0 \\ 6 & -9 & 10 \mid 0 \end{bmatrix} \xrightarrow{R_2 \to R_2 - 2R_1, \\ R_4 \to R_4 - 6R_1} \begin{bmatrix} 1 & 1 & -2 \mid 0 \\ 0 & -5 & 5 \mid 0 \\ 0 & -10 & 16 \mid 0 \\ 0 & -15 & 22 \mid 0 \end{bmatrix} \xrightarrow{R_2 \to R_2 \div (-5), \\ R_3 \to R_3 \div (-2)} \begin{bmatrix} 1 & 1 & -2 \mid 0 \\ 0 & 1 & -1 \mid 0 \\ 0 & 5 & -8 \mid 0 \\ 0 & -15 & 22 \mid 0 \end{bmatrix} \xrightarrow{R_3 \to R_3 \div (-2)} \begin{bmatrix} 1 & 1 & -2 \mid 0 \\ 0 & 5 & -8 \mid 0 \\ 0 & -15 & 22 \mid 0 \end{bmatrix}$$
$$\xrightarrow{R_3 \to R_3 - 5R_2, \\ R_4 \to R_4 + 15R_2 \to \\ 0 & 0 & -3 \mid 0 \\ 0 & 0 & 7 \mid 0 \end{bmatrix}} \xrightarrow{R_3 \to R_3 \div (-3), \\ R_4 \to R_4 \div 7 \to \\ 0 & 0 & 1 \mid 0 \\ 0 & 0 & 1 \mid 0 \end{bmatrix}} \xrightarrow{R_4 \to R_4 - R_3} \xrightarrow{R_4 \to R_4 - R_3} \begin{bmatrix} 1 & 1 & -2 \mid 0 \\ 0 & 1 & -1 \mid 0 \\ 0 & 1 & -1 \mid 0 \\ 0 & 0 & 1 \mid 0 \\ 0 & 0 & 1 \mid 0 \end{bmatrix}}$$

So, $\rho(A) = \rho([A | O]) = 3$ = Number of unknowns. Hence the system has trivial solution only.

Example 1.38

Determine the values of λ for which the following system of equations $(3\lambda - 8)x + 3y + 3z = 0$, $3x + (3\lambda - 8)y + 3z = 0$, $3x + 3y + (3\lambda - 8)z = 0$

has a non-trivial solution.

Solution

Here the number of unknowns is 3. So, if the system is consistent and has a non-trivial solution, then the rank of the coefficient matrix is equal to the rank of the augmented matrix and is less than 3. So the determinant of the coefficient matrix should be 0.

Hence we get

 $\begin{vmatrix} 3\lambda - 8 & 3 & 3 \\ 3 & 3\lambda - 8 & 3 \\ 3 & 3 & 3\lambda - 8 \end{vmatrix} = 0 \text{ or } \begin{vmatrix} 3\lambda - 2 & 3\lambda - 2 & 3\lambda - 2 \\ 3 & 3\lambda - 8 & 3 \\ 3 & 3 & 3\lambda - 8 \end{vmatrix} = 0 \text{ (by applying } R_1 \to R_1 + R_2 + R_3)$ or $(3\lambda - 2) \begin{vmatrix} 1 & 1 & 1 \\ 3 & 3\lambda - 8 & 3 \\ 3 & 3 & 3\lambda - 8 \end{vmatrix} = 0 \text{ (by taking out } (3\lambda - 2) \text{ from } R_1)$

45

Applications of Matrices and Determinants

or
$$(3\lambda - 2) \begin{vmatrix} 1 & 1 & 1 \\ 0 & 3\lambda - 11 & 0 \\ 0 & 0 & 3\lambda - 11 \end{vmatrix} = 0$$
 (by applying $R_2 \to R_2 - 3R_1, R_3 \to R_3 - 3R_1$)
or $(3\lambda - 2)(3\lambda - 11)^2 0$. So $\lambda = \frac{2}{3}$ and $\lambda = \frac{11}{3}$.

We now give an application of system of linear homogeneous equations to chemistry. You are already aware of balancing chemical reaction equations by inspecting the number of atoms present on both sides. A direct method is explained as given below.

۲

Example 1.39

By using Gaussian elimination method, balance the chemical reaction equation:

 $C_5H_8 + O_2 \rightarrow CO_2 + H_2O.$

(The above is the reaction that is taking place in the burning of organic compound called isoprene.)

Solution

We are searching for positive integers x_1, x_2, x_3 and x_4 such that

$$x_1 C_5 H_8 + x_2 O_2 = x_3 C O_2 + x_4 H_2 O.$$
 (1)

The number of carbon atoms on the left-hand side of (1) should be equal to the number of carbon atoms on the right-hand side of (1). So we get a linear homogenous equation

$$5x_1 = x_3 \implies 5x_1 - x_3 = 0$$
 ... (2)

Similarly, considering hydrogen and oxygen atoms, we get respectively,

$$8x_1 = 2x_4 \implies 4x_1 - x_4 = 0, \qquad \dots (3)$$

$$2x_2 = 2x_3 + x_4 \implies 2x_2 - 2x_3 - x_4 = 0. \qquad \dots (4)$$

Equations (2), (3), and (4) constitute a homogeneous system of linear equations in four unknowns.

The augmented matrix is $[A | B] = \begin{bmatrix} 5 & 0 & -1 & 0 & | & 0 \\ 4 & 0 & 0 & -1 & | & 0 \\ 0 & 2 & -2 & -1 & | & 0 \end{bmatrix}$.

By Gaussian elimination method, we get

$$\begin{bmatrix} A \mid B \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} 4 & 0 & 0 & -1 & | & 0 \\ 5 & 0 & -1 & 0 & | & 0 \\ 0 & 2 & -2 & -1 & | & 0 \end{bmatrix} \xrightarrow{R_2 \leftrightarrow R_3} \begin{bmatrix} 4 & 0 & 0 & -1 & | & 0 \\ 0 & 2 & -2 & -1 & | & 0 \\ 5 & 0 & -1 & 0 & | & 0 \end{bmatrix}$$
$$\xrightarrow{R_3 \rightarrow 4R_3 - 5R_1} \begin{bmatrix} 4 & 0 & 0 & -1 & | & 0 \\ 0 & 2 & -2 & -1 & | & 0 \\ 0 & 0 & -4 & 5 & | & 0 \end{bmatrix}.$$

Therefore, $\rho(A) = \rho([A | B]) = 3 < 4 =$ Number of unknowns.

The system is consistent and has infinite number of solutions.

Writing the equations using the echelon form, we get $4x_1 - x_4 = 0$, $2x_2 - 2x_3 - x_4 = 0$, $-4x_3 + 5x_4 = 0$. So, one of the unknowns should be chosen arbitrarily as a non-zero real number.

Let us choose
$$x_4 = t, t \neq 0$$
. Then, by back substitution, we get $x_3 = \frac{5t}{4}, x_2 = \frac{7t}{4}, x_1 = \frac{t}{4}$.

XII - Mathematics

Since x_1, x_2, x_3 , and x_4 are positive integers, let us choose t = 4.

Then, we get $x_1 = 1, x_2 = 7, x_3 = 5$ and $x_4 = 4$.

So, the balanced equation is $C_5H_8 + 7O_2 \rightarrow 5CO_2 + 4H_2O$.

Example 1.40

If the system of equations px + by + cz = 0, ax + qy + cz = 0, ax + by + rz = 0 has a non-trivial

۲

solution and $p \neq a, q \neq b, r \neq c$, prove that $\frac{p}{p-a} + \frac{q}{q-b} + \frac{r}{r-c} = 2$.

Solution

Assume that the system px + by + cz = 0, ax + qy + cz = 0, ax + by + rz = 0 has a non-trivial solution.

So, we have
$$\begin{vmatrix} p & b & c \\ a & q & c \\ a & b & r \end{vmatrix} = 0$$
. Applying $R_2 \to R_2 - R_1$ and $R_3 \to R_3 - R_1$ in the above equation,

we get

$$\begin{vmatrix} p & b & c \\ a-p & q-b & 0 \\ a-p & 0 & r-c \end{vmatrix} = 0. \text{ That is, } \begin{vmatrix} p & b & c \\ -(p-a) & q-b & 0 \\ -(p-a) & 0 & r-c \end{vmatrix} = 0.$$

Since
$$p \neq a, q \neq b, r \neq c$$
, we get $(p-a)(q-b)(r-c) \begin{vmatrix} \frac{p}{p-a} & \frac{b}{q-b} & \frac{c}{r-c} \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{vmatrix} = 0$
So, we have $\begin{vmatrix} \frac{p}{p-a} & \frac{b}{q-b} & \frac{c}{r-c} \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{vmatrix} = 0.$

Expanding the determinant, we get $\frac{p}{p-a} + \frac{b}{q-b} + \frac{c}{r-c} = 0.$

That is,
$$\frac{p}{p-a} + \frac{q-(q-b)}{q-b} + \frac{r-(r-c)}{r-c} = 0 \implies \frac{p}{p-a} + \frac{q}{q-b} + \frac{r}{r-c} = 2$$
.

EXERCISE 1.7

1. Solve the following system of homogenous equations.

- (i) 3x + 2y + 7z = 0, 4x 3y 2z = 0, 5x + 9y + 23z = 0
- (ii) 2x + 3y z = 0, x y 2z = 0, 3x + y + 3z = 0
- 2. Determine the values of λ for which the following system of equations x + y + 3z = 0, 4x + 3y + λz = 0, 2x + y + 2z = 0 has
 (i) a unique solution (ii) a non-trivial solution.
- 3. By using Gaussian elimination method, balance the chemical reaction equation: $C_2H_6 + O_2 \rightarrow H_2O + CO_2$

47

Applications of Matrices and Determinants



Choose the Correct of	or the most suitable answer fro	om the given four alt	ernatives :
1. If $ \operatorname{adj}(\operatorname{adj} A) =$	$ A ^9$, then the order of the squar	e matrix A is	
(1) 3	(2) 4	(3) 2	(4) 5
2. If A is a 3×3	non-singular matrix such that A	$A^T = A^T A$ and $B = A^{-1} A$	A^T , then $BB^T =$
(1) A	(2) B	(3) I_{3}	(4) B^T
3. If $A = \begin{bmatrix} 3 & 5 \\ 1 & 2 \end{bmatrix}$,	$B = \operatorname{adj} A$ and $C = 3A$, then $\frac{ \operatorname{adj}}{ C }$	$\frac{B }{C } =$	
(1) $\frac{1}{3}$	(2) $\frac{1}{9}$	(3) $\frac{1}{4}$	(4) 1
$4. \text{ If } A \begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 & 4 \end{bmatrix}$	$\begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix}$, then $A =$		
$(1)\begin{bmatrix}1 & -2\\1 & 4\end{bmatrix}$	$(2)\begin{bmatrix}1&2\\-1&4\end{bmatrix}\qquad(3)\begin{bmatrix}4&2\\-1&1\end{bmatrix}$	$(4)\begin{bmatrix} 4 & -1\\ 2 & 1 \end{bmatrix}$	
5. If $A = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$,	then $9I_2 - A =$		M38JTX
(1) A^{-1}	(2) $\frac{A^{-1}}{2}$	(3) $3A^{-1}$	$(4) 2A^{-1}$
$6. If A = \begin{bmatrix} 2 & 0 \\ 1 & 5 \end{bmatrix} a$	and $B = \begin{bmatrix} 1 & 4 \\ 2 & 0 \end{bmatrix}$ then $ \operatorname{adj}(AB) =$		
(1) -40	(2) -80	(3) -60	(4) -20
7. If $P = \begin{bmatrix} 1 & x \\ 1 & 3 \\ 2 & 4 \end{bmatrix}$	$\begin{bmatrix} 0 \\ 0 \\ -2 \end{bmatrix}$ is the adjoint of 3×3 matrix	A and A = 4, then x	: is
(1) 15	(2) 12	(3) 14	(4) 11
8. If $A = \begin{bmatrix} 3 & 1 \\ 2 & -2 \\ 1 & 2 \end{bmatrix}$	$\begin{bmatrix} -1\\0\\-1 \end{bmatrix} \text{ and } A^{-1} = \begin{bmatrix} a_{11} & a_{12} & a_{13}\\a_{21} & a_{22} & a_{23}\\a_{31} & a_{32} & a_{33} \end{bmatrix} t$	hen the value of a_{23} is	
(1) 0	(2) -2	(3) -3	(4) -1
9. If A, B and C	are invertible matrices of some	e order, then which or	ne of the following is not
true?			C C
(1) $\operatorname{adj} A = A $	$A^{-1} \qquad (2) \operatorname{adj}(AB) = (a)$	$\operatorname{adj} A$)($\operatorname{adj} B$)	
(3) det $A^{-1} = (d)$	et A) ⁻¹ (4) $(ABC)^{-1} = 0$	$C^{-1}B^{-1}A^{-1}$	
10. If $(AB)^{-1} = \begin{bmatrix} 12\\ -1 \end{bmatrix}$	$\begin{bmatrix} 2 & -17 \\ 9 & 27 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix}$, the	$B^{-1} =$	
$(1)\begin{bmatrix}2 & -5\\-3 & 8\end{bmatrix}$	$(2)\begin{bmatrix} 8 & 5\\ 3 & 2 \end{bmatrix}$	$(3)\begin{bmatrix}3 & 1\\2 & 1\end{bmatrix}$	$(4)\begin{bmatrix} 8 & -5\\ -3 & 2 \end{bmatrix}$

11. If $A^T A^{-1}$ is symmetric, then $A^2 =$ (1) A^{-1} (2) $(A^T)^2$ (3) A^{T} $(4) (A^{-1})^2$ 12. If A is a non-singular matrix such that $A^{-1} = \begin{bmatrix} 5 & 3 \\ -2 & -1 \end{bmatrix}$, then $(A^T)^{-1} =$ $(2)\begin{bmatrix}5&3\\-2&-1\end{bmatrix} \qquad (3)\begin{bmatrix}-1&-3\\2&5\end{bmatrix} \qquad (4)\begin{bmatrix}5&-2\\3&-1\end{bmatrix}$ $(1) \begin{vmatrix} -5 & 3 \\ 2 & 1 \end{vmatrix}$ 13. If $A = \begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ x & \frac{3}{5} \end{bmatrix}$ and $A^{T} = A^{-1}$, then the value of x is $(1) \frac{-4}{5}$ $(3) \frac{3}{5}$ $(2) \frac{-3}{5}$ $(4) \frac{4}{5}$ 14. If $A = \begin{bmatrix} 1 & \tan\frac{\theta}{2} \\ -\tan\frac{\theta}{2} & 1 \end{bmatrix}$ and $AB = I_2$, then B =(1) $\left(\cos^2\frac{\theta}{2}\right)A$ (2) $\left(\cos^2\frac{\theta}{2}\right)A^T$ (3) $\left(\cos^2\theta\right)I$ (4) $\left(\sin^2\frac{\theta}{2}\right)A$ 15. If $A = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$ and $A(\operatorname{adj} A) = \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix}$, then $k = \begin{bmatrix} -\sin\theta & \cos\theta \\ 0 & k \end{bmatrix}$ (1) 0(2) $\sin\theta$ (3) $\cos\theta$ (4) 116. If $A = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$ be such that $\lambda A^{-1} = A$, then λ is (3) 19(1) 17(2) 14(4) 2117. If adj $A = \begin{bmatrix} 2 & 3 \\ 4 & -1 \end{bmatrix}$ and adj $B = \begin{bmatrix} 1 & -2 \\ -3 & 1 \end{bmatrix}$ then adj (AB) is $(1) \begin{vmatrix} -7 & -1 \\ 7 & -9 \end{vmatrix}$ $(2)\begin{bmatrix} -6 & 5 \\ -2 & -10 \end{bmatrix} \qquad (3)\begin{bmatrix} -7 & 7 \\ -1 & -9 \end{bmatrix} \qquad (4)\begin{bmatrix} -6 & -2 \\ 5 & -10 \end{bmatrix}$ 18. The rank of the matrix $\begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 4 & 6 & 8 \\ -1 & -2 & -3 & -4 \end{bmatrix}$ is (1) 1(4) 3 (2) 2(3) 419. If $x^a y^b = e^m, x^c y^d = e^n, \Delta_1 = \begin{vmatrix} m & b \\ n & d \end{vmatrix}, \Delta_2 = \begin{vmatrix} a & m \\ c & n \end{vmatrix}, \Delta_3 = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$, then the values of x and y are respectively, (1) $e^{(\Delta_2/\Delta_1)} \cdot e^{(\Delta_3/\Delta_1)}$ (2) $\log(\Delta_1 / \Delta_3), \log(\Delta_2 / \Delta_3)$ (4)) $e^{(\Delta_1/\Delta_3)}, e^{(\Delta_2/\Delta_3)}$ (3) $\log(\Delta_2 / \Delta_1), \log(\Delta_3 / \Delta_1)$ 49 **Applications of Matrices and Determinants**

۲

۲

۲

۲ 20. Which of the following is/are correct? (i) Adjoint of a symmetric matrix is also a symmetric matrix. (ii) Adjoint of a diagonal matrix is also a diagonal matrix. (iii) If A is a square matrix of order n and λ is a scalar, then $adj(\lambda A) = \lambda^n adj(A)$. (iv) A(adjA) = (adjA)A = |A|I(1) Only (i) (2) (ii) and (iii) (3) (iii) and (iv) (4) (i), (ii) and (iv) 21. If $\rho(A) = \rho([A | B])$, then the system AX = B of linear equations is (1) consistent and has a unique solution (2) consistent (3) consistent and has infinitely many solution (4) inconsistent 22. If $0 \le \theta \le \pi$ and the system of equations $x + (\sin \theta)y - (\cos \theta)z = 0, (\cos \theta)x - y + z = 0$, $(\sin\theta)x + y - z = 0$ has a non-trivial solution then θ is (2) $\frac{3\pi}{4}$ (3) $\frac{5\pi}{6}$ (4) $\frac{\pi}{4}$ (1) $\frac{2\pi}{2}$ 23. The augmented matrix of a system of linear equations is $\begin{bmatrix} 1 & 2 & 7 & 3 \\ 0 & 1 & 4 & 6 \\ 0 & 0 & \lambda - 7 & \mu + 5 \end{bmatrix}$. The system has infinitely many solutions if (2) $\lambda = -7, \mu = 5$ (3) $\lambda \neq 7, \mu \neq -5$ (4) $\lambda = 7, \mu = -5$ (1) $\lambda = 7, \mu \neq -5$ 24. Let $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$ and $4B = \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & x \\ -1 & 1 & 3 \end{bmatrix}$. If B is the inverse of A, then the value of x is (1) 2(2) 4 (3) 3 (4) 125. If $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$, then $\operatorname{adj}(\operatorname{adj} A)$ is $(1) \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$ $(2) \begin{bmatrix} 6 & -6 & 8 \\ 4 & -6 & 8 \\ 0 & -2 & 2 \end{bmatrix}$ $(3) \begin{bmatrix} -3 & 3 & -4 \\ -2 & 3 & -4 \\ 0 & 1 & -1 \end{bmatrix}$ $(4) \begin{bmatrix} 3 & -3 & 4 \\ 0 & -1 & 1 \\ 2 & -3 & 4 \end{bmatrix}$

SUMMARY

(1) Adjoint of a square matrix A = Transpose of the cofactor matrix of A.

(2)
$$A(\operatorname{adj} A) = (\operatorname{adj} A)A = |A|I_n$$
.

(3)
$$A^{-1} = \frac{1}{|A|} \operatorname{adj} A.$$

(4) (i) $|A^{-1}| = \frac{1}{|A|}$ (ii) $(A^T)^{-1} = (A^{-1})^T$ (iii) $(\lambda A)^{-1} = \frac{1}{\lambda} A^{-1}$, where λ is a non-zero scalar.

XII - Mathematics

(5) (i) $(AB)^{-1} = B^{-1}A^{-1}$. (ii) $(A^{-1})^{-1} = A$ (6) If A is a non-singular square matrix of order n, then (i) $(adj A)^{-1} = adj(A^{-1}) = \frac{1}{|A|}A$ (ii) $|adj A| = |A|^{n-1}$ (iii) $adj (adj A) = |A|^{n-2}A$ (iv) $adj(\lambda A) = \lambda^{n-1}adj(A)$, λ is a nonzero scalar (v) $|adj (adj A)| = |A|^{(n-1)^2}$ (vi) $(adj A)^T = adj(A^T)$ (vii) adj(AB) = (adjB)(adjA)(7) (i) $A^{-1} = \pm \frac{1}{\sqrt{|adj A|}}adjA$. (ii) $A = \pm \frac{1}{\sqrt{|adj A|}}adj(adjA)$. (8) (i) A matrix A is orthogonal if $AA^T = A^T A = I$ (ii) A matrix A is orthogonal if and only if A is non-singular and $A^{-1} = A^T$ (8) Methods to solve the system of linear equations AX = B(i) By matrix inversion method $X = A^{-1}B$, $|A| \neq 0$

(ii) By Cramer's rule
$$x = \frac{\Delta_1}{\Delta}, y = \frac{\Delta_2}{\Delta}, z = \frac{\Delta_3}{\Delta}, \Delta \neq 0$$

(iii) By Gaussian elimination method

(9) (i) If $\rho(A) = \rho([A | B])$ = number of unknowns, then the system has unique solution.

۲

- (ii) If $\rho(A) = \rho([A | B])$ < number of unknowns, then the system has infinitely many solutions.
- (iii) If $\rho(A) \neq \rho([A | B])$ then the system is inconsistent and has no solution.
- (10) The homogenous system of linear equations AX = 0
 - (i) has the trivial solution, if $|A| \neq 0$.
 - (ii) has a non trivial solution, if |A| = 0.

Scan the QR Code

Scan the QR code. GeoGebra work book named "12th Standard Mathematics" will open. In the left side of the work book there are many chapters related to your text book. Click on the chapter named "Applications of Matrices and Determinants". You can see several work sheets related to the chapter. Select the work sheet "Application Matrices–Problem"



Applications of Matrices and Determinants