

# NBSE Class 12 Maths Question Paper 2019

Total number of printed pages : 5

2019/XII/MAT

## 2019 MATHEMATICS

Full marks: 100

Time: 3 hours

### General instructions:

- i) Approximately 15 minutes is allotted to read the question paper and revise the answers.
  - ii) The question paper consists of 26 questions. All questions are compulsory.
  - iii) Marks are indicated against each question.
  - iv) Internal choice has been provided in some questions.
  - v) Use of simple calculators (non-scientific and non-programmable) only is permitted.
- N.B:** Check that all pages of the question paper is complete as indicated on the top left side.

### Section – A

#### 1. Choose the correct answer from the given alternatives:

- (a) If  $f(x)=|x|$  and  $g(x)=[x]$ , then  $g \circ f(-3.7)$  is equal to 1  
(i) -3.7                      (ii) 3                      (iii) 3.7                      (iv) 4
- (b) Consider the set  $\mathbf{Q}$  with the binary operation  $*$  as  $a * b = \frac{ab}{4}$ . Then the identity element is 1  
(i)  $\frac{1}{4}$                       (ii) 1                      (iii) 4                      (iv) 16
- (c) If a matrix A is both symmetric and skew-symmetric matrix, then 1  
(i) A is a diagonal matrix                      (ii) A is a zero matrix  
(iii) A is a square matrix                      (iv) none of these
- (d) If  $y = a^x x^a$  then  $\frac{dy}{dx}$  is equal to 1  
(i)  $a^x x^{a-1}(a - x \log a)$                       (ii)  $a^x x^{a-1}(a + x \log a)$   
(iii)  $a^x x^a(a + x \log a)$                       (iv)  $a^x x^{a-1}(x + a \log a)$
- (e) The point on the curve  $y = 2x^2$ , where the slope of the tangent is 8, is 1  
(i) (0, 2)                      (ii) (0, 8)                      (iii) (2, 8)                      (iv) (8, 2)
- (f) The value of  $\int \tan^2 x dx$  is 1  
(i)  $x - \tan x + C$                       (ii)  $\tan x + x + C$                       (iii)  $\tan x - x + C$                       (iv)  $x \tan x + C$

- (g) The value of  $\int_{-1}^1 \log\left(\frac{2-x}{2+x}\right) dx$  is 1
- (i) -1                      (ii) 0                      (iii) 1                      (iv) 2
- (h) If  $p\hat{i} + 3\hat{j}$  is a vector of magnitude 5, then the value of  $p$  is 1
- (i) 0                      (ii) 1                      (iii)  $\pm 3$                       (iv)  $\pm 4$
- (i) Let A and B be events such that  $P(A) = \frac{7}{13}$ ,  $P(B) = \frac{9}{13}$  and  $P(A \cap B) = \frac{4}{13}$ , then  $P(A|B)$  is equal to 1
- (i)  $\frac{4}{9}$                       (ii)  $\frac{7}{13}$                       (iii)  $\frac{2}{3}$                       (iv)  $\frac{9}{4}$
- (j) If A & B are two events such that  $P(A) = \frac{1}{4}$ ,  $P(B) = \frac{1}{3}$ ,  $P(A \cup B) = \frac{1}{2}$ , then the events A and B are 1
- (i) independent                      (ii) dependent  
(iii) mutually exclusive                      (iv) none of these

### Section – B

2. Consider the set of real numbers  $\mathbf{R}$ . Define the relation R on  $\mathbf{R}$  as “ $a R b$  if and only if  $a^2 + b^2 = 1$ ”. Write the domain of R. Also, prove that R is not transitive. 2
3. Find  $f \circ g$  and  $g \circ f$  if  $f(x) = |x|$  and  $g(x) = |4x + 3|$ . Are they equal? 2
4. Find the value of  $\tan\left(\tan^{-1}\sqrt{3} + \sin^{-1}\frac{1}{\sqrt{2}} - \cot^{-1}1\right)$  2
5. Solve the following equation for  $x$ :  $\sin\left(\sin^{-1}\frac{1}{5} + \cos^{-1}x\right) = 1$  2
6. If  $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$ , show that  $A'A = I_2$  2
7. Differentiate  $\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$  with respect to  $x$ . 2
8. If  $y = 5\cos x - 3\sin x$ , prove that  $\frac{d^2y}{dx^2} + y = 0$  2
9. Evaluate  $\int \sin^4 x dx$  2

10. Form a differential equation representing the given curve,  $y = ae^{bx}$ , where  $a$  &  $b$  are arbitrary constants. 2
11. Find the value of  $\lambda$  for which  $\vec{a}$  and  $\vec{b}$  are perpendicular if  $\vec{a} = 7\hat{i} - \lambda\hat{j} - 7\hat{k}$  and  $\vec{b} = 4\hat{i} + 5\hat{j} - \hat{k}$  2

### Section – C

12. a. If  $A = \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix}$ , show that  $2A^{-1} = 9I - A$ .

**Or**

- b. Using properties of determinants, prove that:

$$\begin{vmatrix} a+b & b+c & c+a \\ b+c & c+a & a+b \\ c+a & a+b & b+c \end{vmatrix} = 2 \begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix}$$

13. a. If  $y = a \cos(\log x) + b \sin(\log x)$ , prove that  $x^2 y_2 + x y_1 + y = 0$

**Or**

- b. Find the coordinates of the point at which the tangent to the curve  $f(x) = x^2 - 6x + 1$  is parallel to the chord joining the points  $(1, -4)$  and  $(3, -8)$  4

14. a. If  $x = a \sin 2t (1 + \cos 2t)$ ,  $y = b \cos 2t (1 - \cos 2t)$ , show that  $\frac{dy}{dx} = \frac{b}{a}$  at  $t = \frac{\pi}{4}$

**Or**

- b. If  $f(x) = \left(\frac{3+x}{1+x}\right)^{2+3x}$ , find  $f'(0)$ .

15. Evaluate  $\int \frac{\cos^5 x}{\sin x} dx$  4

16. a. Evaluate  $\int_0^{\frac{\pi}{2}} \frac{\cos x}{(1 + \sin x)(2 + \sin x)} dx$

**Or**

- b. Evaluate  $\int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{x}{1 + \sin x} dx = \pi(\sqrt{2} - 1)$  4

17. Solve the differential equation  $x \sin \frac{y}{x} \frac{dy}{dx} + x - y \sin \frac{y}{x} = 0$ , given that  $y(1) = \frac{\pi}{2}$  4
18. **a.** If  $\vec{a} = \hat{i} - \hat{j}$ ,  $\vec{b} = 3\hat{j} - \hat{k}$  and  $\vec{c} = 7\hat{i} - \hat{k}$ , find the vector  $\vec{d}$  which is perpendicular to both  $\vec{a}$  and  $\vec{b}$  and  $\vec{c} \cdot \vec{d} = 4$  4
- Or** 4
- b.** Show that the points A, B, C, D with position vectors  $4\hat{i} + 8\hat{j} + 12\hat{k}$ ,  $2\hat{i} + 4\hat{j} + 6\hat{k}$ ,  $3\hat{i} + 5\hat{j} + 4\hat{k}$  and  $5\hat{i} + 8\hat{j} + 5\hat{k}$  respectively are coplanar.
19. Find the foot and the length of the perpendicular drawn from the point (3, 4, 5) to the plane  $2x - 5y + 3z = 39$  4
20. In a bulb factory, machines A, B and C manufacture 60%, 30% and 10% bulbs respectively. Out of these bulbs, 1%, 2% and 3% of the bulbs produced respectively by A, B and C are found to be defective. A bulb is picked up at random from the total production and found to be defective. Find the probability that this bulb was produced by the machine A. 4
21. A die is tossed once. If the random variable X is defined as:
- $$X = \begin{cases} 1, & \text{if the die results in an even number} \\ 0, & \text{if the die results in an odd number} \end{cases}$$
- 4
- Then, find the mean and variance of X.

#### Section – D

22. **a.** Using elementary row transformations, find the inverse of the matrix  $\begin{bmatrix} 1 & 3 & -2 \\ -3 & 0 & -5 \\ 2 & 5 & 0 \end{bmatrix}$  6
- Or**
- b.** Solve the following system of linear equations using matrix method:
- $$\begin{aligned} 2x + 3y + 3z &= 5 \\ x - 2y + z &= -4 \\ 3x - y - 2z &= 3 \end{aligned}$$
23. **a.** Show that the semi-vertical angle of a cone maximum volume and given slant height is  $\tan^{-1} \sqrt{2}$  6

**Or**

6

- b. Prove that the volume of the largest cone that can be inscribed in a sphere of radius  $R$  is  $\frac{8}{27}$  of the volume of the sphere.

24. a. Find the area of the smaller region bounded by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and the straight line  $\frac{x}{a} + \frac{y}{b} = 1$

**Or**

**6**

- b. Using the method of integration, find the area of the region bounded by the lines  $2x + y = 4$ ,  $3x - 2y = 6$  and  $x - 3y + 5 = 0$

25. a. Find the image of the point  $(1, 3, 4)$  in the plane  $2x - y + z + 3 = 0$ . Also, find the distance of the point from its image.

**Or**

**6**

- b. Show that the lines  $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$  and  $\frac{x-4}{5} = \frac{y-1}{2} = z$  intersect. Also, find their point of intersection.

26. a. A housewife wishes to mix two types of food X and Y in such a way that the mixture contains at least 8 units of vitamin A and 10 units of vitamin B. X contains 2 units/kg of vitamin A and 1 unit/kg of vitamin B. While Y contains 1 unit/kg of vitamin A and 2 units/kg of vitamin B. It costs Rs 60/kg of X and Rs 80/kg of Y. Formulate this problem as a linear programming problem to minimize the cost of such a mixture and solve it.

**Or**

**6**

- b. A shopkeeper wants to invest Rs 5400 on two types of pens. Type A costs Rs 180 per packet and type B costs Rs 60 per packet. He can get a profit of Rs 15 on type A and Rs 10 on type B. He has a space for 50 packets only. Formulate this as an LPP so as to get the number of each type of packets and the maximum profit. Also, find the maximum profit.

\*\*\*\*\*