

EXERCISE 5.3

PAGE NO: 5.41

1. Compute the indicated products:

(<i>i</i>)	$a \\ -b$	$b \\ a$	$\begin{bmatrix} a \\ b \end{bmatrix}$	-b a		
(ii)	$\begin{bmatrix} 1\\ 2 \end{bmatrix}$	$-2 \\ 3$] [1 1 -3 1	$\begin{bmatrix} 2 & 3 \\ 2 & 1 \end{bmatrix}$	
	2	3	4	1	-3^{-1}	5
(iii)	3	4	5	0	2	4
	4	5	6	3	0	5

Solution:

(i) Consider

$$\begin{bmatrix} a & b \\ -b & a \end{bmatrix} \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} a \times a \times +b + b & a \times (-b) + b \times a \\ (-b) \times a + a \times b & (-b) \times (-b) + a \times a \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} a^2 + b^2 & -ab^2 + ab \\ -ab + ab & b^2 + a^2 \end{bmatrix}$$
On simplification we get

On simplification we get,

$$\Rightarrow \begin{bmatrix} a^2 + b^2 & 0 \\ 0 & a^2 + b^2 \end{bmatrix}$$

(ii) Consider

$$\begin{bmatrix} 1 & -2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ -3 & 2 & -1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 \times 1 + (-2 \times (-3)) & 1 \times 2 + (-2) \times 2 & 1 \times 3 + (-2) \times (-1) \\ 2 \times 1 + 3 \times (-3) & 2 \times 2 + 3 \times 2 & 2 \times 3 + 3 \times (-1) \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 + 6 & 2 - 4 & 3 + 2 \\ 2 - 9 & 4 + 6 & 6 - 3 \end{bmatrix}$$

On simplification we get,

$$\Rightarrow \left[egin{array}{ccc} 7 & -2 & 5 \ -7 & 10 & 3 \end{array}
ight]$$



(iii) Consider

 $\begin{bmatrix} 2 & 3 & 4 \\ 3 & 4 & 5 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 1 & -3 & 5 \\ 0 & 2 & 4 \\ 3 & 0 & 5 \end{bmatrix}$ $\Rightarrow \begin{bmatrix} 2 \times 1 + 3 \times 0 + 4 \times 3 & 2 \times (-3) + 3 \times 2 + 4 \times 0 & 2 \times 5 + 3 \times 4 + 4 \times 5 \\ 3 \times 1 + 4 \times 0 + 5 \times 3 & 3 \times (-3) + 4 \times 2 + 5 \times 0 & 3 \times 5 + 4 \times 4 + 5 \times 5 \\ 4 \times 1 + 5 \times 0 + 6 \times 3 & 4 \times (-3) + 5 \times 2 + 6 \times 0 & 4 \times 5 + 5 \times 4 + 6 \times 5 \end{bmatrix}$ $\Rightarrow \begin{bmatrix} 2+0+12 & -6+6+0 & 10+12+20 \\ 3+0+15 & -9+8+0 & 15+16+25 \\ 4+0+18 & -12+10+0 & 20+20+30 \end{bmatrix}$

On simplification we get,

	14	0	42
\Rightarrow	18	-1	56
	22	-2	70

:در ال **2.** Show that AB \neq BA in each of the following cases:

$$(i)A = \begin{bmatrix} 5 & -1 \\ 6 & 7 \end{bmatrix} andB = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$$
$$(ii)A = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \\ 2 & 3 & 4 \end{bmatrix} andB = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \\ 4 & 1 & 0 \end{bmatrix}$$
$$(iii)A = \begin{bmatrix} 1 & 3 & 0 \\ 1 & 3 & 0 \\ 1 & 1 & 0 \\ 4 & 1 & 0 \end{bmatrix} andB = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 5 & 1 \end{bmatrix}$$

Solution:

(i) Consider,

Again consider,



From equation (1) and (2), it is clear that $AB \neq BA$

(ii) Consider,

$$AB = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \\ 2 & 3 & 4 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$$
$$\Rightarrow AB = \begin{bmatrix} -1+0+0 & -2+1+0 & -3+0+0 \\ 0+0+1 & 0-1+1 & 0+0+0 \\ 2+0+4 & 4+3+4 & 6+0+0 \end{bmatrix}$$
$$\Rightarrow AB = \begin{bmatrix} -1 & -1 & -3 \\ 1 & 0 & 0 \\ 6 & 11 & 6 \end{bmatrix} \dots \dots \dots \dots (1)$$



Now again consider,

From equation (1) and (2), it is clear that $AB \neq BA$

(iii) Consider,



Now again consider,

$$BA = egin{bmatrix} 0 & 1 & 0 \ 1 & 0 & 0 \ 0 & 5 & 1 \end{bmatrix} egin{bmatrix} 1 & 3 & 0 \ 1 & 1 & 0 \ 4 & 1 & 0 \end{bmatrix}$$



From equation (1) and (2), it is clear that $AB \neq BA$

3. Compute the products AB and BA whichever exists in each of the following cases:

$$(i)A = \begin{bmatrix} 1 & -2 \\ 2 & 3 \end{bmatrix} andB = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{bmatrix}$$
$$(ii)A = \begin{bmatrix} 3 & 2 \\ -1 & 0 \\ -1 & 1 \end{bmatrix} andB = \begin{bmatrix} 4 & 5 & 6 \\ 0 & 1 & 2 \end{bmatrix}$$
$$(iii)A = \begin{bmatrix} 1 & -1 & 2 & 3 \end{bmatrix} andB = \begin{bmatrix} 0 \\ 1 \\ 3 \\ 2 \end{bmatrix}$$



$$(iv) \begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} + \begin{bmatrix} a & b & c & d \end{bmatrix} \begin{vmatrix} a \\ b \\ c \\ d \end{vmatrix}$$

Solution:

(i) Consider,

$$AB = \begin{bmatrix} 1 & -2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{bmatrix}$$
$$\Rightarrow AB = \begin{bmatrix} 1-4 & 2-6 & 3-2 \\ 2+6 & 4+9 & 6+3 \end{bmatrix}$$
$$\Rightarrow AB = \begin{bmatrix} -3 & -4 & 1 \\ 8 & 13 & 9 \end{bmatrix}$$

BA does not exist

Because the number of columns in B is greater than the rows in A

(ii) Consider,

$$AB = \begin{bmatrix} 3 & 2 \\ -1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 & 6 \\ 0 & 1 & 2 \end{bmatrix}$$

$$\Rightarrow AB = \begin{bmatrix} 12+0 & 15+2 & 18+4 \\ -4+0 & -5+0 & -6+0 \\ -4+0 & -5+1 & -6+2 \end{bmatrix}$$

$$\Rightarrow AB = \begin{bmatrix} 12 & 17 & 22 \\ -4 & -5 & -6 \\ -4 & -4 & -4 \end{bmatrix}$$

Again consider,

$$BA = \begin{bmatrix} 4 & 5 & 6 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ -1 & 0 \\ -1 & 1 \end{bmatrix}$$
$$\Rightarrow BA = \begin{bmatrix} 12 - 5 - 6 & 8 + 0 + 6 \\ 0 - 1 - 2 & 0 + 0 + 2 \end{bmatrix}$$
$$\Rightarrow BA = \begin{bmatrix} 1 & 14 \\ -3 & 2 \end{bmatrix}$$



(iii) Consider,

$$AB = \begin{bmatrix} 1 & -1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 3 \\ 2 \end{bmatrix}$$

$$AB = \begin{bmatrix} 0 + (-1) + 6 + 6 \end{bmatrix}$$

$$AB = 11$$

$$Again consider,$$

$$BA = \begin{bmatrix} 0 \\ 1 \\ 3 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & -1 & 2 & 3 \end{bmatrix}$$

$$\Rightarrow BA = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & -1 & 2 & 3 \\ 3 & -3 & 6 & 9 \\ 2 & -2 & 4 & 6 \end{bmatrix}$$
(iv) Consider,

$$\begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} + \begin{bmatrix} a & b & c & d \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$$

$$\Rightarrow [ac + bd] + [a^2 + b^2 + c^2 + d^2]$$

$$[a^2 + b^2 + c^2 + d^2 + ac + bd]$$

4. Show that AB ≠ BA in each of the following cases:

$$\begin{aligned} (i)A &= \begin{bmatrix} 1 & 3 & -1 \\ 2 & -1 & -1 \\ 3 & 0 & -1 \end{bmatrix} and B = \begin{bmatrix} -2 & 3 & -1 \\ -1 & 2 & -1 \\ -6 & 9 & -4 \end{bmatrix} \\ (ii)A &= \begin{bmatrix} 10 & -4 & -1 \\ -11 & 5 & 0 \\ 9 & -5 & 1 \end{bmatrix} and B = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 4 & 2 \\ 1 & 3 & 2 \end{bmatrix} \end{aligned}$$



Solution:

(i) Consider,

Again consider,

From equation (1) and (2), it is clear that AB ≠ BA

(ii) Consider,

$$AB = \begin{bmatrix} 10 & -4 & -1 \\ -11 & 5 & 0 \\ 9 & -5 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 3 & 4 & 2 \\ 1 & 3 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 10 - 12 - 1 & 20 - 16 - 3 & 10 - 8 - 2 \\ -11 + 15 + 0 & -22 + 20 + 0 & -11 + 10 + 0 \\ 9 - 15 + 1 & 18 - 20 + 3 & 9 - 10 + 2 \end{bmatrix}$$



$$AB = \begin{bmatrix} -3 & 1 & 0\\ 4 & -2 & -1\\ -5 & 1 & 1 \end{bmatrix} \dots \dots (1)$$

Again consider,

$$BA = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 4 & 2 \\ 1 & 3 & 2 \end{bmatrix} \begin{bmatrix} 10 & -4 & -1 \\ -11 & 5 & 0 \\ 9 & -5 & 1 \end{bmatrix}$$

 $= \begin{bmatrix} 10 - 22 + 9 & -4 + 10 - 5 & -9 + 0 + 1 \\ 30 - 44 + 10 & -12 + 20 - 10 & -3 + 0 + 2 \\ 10 - 33 + 18 & -4 + 15 - 10 & -1 + 0 + 2 \end{bmatrix}$ ining AP

$$BA = \begin{bmatrix} -3 & 1 & 0\\ 4 & -2 & -1\\ -5 & 1 & 1 \end{bmatrix} \dots \dots (2)$$

From equation (1) and (2) it is clear that, AB ≠ BA

5. Evaluate the following:

(i)
$$\begin{pmatrix} 1 & 3 \\ -1 & -4 \end{pmatrix} + \begin{pmatrix} 3 & -2 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{pmatrix}$$

(ii) $\begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 2 & 0 & 1 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$
(ii) $\begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 2 & 3 \end{bmatrix} \begin{pmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 2 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 1 & 2 \\ 1 & 0 & 2 \end{bmatrix} \end{pmatrix}$

Solution:

(i) Given

 $\left(\begin{bmatrix} 1 & 3 \\ -1 & -4 \end{bmatrix} + \begin{bmatrix} 3 & -2 \\ -1 & 1 \end{bmatrix} \right) \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix}$

First we have to add first two matrix,



$$\Rightarrow \left(\begin{bmatrix} 1+3 & 3-2\\ -1-1 & -4+1 \end{bmatrix} \right) \begin{bmatrix} 1 & 3 & 5\\ 2 & 4 & 6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 4 & 1\\ -2 & -3 \end{bmatrix} \begin{bmatrix} 1 & 3 & 5\\ 2 & 4 & 6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 4+2 & 12+4 & 20+6\\ -2-6 & -6-12 & -10-18 \end{bmatrix}$$

On simplifying, we get

$$\Rightarrow \begin{bmatrix} 6 & 16 & 26\\ -8 & -18 & -28 \end{bmatrix}$$

(ii) Given,

$$\begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2\\ 2 & 0 & 1\\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 2\\ 4\\ 6 \end{bmatrix}$$

First we have to multiply first two given matrix,

$$\Rightarrow \begin{bmatrix} 1+4+0 & 0+0+3 & 2+2+6 \end{bmatrix} \begin{bmatrix} 2\\ 4\\ 6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 5 & 3 & 10 \end{bmatrix} \begin{bmatrix} 2\\ 4\\ 6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 10+12+60 \end{bmatrix} = 82$$

(iii) Given

$$\begin{bmatrix} 1 & -1\\ 0 & 2\\ 2 & 3 \end{bmatrix} \left(\begin{bmatrix} 1 & 0 & 2\\ 2 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 1 & 2\\ 1 & 0 & 2 \end{bmatrix} \right)$$

First we have subtract the matrix which is inside the bracket,

$$\Rightarrow \begin{bmatrix} 1 & -1\\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1-0 & 0-1 & 2-2\\ 1 & 0 & 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1-0 & 0-1 & 2-2 \\ 2-1 & 0-0 & 1-2 \end{bmatrix}$$

=

https://byjus.com



$$\Rightarrow \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \end{bmatrix}$$
$$\Rightarrow \begin{bmatrix} 1-1 & -1+0 & 0+1 \\ 0+2 & 0+0 & 0-2 \\ 2+3 & -2+0 & 0-3 \end{bmatrix}$$
$$\Rightarrow \begin{bmatrix} 0 & -1 & 1 \\ 2 & 0 & -2 \\ 5 & -2 & -3 \end{bmatrix}$$

6.If
$$\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
, $\mathbf{B} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ and $\mathbf{C} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, show that $\mathbf{A}^2 = \mathbf{B}^2 = \mathbf{C}^2 = \mathbf{I}_2$
Solution:
Given
 $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ and $C = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$
We know that,
 $A^2 = AA$
 $\Rightarrow A^2 = \begin{bmatrix} 1 & 0 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 \end{bmatrix}$

Solution:

Given

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \text{ and } C = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

We know that,

Again we know that,

Now, consider,



 $C^2 = C C$

We have,

$$I_2=egin{bmatrix} 1&0\0&1 \end{bmatrix}$$
.....(4)

Now, from equation (1), (2), (3) and (4), it is clear that $A^2 = B^2 = C^2 = I_2$ ning Apt

7.If
$$\mathbf{A} = \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix}$$
, $\mathbf{B} = \begin{bmatrix} 0 & 4 \\ -1 & 7 \end{bmatrix}$, find $3\mathbf{A^2} - 2\mathbf{B} + \mathbf{I}$

Solution:

Given

$$A = \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix}, B = \begin{bmatrix} 0 & 4 \\ -1 & 7 \end{bmatrix}$$

Consider,

$$A^2 = A A$$

$$\Rightarrow A^{2} = \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix}$$
$$\Rightarrow A^{2} = \begin{bmatrix} 4-3 & -2-2 \\ 6+6 & -3+4 \end{bmatrix}$$
$$\Rightarrow A^{2} = \begin{bmatrix} 1 & -4 \\ 12 & 1 \end{bmatrix}$$

Now we have to find, $3A^2 - 2B + I$

$$\Rightarrow 3A^2 - 2B + I = 3\begin{bmatrix} 1 & -4 \\ 12 & 1 \end{bmatrix} - 2\begin{bmatrix} 0 & 4 \\ -1 & 7 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
$$\Rightarrow 3A^2 - 2B + I = \begin{bmatrix} 3 & -12 \\ 36 & 3 \end{bmatrix} - \begin{bmatrix} 0 & 8 \\ -2 & 14 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$



$$egin{array}{lll} \Rightarrow 3A^2-2B+I = egin{bmatrix} 3-0+1 & -12-8+0\ 36+2+0 & 3-14+1 \end{bmatrix} \ \Rightarrow 3A^2-2B+I = egin{bmatrix} 4 & -20\ 38 & -10 \end{bmatrix} \end{array}$$

8.If
$$\mathbf{A} = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$$
, prove that $(\mathbf{A} - 2\mathbf{I})(\mathbf{A} - 3\mathbf{I}) = \mathbf{0}$.

Solution:

Given $A = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$ Consider, $\Rightarrow (A-2I)(A-3I) = \left(\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} - 2 \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \right) \left(\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} - 3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right)$ $\Rightarrow (A-2I)(A-3I) = \left(\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} - \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \right) \left(\begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix} - \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} \right)$ $\Rightarrow (A-2I)(A-3I) = \begin{bmatrix} 4-2 & 2-0 \\ -1-0 & 1-2 \end{bmatrix} \begin{bmatrix} 4-3 & 2-0 \\ -1-0 & 1-3 \end{bmatrix}$ $\Rightarrow (A-2I)(A-3I) = \begin{bmatrix} 2 & 2 \\ -1 & -1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ -1 & -2 \end{bmatrix}$ $\Rightarrow (A-2I)(A-3I) = \begin{bmatrix} 2-2 & 4-4 \\ -1+1 & -2+2 \end{bmatrix}$ $\Rightarrow (A-2I)(A-3I) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ $\Rightarrow (A-2I)(A-3I) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ $\Rightarrow (A-2I)(A-3I) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

Hence the proof.

9.If
$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$
, show that $\mathbf{A}^2 = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ and $\mathbf{A}^3 = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$

Solution:

Given,



$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

Consider,
$$A^{2} = A A$$

$$A^{2} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

$$A^{2} = \begin{bmatrix} 1+0 & 1+1 \\ 0+0 & 0+1 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

Again consider,
$$A^{3} = A^{2}A$$

$$A^{3} = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

A

$$egin{aligned} &A^3 = A^2 A\ &A^3 = egin{bmatrix} 1 & 2\ 0 & 1 \end{bmatrix} egin{bmatrix} 1 & 1\ 0 & 1 \end{bmatrix}\ &\Rightarrow A^3 = egin{bmatrix} 1+0 & 1+2\ 0+0 & 0+1 \end{bmatrix}\ &A^3 = egin{bmatrix} 1 & 3\ 0 & 1 \end{bmatrix} \end{aligned}$$

Hence the proof.

10.If
$$\mathbf{A} = \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix}$$
, show that $\mathbf{A}^2 = \mathbf{0}$

Solution:

Given,

$$A = \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix}$$

Consider,

$$A^2 = A A$$

$$egin{array}{lll} \Rightarrow A^2 = egin{bmatrix} ab & b^2 \ -a^2 & -ab \end{bmatrix} egin{bmatrix} ab & b^2 \ -a^2 & -ab \end{bmatrix} \ \Rightarrow A^2 = egin{bmatrix} a^2b^2 - a^2b^2 & ab^3 - ab^3 \ -a^3b + a^3b & -a^2b^2 + a^2b^2 \end{bmatrix}$$



$$\Rightarrow A^2 egin{bmatrix} 0 & 0 \ 0 & 0 \end{bmatrix}$$

$$\Rightarrow A^2 = 0$$

Hence the proof.

11.If
$$\mathbf{A} = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$
, find \mathbf{A}^2

Solution:

Given,

$$A = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$

Consider,

 $egin{aligned} &A^2 = A \ &\Rightarrow A^2 = \left[egin{aligned} &\cos 2 heta \,\sin 2 heta \ &-\sin 2 heta \,\cos 2 heta \end{array}
ight] \left[egin{aligned} &\cos 2 heta \,\sin 2 heta \ &-\sin 2 heta \,\cos 2 heta \end{array}
ight] \ &\Rightarrow A^2 = \left[egin{aligned} &\cos 2 heta \,\sin 2 heta \ &-\sin 2 heta \,\cos 2 heta \end{array}
ight] &\cos 2 heta \,\sin 2 heta \,\cos 2 heta \ &-\sin 2 heta \,\cos 2 heta \ &-\sin^2(2 heta) + \cos^2(2 heta) \end{array}
ight] \end{aligned}$

We know that, $\cos^2\theta - \sin^2\theta = \cos^2(2\theta)$

$$\Rightarrow A^2 = egin{bmatrix} \cos artheta & = \cos (2 artheta) \ -2\sin 2 artheta \cos (2 artheta) \ -2\sin 2 artheta \cos (2 artheta) \ \cos (2 imes 2 artheta) \end{bmatrix}$$

Again we have,

 $\sin 2\theta = 2\sin\theta\cos\theta$

$$\Rightarrow A^2 = egin{bmatrix} \cos 4 heta \, \sin(2 imes 2 heta) \ -\sin(2 imes 2 heta) \, \cos 4 heta \end{bmatrix}$$
 $\Rightarrow A^2 = egin{bmatrix} \cos 4 heta \, \sin 4 heta \ -\sin 4 heta \, \cos 4 heta \end{bmatrix}$

12.If
$$\mathbf{A} = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$$
 and $\mathbf{B} = \begin{bmatrix} -1 & 3 & 5 \\ 1 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$ show that $\mathbf{AB} = \mathbf{BA} = \mathbf{0}_{\mathbf{3} \times \mathbf{3}}$

Solution:



Given, $A = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix} \text{ and } B = \begin{bmatrix} -1 & 3 & 5 \\ 1 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$ Consider. $AB = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix} \begin{bmatrix} -1 & 3 & 5 \\ 1 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$ $= \begin{bmatrix} -2-3+5 & 6+9-15 & 5+15-20\\ 1+4-5 & -3-12+15 & -5-15+20\\ -1-3+4 & 3+9-12 & 5+15-20 \end{bmatrix}$ arning Apr $= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ $AB = 0_{3 \times 3} \dots (1)$ Again consider, $\mathsf{BA} = \begin{bmatrix} -1 & 3 & 5\\ 1 & -3 & -5\\ -1 & 2 & 5 \end{bmatrix} \begin{bmatrix} 2 & -3 & -5\\ -1 & 4 & 5\\ 1 & -3 & -5 \end{bmatrix}$ $= \begin{bmatrix} -2-3+5 & 3+12-15 & 5+15-20\\ 2+3-5 & -3-12+15 & -5-15+20\\ -2-3+5 & 3+9-12 & 5+15-20 \end{bmatrix}$ $= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

 $BA = 0_{3\times3}$ (2) From equation (1) and (2) $AB = BA = 0_{3\times3}$

$$13.If \ A = egin{bmatrix} 0 & c & -b \ -c & 0 & a \ b & -a & 0 \end{bmatrix} \ and \ B = egin{bmatrix} a^2 & ab & ac \ ab & b^2 & bc \ ac & bc & c^2 \end{bmatrix} \ show \ that \ AB = BA = 0_{3 imes 3}$$

Solution:

Given

$$A = \begin{bmatrix} 0 & c & -b \\ -c & 0 & a \\ b & -a & 0 \end{bmatrix} \text{ and } B = \begin{bmatrix} a^2 & ab & ac \\ ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix}$$

Consider,

$$AB = egin{bmatrix} 0 & c & -b \ -c & 0 & a \ b & -a & 0 \end{bmatrix} egin{bmatrix} a^2 & ab & ac \ ab & b^2 & bc \ ac & bc & c^2 \end{bmatrix}$$

$$AB = \begin{bmatrix} -c & 0 & a \\ b & -a & 0 \end{bmatrix} \begin{bmatrix} ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix}$$
$$\Rightarrow AB = \begin{bmatrix} 0 + abc - abc & 0 + b^2c - b^2c & 0 + bc^2 - bc^2 \\ -a^2c + 0 + a^2c & -abc + 0 + abc & -ac^2 + 0 + ac^2 \\ a^2b - a^2b + 0 & ab^2 - ab^2 + 0 & abc - abc + 0 \end{bmatrix}$$

$$\Rightarrow AB = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

 $\Rightarrow AB = O_{3 \times 3} \dots (1)$ Again consider, $BA = \begin{bmatrix} a^2 & ab & ac \\ ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix} \begin{bmatrix} 0 & c & -b \\ -c & 0 & a \\ b & -a & 0 \end{bmatrix}$

$$\Rightarrow BA = \begin{bmatrix} 0 - abc + abc & a^2c + 0 - a^2c & -a^2b + a^2b + 0\\ 0 - b^2c + b^2c & abc + 0 - abc & -ab^2 + ab^2 + 0\\ 0 - bc^2 + bc^2 & ac^2 + 0 - ac^2 & -abc + abc + 0 \end{bmatrix}$$



$$\Rightarrow BA = egin{bmatrix} 0 & 0 & 0 \ 0 & 0 & 0 \ 0 & 0 & 0 \end{bmatrix}$$

$$\Rightarrow BA = O_{3 \times 3} \dots (2)$$

From equation (1) and (2) $AB = BA = O_{3x3}$

$$14.If \ A = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix} \ and \ B = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix} \ show \ that \ AB = A \ and \ BA = B.$$

Solution:

Given

Solution:
Given

$$A = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix} and B = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$$
Now consider,

$$\begin{bmatrix} 2 & -3 & -5 \\ -1 & -3 & -4 \end{bmatrix} \begin{bmatrix} 2 & -2 & -4 \\ -1 & -2 & -3 \end{bmatrix}$$

Now consider,

$$AB = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix} \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$$

$$= \begin{bmatrix} 4+3-5 & -4-9+10 & -8-12+15\\ -2-4+5 & 2+12-10 & 4+16-15\\ 2+3-4 & -2-9+18 & -4-12+12 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$$

Therefore AB = AAgain consider, BA we get,

$$\mathsf{BA} = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix} \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$$



$$= \begin{bmatrix} 4+2-4 & -6-8+12 & -10-10+16\\ -2-3+4 & 3+12-12 & 5+15-16\\ 2+2-3 & -3-8+9 & -5-10+12 \end{bmatrix}$$

 $= \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$

Hence BA = B Hence the proof.

15.Let
$$A = \begin{bmatrix} -1 & 1 & -1 \\ 3 & -3 & 3 \\ 5 & 5 & 5 \end{bmatrix}$$
 and $B = \begin{bmatrix} 0 & 4 & 3 \\ 1 & -3 & -3 \\ -1 & 4 & 4 \end{bmatrix}$, compute $A^2 - B^2$.
Solution:
Given,
 $A = \begin{bmatrix} -1 & 1 & -1 \\ 3 & -3 & 3 \\ 5 & 5 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 4 & 3 \\ 1 & -3 & -3 \\ -1 & 4 & 4 \end{bmatrix}$
Consider,
 $A = \begin{bmatrix} -1 & 1 & -1 \\ -1 & 1 & -1 \end{bmatrix} \begin{bmatrix} -1 & 1 & -1 \end{bmatrix}$

Solution:

Given,

$$A = \begin{bmatrix} -1 & 1 & -1 \\ 3 & -3 & 3 \\ 5 & 5 & 5 \end{bmatrix} and B = \begin{bmatrix} 0 & 4 & 3 \\ 1 & -3 & -3 \\ -1 & 4 & 4 \end{bmatrix}$$

Consider,

$$A^{2} = \begin{bmatrix} -1 & 1 & -1 \\ 3 & -3 & 3 \\ 5 & 5 & 5 \end{bmatrix} \begin{bmatrix} -1 & 1 & -1 \\ 3 & -3 & 3 \\ 5 & 5 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 1+3+5 & -1-3-5 & 1+3-5 \\ -3-9+15 & 3+9+15 & -3-9+15 \\ -5+15+25 & 5-15+25 & -5+15+25 \end{bmatrix}$$

$$A^{2} = \begin{bmatrix} -1 & 9 & -1 \\ 3 & 27 & 3 \\ 35 & 15 & 35 \end{bmatrix} \dots \dots (1)$$

Now again consider, B²



$$B^{2} = \begin{bmatrix} 0 & 4 & 3 \\ 1 & -3 & -3 \\ -1 & 4 & 4 \end{bmatrix} \begin{bmatrix} 0 & 4 & 3 \\ 1 & -3 & -3 \\ -1 & 4 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} 0 + 4 + 3 & 0 - 12 + 12 & 0 - 12 + 12 \\ 0 - 3 + 3 & 4 + 9 - 12 & 3 + 9 - 12 \\ 0 + 4 - 4 & -4 - 12 + 16 & -3 - 12 + 16 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \dots \dots (2)$$

Now by subtracting equation (2) from equation (1) we get,

 $A^{2} - B^{2} = \begin{bmatrix} -1 & 9 & -1 \\ 3 & 27 & 3 \\ 35 & 15 & 35 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

 $= \begin{bmatrix} -2 & 9 & -1 \\ 3 & 26 & 3 \\ 35 & 15 & 34 \end{bmatrix}$

16. For the following matrices verify the associativity of matrix multiplication i.e. (AB) C = A (BC)

$$(i)A = \begin{bmatrix} 1 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ -1 & 2 \\ 0 & 3 \end{bmatrix}, and C = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$
$$(ii)A = \begin{bmatrix} 4 & 2 & 3 \\ 1 & 1 & 2 \\ 3 & 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix}, and C = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

Solution:

(i) Given

$$A = \begin{bmatrix} 1 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ -1 & 2 \\ 0 & 3 \end{bmatrix}, and C = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Consider,





$$(AB)C = \left(\begin{bmatrix} 1 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 2 \\ 0 & 3 \end{bmatrix} \right) \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$
$$= \begin{bmatrix} 1-2+0 & 0+4+0 \\ -1+0+0 & 0+0+3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$
$$= \begin{bmatrix} -1 & 4 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$
$$= \begin{bmatrix} -1-4 \\ -1 & 3 \end{bmatrix}$$
$$(AB)C = \begin{bmatrix} -5 \\ -4 \end{bmatrix} \dots (1)$$
Now consider RHS,
$$A(BC) = \begin{bmatrix} 1 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix} \left(\begin{bmatrix} 1 & 0 \\ -1 & 2 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right)$$
$$= \begin{bmatrix} 1 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1+0 \\ -1-2 \\ 0-3 \end{bmatrix}$$
$$= \begin{bmatrix} 1 & 2 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -3 \\ -3 \end{bmatrix}$$
$$= \begin{bmatrix} 1-6+0 \\ -1+0-3 \end{bmatrix}$$

$$A(BC) = \begin{bmatrix} -5 \\ -4 \end{bmatrix} \dots \dots (2)$$

From equation (1) and (2), it is clear that (AB) C = A (BC)

https://byjus.com



(ii) Given,

$$A = \begin{bmatrix} 4 & 2 & 3 \\ 1 & 1 & 2 \\ 3 & 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix}, and C = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$
Consider the LHS,

$$(AB)C = \begin{bmatrix} 4 & 2 & 3 \\ 1 & 1 & 2 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 4 + 0 + 6 & -4 + 2 - 3 & 4 + 4 + 3 \\ 1 + 0 + 4 & -1 + 1 - 2 & 1 + 2 + 2 \\ 3 + 0 + 2 & -3 + 0 - 1 & 3 + 0 + 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 10 & -5 & 11 \\ 5 & -2 & 5 \\ 5 & -4 & 4 \end{bmatrix} \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 10 - 15 + 0 & 20 + 0 + 0 & -10 + 5 + 11 \\ 5 - 6 + 0 & 10 + 0 + 0 & -5 - 2 + 5 \\ 5 - 12 + 0 & 10 + 0 + 0 & -5 - 4 + 4 \end{bmatrix}$$

$$(AB)C = \begin{bmatrix} -5 & 20 & -4 \\ -1 & 10 & -2 \\ -7 & 10 & -5 \end{bmatrix} \dots \dots (1)$$
Now consider RHS,

$$A(BC) = \begin{bmatrix} 4 & 2 & 3 \\ 1 & 1 & 2 \\ 3 & 0 & 1 \end{bmatrix} \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 2 & 3 \\ 1 & 1 & 2 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 - 3 + 0 & 2 + 0 + 0 & -1 - 1 + 1 \\ 0 + 3 + 0 & 0 + 0 + 0 & -2 - 1 + 1 \end{bmatrix}$$



$$= \begin{bmatrix} 4 & 2 & 3 \\ 1 & 1 & 2 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} -2 & 2 & 1 \\ 3 & 0 & 3 \\ -1 & 4 & -2 \end{bmatrix}$$

$$=\begin{bmatrix} -8 + 6 - 3 & 8 + 0 + 12 & -4 + 6 - 6 \\ -2 + 3 - 2 & 2 + 0 + 8 & -1 + 3 - 4 \\ -6 + 0 - 1 & 6 + 0 + 4 & -3 + 0 - 2 \end{bmatrix}$$

$$A(BC) = \begin{bmatrix} -5 & 20 & -4 \\ -1 & 10 & -2 \\ -7 & 10 & -5 \end{bmatrix} \dots \dots (2)$$

From equation (1) and (2), it is clear that (AB) C = A (BC)

17. For the following matrices verify the distributivity of matrix multiplication over matrix addition i.e. A (B + C) = AB + AC.

$$\begin{aligned} (i)A &= \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix}, \ B &= \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, \ and \ C &= \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \\ (ii)A &= \begin{bmatrix} 2 & -1 \\ 1 & 1 \\ -1 & 2 \end{bmatrix}, \ B &= \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}, \ and \ C &= \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

Solution:

(i) Given

$$A = \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix}, B = \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix}, and C = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}$$

Consider LHS,

 $A(B + C) = \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \left(\begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \right)$

$$= \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} -1 + 0 & 0 + 1 \\ 2 + 1 & 1 - 1 \end{bmatrix}$$

 $= \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 3 & 0 \end{bmatrix}$





$$= \begin{bmatrix} -1 - 3 & 1 + 0 \\ 0 + 6 & 0 + 0 \end{bmatrix}$$

$$A(B + C) = \begin{bmatrix} -4 & 1 \\ 6 & 0 \end{bmatrix} \dots \dots (1)$$

Now consider RHS,
$$AB = AC = \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 2 & 1 \end{bmatrix} + \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}$$

 $= \begin{bmatrix} -1-2 & 0-1 \\ 0+4 & 0+2 \end{bmatrix} + \begin{bmatrix} 0+-1 & 1+1 \\ 0+2 & 0-2 \end{bmatrix}$

$$= \begin{bmatrix} -3 & -1 \\ 4 & 2 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 2 & -2 \end{bmatrix}$$

 $= \begin{bmatrix} -3-1 & -1+2\\ 4+2 & 2-2 \end{bmatrix}$

$$AB + AC = \begin{bmatrix} -4 & 1 \\ 6 & 0 \end{bmatrix} \dots \dots (2)$$

From equation (1) and (2), it is clear that A (B + C) = AB + AC

(ii) Given,

$$A = \begin{bmatrix} 2 & -1 \\ 1 & 1 \\ -1 & 2 \end{bmatrix}, B = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}, and C = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

Consider the LHS

$$A(B + C) = \begin{bmatrix} 2 & -1 \\ 1 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & -1 \\ 1 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 0 + 1 & 1 - 1 \\ 1 + 0 & 1 + 1 \end{bmatrix}$$

RD Sharma Solutions for Class 12 Maths Chapter 5 **Algebra of Matrices**



$$= \begin{bmatrix} 2 & -1 \\ 1 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 2-1 & 0+2\\ 1+1 & 0+2\\ -1+2 & 0+4 \end{bmatrix}$$

$$A(B + C) = \begin{bmatrix} 1 & -2 \\ 2 & 2 \\ 1 & 4 \end{bmatrix} \dots \dots (1)$$

Now consider RHS,

$$A(B + C) = \begin{bmatrix} 2 & 2 \\ 1 & 4 \end{bmatrix} \dots \dots (1)$$

Now consider RHS,
$$AB + AC = \begin{bmatrix} 2 & -1 \\ 1 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 2 & -1 \\ 1 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 0 + 1 & 2 - 1 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 + 0 & -2 - 1 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 2 + 0 & -2 - 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 + 1 & 2 - 1 \\ 0 + 1 & 1 + 1 \\ 0 + 2 & -1 + 2 \end{bmatrix} + \begin{bmatrix} 2 + 0 & -2 - 1 \\ 1 + 0 & -1 + 1 \\ -1 + 0 & 1 + 2 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & 1 \\ 1 & 2 \\ 2 & 1 \end{bmatrix} + \begin{bmatrix} 2 & -3 \\ 1 & 0 \\ -1 & 3 \end{bmatrix}$$

 $= \begin{bmatrix} -1 + 2 & 1 - 3 \\ 1 + 1 & 2 + 0 \\ 2 - 1 & 1 + 3 \end{bmatrix}$

$$AB + AC = \begin{bmatrix} 1 & -2 \\ 2 & 2 \\ 1 & 4 \end{bmatrix} \dots \dots (2)$$

$$18.IfA = \begin{bmatrix} 1 & 0 & -2 \\ 3 & -1 & 0 \\ -2 & 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 0 & 5 & -4 \\ -2 & 1 & 3 \\ -1 & 0 & 2 \end{bmatrix}, and C = \begin{bmatrix} 1 & 5 & 2 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix},$$



verify that A(B-C) = AB - AC.

Solution:

Given,

$$A = \begin{bmatrix} 1 & 0 & -2 \\ 3 & -1 & 0 \\ -2 & 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 0 & 5 & -4 \\ -2 & 1 & 3 \\ -1 & 0 & 2 \end{bmatrix}$$

 $C = \begin{bmatrix} 1 & 5 & 2 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$

Consider the LHS,

Consider the LHS,

$$A(B - C) = \begin{bmatrix} 1 & 0 & -2 \\ 3 & -1 & 0 \\ -2 & 1 & 1 \end{bmatrix} \left(\begin{bmatrix} 0 & 5 & -4 \\ -2 & 1 & 3 \\ -1 & 0 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 5 & 2 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \right)$$

$$= \begin{bmatrix} 1 & 0 & -2 \\ 3 & -1 & 0 \\ -2 & 1 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 & 6 \\ -1 & 0 & 3 \\ -1 & 1 & 1 \end{bmatrix}$$

$$A(B - C) = \begin{bmatrix} 1 & -2 & -8 \\ -2 & 0 & -21 \\ 0 & 1 & 16 \end{bmatrix}$$

Now consider RHS

$$AB - AC = \begin{bmatrix} 1 & 0 & -2 \\ 3 & -1 & 0 \\ -2 & 1 & 1 \end{bmatrix} \begin{bmatrix} 0 & 5 & -4 \\ -2 & 1 & 3 \\ -1 & 0 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 0 & -2 \\ 3 & -1 & 0 \\ -2 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 5 & 2 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 5 & -8 \\ 2 & 14 & -15 \\ -3 & -9 & 13 \end{bmatrix} - \begin{bmatrix} 1 & 7 & 0 \\ 4 & 14 & 6 \\ -3 & -10 & -3 \end{bmatrix}$$

 $= \begin{bmatrix} 1 & -2 & -8 \\ -2 & 0 & -21 \\ 0 & 1 & 16 \end{bmatrix}$



From the above equations LHS = RHS Therefore, A (B - C) = AB - AC.

19. Compute the elements a_{43} and a_{22} of the matrix:

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 2 \\ 0 & 3 & 2 \\ 4 & 0 & 4 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -3 & 2 \\ 4 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 & -1 & 2 & -2 \\ 3 & -3 & 4 & -4 & 0 \end{bmatrix}$$

Solution:

Given

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 2 \\ 0 & 3 & 2 \\ 4 & 0 & 4 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -3 & 2 \\ 4 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 & -1 & 2 & -2 \\ 3 & -3 & 4 & -4 & 0 \end{bmatrix}$$

$$A = \begin{bmatrix} -3 & 2 \\ 12 & 4 \\ -1 & 12 \\ 24 & 8 \end{bmatrix} \begin{bmatrix} 0 & 1 & -1 & 2 & -2 \\ 3 & -3 & 4 & -4 & 0 \end{bmatrix}$$

$$A = \begin{bmatrix} 6 & -9 & 11 & -14 & 6 \\ 12 & 0 & 4 & 8 & -24 \\ 36 & -37 & 49 & -50 & 2 \\ 24 & 0 & 8 & 16 & -48 \end{bmatrix}$$

From the above matrix, $a_{43} = 8$ and $a_{22} = 0$

 $20.IfA = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ p & q & r \end{bmatrix} \text{ and } I \text{ is the identity matrix of order } 3, that \ A^3 = pI + qA + rA^2$

Solution:

Given

 $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ p & q & r \end{bmatrix}$

Consider, $A^2 = A.A$

$$= \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ p & q & r \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ p & q & r \end{bmatrix}$$

$$= \begin{bmatrix} 0+0+0 & 0+0+0 & 0+1+0\\ 0+0+p & 0+0+q & 0+0+r\\ 0+0+pr & p+0+qr & 0+q+r^2 \end{bmatrix}$$

Again conside

Now, consider the RHS

 $pI + qA + rA^2$

 $A^{3} = A^{2}.A$

$$= \begin{bmatrix} 0+0+0 & 0+0+0 & 0+1+0\\ 0+0+p & 0+0+q & 0+0+r\\ 0+0+pr & p+0+ar & 0+a+r^2 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0\\ 0 & 0 & 1\\ p & q & r \end{bmatrix}$$

 $= p \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + q \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ p & q & r \end{bmatrix} + r \begin{bmatrix} 0 & 0 & 1 \\ p & q & r \\ pr & p + qr & q + r^2 \end{bmatrix}$

$$= \begin{bmatrix} 0+0+p & 0+0+q & 0+0+r \\ 0+0+pr & p+0+qr & 0+q+r^2 \\ 0+0+pq+pr^2 & pr+0+q^2+qr^2 & 0+p+qr+qr+r^2 \end{bmatrix}$$

$$= \begin{bmatrix} 0+0+p & 0+0+q & 0+0+r \\ 0+0+pr & p+0+qr & 0+q+r^2 \\ 0+0+pq+pr^2 & pr+0+q^2+qr^2 & 0+p+qr+qr+r^2 \end{bmatrix}$$

$$= \begin{bmatrix} 0+0+p & 0+0+q & 0+0+r \\ 0+0+pr & p+0+qr & 0+q+r^2 \end{bmatrix}$$

$$= \begin{bmatrix} 0 + 0 + 0 & 0 + 0 + 0 & 0 + 1 + 0 \\ 0 + 0 + p & 0 + 0 + q & 0 + 0 + r \\ 0 + 0 + p r & p + 0 + q r & 0 + q + r^{2} \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ p & q & r \end{bmatrix}$$

 $= \begin{bmatrix} p & q & r \\ pr & p+qr & q+r^2 \\ pq+pr^2 & pr+q^2+qr^2 & p+2qr+r^2 \end{bmatrix}$

$$\begin{bmatrix} 0+0+0 & 0+0+0 & 0+1+0 \\ 0+0+p & 0+0+q & 0+0+r \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{array}{c} + & 0 \\ 0 + & r \\ + & r^2 \end{array}$$

$$\begin{pmatrix} 1 + 0 \\ 0 + r \\ q + r^2 \end{bmatrix}$$



$$= \begin{bmatrix} p & q & r \\ pr & p+qr & q+r^2 \\ pq+pr^2 & pr+q^2+qr^2 & p+2qr+r^2 \end{bmatrix}$$

Therefore, $A^3 = p I + q A + rA^2$ Hence the proof.

21. If ω is a complex cube root of unity, show that

$$\left(\begin{bmatrix}1&\omega&\omega^2\\\omega&\omega^2&1\\\omega^2&1&\omega\end{bmatrix}+\begin{bmatrix}\omega&\omega^2&1\\\omega^2&1&\omega\\\omega&\omega^2&1\end{bmatrix}\right)\begin{bmatrix}1\\\omega\\\omega^2\end{bmatrix}=\begin{bmatrix}0\\0\\0\end{bmatrix}$$

Solution:

Given

$$\left(\begin{bmatrix}1 & \omega & \omega^2\\ \omega & \omega^2 & 1\\ \omega^2 & 1 & \omega\end{bmatrix} + \begin{bmatrix}\omega & \omega^2 & 1\\ \omega^2 & 1 & \omega\\ \omega & \omega^2 & 1\end{bmatrix}\right)\begin{bmatrix}1\\ \omega\\ \omega^2\end{bmatrix} = \begin{bmatrix}0\\ 0\\ 0\end{bmatrix}$$

It is also given that $\boldsymbol{\omega}$ is a complex cube root of unity, Consider the LHS,

$$= \begin{bmatrix} 1+\omega & \omega+\omega^2 & \omega^2+1\\ \omega+\omega^2 & \omega^2+1 & 1+\omega\\ \omega^2+\omega & 1+\omega^2 & \omega+1 \end{bmatrix} \begin{bmatrix} 1\\ \omega\\ \omega^2 \end{bmatrix}$$

We know that $1 + \omega + \omega^2 = 0$ and $\omega^3 = 1$

$$= \begin{bmatrix} -\omega^2 & -1 & -\omega \\ -1 & -\omega & -\omega^2 \\ -1 & -\omega & -\omega^2 \end{bmatrix} \begin{bmatrix} 1 \\ \omega \\ \omega^2 \end{bmatrix}$$

Now by simplifying we get,

$$= \begin{bmatrix} -\omega^2 & -\omega & -\omega^3 \\ -1 & -\omega^2 & -\omega^4 \\ -1 & -\omega^2 & -\omega^4 \end{bmatrix}$$

Again by substituting $1 + \omega + \omega^2 = 0$ and $\omega^3 = 1$ in above matrix we get,

 $egin{array}{c} 0 \\ 0 \\ 0 \end{array}$

[0]

Therefore LHS = RHS Hence the proof.

https://byjus.com



22.If
$$A = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$$
, show that $A^2 = A$

Solution:

Given,

$$A = \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$$

Consider A²

 $A^2 = A.A$

$$= \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix} \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$$

$$A^{2} = A A$$

$$= \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix} \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix}$$

$$= \begin{bmatrix} 4 + 3 - 5 & -6 - 12 + 15 & -10 - 15 + 20 \\ -2 - 4 + 5 & 3 + 16 - 15 & 5 + 20 - 20 \\ 2 + 3 - 4 & -3 - 12 + 12 & -5 - 15 + 16 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & -3 & -5 \\ -1 & 4 & 5 \\ 1 & -3 & -4 \end{bmatrix} = \mathbf{A}$$

Therefore $A^2 = A$

23.If
$$A = \begin{bmatrix} 4 & -1 & -4 \\ 3 & 0 & -4 \\ 3 & -1 & -3 \end{bmatrix}$$
, show that $A^2 = I_3$

Solution:

Given

 $A = \begin{bmatrix} 4 & -1 & -4 \\ 3 & 0 & -4 \\ 3 & -1 & -3 \end{bmatrix}$ Consider A²,



 $A^2 = A. A$

$$= \begin{bmatrix} 4 & -1 & -4 \\ 3 & 0 & -4 \\ 3 & -1 & -3 \end{bmatrix} \begin{bmatrix} 4 & -1 & -4 \\ 3 & 0 & -4 \\ 3 & -1 & -3 \end{bmatrix}$$
$$= \begin{bmatrix} 16 - 3 - 12 & -4 + 0 + 4 & 16 + 4 + 12 \\ 12 + 0 - 12 & -3 + 0 + 4 & -12 + 0 + 12 \\ 12 - 3 - 9 & -3 + 0 + 3 & -12 + 4 + 9 \end{bmatrix}$$
$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_3$$
Hence $A^2 = I_3$
Hence $A^2 = I_3$
(ii) If $\begin{bmatrix} 1 & 1 & x \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} -4 & 6 \\ -9 & x \end{bmatrix}$, find x.
(iii) If $\begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix} \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} -4 & 6 \\ -9 & x \end{bmatrix}$, find x.
Solution:
(i) Given
 $\begin{bmatrix} 1 & 1 & x \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 \\ -9 & x \end{bmatrix}$, find x.
$$= \begin{bmatrix} 1 + 2x + 0 & x + 0 + 2 & 2 + 1 + 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = 0$$
$$\Rightarrow \begin{bmatrix} 2x + 4 & x + 2 & 2x + 4 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = 0$$
$$= \begin{bmatrix} 2x + 1 + 2 + x + 3 \end{bmatrix} = 0$$
$$= \begin{bmatrix} 3x + 6 \end{bmatrix} = 0$$
$$= 3x = -6$$



x = -6/3 x = -2

(ii) Given,

$$\begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix} \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} -4 & 6 \\ -9 & x \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 2-6 & -6+12 \\ 5-14 & -15+28 \end{bmatrix} = \begin{bmatrix} -4 & 6 \\ -9 & x \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -4 & 6 \\ -9 & 13 \end{bmatrix} = \begin{bmatrix} -4 & 6 \\ -9 & x \end{bmatrix}$$

On comparing the above matrix we get, x = 13

25. If
$$\begin{bmatrix} x & 4 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 2 \\ 1 & 0 & 2 \\ 0 & 2 & -4 \end{bmatrix} \begin{bmatrix} x \\ 4 \\ -1 \end{bmatrix} = 0$$
, find x.
Solution:
Given

Solution:

Given

$$\begin{bmatrix} x & 4 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 2 \\ 1 & 0 & 2 \\ 0 & 2 & -4 \end{bmatrix} \begin{bmatrix} x \\ 4 \\ -1 \end{bmatrix}$$

$$\Rightarrow [2x + 4 + 0 \quad x + 0 + 2 \quad 2x + 8 - 4] \begin{bmatrix} x \\ 4 \\ -1 \end{bmatrix} = 0$$

$$\Rightarrow [2x + 4 \quad x + 2 \quad 2x + 4] \begin{bmatrix} x \\ 4 \\ -1 \end{bmatrix} = 0$$

$$\Rightarrow [(2x + 4) x + 4 (x + 2) - 1(2x + 4)] = 0$$

$$\Rightarrow 2x^{2} + 4x + 4x + 8 - 2x - 4 = 0$$

$$\Rightarrow 2x^{2} + 6x + 4 = 0$$

$$\Rightarrow 2x^{2} + 2x + 4x + 4 = 0$$

$$\Rightarrow 2x^{2} + 2x + 4x + 4 = 0$$

$$\Rightarrow 2x (x + 1) + 4 (x + 1) = 0$$

$$\Rightarrow (x + 1) (2x + 4) = 0$$

$$\Rightarrow x = -1 \text{ or } x = -2$$

https://byjus.com





Hence, x = -1 or x = -2

26. If
$$\begin{bmatrix} 1 & -1 & x \end{bmatrix} \begin{bmatrix} 0 & 1 & -1 \\ 2 & 1 & 3 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = 0$$
, find x.

Solution:

Given

$$\begin{bmatrix} 1 & -1 & x \end{bmatrix} \begin{bmatrix} 0 & 1 & -1 \\ 2 & 1 & 3 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = 0$$

By multiplying we get,

$$\Rightarrow \begin{bmatrix} 0 - 2 + x & x & (-1) - 3 + x \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = 0$$

$$\Rightarrow \begin{bmatrix} x - 2 & x & x - 4 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = 0$$

$$\begin{bmatrix} (x - 2) \times 0 + x \times 1 + (x - 4) \times 1 \end{bmatrix} = 0$$

$$\Rightarrow \begin{bmatrix} x - 2 & x & x - 4 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = 0$$

$$[(x-2) \times 0 + x \times 1 + (x-4) \times 1] = 0$$

$$\Rightarrow x + x - 4 = 0$$

 $\Rightarrow 2x = 4 \Rightarrow x = 2$

27. If
$$A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$$
 and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then prove that $A^2 - A + 2I = 0$.

Solution:

Given

 $A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix} and I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ Now we have to prove $A^2 - A + 2I = 0$



Now, we will find the matrix for A^2 , we get

$$A^{2} = A \times A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 3 \times 3 + (-2 \times 4) & 3 \times (-2) + (-2 \times -2) \\ 4 \times 3 + (-2 \times 4) & 4 \times (-2) + (-2 \times -2) \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 9 - 8 & -6 + 4 \\ 12 - 8 & -8 + 4 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 1 & -2 \\ 4 & -4 \end{bmatrix} \dots \dots \dots (i)$$

Now, we will find the matrix for 2l, we get

$$2I = 2\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow 2I = \begin{bmatrix} 2 \times 1 & 2 \times 0 \\ 2 \times 0 & 2 \times 1 \end{bmatrix}$$

$$\Rightarrow 2I = \begin{bmatrix} 2 & 1 & 2 \times 0 \\ 2 \times 0 & 2 \times 1 \end{bmatrix}$$

$$\Rightarrow 2I = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \dots \dots \dots (ii)$$

Substitute corresponding values from eqn (i) and eqn (ii), we get

 $\Rightarrow = \begin{bmatrix} 1 & -2 \\ 4 & -4 \end{bmatrix} - \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix} + \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$ $\Rightarrow = \begin{bmatrix} 1 - 3 + 2 & -2 - (-2) + 0 \\ 4 - 4 + 0 & -4 - (-2) + 2 \end{bmatrix}$ $\Rightarrow = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$ Therefore, $A^2 - A + 2I = 0$

Hence proved

28. If
$$A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$$
 and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then prove that $A^2 = 5A + \lambda I$.



Solution:

Given

$$A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} and I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Now, we have to find A²,

$$A^{2} = A \times A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 3 \times 3 + (1 \times -1) & 3 \times 1 + 1 \times 2 \\ (-1 \times 3) + 2 \times (-1) & (-1 \times 1) + 2 \times 2 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 9 - 1 & 3 + 2 \\ -3 - 2 & -1 + 4 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} \dots \dots \dots (i)$$

Now, we will find the matrix for 5A, we get

$$5A = 5\begin{bmatrix} 3 & 1\\ -1 & 2 \end{bmatrix}$$

$$\Rightarrow 5A = \begin{bmatrix} 5 \times 3 & 5 \times 1\\ 5 \times (-1) & 5 \times 2 \end{bmatrix}$$

$$\Rightarrow 5A = \begin{bmatrix} 15 & 5\\ -5 & 10 \end{bmatrix} \dots \dots \dots \dots \dots (ii)$$

So,

 $A^2~=~5A~+~\lambda I$

Substitute corresponding values from eqn (i) and eqn (ii), we get

$$\Rightarrow \begin{bmatrix} 8 & 5\\ -5 & 3 \end{bmatrix} = \begin{bmatrix} 15 & 5\\ -5 & 10 \end{bmatrix} + \lambda \begin{bmatrix} 1 & 0\\ 0 & 1 \end{bmatrix}$$
$$\Rightarrow \begin{bmatrix} 8 & 5\\ -5 & 3 \end{bmatrix} = \begin{bmatrix} 15 & 5\\ -5 & 10 \end{bmatrix} + \begin{bmatrix} \lambda & 0\\ 0 & \lambda \end{bmatrix}$$





 $\begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} = \begin{bmatrix} 15 + \lambda & 5 + 0 \\ -5 + 0 & 10 + \lambda \end{bmatrix}$

And to satisfy the above condition of equality, the corresponding entries of the matrices should be equal,

Hence,

 $8 = 15 + \lambda \Rightarrow \lambda = -7$

 $3 = 10 + \lambda \Rightarrow \lambda = -7$

29. If
$$A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$$
 show that $A^2 - 5A + 7I_2 = 0$.

Solution:

Given

$$A = \begin{bmatrix} 3 & 1\\ -1 & 2 \end{bmatrix}$$

 I_2 is an identity matrix of size 2, so

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

To show that

 $A^2 - 5A + 7I_2 = 0$ Now, we will find the matrix for A^2 , we get $A^2 = A \times A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$

$$\Rightarrow A^{2} = \begin{bmatrix} 3 \times 3 + (1 \times -1) & 3 \times 1 + 1 \times 2 \\ (-1 \times 3) + 2 \times (-1) & (-1 \times 1) + 2 \times 2 \end{bmatrix}$$
$$\Rightarrow A^{2} = \begin{bmatrix} 9 - 1 & 3 + 2 \\ -3 - 2 & -1 + 4 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} \dots \dots \dots \dots (i)$$



Now, we will find the matrix for 5A, we get

Now,

$$7I_{2} = 7\begin{bmatrix} 1 & 0\\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 7 & 0\\ 0 & 7 \end{bmatrix} \dots \dots (iii)$$

So,
$$A^{2} - 5A + 7I_{2}$$

Substitute corresponding values from eqn (i), (ii) and (iii), we get

$$\Rightarrow = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix} + \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$$
$$\Rightarrow = \begin{bmatrix} 8 - 15 + 7 & 5 - 5 + 0 \\ -5 - (-5) + 0 & 3 - 10 + 7 \end{bmatrix}$$
$$\Rightarrow = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$$
Hence the proof.

30. If
$$A = \begin{bmatrix} 2 & 3 \\ -1 & 0 \end{bmatrix}$$
 show that $A^2 - 2A + 3I_2 = 0$.

Solution:

Given

$$A = \begin{bmatrix} 2 & 3\\ -1 & 0 \end{bmatrix}$$

 $I_{\rm 2}$ is an identity matrix of size 2, so

 $I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$



Now we have to show, $A^2 - 2A + 3I_2 = 0$

Now, we will find the matrix for A², we get

$$A^{2} = A \times A = \begin{bmatrix} 2 & 3 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ -1 & 0 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 2 \times 2 + (3 \times -1) & 2 \times 3 + 3 \times 0 \\ (-1 \times 2) + 0 \times (-1) & (-1 \times 3) + 0 \times 0 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 4 - 3 & 6 + 0 \\ -2 + 0 & -3 + 0 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 1 & 6 \\ -2 & -3 \end{bmatrix} \dots \dots \dots (i)$$

Now, we will find the matrix for 2A, we get

$$2A = 2 \begin{bmatrix} 2 & 3 \\ -1 & 0 \end{bmatrix}$$

$$\Rightarrow 2A = \begin{bmatrix} 2 \times 2 & 2 \times 3 \\ 2 \times (-1) & 2 \times 0 \end{bmatrix}$$

$$\Rightarrow 2A = \begin{bmatrix} 4 & 6 \\ -2 & 0 \end{bmatrix} \dots \dots \dots (ii)$$

Now,

$$3I_{2} = 3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} \dots \dots (iii)$$

So,

 $A^2 - 2A + 3I_2$

Substitute corresponding values from eqn (j), (ii) and (iii), we get

$$\Rightarrow = \begin{bmatrix} 1 & 6 \\ -2 & -3 \end{bmatrix} - \begin{bmatrix} 4 & 6 \\ -2 & 0 \end{bmatrix} + \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$$
$$\Rightarrow = \begin{bmatrix} 1-4+3 & 6-6+0 \\ -2-(-2)+0 & -3-0+3 \end{bmatrix}$$
$$\Rightarrow = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$$

Hence the proof.



31. Show that the matrix
$$A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$$
 satisfies the equation $A^3 - 4A^2 + A = 0$.

Solution:

Given

 $A = \begin{bmatrix} 2 & 3\\ 1 & 2 \end{bmatrix}$

To show that $A^3 - 4A^2 + A = 0$

Now, we will find the matrix for A², we get $A^{2} = (A \times A) = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ $\Rightarrow A^{2} = \begin{bmatrix} 2 \times 2 + (3 \times 1) & 2 \times 3 + 3 \times 2 \\ 1 \times 2 + 2 \times 1 & 1 \times 3 + 2 \times 2 \end{bmatrix}$ $\Rightarrow A^{2} = \begin{bmatrix} 4 + 3 & 6 + 6 \\ 2 + 2 & 3 + 4 \end{bmatrix}$ $\Rightarrow A^{2} = \begin{bmatrix} 7 & 12 \\ 4 & 7 \end{bmatrix} \dots \dots \dots \dots (i)$ Now, we will find the matrix for A³, we get $A^{3} = A^{2} \times A = \begin{bmatrix} 7 & 12 \\ 4 & 7 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ $\Rightarrow A^{3} = \begin{bmatrix} 7 \times 2 + 12 \times 1 & 7 \times 3 + 12 \times 2 \\ 4 \times 2 + 7 \times 1 & 4 \times 3 + 7 \times 2 \end{bmatrix}$ $\Rightarrow A^{3} = \begin{bmatrix} 14 + 12 & 21 + 24 \\ 8 + 7 & 12 + 14 \end{bmatrix}$ $\Rightarrow A^{3} = \begin{bmatrix} 26 & 45 \\ 15 & 26 \end{bmatrix} \dots \dots \dots \dots (ii)$ So, $A^{3} - 4A^{2} + A$

Substitute corresponding values from eqn (i) and (ii), we get

$$\Rightarrow = \begin{bmatrix} 26 & 45\\ 15 & 26 \end{bmatrix} - 4 \begin{bmatrix} 7 & 12\\ 4 & 7 \end{bmatrix} + \begin{bmatrix} 2 & 3\\ 1 & 2 \end{bmatrix}$$



$$\Rightarrow = \begin{bmatrix} 26 & 45 \\ 15 & 26 \end{bmatrix} - \begin{bmatrix} 4 \times 7 & 4 \times 12 \\ 4 \times 4 & 4 \times 7 \end{bmatrix} + \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$$

$$\Rightarrow = \begin{bmatrix} 26 & 45 \\ 15 & 26 \end{bmatrix} - \begin{bmatrix} 28 & 48 \\ 16 & 28 \end{bmatrix} + \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$$

$$\Rightarrow = \begin{bmatrix} 26 - 28 + 2 & 45 - 48 + 3 \\ 15 - 16 + 1 & 26 - 28 + 2 \end{bmatrix}$$

$$\Rightarrow = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$$
Therefore,
$$A^{3} - 4A^{2} + A = 0$$

Hence matrix A satisfies the given equation.

32. Show that the matrix $A = \begin{bmatrix} 5 & 3 \\ 12 & 7 \end{bmatrix}$ satisfies the equation $A^2 - 12A - I = 0$.

Solution:

Given

 $A = \begin{bmatrix} 5 & 3\\ 12 & 7 \end{bmatrix}$

I is an identity matrix so $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

To show that $A^2 - 12A - I = 0$

Now, we will find the matrix for A^2 , we get

$$A^{2} = A \times A = \begin{bmatrix} 5 & 3\\ 12 & 7 \end{bmatrix} \begin{bmatrix} 5 & 3\\ 12 & 7 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 5 \times 5 + 3 \times 12 & 5 \times 3 + 3 \times 7\\ 12 \times 5 + 7 \times 12 & 12 \times 3 + 7 \times 7 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 25 + 36 & 15 + 21\\ 60 + 84 & 36 + 49 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 61 & 36\\ 144 & 85 \end{bmatrix} \dots \dots \dots (i)$$





Now, we will find the matrix for 12A, we get

So,

$$A^2 - 12A - I$$

APF Substitute corresponding values from eqn (i) and (ii), we get

 $\Rightarrow = \begin{bmatrix} 61 & 36 \\ 144 & 85 \end{bmatrix} - \begin{bmatrix} 60 & 36 \\ 144 & 84 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ $\Rightarrow = \begin{bmatrix} 61 - 60 - 1 & 36 - 36 - 0 \\ 144 - 144 - 0 & 85 - 84 - 1 \end{bmatrix}$ $\Rightarrow = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$

Therefore, $A^2 - 12A - I = 0$

Hence matrix A is the root of the given equation.

33. If
$$A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$$
 find $A^2 - 5A - 14I$.

Solution:

Given

$$A = \begin{bmatrix} 3 & -5\\ -4 & 2 \end{bmatrix}$$

I is identity matrix so

$$14I = 14\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 14 & 0 \\ 0 & 14 \end{bmatrix}$$



To find $A^2 - 5A - 14I$

Now, we will find the matrix for A², we get

$$A^{2} = A \times A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 3 \times 3 + (-5 \times -4) & 3 \times (-5) + (-5 \times 2) \\ (-4 \times 3) + (2 \times -4) & (-4 \times -5) + 2 \times 2 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 9 + 20 & -15 - 10 \\ -12 - 8 & 20 + 4 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 29 & -25 \\ -20 & 24 \end{bmatrix} \dots \dots \dots (i)$$

Now, we will find the matrix for 5A, we get

$$5A = 5\begin{bmatrix} 3 & -5\\ -4 & 2 \end{bmatrix}$$
$$\Rightarrow 5A = \begin{bmatrix} 5 \times 3 & 5 \times (-5)\\ 5 \times (-4) & 5 \times 2 \end{bmatrix}$$
$$\Rightarrow 5A = \begin{bmatrix} 15 & -25\\ -20 & 10 \end{bmatrix} \dots \dots \dots \dots \dots (ii)$$



So,

 $A^2 - 5A - 14I$

Substitute corresponding values from eqn (i) and (ii), we get

$$\Rightarrow = \begin{bmatrix} 29 & -25 \\ -20 & 24 \end{bmatrix} - \begin{bmatrix} 15 & -25 \\ -20 & 10 \end{bmatrix} - \begin{bmatrix} 14 & 0 \\ 0 & 14 \end{bmatrix}$$
$$\Rightarrow = \begin{bmatrix} 29 - 15 - 14 & -25 + 25 - 0 \\ -20 + 20 - 0 & 24 - 10 - 14 \end{bmatrix}$$
$$\Rightarrow = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$$

34. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$ show that $A^2 - 5A + 7I = 0$. Use this to find A^4 .

https://byjus.com



Solution:

Given

$$A = \begin{bmatrix} 3 & 1\\ -1 & 2 \end{bmatrix}$$

I is identity matrix so

$$7I = 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$$

To show that $A^2 - 5A + 7I = 0$

Now, we will find the matrix for A^2 , we get

$$A^{2} = A \times A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 3 \times 3 + (1 \times -1) & 3 \times 1 + 1 \times 2 \\ (-1 \times 3) + (2 \times -1) & (-1 \times 1) + 2 \times 2 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 9 - 1 & 3 + 2 \\ -3 - 2 & -1 + 4 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} \dots \dots \dots (i)$$

Now, we will find the matrix for 5A, we get

So,

 $A^2 - 5A + 7I$

Substitute corresponding values from eqn (i) and (ii), we get

 $\Rightarrow = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix} - \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$





$$\Rightarrow = \begin{bmatrix} 8 - 15 - 7 & 5 - 5 - 0 \\ -5 + 5 - 0 & 3 - 10 - 7 \end{bmatrix}$$
$$\Rightarrow = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$$

Therefore

 $A^2 - 5A + 7I = 0$

Hence proved

We will find A⁴

$$A^2 - 5A + 7I = 0$$

Multiply both sides by A², we get

$$A^{2}(A^{2} - 5A + 7I) = A^{2}(0)$$

$$\Rightarrow A^{4} - 5A^{2}.A + 7I.A^{2}$$

$$\Rightarrow A^{4} = 5A^{2}.A - 7I.A^{2}$$

$$\Rightarrow A^{4} = 5A^{2}A - 7A^{2}$$

As multiplying by the identity matrix, I don't change anything. Now will substitute the corresponding values we get

$$\Rightarrow A^{4} = 5 \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} - 7 \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix}
\Rightarrow A^{4} = 5 \begin{bmatrix} 24-5 & 8+10 \\ -15-3 & -5+6 \end{bmatrix} - 7 \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix}
\Rightarrow A^{4} = 5 \begin{bmatrix} 19 & 18 \\ -18 & 1 \end{bmatrix} - 7 \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix}
\Rightarrow A^{4} = \begin{bmatrix} 5 \times 19 & 5 \times 18 \\ 5 \times (-18) & 5 \times 1 \end{bmatrix} - \begin{bmatrix} 7 \times 8 & 7 \times 5 \\ 7 \times (-5) & 7 \times 3 \end{bmatrix}
\Rightarrow A^{4} = \begin{bmatrix} 95 & 90 \\ -90 & 5 \end{bmatrix} - \begin{bmatrix} 56 & 35 \\ -35 & 21 \end{bmatrix}
\Rightarrow A^{4} = \begin{bmatrix} 95-56 & 90-35 \\ -90 + 35 & 5-21 \end{bmatrix}$$



$$\Rightarrow A^4 = \begin{bmatrix} 39 & 55 \\ -55 & -16 \end{bmatrix}$$

35. If
$$A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$$
 find k such that $A^2 = kA - 2I_2$.

Solution:

Given

$$A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$$

 $I_{\rm 2}$ is an identity matrix of size 2, so

$$2I_2 = 2\begin{bmatrix} 1 & 0\\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0\\ 0 & 2 \end{bmatrix}$$

Also given,

$$A^2 = kA - 2I_2$$

Now, we will find the matrix for A², we get

$$A^{2} = A \times A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 3 \times 3 + (-2 \times 4) & 3 \times (-2) + (-2 \times -2) \\ (4 \times 3) + (-2 \times 4) & (4 \times -2) + (-2 \times -2) \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 9 - 8 & -6 + 4 \\ 12 - 8 & -8 + 4 \end{bmatrix}$$

$$\Rightarrow A^{2} = \begin{bmatrix} 1 & -2 \\ 4 & -4 \end{bmatrix} \dots \dots \dots (i)$$

Now, we will find the matrix for kA, we get

$$kA = k \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$$
$$\Rightarrow kA = \begin{bmatrix} k \times 3 & k \times (-2) \\ k \times 4 & k \times (-2) \end{bmatrix}$$





So,

$$A^2 = kA - 2I_2$$

Substitute corresponding values from eqn (i) and (ii), we get

$$\Rightarrow \begin{bmatrix} 1 & -2\\ 4 & -4 \end{bmatrix} = \begin{bmatrix} 3k & -2k\\ 4k & -2k \end{bmatrix} - \begin{bmatrix} 2 & 0\\ 0 & 2 \end{bmatrix}$$
$$\Rightarrow \begin{bmatrix} 1 & -2\\ 4 & -4 \end{bmatrix} = \begin{bmatrix} 3k-2 & -2k-0\\ 4k-0 & -2k-2 \end{bmatrix}$$

And to satisfy the above condition of equality, the corresponding entries of the matrices should be equal

Hence, $3k-2 = 1 \Rightarrow k = 1$

Therefore, the value of k is 1

36. If
$$A = \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix}$$
 find k such that $A^2 - 8A + kI = 0$.

Solution:

Given

 $A = \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix}$ I is identity matrix, so $kI = k \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix}$ Also given, $A^2 - 8A + kI = 0$ Now, we have to find A^2 , we get $A^2 = A \times A = \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix}$ $\Rightarrow A^2 = \begin{bmatrix} 1 \times 1 + 0 & 0 + 0 \\ (-1 \times 1) + 7 \times (-1) & 0 + 7 \times 7 \end{bmatrix}$ $\Rightarrow A^2 = \begin{bmatrix} 1 & 0 \\ -8 & 49 \end{bmatrix} \dots \dots (i)$ Now, we will find the matrix for 8A, we get $8A = 8 \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix}$



$$\Rightarrow 8A = \begin{bmatrix} 8 \times 1 & 8 \times 0 \\ 8 \times (-1) & 8 \times 7 \end{bmatrix}$$

$$\Rightarrow 8A = \begin{bmatrix} 8 & 0 \\ -8 & 56 \end{bmatrix} \dots \dots \dots \dots \dots (ii)$$

So,

$$A^{2} - 8A + kI = 0$$

Substitute corresponding values from eqn (i) and (ii), we get

$$\Rightarrow \begin{bmatrix} 1 & 0 \\ -8 & 49 \end{bmatrix} - \begin{bmatrix} 8 & 0 \\ -8 & 56 \end{bmatrix} + \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix} = 0$$

$$\Rightarrow \begin{bmatrix} 1 - 8 + k & 0 - 0 + 0 \\ -8 + 8 + 0 & 49 - 56 + k \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

And to satisfy the above condition of equality, the corresponding entries of the matrix o

And to satisfy the above condition of equality, the corresponding entries of the matrices should be equal

Hence,

$$1-8 + k = 0 \Rightarrow k = 7$$

Therefore, the value of k is 7

37. If
$$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$
 and $f(x) = x^2 - 2x - 3$ show that $f(A) = 0$

Solution:

Given

 $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$

To show that f(A) = 0Substitute x = A in f(x), we get $f(A) = A^2 - 2A - 3I \dots \dots (i)$ I is identity matrix, so $3I = 3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$ Now, we will find the matrix for A^2 , we get $A^2 = A \times A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ $\Rightarrow A^2 = \begin{bmatrix} 1 \times 1 + 2 \times 2 & 1 \times 2 + 2 \times 1 \\ 2 \times 1 + 1 \times 2 & 2 \times 2 + 1 \times 1 \end{bmatrix}$ $\Rightarrow A^2 = \begin{bmatrix} 1 + 4 & 2 + 2 \\ 2 + 2 & 4 + 1 \end{bmatrix}$



$$\Rightarrow A^2 = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix} \dots \dots \dots (ii)$$

Now, we will find the matrix for 2A, we get

$$2A = 2\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$

$$\Rightarrow 2A = \begin{bmatrix} 2 \times 1 & 2 \times 2 \\ 2 \times 2 & 2 \times 1 \end{bmatrix}$$

$$\Rightarrow 2A = \begin{bmatrix} 2 & 4 \\ 4 & 2 \end{bmatrix} \dots \dots \dots \dots (iii)$$

Substitute corresponding values from eqn (ii) and (iii) in eqn (i), we get

$$f(A) = A^{2} - 2A - 3I$$

$$\Rightarrow f(A) = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix} - \begin{bmatrix} 2 & 4 \\ 4 & 2 \end{bmatrix} - \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$$

$$\Rightarrow f(A) = \begin{bmatrix} 5 - 2 - 3 & 4 - 4 - 0 \\ 4 - 4 - 0 & 5 - 2 - 3 \end{bmatrix}$$

$$\Rightarrow f(A) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

So,

 $\Rightarrow f(A) = 0$ Hence Proved

38. If
$$A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$$
 and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ then find λ, μ so that $A^2 = \lambda A + \mu I$

Solution:

Given

$$A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} and I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

So

$$\mu \mathbf{I} = \mu \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \mu & 0 \\ 0 & \mu \end{bmatrix}$$

Now, we will find the matrix for A^2 , we get

$$A^{2} = A \times A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$$
$$\Rightarrow A^{2} = \begin{bmatrix} 2 \times 2 + 3 \times 1 & 2 \times 3 + 3 \times 2 \\ 1 \times 2 + 2 \times 1 & 1 \times 3 + 2 \times 2 \end{bmatrix}$$
$$\Rightarrow A^{2} = \begin{bmatrix} 7 & 12 \\ 4 & 7 \end{bmatrix} \dots \dots \dots (i)$$

Now, we will find the matrix for λ A, we get



 $\lambda A = \lambda \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ $\Rightarrow \lambda A = \begin{bmatrix} \lambda \times 2 & \lambda \times 3 \\ \lambda \times 1 & \lambda \times 2 \end{bmatrix}$ $\Rightarrow \lambda A = \begin{bmatrix} 2\lambda & 3\lambda \\ \lambda & 2\lambda \end{bmatrix} \dots \dots \dots \dots (ii)$ But given, $A^2 = \lambda A + \mu I$ Substitute corresponding values from equation (i) and (ii), we get $\Rightarrow \begin{bmatrix} 7 & 12 \\ 4 & 7 \end{bmatrix} = \begin{bmatrix} 2\lambda & 3\lambda \\ 2\lambda & 2\lambda \end{bmatrix} + \begin{bmatrix} \mu & 0 \\ 0 & \mu \end{bmatrix}$

- l4	7 J	_	Lλ	2λ]	' [O	μ
[7	12]		[2λ	+μ	3λ+	0]
⇒l4	7]	=	lλ·	+ 0	2λ +	μ

And to satisfy the above condition of equality, the corresponding entries of the matrices should be equal

Hence,
$$\lambda + 0 = 4 \Rightarrow \lambda = 4$$

And also, $2\lambda + \mu = 7$

Substituting the obtained value of λ in the above equation, we get

 $2(4) + \mu = 7 \Rightarrow 8 + \mu = 7 \Rightarrow \mu = -1$

Therefore, the value of λ and μ are 4 and – 1 respectively

39. Find the value of x for which the matrix product

2	0	7	$\left[-x\right]$	14x	7x	
0	1	0	0	1	0	equal to an identity matrix.
1	$^{-2}$	1	x	-4x	-2x	

Solution:

We know,

 $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

is identity matrix of size 3.

So according to the given criteria

 $\begin{bmatrix} 2 & 0 & 7 \\ 0 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix} \begin{bmatrix} -x & 14x & 7x \\ 0 & 1 & 0 \\ x & -4x & -2x \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Now we will multiply the two matrices on LHS using the formula $c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + ... + a_{in}b_{nj}$, we get

BYJU'S

RD Sharma Solutions for Class 12 Maths Chapter 5 Algebra of Matrices



And to satisfy the above condition of equality, the corresponding entries of the matrices should be equal

So we get

 $5x = 1 \Rightarrow x = \frac{1}{5}$ So the value of x is $\frac{1}{5}$

https://byjus.com