

EXERCISE 27.2

PAGE NO: 27.15

1. The following table shows the number of patients discharged from a hospital with HIV diagnosis in different years:

Years	2002	2003	2004	2005	2006
Number of	150	170	195	225	230
Patients					

Represent the information by a graph. Solution:

Here we know, year is an independent variable and the number of patients is a dependent variable. So let us take years on the x-axis and the number of patients on the y-axis. Let us choose the following scale:

On x-axis: 2 cm = 1 year

On y-axis: 1 cm = 10 patients

Let us assume that on the x-axis, origin (O) represents 2001 and on the y-axis, origin (O) represents 120, i.e. O (2001, 120).

Now, let us plot (2002, 150), (2003, 170), (2004, 195), (2005, 225), (2006, 230). These points are joined to get the graph representing the given information as shown in the figure below.





2. The following table shows the amount of rice grown by a farmer in different vears:

Jearse							
Years	2000	2001	2002	2003	2004	2005	2006
Rice	200	180	240	260	250	200	270
grown (in							
quintals)							

Plot a graph to illustrate this information.

Solution:

We know that, the year is an independent variable and quantity of rice grown is a dependent variable. So let us take years on the x-axis and quantity of rice grown on the y-axis.

By choosing the following scale:

On x-axis: 2 cm = 1 year

On y-axis: 1 cm = 20 quintals

Let us assume that the origin O represents the coordinates (1999, 160).

Now, let us plot (2000, 200), (2001, 180), (2002, 240), (2003, 260), (2004, 250), (2005, 200), (2006, 270).

These points are joined to get the graph representing the given information as shown in the figure below.



3. The following table gives the information regarding the number of persons employed to a piece of work and time taken to complete the work:

Number of	2	4	6	8
persons				
Time taken (in	12	6	4	3
days)				

Plot the graph of this information. Solution:

We know, the number of persons is an independent variable and time taken is a dependent variable.

So, we take the number of persons on the x-axis and time taken on the y-axis.

By choosing the following scale:

On x-axis: 2 cm = 2 persons

On y-axis: 2 cm = 2 days

Now, let us plot (2, 12), (4, 6), (6, 4), (8, 3).

These points are joined to get the graph representing the given information as shown in the figure below.





4. The following table gives the information regarding length of a side of a square and its area

Length of a side (in cm)	1	2	3	4	5
Area of	1	4	9	16	25
square (in cm ²)					

Draw a graph to illustrate this information.

Solution:

We know, length of a side is an independent variable and area of the square is a dependent variable.

So, let us take the length of a side on the x-axis and area of the square on the y-axis.

By choosing the following scale:

On x-axis: 2 cm = 1 cm

On y-axis: $1 \text{ cm} = 2 \text{ cm}^2$

Now, let us plot (1, 1), (2, 4), (3, 9), (4, 16), (5, 25).





5. The following table shows the sales of a commodity during the years 2000 to 2006.

Years	2000	2001	2002	2003	2004	2005	2006
Sales (in	1.5	1.8	2.4	3.2	5.4	7.8	8.6
lakhs of							
Rs)							

Draw a graph of this information.

Solution:

We know, year is an independent variable and sales is a dependent variable. So, we take year on the x-axis and sales on the y-axis.

By choosing the following scale:

On x-axis: 2 cm = 1 year

On y-axis: 2 cm = 1 lakh rupees

Assume that on x-axis, origin (O) represents 1991.

So, the coordinates of O are (1991, 0).

Now, let us plot (2000, 1.5), (2001, 1.8), (2002, 2.4), (2003, 3.2), (2004, 5.4), (2005, 7.8) and (2006, 8.6).



9 -			Scale : X axis : 2 cm = 1 year		
			Y axis : 2 cm = 1 lakh	G(2006, 8.6)	
8 -	8			F(2005, 7.8)	
7 -	9			/	
6-			/		
5 -			E(20	04, 5.4)	
4 -			/		
3 -			D(2003, 3.2)		
2 -			C(2002, 2.4)		
1-	A(2000, 1.	B(2001, 5)	1.8)		
				X - axis	
0	91 2000	2001 20	02 2003 2004	2005 2006	

6. Draw the temperature-time graph in each of the following cases: (i)

Time (in hours)	7:00	9:00	11:00	13:00	15:00	17:00	19:00	21:00
Temperature (in ^o F)	100	101	104	102	100	99	100	98

(ii) Time (in 8:00 10:00 12:00 14:00 16:00 18:00 20:00 hours) Temperature 100 101 104 103 99 **98** 100 (in ^oF)

Solution:

(i) We know that, time is an independent variable and temperature is a dependent variable. So, we take time on the x-axis and temperature on the y-axis.



Let us choose the following scale:

On x-axis: 1 cm = 1 hour

On y-axis: $1 \text{ cm} = 1^{\circ}\text{F}$

Let us assume that on the x-axis, the coordinate of origin is 6:00.

On y-axis, the coordinate of origin is 94°F.

So, the coordinates of 0 are (6:00, 94).

Now, let us plot (7:00, 100), (9:00, 101), (11:00, 104)...(21:00, 98)

These points are joined to get the graphs representing the given information as shown in the figures below.



(ii) We know that, time is an independent variable and temperature is a dependent variable. So, we take time on the x-axis and temperature on the y-axis.

Let us choose the following scale:

On x-axis: 2 cm = 2 hours

On y-axis: $1 \text{ cm} = 1^{\circ}\text{F}$

Let us assume that on the x-axis, the coordinate of origin is 6:00.

On y-axis, the coordinate of origin is 94°F.

So, the coordinates of 0 are (6:00, 94).

Now, let us plot (8:00, 100), (10:00, 101), (12:00, 104) (20:00, 100).

These points are joined to get the graphs representing the given information as shown in the figures below.





7. Draw the velocity-time graph from the following data:

Time (in hours)	7:00	8:00	9:00	10:00	11:00	12:00	13:00	14:00
Speed (in km/hr)	30	45	60	50	70	50	40	45

Solution:

We know that, time is an independent variable and speed is a dependent variable. So, we take time on the x-axis and speed on the y-axis.

By choosing the following scale:

On x-axis: 2 big division = 1 hour

On y-axis: 1 big division = 10 km/hr

Let us assume that on the x-axis, the coordinate of origin (O) is 7:00.

So, the coordinates of O are (7:00, 0).

Now, let us plot (7:00, 30), (8:00, 45), (9:00, 60), (10:00, 50), (11:00, 70), (12:00, 50), (13:00, 40), (14:00, 45).





8. The runs scored by a cricket team in first 15 overs are given below:

Overs:	Ι	Π	III	IV	V	VI	VII	VIII	IX	X	XI	XII	XIII	XIV	XV
Runs:	2	1	4	2	6	8	10	21	5	8	3	2	6	8	12

Draw the graph representing the above data in two different ways as a graph and as a bar chart.

Solution:

We know that, over is an independent variable and run is a dependent variable. So, we take overs on the x-axis and runs on the y-axis.

By choosing the following scale:

On x-axis: 1 cm = 1 over

On y-axis: 1 cm = 2 runs

Now, let us plot (I, 2), (II, 1), (III, 4)... (XV, 12).





The bar graph is as follows:



9. The runs scored by two teams A and B in first 10 overs are given below:

Overs	Ι	II	III	IV	V	VI	VII	VIII	IX	X
Team	2	1	8	9	4	5	6	10	6	2
Α										



Team	5	6	2	10	5	6	3	4	8	10
B										

Draw a graph depicting the data, making the graphs on the same axes in each case in two different ways as a graph and as a bar chart.

Solution:

We know that, over is an independent variable and run is a dependent variable. So, we take overs on x-axis and runs on the y-axis.

By choosing the following scale:

On x-axis: 1 cm = 1 over

On y-axis: 1 cm = 1 run

Now, let us plot (I, 2), (II, 1), (III, 8)... (X, 2) for team A and (I, 5), (II, 6), (III, 8) (X, 10) for team B.

These points are joined to get the graph representing the given information as shown in the figure below.





The bar graph is as follows:

