Date: 7<sup>th</sup> January 2020

Time: 02:30 PM - 05:30 PM

**Subject: Chemistry** 



## Answer: D

## Solution:

In aryl halides, due to the partial double bond character generated by chlorine, the aryl cation is not formed.

Vinyl halides do not give Friedel-Crafts reaction, because the intermediate that is generated (vinyl cation) is not stable.

 $H_2C = CH$ 

vinyl cation

2. In the following reaction sequence,



### The major product B is:



### Solution:

During trisubstitution, the acetanilide group attached to the benzene ring is more electron donating than the methyl group attached, owing to +M effect, and therefore, the incoming electrophile would prefer ortho w.r.t the acetanilide group.





3. For the following reactions,

$$CH_{3}CH_{2}CH_{2}Br + Z^{\ominus} - \underbrace{\begin{vmatrix} k_{s} \\ Substitution \end{vmatrix}}_{k_{e}} CH_{3}CH_{2}CH_{2}Z + Br^{\ominus}$$

$$k_{e} \\ Elimination CH_{3}CH=CH_{2} + Br^{\ominus}$$

Where,

$$\stackrel{\bigcirc}{=} Z = CH_3CH_2O \stackrel{\bigcirc}{(A)} or H_3C \stackrel{CH_3}{-C-O} (B)$$

 $k_s$  and  $k_e$ , are, respectively, the rate constants for substitution and elimination, and  $\mu = \frac{k_s}{k_e}$ , the correct option is \_\_\_\_\_.

a.  $\mu_A > \mu_B$  and  $k_e(A) > k_e(B)$ 

b.  $\mu_B > \mu_A$  and  $k_e(A) > k_e(B)$ 

c.  $\mu_A > \mu_B$  and  $k_e(B) > k_e(A)$ 

d.  $\mu_B > \mu_A$  and  $k_e(B) > k_e(A)$ 

### Answer: c

The base  $CH_3CH_2O^-$  favours substitution reaction over elimination reaction and thus  $k_e(A) < k_s(A)$ .

Since the bulkier base tertiary butoxide (B) favours elimination reaction over substitution reaction,  $k_e(B)$  is higher. Since  $\mu = \frac{k_s}{k_o}$ ,  $\mu_A$  will be greater as it is having lower  $k_e$  value.

### Solution:

- 4. Which of the following statements is correct?
  - a. Gluconic acid can form cyclic (acetal/hemiacetal) structure
  - b. Gluconic acid is a dicarboxylic acid
  - c. Gluconic acid is obtained by oxidation of glucose with HNO<sub>3</sub>
  - d. Gluconic acid is a partial oxidation product of glucose

### Answer: d

# B

## Solution:

The gluconic acid formed is a monocarboxylic acid which is formed during the partial oxidation of glucose. Glucose on reaction with  $HNO_3$  will give glucaric acid:



7th January 2020 (Shift- 2), Chemistry



6. In the following reaction sequence, structures of A and B, respectively will be:



- 7. A chromatography column, packed with silica gel as stationary phase, was used to separate a mixture of compounds consisting of (A) benzanilide, (B) aniline and (C) acetophenone. When the column is eluted with a mixture of solvents, hexane : ethyl acetate (20:80), the sequence of obtained compounds is:
  - a. (B), (A) and (C)
  - b. (C), (A) and (B)
  - c. (B), (C) and (A)
  - d. (A), (B) and (C)

## Answer: b



### Solution:

Since the column is eluted with the solvent having more proportion of ethyl acetate, the more polar compound will come out first.

The order of polarity in the given compounds is ; acetophenone > benzanilide > aniline.

(Dipolemoment ( $\mu$ ) of acetophenone=3.05 D, benzanilide=2.71 D and aniline=1.59 D).

8. The number of possible optical isomers for the complexes  $[MA_2B_2]$  with sp<sup>3</sup> or dsp<sup>2</sup> hybridized metal atom, respectively, is:

Note: A and B are unidentate neutral and unidentate monoanionic ligands, respectively.

- a. 0 and 1
- b. 0 and 0

c. 2 and 2

d. 0 and 2

### Answer: b

### Solution:

Case 1: If M is sp<sup>3</sup> hybridized, the geometry will be tetrahedral. There will be a plane of symmetry and thus it does not show optical activity.



Case 2: If M is dsp<sup>2</sup> hybridized, the geometry will be square planar. Due to the presence of a plane of symmetry, it does not show optical activity.

- 9. The bond order and magnetic characteristics of CN<sup>-</sup> are:
  - a. 3, paramagnetic
  - b. 2.5, diamagnetic

- c. 3, diamagnetic
- d. 2.5, paramagnetic

Answer: c

Solution:

 $CN^{-}$  is a 14 electron system. The bond order and magnetism can be predicted using MOT.

The MOT electronic configuration of CN<sup>-</sup> is:

 $\sigma_{1s}^2 \sigma_{1s}^{*2} \sigma_{2s}^2 \sigma_{2s}^{*2} \pi_{2p_x}^2 \pi_{2p_y}^2 \sigma_{2p_z}^2$ 

Bond order =  $\frac{1}{2} \times (N_{\text{bonding}} - N_{\text{antibonding}}) = 3$ 

As CN<sup>-</sup> does not have any unpaired electrons, and hence it is diamagnetic.

### 10. The equation that is incorrect is:

- a.  $\Lambda_m^0 \text{NaBr} \Lambda_m^0 \text{NaI} = \Lambda_m^0 \text{KBr} \Lambda_m^0 \text{NaBr}$ b.  $\Lambda_m^0 \text{NaBr} \Lambda_m^0 \text{NaCl} = \Lambda_m^0 \text{KBr} \Lambda_m^0 \text{KCl}$ c.  $\Lambda_m^0 \text{KCl} \Lambda_m^0 \text{NaCl} = \Lambda_m^0 \text{KBr} \Lambda_m^0 \text{NaBr}$ d.  $\Lambda_m^0 \text{H}_2 \text{O} = \Lambda_m^0 \text{HCl} + \Lambda_m^0 \text{NaOH} \Lambda_m^0 \text{NaBr}$

### Answer: a

### Solution:

 $\Lambda_m^0 \text{NaI} - \Lambda_m^0 \text{NaBr} = \Lambda_m^0 \text{NaBr} - \Lambda_m^0 \text{KBr}$ 

$$\begin{split} [\lambda_m^0 Na^+ + \lambda_m^0 I^-] - [\lambda_m^0 Na^+ + \lambda_m^0 Br^-] &= [\lambda_m^0 Na^+ + \lambda_m^0 Br^-] - [\lambda_m^0 K^+ + \lambda_m^0 Br^-] \\ \lambda_m^0 I^- - \lambda_m^0 Br^- \neq \lambda_m^0 Na^+ - \lambda_m^0 K^+ \end{split}$$

11. In the following reactions, product (A) and (B), respectively, are:

 $NaOH + Cl_2 \rightarrow (A) + side products$ 

(hot & conc.)

 $Ca(OH)_2 + Cl_2 \rightarrow (B) + side products$ (dry)

- a. NaClO<sub>3</sub> and Ca(ClO<sub>3</sub>)<sub>2</sub>
- b. NaOCl and  $Ca(ClO_3)_2$
- c. NaOCl and Ca(OCl)<sub>2</sub>
- d. NaClO<sub>3</sub> and Ca(OCl)<sub>2</sub>

#### Answer: d

### Solution:

 $6NaOH + 3Cl_2 \rightarrow 5NaCl + NaClO_3 + 3H_2O$  $2Ca(OH)_2 + Cl_2 \rightarrow Ca(OCl)_2 + CaCl_2 + H_2O$ 

12. Two open beakers one containing a solvent and the other containing a mixture of that solvent

- with a non-volatile solute are together sealed in a container. Over time:
- a. the volume of the solution and the solvent does not change
- b. the volume of the solution increases and the volume of the solvent decreases
- c. the volume of the solution decreases and the volume of the solvent increases
- d. the volume of the solution does not change and the volume of the solvent decreases

#### Answer: b



### Solution:

Consider beaker I contains the solvent and beaker 2 contains the solution. Let the vapour pressure of the beaker I be  $P^o$  and the vapour pressure of beaker II be  $P^s$ . According to Raoult's law, the vapour pressure of the solvent ( $P^o$ ) is greater than the vapour pressure of the solution ( $P^s$ )

 $(P_0 > P_s)$ 

Due to a higher vapour pressure, the solvent flows into the solution. So volume of beaker II would increase.

In a closed beaker, both the liquids on attaining equilibrium with the vapour phase will end up having the same vapour pressure. Beaker II attains equilibrium at a lower vapour pressure and so in its case, condensation will occur so as to negate the increased vapour pressure from beaker I, which results in an increase in its volume.

On the contrary, since particles are condensing from the vapour phase in beaker II, the vapour pressure will decrease. Since beaker I at equilibrium attains a higher vapour pressure evaporation will be favoured more so as to compensate for the decreased vapour pressure, as mentioned in the previous statement.

- 13. The refining method used when the metal and the impurities have low and high melting
  - temperatures, respectively, is: a. vapour phase refining

- b. distillation
- d. zone refining

### Answer: c

c. liquation

### Solution:

Liquation is the process of refining a metal with a low melting point containing impurities of high melting point

- 14. Among statements I-IV, the correct ones are:
  - I. Decomposition of hydrogen peroxide gives dioxygen
  - II. Like hydrogen peroxide, compounds, such as  $KClO_3$ ,  $Pb(NO_3)_2$  and  $NaNO_3$  when heated liberate dioxygen.
  - III. 2-Ethylanthraquinone is useful for the industrial preparation of hydrogen peroxide.
  - IV. Hydrogen peroxide is used for the manufacture of sodium perborate
  - a. I,II, III and IV
  - c. I, III and IV only

- b. I, II and III only
- d. I and III only

AllSwella	Answer:	а
-----------	---------	---

### Solution:

Decomposition of  $H_2O_2 : 2H_2O_2(l) \rightarrow O_2(g) + 2H_2O(l)$ 

Industrially,  $H_2O_2$  is prepared by the auto-oxidation of 2-alklylanthraquinols. 2KClO<sub>3</sub>  $\xrightarrow{150-300^{\circ}C}$  2KCl + 3O<sub>2</sub>

 $2Pb(NO_3)_2 \xrightarrow{200-470^{\circ}C} 2PbO + 4NO_2 + O_2$ 

 $2NaNO_3 \rightarrow 2NaNO_2 + O_2$ 

Synthesis of sodium perborate:

 $\mathrm{Na_2B_4O_7} + 2\mathrm{NaOH} + 4\mathrm{H_2O_2} \rightarrow 2\mathrm{NaBO_3} + 5\mathrm{H_2O}$ 

### 15. The redox reaction among the following is:

- a. formation of ozone from atmospheric oxygen in the presence of sunlight
- b. reaction of H<sub>2</sub>SO<sub>4</sub> with NaOH
- c. combination of dinitrogen with dioxygen at 2000 K
- d. Reaction of  $[Co(H_2O)_6]Cl_3$  with AgNO<sub>3</sub>

#### Answer: c

#### Solution:

 $N_2 + O_2 \xrightarrow{2000 \text{ K}} 2\text{NO}$ : The oxidation state of N changes from 0 to +2, and the oxidation state of O changes from 0 to -2

In all the remaining reactions, there is no change in oxidation states of the elements participating in the reaction.

 $30_2 \rightarrow 20_3$ 

 $2NaOH + H_2SO_4 \rightarrow Na_2SO_4 + 2H_2O$  (Neutralisation reaction)

 $3 \text{ AgNO}_3 + [\text{Co}(\text{H}_2\text{O})_6]\text{Cl}_3 \rightarrow [\text{Co}(\text{H}_2\text{O})_6](\text{NO}_3)_3 + 3 \text{ AgCl}$  (Double displacement)



16. Identify the correct labels of A, B and C in the following graph from the options given below:



Root mean square speed ( $V_{rms}$ ); most probable speed ( $V_{mp}$ ); average speed ( $V_{av}$ )

a.  $A = V_{mp}$ ,  $B = V_{av}$ ,  $C = V_{rms}$ 

c. 
$$A = V_{av}, B = V_{rms}, C = V_{mp}$$

# Answer: a

### Solution:

 $C_{RMS} = \sqrt{\frac{3RT}{M}}$  $C_{Average} = \sqrt{\frac{8RT}{\pi M}}$ 

$$C_{MPS} = \sqrt{\frac{2RT}{M}}$$

$$\sqrt{3} > \sqrt{\frac{8}{\pi}} > \sqrt{2}$$

 $C_{RMS} > C_{Average} > C_{MP}$ 

17. For the reaction,

 $2H_2(g) + 2NO(g) \rightarrow N_2(g) + 2H_2O(g)$ The observed rate expression is, rate =  $k_f[NO]^2[H_2]$ . The rate expression for the reverse reaction is:

a. 
$$k_b [N_2] [H_2 O]^2$$

b. 
$$k_b[N_2][H_2O]$$

c. 
$$k_b[N_2][H_2O]^2/[H_2]$$

d.  $k_b[N_2][H_2O]^2/[NO]$ 

# Answer: c



b. 
$$A = V_{mp}, B = V_{rms}, C = V_{av}$$

I. 
$$A = V_{rms}$$
,  $B = V_{mp}$ ,  $C = V_{av}$ 

### Solution:

The reverse reaction is:

 $2H_2(g) + 2NO(g) \rightleftharpoons N_2(g) + 2H_2O(g)$ 

The equilibrium constant,  $K_c = \frac{k_f}{k_b} = \frac{[N_2][H_2O]^2}{[H_2]^2[NO]^2}$ 

At equilibrium, rate of forward reaction=rate of backward reaction

i.e. ,  $k_{\rm f}[\rm NO]^2[\rm H_2] = k_b \frac{[N_2][H_2O]^2}{[H_2]}$ 

Hence, rate of reverse reaction= $k_b \frac{[N_2][H_2O]^2}{[H_2]}$ 

18. Within each pair of elements F & Cl, S & Se and Li & Na, respectively, the elements that release more energy upon an electron gain are:

a. Cl, Se and Na

c. F, S and Li

b. Cl, S and Li

d. F, Se and Na

c. A and D only d. C and D only

### Answer: b

#### Solution:

Element	First Ele	ectron gain enthalpy(kJ/mol)
Li	-60	
Na	-53	
F	-320	
S	-200	
Cl	-340	
Se	-195	

Despite F being more electronegative than Cl, due to the small size of F, Cl would have a more negative value of electron gain enthalpy because of inter-electronic repulsions.

As we go down the group, the negative electron gain enthalpy decreases.

19. Among the following statements A-D, the incorrect ones are:

A. Octahedral Co(III) complexes with strong field ligands have high magnetic moments

- B. When  $\Delta_0$  < P, the d- electron configuration of Co(III) in an octahedral complex is  $t_{2g}^4$ ,  $e_g^2$ .
- C. Wavelength of light absorbed by  $[Co(en)_3]^{3+}$  is lower than that of  $[CoF_6]^{3-}$ .
- D. If the  $\Delta_0$  for an octahedral complex of Co(III) is 18000 cm<sup>-1</sup>, the  $\Delta_t$  for its tetrahedral complex with the same ligand will be16000 cm<sup>-1</sup>.
- a. B and C only b. A and B only

Answer: c



### Solution:

 $\text{Co}^{3+}$  has d<sup>6</sup> electronic configuration. In the presence of strong field ligand,  $\Delta_o > P$ . Thus the splitting occurs as:  $t_{2g}^6$ ,  $e_g^0$ ; so the magnetic moment is zero.

According to the spectrochemical series, en is a stronger ligand than F and therefore promotes pairing. This implies that the  $\Delta_0$  of en is more than the  $\Delta_0$  of F.

$$\Delta_{o} = \frac{hc}{\lambda_{abs}}$$
$$\Delta_{t} = \frac{4}{9}\Delta_{o} = 8000 \text{ cm}^{-1}$$

20. The ammonia  $(NH_3)$  released on quantitative reaction of 0.6 g urea  $(NH_2CONH_2)$  with sodium hydroxide (NaOH) can be neutralized by:

a. 200 mL of 0.2 N HCl

b. 200 mL of 0.4 N HCl

#### Answer: d

### Solution:

Moles of urea=  $\left(\frac{0.6}{60}\right)$  = 0.01

 $NH_2CONH_2 + 2NaOH \rightarrow Na_2CO_3 + 2NH_3$ 

0.01

0.02

c. 100 mL of 0.1 N HCl

d. 100 mL of 0.2 N HCl

```
0.02 moles of NH<sub>3</sub> reacts with 0.02 moles of HCl.
```

Moles of HCl in option a=  $0.2 \times \frac{100}{1000} = 0.02$ 

21. Number of sp<sup>2</sup> hybrid carbon atoms present in aspartame is \_\_\_\_.

Answer: 9

Solution:



The marked carbons are sp<sup>2</sup> hybridised.





22. 3 grams of acetic acid is added to 250 mL of 0.1 M HCl and the solution is made up to 500 mL. To 20 mL of this solution  $\frac{1}{2}$  mL of 5 M NaOH is added. The pH of this solution is\_\_\_\_\_.

(Given:  $\log 3 = 0.4771$ , pK<sub>a</sub> of acetic acid = 4.74, molar mass of acetic acid = 60 g/mole).

### **Answer: 5.22**

### Solution:

-

mmole of acetic acid in 20 mL = 2mmole of HCl in 20 mL = 1mmole of NaOH = 2.5

1

1

 $HCl + NaOH \rightarrow NaCl + H_2O$ 

2.5 1 1.5

 $CH_3COOH + NaOH (remaining) \longrightarrow CH_3COONa + water$ 

2 1.5 0.5 0

1.5

 $pH = pK_a + \log \frac{1.5}{0.5} = 4.74 + \log 3 = 4.74 + 0.48 = 5.22$ 

23. The flocculation value of HCl for As<sub>2</sub>S<sub>3</sub> sol is 30 mmolL<sup>-1</sup>. If H<sub>2</sub>SO<sub>4</sub> is used for the flocculation of arsenic sulphide, the amount, in grams, of H<sub>2</sub>SO<sub>4</sub> in 250 mL required for the above purpose is

### Answer: 0.3675 g

### Solution:

For 1L sol 30 mmol of HCl is required For 1L sol 15 mmol of H<sub>2</sub>SO<sub>4</sub> is required For 250 mL of sol,  $\frac{15}{4} \times 98 \times 10^{-3}$  g of H<sub>2</sub>SO<sub>4</sub>= 0.3675 g

24. Consider the following reactions :

 $NaCl + K_2Cr_2O_7 + H_2SO_4 \rightarrow (A) + side products$ 

(A) + NaOH  $\rightarrow$  (B) + side products

 $(B) + H_2SO_4(dil.) + H_2O_2 \rightarrow (C) + side products$ 

The sum of the total number of atoms in one molecule of (A), (B) & (C) is \_\_\_\_\_.

### Answer: 18



Solution:

 $\begin{aligned} &\text{NaCl} \xrightarrow{\text{K}_2\text{Cr}_2\text{O}_7/\text{Conc.H}_2\text{SO}_4} &\text{CrO}_2\text{Cl}_2 \xrightarrow{\text{NaOH}} \text{Na}_2\text{CrO}_4 + \text{NaCl} \\ &\text{Na}_2\text{CrO}_4 \xrightarrow{\text{Dil.H}_2\text{SO}_4} &\text{Na}_2\text{Cr}_2\text{O}_7 \xrightarrow{\text{Dil.H}_2\text{O}_2} &\text{CrO}_5 \\ &\text{(A)} &= \text{CrO}_2\text{Cl}_2, (B) = \text{Na}_2\text{CrO}_4 \text{ and } (C) = \text{CrO}_5 \end{aligned}$ 

25. The standard heat of formation ( $\Delta_{\rm f}H_{298}^{\circ}$ ) of ethane (in kJ/mol), if the heat of combustion of ethane, hydrogen and graphite are -1560, -393.5 and -286 kJ/mol, respectively, is\_\_\_\_\_.

#### Answer: -192.5 kJ/mol

#### Solution:

 $C_{(\text{graphite})} + O_2(g) \rightarrow CO_2(g) \ \Delta H_c^{\circ} = -286 \text{ kJ/mol.....(1)}$   $H_2(g) + 0.5O_2(g) \rightarrow H_2O(l) \ \Delta H_c^{\circ} = -393.5 \text{ kJ/mol.....(2)}$   $C_2H_6(g) + 3.5O_2(g) \rightarrow 2CO_2(g) + 3H_2O(l) \ \Delta H_c^{\circ} = -1560 \text{ kJ/mol....(3)}$   $2C_{(\text{graphite})} + 3H_2(g) \rightarrow C_2H_6(g); \ \Delta H_f^{\circ} = ?$ 

By inverting (3) and multiplying (1) by 2 and (2) by 3 and adding, we get,

2 × (-286) + 3 × (-393.5) - (-1560) = -192.5 kJ/mol