

## EXERCISE 7.1

PAGE NO: 7.22

1. Find the adjoint of each of the following matrices:

(i)  $\begin{bmatrix} -3 & 5 \\ 2 & 4 \end{bmatrix}$

(ii)  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$

(iii)  $\begin{bmatrix} \cos\alpha & \sin\alpha \\ \sin\alpha & \cos\alpha \end{bmatrix}$

(iv)  $\begin{bmatrix} 1 & \tan\frac{\alpha}{2} \\ -\tan\frac{\alpha}{2} & 1 \end{bmatrix}$

 Verify that  $(\text{adj } A) A = |A| I = A (\text{adj } A)$  for the above matrices.

**Solution:**

(i) Let

$$A = \begin{bmatrix} -3 & 5 \\ 2 & 4 \end{bmatrix}$$

Cofactors of A are

$$C_{11} = 4$$

$$C_{12} = -2$$

$$C_{21} = -5$$

$$C_{22} = -3$$

$$\text{Since, adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$$

$$(\text{adj } A) = \begin{bmatrix} 4 & -2 \\ -5 & -3 \end{bmatrix}^T$$

$$= \begin{bmatrix} 4 & -5 \\ -2 & -3 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) A = \begin{bmatrix} 4 & -5 \\ -2 & -3 \end{bmatrix} \begin{bmatrix} -3 & 5 \\ 2 & 4 \end{bmatrix} = \begin{bmatrix} -12 - 10 & 20 - 20 \\ 6 - 6 & -10 - 12 \end{bmatrix}$$

$$(\text{adj } A) A = \begin{bmatrix} -22 & 0 \\ 0 & -22 \end{bmatrix}$$

$$\text{And, } |A| I = \begin{vmatrix} -3 & 5 \\ 2 & 4 \end{vmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = (-22) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} -22 & 0 \\ 0 & -22 \end{bmatrix}$$

$$\text{Also, } A (\text{adj } A) = \begin{bmatrix} -3 & 5 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -5 \\ -2 & -3 \end{bmatrix} = \begin{bmatrix} -12 - 10 & 20 - 20 \\ 6 - 6 & -10 - 12 \end{bmatrix}$$

$$A (\text{adj } A) = \begin{bmatrix} -22 & 0 \\ 0 & -22 \end{bmatrix}$$

Hence,  $(\text{adj } A) A = |A| I = A (\text{adj } A)$

(ii) Let

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Therefore cofactors of A are

$$C_{11} = d$$

$$C_{12} = -c$$

$$C_{21} = -b$$

$$C_{22} = a$$

$$\text{We know that, } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$$

Therefore by substituting these values we get,

$$(\text{adj } A) = \begin{bmatrix} d & -c \\ -b & a \end{bmatrix}^T$$

$$= \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) A = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} ad - bc & bd - bd \\ -ac + ac & -bc + ad \end{bmatrix}$$

$$(\text{adj } A) A = \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix}$$

$$\text{And, } |A| \cdot I = \begin{vmatrix} a & b \\ c & d \end{vmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = (ad - bc) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix}$$

Also,

$$A (\text{adj } A) = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix}$$

Hence,  $(\text{adj } A) A = |A| I = A (\text{adj } A)$

(iii) Let

$$A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

Therefore cofactors of A are

$$C_{11} = \cos \alpha$$

$$C_{12} = -\sin \alpha$$

$$C_{21} = -\sin \alpha$$

$$C_{22} = \cos \alpha$$

$$\text{We know that, } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$$

$$\begin{aligned} (\text{adj } A) &= \begin{bmatrix} \cos \alpha & -\sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}^T \\ &= \begin{bmatrix} \cos \alpha & -\sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix} \end{aligned}$$

$$\text{Now, } (\text{adj } A) A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

$$= \begin{bmatrix} -\sin^2 \alpha + \cos^2 \alpha & \cos \alpha \sin \alpha - \sin \alpha \cos \alpha \\ -\cos \alpha \sin \alpha + \sin \alpha \cos \alpha & -\sin^2 \alpha + \cos^2 \alpha \end{bmatrix}$$

$$(\text{adj } A) A = \begin{bmatrix} \cos 2\alpha & 0 \\ 0 & \cos 2\alpha \end{bmatrix}$$

$$\text{And, } |A| = \begin{vmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & \cos \alpha \end{vmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= (\cos^2 \alpha - \sin^2 \alpha) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \alpha - \sin^2 \alpha & 0 \\ 0 & \cos^2 \alpha - \sin^2 \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos 2\alpha & 0 \\ 0 & \cos 2\alpha \end{bmatrix}$$

Also,  $A (\text{adj } A)$

$$= \begin{bmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \alpha & -\sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix} = \begin{bmatrix} \cos^2 \alpha - \sin^2 \alpha & 0 \\ 0 & \cos^2 \alpha - \sin^2 \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos 2\alpha & 0 \\ 0 & \cos 2\alpha \end{bmatrix}$$

Hence,  $(\text{adj } A) A = |A| I = A (\text{adj } A)$

(iv) Let

$$A = \begin{bmatrix} 1 & \tan \frac{\alpha}{2} \\ -\tan \frac{\alpha}{2} & 1 \end{bmatrix}$$

Therefore cofactors of A are

$$C_{11} = 1$$

$$C_{12} = \tan \frac{\alpha}{2}$$

$$C_{21} = -\tan \frac{\alpha}{2}$$

$$C_{22} = 1$$

We know that,  $\text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$

$$(\text{adj } A) = \begin{bmatrix} 1 & \tan \frac{\alpha}{2} \\ -\tan \frac{\alpha}{2} & 1 \end{bmatrix}^T$$

$$= \begin{bmatrix} 1 & -\tan \frac{\alpha}{2} \\ \tan \frac{\alpha}{2} & 1 \end{bmatrix}$$

Now,  $(\text{adj } A) A = \begin{bmatrix} 1 & -\tan \frac{\alpha}{2} \\ \tan \frac{\alpha}{2} & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan \frac{\alpha}{2} \\ -\tan \frac{\alpha}{2} & 1 \end{bmatrix}$

$$= \begin{bmatrix} 1 + \tan^2 \frac{\alpha}{2} & \tan \frac{\alpha}{2} - \tan \frac{\alpha}{2} \\ \tan \frac{\alpha}{2} - \tan \frac{\alpha}{2} & 1 + \tan^2 \frac{\alpha}{2} \end{bmatrix}$$

$$(\text{adj } A)A = \begin{bmatrix} 1 + \tan^2 \frac{\alpha}{2} & 0 \\ 0 & 1 + \tan^2 \frac{\alpha}{2} \end{bmatrix}$$

$$\text{And, } |A| \cdot I = \begin{vmatrix} 1 & \tan \frac{\alpha}{2} \\ -\tan \frac{\alpha}{2} & 1 \end{vmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = (1 + \tan^2 \frac{\alpha}{2}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 + \tan^2 \frac{\alpha}{2} & 0 \\ 0 & 1 + \tan^2 \frac{\alpha}{2} \end{bmatrix}$$

$$\text{Also, } A (\text{adj } A) = \begin{bmatrix} 1 & \tan \frac{\alpha}{2} \\ -\tan \frac{\alpha}{2} & 1 \end{bmatrix} \begin{bmatrix} 1 & -\tan \frac{\alpha}{2} \\ \tan \frac{\alpha}{2} & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 + \tan^2 \frac{\alpha}{2} & \tan \frac{\alpha}{2} - \tan \frac{\alpha}{2} \\ \tan \frac{\alpha}{2} - \tan \frac{\alpha}{2} & 1 + \tan^2 \frac{\alpha}{2} \end{bmatrix}$$

$$= \begin{bmatrix} 1 + \tan^2 \frac{\alpha}{2} & 0 \\ 0 & 1 + \tan^2 \frac{\alpha}{2} \end{bmatrix}$$

Hence,  $(\text{adj } A) A = |A| I = A (\text{adj } A)$

**2. Compute the adjoint of each of the following matrices.**

$$(i) \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$(ii) \begin{bmatrix} 1 & 2 & 5 \\ 2 & 3 & 1 \\ -1 & 1 & 1 \end{bmatrix}$$

$$(iii) \begin{bmatrix} 2 & -1 & 3 \\ 4 & 2 & 5 \\ 0 & 4 & -1 \end{bmatrix}$$

$$(iv) \begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 1 & 1 & 3 \end{bmatrix}$$

**Solution:**

(i) Let

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

Therefore cofactors of A are

$$C_{11} = -3$$

$$C_{21} = 2$$

$$C_{31} = 2$$

$$C_{12} = 2$$

$$C_{22} = -3$$

$$C_{23} = 2$$

$$C_{13} = 2$$

$$C_{23} = 2$$

$$C_{33} = -3$$

$$\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) A = \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} -3 + 4 + 4 & -6 + 2 + 4 & -6 + 4 + 2 \\ 2 - 3 + 4 & 4 - 3 + 4 & 4 - 6 + 2 \\ 2 + 4 - 6 & 4 + 2 - 6 & 4 + 4 - 3 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

$$\text{Also, } |A| = \begin{vmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{vmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = (-3 + 4 + 4) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

$$\text{Then, } A (\text{adj } A) = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

$$= \begin{bmatrix} -3 + 4 + 4 & -6 + 2 + 4 & -6 + 4 + 2 \\ 2 - 3 + 4 & 4 - 3 + 4 & 4 - 6 + 2 \\ 2 + 4 - 6 & 4 + 2 - 6 & 4 + 4 - 3 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

Since,  $(\text{adj } A) A = |A| I = A (\text{adj } A)$

(ii) Let

$$A = \begin{bmatrix} 1 & 2 & 5 \\ 2 & 3 & 1 \\ -1 & 1 & 1 \end{bmatrix}$$

Cofactors of A

$$C_{11} = 2$$

$$C_{21} = 3$$

$$C_{31} = -13$$

$$C_{12} = -3$$

$$C_{22} = 6$$

$$C_{32} = 9$$

$$C_{13} = 5$$

$$C_{23} = -3$$

$$C_{33} = -1$$

$$\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} 2 & -3 & 5 \\ 3 & 6 & -3 \\ -13 & 9 & -1 \end{bmatrix}^T$$

$$\text{adj } A = \begin{bmatrix} 2 & 3 & -13 \\ -3 & 6 & 9 \\ 5 & -3 & -1 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) A = \begin{bmatrix} 2 & 3 & -13 \\ -3 & 6 & 9 \\ 5 & -3 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 5 \\ 2 & 3 & 1 \\ -1 & 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 2 + 6 + 13 & 4 + 9 - 13 & 10 + 3 - 13 \\ -3 + 12 - 9 & -6 + 18 + 9 & -15 + 6 + 9 \\ 5 - 6 + 1 & 10 - 9 - 1 & 25 - 3 - 1 \end{bmatrix}$$

$$= \begin{bmatrix} 21 & 0 & 0 \\ 0 & 21 & 0 \\ 0 & 0 & 21 \end{bmatrix}$$

$$\text{Also, } |A| = \begin{vmatrix} 1 & 2 & 5 \\ 2 & 3 & 1 \\ -1 & 1 & 1 \end{vmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= [1(3 - 1) - 2(2 + 1) + 5(2 + 3)] \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= 21 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 21 & 0 & 0 \\ 0 & 21 & 0 \\ 0 & 0 & 21 \end{bmatrix}$$



$$\begin{aligned} \text{Then, } A(\text{adj } A) &= \begin{bmatrix} 1 & 2 & 5 \\ 2 & 3 & 1 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 3 & -13 \\ -3 & 6 & 9 \\ 5 & -3 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 2 - 6 + 25 & 3 + 12 - 15 & -13 + 18 - 5 \\ 4 - 9 + 5 & 6 + 18 - 3 & -26 + 27 - 1 \\ -2 - 3 + 5 & -3 + 6 - 3 & 13 + 9 - 1 \end{bmatrix} \\ &= \begin{bmatrix} 21 & 0 & 0 \\ 0 & 21 & 0 \\ 0 & 0 & 21 \end{bmatrix} \end{aligned}$$

Hence,  $(\text{adj } A)A = |A|I = A(\text{adj } A)$

(iii) Let

$$A = \begin{bmatrix} 2 & -1 & 3 \\ 4 & 2 & 5 \\ 0 & 4 & -1 \end{bmatrix}$$

Therefore cofactors of A

$$C_{11} = -22$$

$$C_{21} = 11$$

$$C_{31} = -11$$

$$C_{12} = 4$$

$$C_{22} = -2$$

$$C_{32} = 2$$

$$C_{13} = 16$$

$$C_{23} = -8$$

$$C_{33} = 8$$

$$\text{We know that } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

Now by substituting the values in above matrix we get,

$$= \begin{bmatrix} -22 & 4 & 16 \\ 11 & -2 & -8 \\ -11 & 2 & 8 \end{bmatrix}^T$$

$$\text{adj } A = \begin{bmatrix} -22 & 11 & -11 \\ 4 & -2 & 2 \\ 16 & -8 & 8 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) A = \begin{bmatrix} -22 & 11 & -11 \\ 4 & -2 & 2 \\ 16 & -8 & 8 \end{bmatrix} \begin{bmatrix} 2 & -1 & 3 \\ 4 & 2 & 5 \\ 0 & 4 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} -44 + 44 + 0 & 22 + 22 - 44 & -66 + 55 + 11 \\ 8 - 8 + 0 & -4 - 4 + 8 & 12 - 10 - 2 \\ 32 - 32 + 0 & -16 - 16 + 32 & 48 - 40 - -8 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Now, } |A| = \begin{vmatrix} 2 & -1 & 3 \\ 4 & 2 & 5 \\ 0 & 4 & -1 \end{vmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= [2(-2 - 20) + 1(-4 - 0) + 3(16 - 0)] \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= (-44 - 4 + 48) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Then, } A (\text{adj } A) = \begin{bmatrix} 2 & -1 & 3 \\ 4 & 2 & 5 \\ 0 & 4 & -1 \end{bmatrix} \begin{bmatrix} -22 & 11 & -11 \\ 4 & -2 & 2 \\ 16 & -8 & 8 \end{bmatrix}$$

$$= \begin{bmatrix} -44 - 4 + 48 & 22 + 2 - 24 & -22 - 2 + 24 \\ -88 + 8 + 80 & 44 - 4 - 40 & -44 + 4 + 40 \\ 0 + 16 - 16 & 0 - 8 + 8 & 0 + 8 - 8 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Hence,  $(\text{adj } A) A = |A| I = A (\text{adj } A)$

(iv) Let

$$A = \begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 1 & 1 & 3 \end{bmatrix}$$

Therefore cofactors of A

$$C_{11} = 3$$

$$C_{21} = -1$$

$$C_{31} = -1$$

$$C_{12} = -15$$

$$C_{22} = 7$$

$$C_{32} = -5$$

$$C_{13} = 4$$

$$C_{23} = -2$$

$$C_{33} = 2$$

$$\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} 3 & -15 & 4 \\ -3 & 7 & -2 \\ 1 & -5 & 2 \end{bmatrix}^T$$

$$\text{adj } A = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 7 & -5 \\ 4 & -2 & 2 \end{bmatrix}$$

$$\text{Now, } (\text{adj } A) A = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 7 & -5 \\ 4 & -2 & 2 \end{bmatrix} \begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 1 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 6 - 5 + 1 & 0 - 1 + 1 & -3 + 0 + 3 \\ -30 + 35 - 5 & 0 + 7 - 5 & 15 - 0 - 15 \\ 8 - 10 + 2 & 0 - 2 + 2 & -4 - 0 + 6 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

$$\text{Also, } |A| = \begin{vmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 1 & 1 & 3 \end{vmatrix} \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$= [2(3 - 0) + 0(15 - 0) - 1(5 - 1)] \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$= (6 - 4) \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$= \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

$$\text{Then, } A(\text{adj } A) = \begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 3 & -1 & 1 \\ -15 & 7 & -5 \\ 4 & -2 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 6 + 0 - 4 & -2 + 0 + 2 & 2 - 0 - 2 \\ 15 - 15 + 0 & -5 + 7 + 0 & 5 - 5 + 0 \\ 3 - 15 + 12 & -1 + 7 - 6 & 1 - 5 + 6 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

Hence,  $(\text{adj } A)A = |A|I = A(\text{adj } A)$

3. For the matrix  $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 3 & 0 \\ 18 & 2 & 10 \end{bmatrix}$ , show that  $A(\text{adj } A) = 0$

**Solution:**

Given

$$A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 3 & 0 \\ 18 & 2 & 10 \end{bmatrix}$$

Therefore cofactors of A

$$C_{11} = 30$$

$$C_{21} = 12$$

$$C_{31} = -3$$

$$C_{12} = -20$$

$$C_{22} = -8$$

$$C_{32} = 2$$

$$C_{13} = -50$$

$$C_{23} = -20$$

We know that  $\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

By substituting these values in above matrix we get,

$$= \begin{bmatrix} 30 & -20 & -50 \\ 12 & -8 & -20 \\ -3 & 2 & 5 \end{bmatrix}^T$$

$$\text{So, adj } (A) = \begin{bmatrix} 30 & 12 & -3 \\ -20 & -8 & 2 \\ -50 & -20 & 5 \end{bmatrix}$$

$$\text{Now, } A (\text{adj } A) = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 3 & 0 \\ 18 & 2 & 10 \end{bmatrix} \begin{bmatrix} 30 & 12 & -3 \\ -20 & -8 & 2 \\ -50 & -20 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 30 + 20 - 50 & 12 + 8 - 20 & -3 - 2 + 5 \\ 60 - 60 + 0 & 24 - 24 + 0 & -6 + 6 + 0 \\ 540 - 40 - 500 & 216 - 16 - 200 & -54 + 4 + 50 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Hence,  $A (\text{adj } A) = 0$

4. If  $A = \begin{bmatrix} -4 & -3 & -3 \\ 1 & 0 & 1 \\ 4 & 4 & 3 \end{bmatrix}$ , show that  $\text{adj } A = A$

**Solution:**

Given

$$A = \begin{bmatrix} -4 & -3 & -3 \\ 1 & 0 & 1 \\ 4 & 4 & 3 \end{bmatrix}$$

Cofactors of A

$$C_{11} = -4$$

$$C_{21} = -3$$

$$C_{31} = -3$$

$$C_{12} = 1$$

$$C_{22} = 0$$

$$C_{32} = 1$$

$$C_{13} = 4$$

$$C_{23} = 4$$

$$C_{33} = 3$$

We know that  $\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

$$= \begin{bmatrix} -4 & 1 & 4 \\ -3 & 0 & 4 \\ -3 & 1 & 3 \end{bmatrix}^T$$

$$\text{So, adj } A = \begin{bmatrix} -4 & -3 & -3 \\ 1 & 0 & 1 \\ 4 & 4 & 3 \end{bmatrix}$$

Hence,  $\text{adj } A = A$

5. If  $A = \begin{bmatrix} -1 & -2 & -2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix}$ , show that  $\text{adj} A = 3A^T$ .

**Solution:**

Given

$$A = \begin{bmatrix} -1 & -2 & -2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix}$$

Cofactors of A are

$$C_{11} = -3$$

$$C_{21} = 6$$

$$C_{31} = 6$$

$$C_{12} = -6$$

$$C_{22} = 3$$

$$C_{32} = -6$$

$$C_{13} = -6$$

$$C_{23} = -6$$

$$C_{33} = 3$$

$$\text{adj} A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} -3 & -6 & -6 \\ 6 & 3 & -6 \\ 6 & -6 & 3 \end{bmatrix}^T$$

$$\text{So, adj} A = \begin{bmatrix} -3 & 6 & 6 \\ -6 & 3 & -6 \\ -6 & -6 & 3 \end{bmatrix}$$

$$\text{Now, } 3A^T = 3 \begin{bmatrix} -1 & -2 & -2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix} = \begin{bmatrix} -3 & 6 & 6 \\ -6 & 3 & -6 \\ -6 & -6 & 3 \end{bmatrix}$$

$$\text{Hence, adj} A = 3.A^T$$

6. Find  $A(\text{adj}A)$  for the matrix  $A = \begin{bmatrix} 1 & -2 & 3 \\ 0 & 2 & -1 \\ -4 & 5 & 2 \end{bmatrix}$

**Solution:**

Given

$$A = \begin{bmatrix} 1 & -2 & 3 \\ 0 & 2 & -1 \\ -4 & 5 & 2 \end{bmatrix}$$

Cofactors of A are

$$C_{11} = 9$$

$$C_{21} = 19$$

$$C_{31} = -4$$

$$C_{12} = 4$$

$$C_{22} = 14$$

$$C_{32} = 1$$

$$C_{13} = 8$$

$$C_{23} = 3$$

$$C_{33} = 2$$

We know that  $\text{adj} A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

$$= \begin{bmatrix} 9 & 4 & 8 \\ 19 & 14 & 3 \\ -4 & 1 & 2 \end{bmatrix}^T$$

So,  $\text{adj} A = \begin{bmatrix} 9 & 19 & -4 \\ 4 & 14 & 1 \\ 8 & 3 & 2 \end{bmatrix}$

Now,  $A \text{adj} A = \begin{bmatrix} 1 & -2 & 3 \\ 0 & 2 & -1 \\ -4 & 5 & 2 \end{bmatrix} \begin{bmatrix} 9 & 19 & -4 \\ 4 & 14 & 1 \\ 8 & 3 & 2 \end{bmatrix}$

$$= \begin{bmatrix} 9 - 8 + 24 & 19 - 28 + 9 & -4 - 2 + 6 \\ 0 + 8 - 8 & 0 + 28 - 3 & 0 + 2 - 2 \\ -36 + 20 + 16 & -76 + 70 + 6 & 16 + 5 + 4 \end{bmatrix}$$



$$= \begin{bmatrix} 25 & 0 & 0 \\ 0 & 25 & 0 \\ 0 & 0 & 25 \end{bmatrix}$$

Hence,  $A \text{ adj } A = 25 I_3$

**7. Find the inverse of each of the following matrices:**

(i)  $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$

(ii)  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

(iii)  $\begin{bmatrix} a & b \\ c & \frac{1+bc}{a} \end{bmatrix}$

(iv)  $\begin{bmatrix} 2 & 5 \\ -3 & 1 \end{bmatrix}$

**Solution:**

(i) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$\text{Now, } |A| = \cos \theta (\cos \theta) + \sin \theta (\sin \theta) \\ = 1$$

Hence,  $A^{-1}$  exists.

Cofactors of A are

$$C_{11} = \cos \theta$$

$$C_{12} = \sin \theta$$

$$C_{21} = -\sin \theta$$

$$C_{22} = \cos \theta$$

$$\text{Since, } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$$

$$(\text{adj } A) = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}^T$$

$$= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \cdot \text{adj } A$$

$$A^{-1} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

(ii) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

Now,  $|A| = -1 \neq 0$

Hence,  $A^{-1}$  exists.

Cofactors of A are

$$C_{11} = 0$$

$$C_{12} = -1$$

$$C_{21} = -1$$

$$C_{22} = 0$$

$$\text{Since, } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$$

$$(\text{adj } A) = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}^T$$

$$= \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$A^{-1} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

(iii) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$\text{Now, } |A| = \frac{a+abc}{a} - bc = \frac{a+abc-abc}{a} = 1 \neq 0$$

Hence,  $A^{-1}$  exists.

Cofactors of A are

$$C_{11} = \frac{1+bc}{a}$$

$$C_{12} = -c$$

$$C_{21} = -b$$

$$C_{22} = a$$

$$\text{Since, } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$$

$$(\text{adj } A) = \begin{bmatrix} \frac{1+bc}{a} & -c \\ -b & a \end{bmatrix}^T$$

$$= \begin{bmatrix} \frac{1+bc}{a} & -b \\ -c & a \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$A^{-1} = \begin{bmatrix} \frac{1+bc}{a} & -b \\ -c & a \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} \frac{1+bc}{a} & -b \\ -c & a \end{bmatrix}$$

(iv) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$\text{Now, } |A| = 2 + 15 = 17$$

Hence,  $A^{-1}$  exists.

Cofactors of A are

$$C_{11} = 1$$

$$C_{12} = 3$$

$$C_{21} = -5$$

$$C_{22} = 2$$

$$\text{Since, adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$$

$$(\text{adj } A) = \begin{bmatrix} 1 & 3 \\ -5 & 2 \end{bmatrix}^T$$

$$= \begin{bmatrix} 0 & -5 \\ 3 & 2 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \cdot \text{adj } A$$

$$A^{-1} = \frac{1}{17} \begin{bmatrix} 0 & -5 \\ 3 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{1}{17} \begin{bmatrix} 0 & -5 \\ 3 & 2 \end{bmatrix}$$

8. Find the inverse of each of the following matrices.

$$(i) \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix}$$

$$(ii) \begin{bmatrix} 1 & 2 & 5 \\ 1 & -1 & -1 \\ 2 & 3 & -1 \end{bmatrix}$$

$$(iii) \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

$$(iv) \begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$$

$$(v) \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix}$$

$$(vi) \begin{bmatrix} 0 & 0 & -1 \\ 3 & 4 & 5 \\ -2 & -4 & -7 \end{bmatrix}$$

$$(vii) \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos\alpha & \sin\alpha \\ 0 & \sin\alpha & -\cos\alpha \end{bmatrix}$$

**Solution:**

(i) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$|A| = 1 \begin{vmatrix} 3 & 1 \\ 1 & 2 \end{vmatrix} - 2 \begin{vmatrix} 2 & 1 \\ 3 & 2 \end{vmatrix} + 3 \begin{vmatrix} 2 & 3 \\ 3 & 1 \end{vmatrix}$$

$$= 1(6 - 1) - 2(4 - 3) + 3(2 - 9)$$

$$= 5 - 2 - 21$$

$$= -18$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = 5$$

$$C_{21} = -1$$

$$C_{31} = -7$$

$$C_{12} = -1$$

$$C_{22} = -7$$

$$C_{32} = 5$$

$$C_{13} = -7$$

$$C_{23} = 5$$

$$C_{33} = -1$$

We know that  $\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

$$= \begin{bmatrix} 5 & -1 & -7 \\ -1 & -7 & 5 \\ -7 & 5 & -1 \end{bmatrix}^T$$

$$\text{So, adj } A = \begin{bmatrix} 5 & -1 & -7 \\ -1 & -7 & 5 \\ -7 & 5 & -1 \end{bmatrix}$$

Now,  $A^{-1} = \frac{1}{|A|} \text{adj } A$

$$\text{So, } A^{-1} = \frac{1}{(-18)} \begin{bmatrix} 5 & -1 & -7 \\ -1 & -7 & 5 \\ -7 & 5 & -1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} \frac{-5}{18} & \frac{1}{18} & \frac{7}{18} \\ \frac{1}{18} & \frac{7}{18} & \frac{-5}{18} \\ \frac{7}{18} & \frac{-5}{18} & \frac{1}{18} \end{bmatrix}$$

Hence,

(ii) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$\begin{aligned} |A| &= 1 \begin{vmatrix} -1 & -1 \\ 3 & -1 \end{vmatrix} - 2 \begin{vmatrix} 1 & -1 \\ 2 & -1 \end{vmatrix} + 5 \begin{vmatrix} 1 & -1 \\ 2 & 3 \end{vmatrix} \\ &= 1(1 + 3) - 2(-1 + 2) + 5(3 + 2) \\ &= 4 - 2 + 25 \\ &= 27 \end{aligned}$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = 4$$

$$C_{21} = 17$$

$$C_{31} = 3$$

$$C_{12} = -1$$

$$C_{22} = -11$$

$$C_{32} = 6$$

$$C_{13} = 5$$

$$C_{23} = 1$$

$$C_{33} = -3$$

$$\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} 4 & -1 & 5 \\ 17 & -11 & 1 \\ 3 & 6 & -3 \end{bmatrix}^T$$

$$\text{So, adj } A = \begin{bmatrix} 4 & 17 & 3 \\ -1 & -11 & 6 \\ 5 & 1 & -3 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\text{So, } A^{-1} = \frac{1}{(27)} \begin{bmatrix} 4 & 17 & 3 \\ -1 & -11 & 6 \\ 5 & 1 & -3 \end{bmatrix}$$

$$\text{Hence, } A^{-1} = \begin{bmatrix} \frac{4}{27} & \frac{17}{27} & \frac{3}{27} \\ \frac{-1}{27} & \frac{-11}{27} & \frac{6}{27} \\ \frac{5}{27} & \frac{1}{27} & \frac{-3}{27} \end{bmatrix} = \begin{bmatrix} \frac{4}{27} & \frac{17}{27} & \frac{1}{9} \\ \frac{-1}{27} & \frac{-11}{27} & \frac{2}{9} \\ \frac{5}{27} & \frac{1}{27} & \frac{-1}{9} \end{bmatrix}$$

(iii) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$\begin{aligned} |A| &= 2 \begin{vmatrix} 2 & -1 \\ -1 & 2 \end{vmatrix} + 1 \begin{vmatrix} -1 & -1 \\ 1 & 2 \end{vmatrix} + 1 \begin{vmatrix} -1 & 2 \\ 1 & -1 \end{vmatrix} \\ &= 2(4 - 1) + 1(-2 + 1) + 1(1 - 2) \\ &= 6 - 2 \\ &= -4 \end{aligned}$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = 3$$

$$C_{21} = 1$$

$$C_{31} = -1$$

$$C_{12} = +1$$

$$C_{22} = 3$$

$$C_{32} = 1$$

$$C_{13} = -1$$

$$C_{23} = 1$$

$$C_{33} = 3$$

$$\text{We know that } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}^T$$

$$\text{So, } \text{adj } A = \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\text{So, } A^{-1} = \frac{1}{4} \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$$

$$\text{Hence, } A^{-1} = \begin{bmatrix} \frac{3}{4} & \frac{1}{4} & \frac{-1}{4} \\ \frac{1}{4} & \frac{3}{4} & \frac{1}{4} \\ \frac{-1}{4} & \frac{1}{4} & \frac{3}{4} \end{bmatrix}$$

(iv) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$\begin{aligned} |A| &= 2 \begin{vmatrix} 1 & 0 \\ 1 & 3 \end{vmatrix} - 0 \begin{vmatrix} 5 & 0 \\ 0 & 3 \end{vmatrix} - 1 \begin{vmatrix} 5 & 1 \\ 0 & 1 \end{vmatrix} \\ &= 2(3 - 0) - 0 - 1(5) \\ &= 6 - 5 \\ &= 1 \end{aligned}$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = 3$$

$$C_{21} = -1$$

$$C_{31} = 1$$

$$C_{12} = -15$$



$$C_{22} = 6$$

$$C_{32} = -5$$

$$C_{13} = -5$$

$$C_{23} = -2$$

$$C_{33} = 2$$

$$\text{We know that } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} 3 & -15 & 5 \\ -1 & 6 & -2 \\ 1 & -5 & 2 \end{bmatrix}^T$$

$$\text{So, } \text{adj } A = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\text{So, } A^{-1} = \frac{1}{1} \begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix}$$

$$\text{Hence, } A^{-1} = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix}$$

(v) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$|A| = 0 \begin{vmatrix} -3 & 0 \\ -3 & 4 \end{vmatrix} - 1 \begin{vmatrix} 4 & 4 \\ 3 & 4 \end{vmatrix} - 1 \begin{vmatrix} 4 & -3 \\ 3 & -3 \end{vmatrix}$$

$$= 0 - 1(16 - 12) - 1(-12 + 9)$$

$$= -4 + 3$$

$$= -1$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = 0$$

$$C_{21} = -1$$

$$\begin{aligned}C_{31} &= 1 \\C_{12} &= -4 \\C_{22} &= 3 \\C_{32} &= -4 \\C_{13} &= -3 \\C_{23} &= 3 \\C_{33} &= -4\end{aligned}$$

We know that  $\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

$$= \begin{bmatrix} 0 & -4 & -3 \\ -1 & 3 & 3 \\ 1 & -4 & -4 \end{bmatrix}^T$$

$$\text{So, adj } A = \begin{bmatrix} 0 & -1 & 1 \\ -4 & 3 & -4 \\ -3 & 3 & -4 \end{bmatrix}$$

Now,  $A^{-1} = \frac{1}{|A|} \text{adj } A$

$$\text{So, } A^{-1} = \frac{1}{-1} \begin{bmatrix} 0 & -1 & 1 \\ -4 & 3 & -4 \\ -3 & 3 & -4 \end{bmatrix}$$

$$\text{Hence, } A^{-1} = \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix}$$

(vi) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$\begin{aligned}|A| &= 0 \begin{vmatrix} 4 & 5 \\ -4 & -7 \end{vmatrix} - 0 \begin{vmatrix} 3 & 5 \\ -2 & -7 \end{vmatrix} - 1 \begin{vmatrix} 3 & 4 \\ -2 & -4 \end{vmatrix} \\ &= 0 - 0 - 1(-12 + 8) \\ &= 4\end{aligned}$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = -8$$

$$C_{21} = 4$$

$$C_{31} = 4$$

$$C_{12} = 11$$

$$C_{22} = -2$$

$$C_{32} = -3$$

$$C_{13} = -4$$

$$C_{23} = 0$$

$$C_{33} = 0$$

We know that  $\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

$$= \begin{bmatrix} 8 & 11 & -4 \\ 4 & -2 & 0 \\ 4 & -3 & 0 \end{bmatrix}^T$$

$$\text{So, adj } A = \begin{bmatrix} 8 & 4 & 4 \\ 11 & -2 & -3 \\ -4 & 0 & 0 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\text{So, } A^{-1} = \frac{1}{4} \begin{bmatrix} 8 & 4 & 4 \\ 11 & -2 & -3 \\ -4 & 0 & 0 \end{bmatrix}$$

$$\text{Hence, } A^{-1} = \begin{bmatrix} 2 & 1 & 1 \\ \frac{11}{4} & \frac{-1}{2} & \frac{-3}{4} \\ -1 & 0 & 0 \end{bmatrix}$$

(vii) The criteria of existence of inverse matrix is the determinant of a given matrix should not equal to zero.

$$\begin{aligned} |A| &= 1 \begin{vmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{vmatrix} - 0 + 0 \\ &= -(\cos^2 \alpha - \sin^2 \alpha) \\ &= -1 \end{aligned}$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = -1$$

$$C_{21} = 0$$

$$C_{31} = 0$$

$$C_{12} = 0$$

$$C_{22} = -\cos \alpha$$

$$C_{32} = -\sin \alpha$$

$$C_{13} = 0$$

$$C_{23} = -\sin \alpha$$

$$C_{33} = \cos \alpha$$

We know that  $\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

$$= \begin{bmatrix} -1 & 0 & 0 \\ 0 & -\cos \alpha & -\sin \alpha \\ 0 & -\sin \alpha & \cos \alpha \end{bmatrix}^T$$

$$\text{So, adj } A = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -\cos \alpha & -\sin \alpha \\ 0 & -\sin \alpha & \cos \alpha \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \text{adj } A$$

$$\text{So, } A^{-1} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -\cos \alpha & -\sin \alpha \\ 0 & -\sin \alpha & \cos \alpha \end{bmatrix}$$

$$\text{Hence, } A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & \sin \alpha \\ 0 & \sin \alpha & -\cos \alpha \end{bmatrix}$$

9. Find the inverse of each of the following matrices and verify that  $A^{-1}A = I_3$ .

(i)  $\begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$

(ii)  $\begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$

**Solution:**

(i) We have

$$\begin{aligned}
 |A| &= 1 \begin{vmatrix} 4 & 3 \\ 3 & 4 \end{vmatrix} - 3 \begin{vmatrix} 1 & 3 \\ 1 & 4 \end{vmatrix} + 3 \begin{vmatrix} 1 & 4 \\ 1 & 3 \end{vmatrix} \\
 &= 1(16 - 9) - 3(4 - 3) + 3(3 - 4) \\
 &= 7 - 3 - 3 \\
 &= 1
 \end{aligned}$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = 7$$

$$C_{21} = -3$$

$$C_{31} = -3$$

$$C_{12} = -1$$

$$C_{22} = -1$$

$$C_{32} = 0$$

$$C_{13} = -1$$

$$C_{23} = 0$$

$$C_{33} = 1$$

We know that  $\text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$

$$= \begin{bmatrix} 7 & -1 & -1 \\ -3 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix}^T$$

So,  $\text{adj } A = \begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$

Now,  $A^{-1} = \frac{1}{1} \begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$

Also,  $A^{-1}A = \begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$

$$= \begin{bmatrix} 7 - 3 - 3 & 21 - 12 - 9 & 21 - 9 - 12 \\ -1 + 1 + 0 & -3 + 4 + 0 & -3 + 3 + 0 \\ -1 + 0 + 1 & -3 + 0 + 3 & -3 + 0 + 4 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Hence,  $A^{-1}A = I_3$

(ii) We have

$$\begin{aligned} |A| &= 2 \begin{vmatrix} 4 & 1 \\ 7 & 2 \end{vmatrix} - 3 \begin{vmatrix} 3 & 1 \\ 3 & 2 \end{vmatrix} + 1 \begin{vmatrix} 3 & 4 \\ 3 & 7 \end{vmatrix} \\ &= 2(8 - 7) - 3(6 - 3) + 1(21 - 12) \\ &= 2 - 9 + 9 \\ &= 2 \end{aligned}$$

Hence,  $A^{-1}$  exists

Cofactors of A are

$$C_{11} = 1$$

$$C_{21} = 1$$

$$C_{31} = -1$$

$$C_{12} = -3$$

$$C_{22} = 1$$

$$C_{32} = 1$$

$$C_{13} = 9$$

$$C_{23} = -5$$

$$C_{33} = -1$$

$$\text{We know that } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} 1 & -3 & 9 \\ 1 & 1 & -5 \\ -1 & 1 & -1 \end{bmatrix}^T$$

$$\text{So, } \text{adj } A = \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{2} \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix}$$

$$\text{Also, } A^{-1}.A = \frac{1}{2} \begin{bmatrix} 1 & 1 & -1 \\ -3 & 1 & 1 \\ 9 & -5 & -1 \end{bmatrix} \begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 2 + 3 - 3 & 3 + 4 - 7 & 1 + 1 - 2 \\ -6 + 3 + 3 & -9 + 4 + 7 & -3 + 1 + 2 \\ 18 - 15 - 3 & 27 - 20 - 7 & 9 - 5 - 2 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{Hence, } A^{-1}.A = I_3$$

10. For the following pair of matrices verify that  $(AB)^{-1} = B^{-1}A^{-1}$ .

$$(i) A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix} \text{ and } B = \begin{bmatrix} 4 & 6 \\ 3 & 2 \end{bmatrix}$$

$$(ii) A = \begin{bmatrix} 2 & 1 \\ 5 & 3 \end{bmatrix} \text{ and } B = \begin{bmatrix} 4 & 5 \\ 3 & 4 \end{bmatrix}$$

**Solution:**

(i) Given

$$A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix},$$

$$|A| = 1$$

$$\text{Then, } \text{adj } A = \begin{bmatrix} 5 & -2 \\ -7 & 3 \end{bmatrix}$$

$$A^{-1} = \frac{\text{adj } A}{|A|} = \frac{1}{1} \begin{bmatrix} 5 & -2 \\ -7 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 4 & 6 \\ 3 & 2 \end{bmatrix},$$

$$|B| = -10$$

$$\text{Then, adj } B = \begin{bmatrix} 2 & -6 \\ -3 & 4 \end{bmatrix}$$

$$B^{-1} = \frac{1}{10} \begin{bmatrix} 2 & -6 \\ -3 & 4 \end{bmatrix}$$

$$\text{Also, } A \cdot B = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix} \begin{bmatrix} 4 & 6 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} 12 + 6 & 18 + 4 \\ 28 + 15 & 42 + 10 \end{bmatrix}$$

$$AB = \begin{bmatrix} 18 & 22 \\ 43 & 52 \end{bmatrix}$$

$$|AB| = 936 - 946 = -10$$

$$\text{Adj } (AB) = \begin{bmatrix} 52 & -22 \\ -43 & 18 \end{bmatrix}$$

$$(AB)^{-1} = \frac{1}{-10} \begin{bmatrix} 52 & -22 \\ -43 & 18 \end{bmatrix} = \begin{bmatrix} -52 & 22 \\ 43 & -18 \end{bmatrix}$$

$$\text{Now } B^{-1}A^{-1} = \frac{1}{-10} \begin{bmatrix} 2 & -6 \\ -3 & 4 \end{bmatrix} \begin{bmatrix} 5 & -2 \\ -7 & 3 \end{bmatrix}$$

$$= \frac{1}{-10} \begin{bmatrix} 10 + 42 & -4 - 18 \\ -15 - 28 & 6 + 12 \end{bmatrix}$$

$$= \frac{1}{10} \begin{bmatrix} -52 & 22 \\ 43 & -18 \end{bmatrix}$$

$$\text{Hence, } (AB)^{-1} = B^{-1}A^{-1}$$

(ii) Given

$$|A| = 1$$

$$\text{Adj } A = \begin{bmatrix} 3 & -1 \\ -5 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{\text{adj } A}{|A|} = \frac{1}{1} \begin{bmatrix} 3 & -1 \\ -5 & 2 \end{bmatrix}$$

$$B = \begin{bmatrix} 4 & 5 \\ 3 & 4 \end{bmatrix}$$



$$|B| = -1$$

$$B^{-1} = \frac{\text{adj } A}{|A|} = \frac{1}{-1} \begin{bmatrix} 4 & -5 \\ -3 & 4 \end{bmatrix}$$

$$\text{Also, } AB = \begin{bmatrix} 2 & 1 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 3 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} 11 & 14 \\ 29 & 37 \end{bmatrix}$$

$$|AB| = 407 - 406 = 1$$

$$\text{And, } \text{adj } (AB) = \begin{bmatrix} 37 & -14 \\ -29 & 11 \end{bmatrix}$$

$$(AB)^{-1} = \frac{\text{adj } AB}{|AB|}$$

$$= \begin{bmatrix} 37 & -14 \\ -29 & 11 \end{bmatrix}$$

$$\text{Now, } B^{-1}A^{-1} = \begin{bmatrix} 4 & -5 \\ -3 & 4 \end{bmatrix} \begin{bmatrix} 3 & -1 \\ -5 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 37 & -14 \\ -29 & 11 \end{bmatrix}$$

$$\text{Hence, } (AB)^{-1} = B^{-1}A^{-1}$$

11. Let  $A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix}$  and  $B = \begin{bmatrix} 6 & 7 \\ 8 & 9 \end{bmatrix}$ . Find  $(AB)^{-1}$

**Solution:**

Given

$$A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix}$$

$$|A| = 15 - 14 = 1$$

$$\text{Therefore } \text{adj } A = \begin{bmatrix} 5 & -2 \\ -7 & 3 \end{bmatrix}$$

$$A^{-1} = \frac{\text{adj}}{|A|} = \frac{1}{1} \begin{bmatrix} 5 & -2 \\ -7 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 6 & 7 \\ 8 & 9 \end{bmatrix}$$

$$|B| = 54 - 56 = -2 \quad \text{adj } B = \begin{bmatrix} 9 & -7 \\ -8 & 6 \end{bmatrix}$$

$$B^{-1} = \frac{\text{adj}B}{|B|} = \frac{1}{-2} \begin{bmatrix} 9 & -7 \\ -8 & 6 \end{bmatrix}$$

Now,  $(AB)^{-1} = B^{-1}A^{-1}$

$$= \frac{1}{-2} \begin{bmatrix} 9 & -7 \\ -8 & 6 \end{bmatrix} \begin{bmatrix} 5 & -2 \\ -7 & 3 \end{bmatrix}$$

$$= \frac{1}{-2} \begin{bmatrix} 45 + 49 & -18 - 21 \\ -40 - 42 & 16 + 18 \end{bmatrix}$$

$$= \frac{1}{-2} \begin{bmatrix} 94 & -39 \\ -82 & 34 \end{bmatrix}$$

$$(AB)^{-1} = \begin{bmatrix} -47 & \frac{39}{2} \\ 41 & -17 \end{bmatrix}$$

12. Given  $A = \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix}$ , compute  $A^{-1}$  and show that  $2A^{-1} = 9I - A$ .

**Solution:**

Given

$$A = \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix}$$

$$|A| = 14 - 12 = 2 \quad \text{adj } A = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{1}{2} \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$$

To Show:  $2A^{-1} = 9I - A$

We have

$$\text{L.H.S} = 2A^{-1} = 2 \cdot \frac{1}{2} \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix} = \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$$

$$\text{R.H.S} = 9I - A = \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix} - \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix}$$

$$= \begin{bmatrix} 7 & 3 \\ 4 & 2 \end{bmatrix}$$

Hence,  $2A^{-1} = 9I - A$

13. If  $A = \begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix}$ , then show that  $A - 3I = 2(I + 3A^{-1})$ .

**Solution:**

Given

$$A = \begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix}$$

$$|A| = 4 - 10 = -6 \quad \text{adj } A = \begin{bmatrix} 1 & -5 \\ -2 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{1}{-6} \begin{bmatrix} 1 & -5 \\ -2 & 4 \end{bmatrix}$$

To Show:  $A - 3I = 2(I + 3A^{-1})$

We have

$$\text{LHS} = A - 3I$$

$$= \begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix} - 3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 5 \\ 2 & -2 \end{bmatrix}$$

$$\text{R.H.S} = 2(I + 3A^{-1}) = 2I + 6A^{-1}$$

$$= 2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + 6 \frac{1}{6} \begin{bmatrix} -1 & 5 \\ 2 & -4 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} + \begin{bmatrix} -1 & 5 \\ 2 & -4 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 5 \\ 2 & -4 \end{bmatrix}$$

Hence,  $A - 3I = 2(I + 3A^{-1})$

14. Find the inverse of the matrix  $A = \begin{bmatrix} a & b \\ c & \frac{1+bc}{a} \end{bmatrix}$ , and show that  $aA^{-1} = (a^2 + bc + 1)I - aA$ .

**Solution:**

$$A = \begin{bmatrix} a & b \\ c & \frac{1+bc}{a} \end{bmatrix}$$

$$\text{Now, } |A| = \frac{a+abc}{a} - bc = \frac{a+abc-abc}{a} = 1 \neq 0$$

Hence,  $A^{-1}$  exists.

Cofactors of A are

$$C_{11} = \frac{1+bc}{a}$$

$$C_{12} = -c$$

$$C_{21} = -b$$

$$C_{22} = a$$

$$\text{Since, } \text{adj } A = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}^T$$

$$\text{Adj } A = \begin{bmatrix} \frac{1+bc}{a} & -c \\ -b & a \end{bmatrix}^T$$

$$= \begin{bmatrix} \frac{1+bc}{a} & -b \\ -c & a \end{bmatrix}$$

Now,  $A^{-1} = \frac{1}{|A|} \cdot \text{adj } A$

$$A^{-1} = \begin{bmatrix} \frac{1+bc}{a} & -b \\ -c & a \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} \frac{1+bc}{a} & -b \\ -c & a \end{bmatrix}$$

To show  $a A^{-1} = (a^2 + bc + 1) I - aA$ .

LHS =  $a A^{-1}$

$$= a \begin{bmatrix} \frac{1+bc}{a} & -b \\ -c & a \end{bmatrix}$$

$$= \begin{bmatrix} 1+bc & -ab \\ -ac & a^2 \end{bmatrix}$$

RHS =  $(a^2 + bc + 1) I - aA$

$$= \begin{bmatrix} a^2 + bc + 1 & 0 \\ 0 & a^2 + bc + 1 \end{bmatrix} - \begin{bmatrix} a^2 & ab \\ ac & 1 + bc \end{bmatrix} = \begin{bmatrix} 1+bc & -ab \\ -ac & a^2 \end{bmatrix}$$

Hence, LHS = RHS

15. Given  $A = \begin{bmatrix} 5 & 0 & 4 \\ 2 & 3 & 2 \\ 1 & 2 & 1 \end{bmatrix}$ ,  $B^{-1} = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$ . Compute  $(AB)^{-1}$

**Solution:**

Given

$$A = \begin{bmatrix} 5 & 0 & 4 \\ 2 & 3 & 2 \\ 1 & 2 & 1 \end{bmatrix} \text{ and } B^{-1} = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$$

Here,  $(AB)^{-1} = B^{-1} A^{-1}$

$$|A| = -5 + 4 = -1$$

Cofactors of A are

$$C_{11} = -1$$

$$C_{21} = 8$$

$$C_{31} = -12$$

$$C_{12} = 0$$

$$C_{22} = 1$$

$$C_{32} = -2$$

$$C_{13} = 1$$

$$C_{23} = -10$$

$$C_{33} = 15$$

$$\text{Adj } A = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} -1 & 0 & 1 \\ 8 & 1 & -10 \\ -12 & -2 & 15 \end{bmatrix}^T$$

$$\text{So, adj } A = \begin{bmatrix} -1 & 8 & -12 \\ 0 & 1 & -2 \\ 1 & -10 & 15 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{-1} \begin{bmatrix} -1 & 8 & -12 \\ 0 & 1 & -2 \\ 1 & -10 & 15 \end{bmatrix}$$

$$(AB)^{-1} = B^{-1}A^{-1}$$

$$= \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix} \begin{bmatrix} 1 & -8 & 12 \\ 0 & -1 & 2 \\ -1 & 10 & -15 \end{bmatrix}$$

$$= \begin{bmatrix} 1 + 0 - 3 & -8 - 3 + 30 & 12 + 6 - 45 \\ 1 + 0 - 3 & -8 - 4 + 30 & 12 + 8 - 45 \\ 1 + 0 - 4 & -8 - 3 + 40 & 12 + 6 - 60 \end{bmatrix}$$

$$\text{Hence, } = \begin{bmatrix} -2 & 19 & -27 \\ -2 & 18 & -25 \\ -3 & 29 & 42 \end{bmatrix}$$

16. Let  $F(\alpha) = \begin{bmatrix} \cos\alpha & -\sin\alpha & 0 \\ \sin\alpha & \cos\alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$  and  $G(\beta) = \begin{bmatrix} \cos\beta & 0 & \sin\beta \\ 0 & 1 & 0 \\ -\sin\beta & 0 & \cos\beta \end{bmatrix}$ . Show that

(i)  $[F(\alpha)]^{-1} = F(-\alpha)$

(ii)  $[G(\beta)]^{-1} = G(-\beta)$

(iii)  $[F(\alpha) G(\beta)]^{-1} = G(-\beta) F(-\alpha)$

**Solution:**

(i) Given

$$F(\alpha) = \begin{bmatrix} \cos\alpha & -\sin\alpha & 0 \\ \sin\alpha & \cos\alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$|F(\alpha)| = \cos^2\alpha + \sin^2\alpha = 1$$

Cofactors of A are

$$C_{11} = \cos\alpha$$

$$C_{21} = \sin\alpha$$

$$C_{31} = 0$$

$$C_{12} = -\sin\alpha$$

$$C_{22} = \cos\alpha$$

$$C_{32} = 0$$

$$C_{13} = 0$$

$$C_{23} = -10$$

$$C_{33} = 1$$

$$\text{Adj } F(\alpha) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} \cos\alpha & -\sin\alpha & 0 \\ \sin\alpha & \cos\alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}^T$$

$$\text{So, adj } F(\alpha) = \begin{bmatrix} \cos\alpha & \sin\alpha & 0 \\ -\sin\alpha & \cos\alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \dots\dots (i)$$

$$\text{Now, } [F(\alpha)]^{-1} = \frac{1}{1} \begin{bmatrix} \cos\alpha & \sin\alpha & 0 \\ -\sin\alpha & \cos\alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{And, } F(-\alpha) = \begin{bmatrix} \cos(-\alpha) & \sin(-\alpha) & 0 \\ \sin(-\alpha) & \cos(-\alpha) & 0 \\ 0 & 0 & 1 \end{bmatrix} \dots\dots (ii)$$

$$= \begin{bmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Hence,  $[F(\alpha)]^{-1} = F(-\alpha)$

(ii) We have

$$|G(\beta)| = \cos^2 \beta + \sin^2 \beta = 1$$

Cofactors of A are

$$C_{11} = \cos \beta$$

$$C_{21} = \sin \alpha$$

$$C_{31} = \sin \beta$$

$$C_{12} = 0$$

$$C_{22} = 1$$

$$C_{32} = 0$$

$$C_{13} = \sin \beta$$

$$C_{23} = 0$$

$$C_{33} = \cos \beta$$

$$\text{Adj } G(\beta) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}^T$$

$$= \begin{bmatrix} \cos \beta & 0 & \sin \beta \\ 0 & 1 & 0 \\ -\sin \beta & 0 & \cos \beta \end{bmatrix}^T$$

$$\text{So, } \text{adj } G(\beta) = \begin{bmatrix} \cos \beta & 0 & -\sin \beta \\ 0 & 1 & 0 \\ \sin \beta & 0 & \cos \beta \end{bmatrix} \dots\dots (i)$$

$$\text{Now, } [G(\beta)]^{-1} = \frac{1}{1} \begin{bmatrix} \cos \beta & 0 & -\sin \beta \\ 0 & 1 & 0 \\ \sin \beta & 0 & \cos \beta \end{bmatrix}$$



$$\text{So, adj } G(\beta) = \begin{bmatrix} \cos\beta & 0 & -\sin\beta \\ 0 & 1 & 0 \\ \sin\beta & 0 & \cos\beta \end{bmatrix} \dots\dots (j)$$

$$\text{Now, } [G(\beta)]^{-1} = \frac{1}{1} \begin{bmatrix} \cos\beta & 0 & -\sin\beta \\ 0 & 1 & 0 \\ \sin\beta & 0 & \cos\beta \end{bmatrix}$$

$$\text{And, } G(-\beta) = \begin{bmatrix} \cos(-\beta) & 0 & \sin(-\beta) \\ 0 & 1 & 0 \\ \sin(-\beta) & 0 & \cos(-\beta) \end{bmatrix}$$

$$= \begin{bmatrix} \cos\beta & 0 & -\sin\beta \\ 0 & 1 & 0 \\ \sin\beta & 0 & \cos\beta \end{bmatrix}$$

$$\text{Hence, } [G(\beta)]^{-1} = G(-\beta)$$

(iii) Now we have to show that

$$[F(\alpha) G(\beta)]^{-1} = G(-\beta) F(-\alpha)$$

We have already know that

$$[G(\beta)]^{-1} = G(-\beta)$$

$$[F(\alpha)]^{-1} = F(-\alpha)$$

$$\text{And LHS} = [F(\alpha) G(\beta)]^{-1}$$

$$= [G(\beta)]^{-1} [F(\alpha)]^{-1}$$

$$= G(-\beta) F(-\alpha)$$

Hence = RHS

17. If  $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$  verify that  $A^2 - 4A + I = 0$ , where  $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  and  $O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Hence find  $A^{-1}$ .

**Solution:**

Consider,

$$A^2 = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 4 + 3 & 6 + 6 \\ 2 + 2 & 3 + 4 \end{bmatrix}$$

$$= \begin{bmatrix} 7 & 12 \\ 4 & 7 \end{bmatrix}$$

$$4A = 4 \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 8 & 12 \\ 4 & 8 \end{bmatrix}$$

$$I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\text{Now, } A^2 - 4A + I = \begin{bmatrix} 7 & 12 \\ 4 & 7 \end{bmatrix} - \begin{bmatrix} 8 & 12 \\ 4 & 8 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 7 - 8 + 1 & 12 - 2 + 0 \\ 4 - 4 + 0 & 7 - 8 + 1 \end{bmatrix}$$

$$\text{Hence, } = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\text{Now, } A^2 - 4A + I = 0$$

$$A \cdot A - 4A = -I$$

Multiply by  $A^{-1}$  both sides we get

$$A \cdot A (A^{-1}) - 4A A^{-1} = -I A^{-1}$$

$$AI - 4I = -A^{-1}$$

$$A^{-1} = 4I - A = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} - \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix}$$

18. Show that  $A = \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix}$  satisfies the equation  $A^2 + 4A - 42I = 0$ . Hence find  $A^{-1}$ .

**Solution:**

Given

$$A = \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix} = \begin{bmatrix} 64 + 10 & -40 + 20 \\ -16 + 8 & 10 + 16 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix} = \begin{bmatrix} 64 + 10 & -40 + 20 \\ -16 + 8 & 10 + 16 \end{bmatrix}$$

$$= \begin{bmatrix} 74 & -20 \\ -8 & 26 \end{bmatrix}$$

$$4A = 4 \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix} = \begin{bmatrix} -32 & 20 \\ 8 & 16 \end{bmatrix}$$

$$42I = 42 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 42 & 0 \\ 0 & 42 \end{bmatrix}$$

Now,

$$A^2 + 4A - 42I = \begin{bmatrix} 74 & -20 \\ -8 & 26 \end{bmatrix} + \begin{bmatrix} -32 & 20 \\ 8 & 16 \end{bmatrix} - \begin{bmatrix} 42 & 0 \\ 0 & 42 \end{bmatrix}$$

$$= \begin{bmatrix} 74 - 32 - 42 & -20 + 20 \\ -8 + 8 & 26 + 16 - 42 \end{bmatrix}$$

$$\text{Hence, } = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\text{Now, } A^2 + 4A - 42I = 0$$

$$= A^{-1}A \cdot A + 4A^{-1} \cdot A - 42A^{-1}I = 0$$

$$= IA + 4I - 42A^{-1} = 0$$

$$= 42A^{-1} = A + 4I$$

$$= A^{-1} = \frac{1}{42}[A + 4I]$$

$$= \frac{1}{42} \left[ \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix} + \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \right]$$

$$A^{-1} = \frac{1}{42} \begin{bmatrix} -4 & 5 \\ 2 & 8 \end{bmatrix}$$

19. If  $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$  show that  $A^2 - 5A + 7I = 0$ . Hence find  $A^{-1}$ .

**Solution:**

Given

$$A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 9-1 & 3+2 \\ -3-2 & -1+4 \end{bmatrix}$$

$$= \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix}$$

$$\text{Now, } A^2 - 5A + 7I = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} - 5 \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} + 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 8-15+7 & 5-5+0 \\ -5+5+0 & 3-10+7 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\text{So, } A^2 - 5A + 7I = 0$$

Multiply by  $A^{-1}$  both sides

$$= A \cdot A A^{-1} - 5A \cdot A^{-1} + 7I \cdot A^{-1} = 0$$

$$A - 5I + 7A^{-1} = 0$$

$$A^{-1} = \frac{1}{7}[5I - A]$$

$$A^{-1} = \frac{1}{7} \cdot \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix} - \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{1}{7} \cdot \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$$