Circles

Introduction to Circles

Circles

- The **set of all the points** in a plane that is at a **fixed distance** from a **fixed point** makes a circle.

- A **Fixed point** from which the set of points are at fixed distance is called **centre** of the circle.

- A circle divides the plane into 3 parts: **interior** (inside the circle), the **circle** itself and **exterior** (outside the circle)


Radius

- The **distance** between the **center** of the circle and any **point on its edge** is called the **radius**.

Tangent and Secant

A line that touches the circle at exactly one point is called its tangent.

A line that cuts a circle at two points is called as a secant.



In the above figure: PQ is the tangent and AB is the secant.

Chord

-The **line segment** within the circle joining any 2 points on the circle is called the chord.

Diameter

- A Chord passing through the center of the circle is called the diameter.
- The **Diameter is 2 times the radius** and it is the **longest chord**.

Arc

- The **portion** of a circle(curve) **between 2 points** is called an **arc**.

- Among the two pieces made by an arc, the **longer** one is called **major arc** and the **shorter** one is called **minor arc**.

Circumference

The **perimeter** of a circle is the **distance** covered by going around its **boundary once**. The perimeter of a circle has a special name: **Circumference**, which is π times the diameter which is given by the formula $2\pi r$

Segment and Sector

- A circular **segment** is a region of a circle which is "**cut off**" from the rest of the circle by a secant or a chord.

- Smaller region cut off by a chord is called **minor segment** and the **bigger region** is called **major segment**.


- A sector is the portion of a circle enclosed by two radii and an arc, where the smaller area is known as the minor sector and the larger being the major sector.

- For **2 equa**l arcs or for semicircles - both the segment and sector is called the **semicircular** region.

Circles and Their Chords

Theorem of equal chords subtending angles at the center.

- Equal chords subtend equal angles at the center.




Proof : AB and CD are the 2 equal chords. In $\triangle AOB$ and $\triangle COD$

OB = OC [Radii] OA = OD [Radii] AB = CD [Given] $\Delta AOB \cong \Delta COD$ (SSS rule) Hence, $\angle AOB = \angle COD$ [CPCT]

Theorem of equal angles subtended by different chords.


- If the **angles** subtended by the chords of a circle at the center are **equal**, then the **chords are equal**.



Proof : In $\triangle AOB$ and $\triangle COD$ OB = OC [Radii] $\angle AOB = \angle COD$ [Given] OA = OD [Radii] $\triangle AOB \cong \triangle COD$ (SAS rule) Hence, AB = CD [CPCT]

Perpendicular from the center to a chord bisects the chord.


Perpendicular from the center of a circle to a chord bisects the chord.



Proof: AB is a chord and OM is the perpendicular drawn from the center. From ΔOMB and ΔOMA , $\angle OMA = \angle OMB = 90^{0}$ OA = OB (radii) OM = OM (common) Hence, $\Delta OMB \cong \Delta OMA$ (RHS rule) Therefore AM = MB [CPCT]

A Line through the center that bisects the chord is perpendicular to the chord.

- A **line** drawn through the center of a circle to **bisect** a chord, is **perpendicular** to the chord.



Proof: OM drawn from center to bisect chord AB . From ΔOMA and ΔOMB , OA = OB (Radii) OM = OM (common) AM = BM (Given) Therefore, $\Delta OMA \cong \Delta OMB$ (SSS rule) $\Rightarrow \angle OMA = \angle OMB$ (C.P.C.T) But, $\angle OMA + \angle OMB = 180^{0}$ Hence, $\angle OMA = \angle OMB = 90^{0}$ $\Rightarrow OM \bot AB$


Circle through 3 points

- There is **one** and **only** one **circle** passing through **three given noncollinear points**.

- A unique circle passes through 3 vertices of a triangle ABC called as the **circumcircle**. The **centre** and **radius** are called the **circumcenter** and **circumradius** of this triangle, respectively.

Equal chords are at equal distances from the center.


Equal chords of a circle(or of congruent circles) are equidistant from the centre (or centres).



Proof: Given, AB = CD, O is the centre. Join OA and OC. Draw, $OP \perp AB$, $OQ \perp CD$ In $\triangle OAP$ and $\triangle OCQ$, OA = OC (Radii) AP = CQ (AB = CD $\Rightarrow \frac{1}{2}AB = \frac{1}{2}CD$, since OP and OQ bisects the chords AB and CD.) $\triangle OAP \cong \triangle OCQ$ (RHS rule) Hence, OP = OQ (C.P.C.T.C)

Chords equidistant from center are equal

Chords equidistant from the center of a circle are equal in length.



Proof : Given OX = OY (The chords AB and CD are at equidistant) $OX \perp AB, OY \perp CD$ In $\triangle AOX$ and $\triangle DOY$ $\angle OXA = \angle OYD$ (Both 90⁰)

```
OA = OD (Radii)

OX = OY (Given)

\Delta AOX \cong \Delta DOY (RHS rule)

Therefore AX = DY (CPCT)


Similarly XB = YC

So, AB = CD
```

Circles and Quadrilaterals

Angle subtended by an arc of a circle on the circle and at the center

The **angle** subtended by **an arc** at the **centre is double** the angle subtended by it on any **part of the circle**.



Here PQ is the arc of a circle with centre O, that subtends $\angle POQ$ at the centre. Join AO and extend it to B.

```
In \triangle OAQ

OA = OQ..... [Radii]

Hence, \angle OAQ = \angle OQA....[Property of isosceles triangle]

Implies \angle BOQ = 2\angle OAQ .....[Exterior angle of triangle = Sum of 2 interior angles]

Similarly, \angle BOP = 2\angle OAP


\Rightarrow \angle BOQ + \angle BOP = 2\angle OAQ + 2\angle OAP

\Rightarrow \angle POQ = 2\angle PAQ

Hence proved
```

Angles in same segment of a circle.

-Angles in the same segment of a circle are equal.



Consider a circle with centre O.

 $\angle PAQ$ and $\angle PCQ$ are the angles formed in the major segment PACQ with respect to the arc PQ.


Join OP and OQ

 $\angle POQ = 2 \angle PAQ = 2 \angle PCQ$ [Angle subtended by an arc at the centre is double the angle subtended by it in any part of the circle]

 $\Rightarrow \angle PCQ = \angle PAQ$ Hence proved

Angle subtended by diameter on the circle

- **Angle** subtended by **diameter** on a circle is a **right angle**.(Angle in a semicircle is a right angle)



Consider a circle with center O, POQ is the diameter of the circle.

 $\angle PAQ$ is the angle subtended by diameter PQ at the circuference.


 $\angle POQ$ is the angle subtended by diameter PQ at the center.

 $\angle PAQ = \frac{1}{2} \angle POQ...$ [Angle subtended by arc at the centre is double the angle at any other part] $\angle PAQ = \frac{1}{2} \angle 180^0 - 90^0$

 $\angle PAQ = \frac{1}{2} \times 180^0 = 90^0$ Hence proved

Line segment that subtends equal angles at two other points


- If a line segment joining two points subtends equal angles at two other points lying on the same side of the line containing the line segment, the four points lie on a circle.(i.e they are concyclic)



Here $\angle ACB = \angle ADB$ and all 4 points A,B,C,D are concyclic.

Cyclic Quadrilateral

- A Quadrilateral is called a cyclic quadrilateral if all the four vertices lie on a circle.



In a circle, if all **four points** A, B, C and D lie **on the circle**, then quadrilateral ABCD is a **cyclic quadrilateral**.

Sum of opposite angles of a cyclic quadrilateral

- If sum of a pair of opposite angles of a quadrilateral is 180 degree, the quadrilateral is cyclic.

Sum of pair of opposite angles in quadrilateral

- The sum of either pair of opposite angles of a cyclic quadrilateral is 180 degree

S A JUN APP