

Exercise-4.1

Page: 47

1. What are the canal rays?

Solution:

The radiations that are positively charged are canal rays. This discovery was crucial in the discovery of another subatomic particle that was positively charged - proton.

2. If an atom contains one electron and one proton, will it carry any charge or not?

Solution:

Since a proton is a positively charged particle and an electron is a negatively charged particle, the net charge becomes neutral as both the particles neutralizes each other.

Exercise-4.2

Page: 49

3. On the basis of Thompson's model of an atom, explain how the atom is neutral as a whole.

Solution:

As per Thompson's model of an atom,

(i) An atom contains a positively charged sphere in which the negatively charged electrons are implanted.

(ii) Electrons and protons are equal in magnitude hence an atom on the whole is electrically neutral.


4. On the basis of Rutherford's model of an atom, which subatomic particle is present in the nucleus of an atom?

Solution:

As per Rutherford's model of an atom, the positively charged protons are the ones that are present in the atom.

5. Draw a sketch of Bohr's model of an atom with three shells.

Solution:



6. What do you think would be the observation if the α - particle scattering experiment is carried out using a foil of a metal other than gold?

Solution:

In the α – particle scattering experiment, when any other metal foil is used instead of gold, the observation would remain the same. This is because the structure of an atom when considered individually remains the same.

Exercise-4.2.4

Page: 49

7. Name the three subatomic particles of an atom.

Solution:

An atom consists of three subatomic particles:

- Protons - positively charged
- Electrons - negatively charged
- Neutrons - neutral in nature (no charge)

8. Helium atom has an atomic mass of 4 u and two protons in its nucleus. How many neutrons does it have?

Solution:

Given: Atomic mass of helium atom = 4u, 2 protons in helium nucleus

Atomic mass = number of protons + number of neutrons

$$4 = 2 + \text{number of neutrons}$$

$$\text{Number of neutrons} = 4 - 2 = 2$$

Hence, Helium has 2 neutrons.

Exercise-4.3

Page: 50

9. Write the distribution of electrons in Carbon and Sodium atoms.

Solution:

Distribution of electrons in Carbon atoms:

The atomic number of Carbon is 6

Number of electrons is equal to the number of protons in carbon atom i.e., 6

The distribution of electrons in carbon atom is K - 2, L - 4

Distribution of electrons in sodium atoms:

The atomic number of Sodium is 11

Number of electrons is equal to the number of protons in sodium atom i.e., 11

The distribution of electrons in sodium atom is K - 2, L - 8, M - 1

10. If K and L shells of an atom are full, then what would be the total number of electrons in the atom?

Solution:

K shell can hold 2 electrons

L shell can hold 8 electrons

Hence, when both the shells are full, the total number of electrons present in the atom = $2+8 = 10$ electrons.

Exercise-4.4

Page: 52

11. How will you find the valency of chlorine, sulphur and magnesium?

Solution:

The definite combining capacity of the atoms of each element, wherein electrons are lost, gained or shared to make the octet of electrons present in the outermost shell is defined as valency. To measure valency, we can figure out the number of electrons that are required to complete the shell in which it is contained or losing excess electrons if present, once the filling is complete.

To find the valency of chlorine:

The atomic number of chlorine is 17

Number of electrons is equal to the number of protons in chlorine i.e., 17

The distribution of electrons in chlorine atom is K - 2, L - 8, M - 7

Hence, from the distribution of chlorine it is clearly evident that to fill the M shell only one electron is required. Therefore its valency is -1. i.e., one electron less

To find the valency of sulphur:

The atomic number of sulphur is 16

Number of electrons is equal to the number of protons in sulphur i.e., 16

The distribution of electrons in sulphur atom is K - 2, L - 8, M - 6

Hence, from the distribution of sulphur it is clearly evident that to fill the M shell two more electrons are required. Therefore its valency is -2, i.e., two electrons lesser.

To find the valency of magnesium:

The atomic number of magnesium is 12

Number of electrons is equal to the number of protons in magnesium i.e., 12

The distribution of electrons in magnesium atom is K - 2, L - 8, M - 2

Hence, from the distribution of magnesium it is clearly evident that to fill the M shell six more electrons are required. But M shell has two electrons only. It possesses lesser electrons than needed to fill the shell.

Thus, we say that the magnesium atom is not stable as the M shell has 2 electrons. Its valency is +2, meaning it has 2 electrons in excess.

Exercise-4.5

12. If the number of electrons in an atom is 8 and number of protons is also 8, then

- (i) What is the atomic number of the atom? and
- (ii) What is the charge on the atom?

Solution:

Given: Number of electrons = 8

Number of protons = 8

(a) The atomic number of an atom is the same as the number of protons in that atom, hence its atomic number is 8.

(b) In an atom, the number of protons is equal to the number of electrons. Hence both the charges - positive and negative neutralize each other. Therefore, the atom does not possess any charge.

13. With the help of given Table, find out the mass number of oxygen and sulphur atom.

Table: Composition of Atoms of the First Eighteen Elements with Electron Distribution in Various Shells.

Name of Element	Symbol	Atomic number	Number of Protons	Number of Neutrons	Number of electrons	Distribution of electrons				Valency
						K	L	M	N	
Hydrogen	H	1	1	—	1	1	—	—	—	1
Helium	He	2	2	2	2	2	—	—	—	0
Lithium	Li	3	3	4	3	2	1	—	—	1
Beryllium	Be	4	4	5	4	2	2	—	—	2
Boron	B	5	5	6	5	2	3	—	—	3
Carbon	C	6	6	6	6	2	4	—	—	4
Nitrogen	N	7	7	7	7	2	5	—	—	3
Oxygen	O	8	8	8	8	2	6	—	—	2
Fluorine	F	9	9	10	9	2	7	—	—	1
Neon	Ne	10	10	10	10	2	8	—	—	0
Sodium	Na	11	11	12	11	2	8	1	—	1
Magnesium	Mg	12	12	12	12	2	8	2	—	2

NCERT Solution for Class 9 Science - Chapter 4 Structure Of The Atom

Aluminium	Al	13	13	14	13	2	8	3	–	3
Silicon	Si	14	14	14	14	2	8	4	–	4
Phosphorus	P	15	15	16	15	2	8	5	–	3,5
Sulphur	S	16	16	16	16	2	8	6	–	2
Chlorine	Cl	17	17	18	17	2	8	7	–	1
Argon	Ar	18	18	22	18	2	8	8		0

Solution:

(a) To find the mass number of Oxygen:

Number of protons = 8

Number of neutrons = 8

Atomic number = 8

Atomic mass number = Number of protons + number of neutrons = 8 + 8 = 16

Therefore, mass number of oxygen = 16

(b) To find the mass number of Sulphur:

Number of protons = 16

Number of neutrons = 16

Atomic number = 16

Atomic mass number = Number of protons + number of neutrons = 16 + 16 = 32

Exercise-4.6

Page: 53

14. For the symbol H, D and T, tabulate three subatomic particles found in each of them.

Solution:

The following table depicts the subatomic particles in Hydrogen (H), Deuterium (D), and Tritium(T).

Isotope	Symbol	Mass no.	Atomic no.	No. of electrons	No. of protons	No. of neutrons
Hydrogen	H	1	1	1	1	0
Deuterium	D	2	1	1	1	1
Tritium	T	3	1	1	1	2

15. Write the electronic configuration of any one pair of isotopes and isobar.

Solution:

(a) Isotopes: Isotopes are atoms which have the same number of protons but the number of neutrons differs. This leads to the variation in mass number too.

Example: Carbon molecule exists as ${}_6\text{C}^{12}$ and ${}_6\text{C}^{14}$ but when their electronic configuration is noticed, both have K-2; L-4

(b) Isobars: Isobars are atoms which have the same mass number but differ in the atomic number. Electronic configuration of an isobar pair is as follows,

Example: Electronic configuration of ${}_{20}\text{Ca}^{40}$ - K-2; L-8; M-8; N- 2

Electronic configuration of ${}_{18}\text{Ar}^{40}$ - K-2; L-8; M-8

Exercise

Page: 54

1. Compare the properties of electrons, protons and neutrons.

Solution:

Property	Electrons	Protons	Neutrons
Charge	Negatively charged	Positively charged	No charge.
Location	Located outside the nucleus	Located within the nucleus	Located inside the nucleus of an atom
Weight	Mass is negligible	1 a.m.u	1 a.m.u
Affinity	Attracted towards positively charged	Attracted towards negatively charged	Do not get attracted to any charged particle

2. What are the limitations of J.J.Thomson's model of the atom?

Solution:

The following are the limitations of the J.J. Thomson's model of an atom.

NCERT Solution for Class 9 Science - Chapter 4

Structure Of The Atom

- The model failed to explain the outcome of alpha particle scattering which was conducted by Rutherford. The model failed to depict why majority of these alpha particles pass through gold foil while some are diverted through small and big angles, while some others rebound completely, returning back on their path.
- It did not provide any experimental evidence and was established on imagination.

3. What are the limitations of Rutherford's model of the atom?

Solution:

Following are the limitations of Rutherford's model of the atom:


- There is no expected stability in the revolution of the electron in a circular orbit
- Charged particles radiate energy when accelerated thus causing the revolving electrons to lose energy and would fall into the nucleus
- Hence atoms must be highly unstable. Matter would not exist in their known form which clearly is an assumption as atoms are highly stable.

4. Describe Bohr's model of the atom.

Solution:

- An atom holds the nucleus at the centre.
- Negatively charged electrons revolve around the nucleus.
- The atoms in it contains distinct orbits of electrons
- Electrons do not radiate energy when they are in their orbits.
- The distinct orbits are named as K, L, M, N orbits. Numbers used to denote them are $n=1, 2, 3, 4$

NCERT Solution for Class 9 Science - Chapter 4 Structure Of The Atom



5. Compare all the proposed models of an atom given in this chapter.


Solution:

Thomson	Rutherford	Bohr


NCERT Solution for Class 9 Science - Chapter 4 Structure Of The Atom

<ul style="list-style-type: none">● Sphere is positively charged● Electrons are negatively charged and scattered all through the inside of the sphere.● Positively charged = negatively charged● The net charge in the atom is zero.	<ul style="list-style-type: none">● The nucleus is at the centre and is positively charged holding the entire mass.● Electrons are negatively charged revolving in a well-defined path● In comparison with the nucleus, the size of the atom is very large.	<ul style="list-style-type: none">● Nucleus is present at the centre and is positively charged● Electrons are negatively charged, revolving around but do not radiate energy.● The distinct orbits are labelled as K, L, M, N
---	---	---


Thomson's Model of Atom



Rutherford's Model of Atoms



Bohr's model of the atom



6. Summarise the rules for writing of distribution of electrons in various shells for the first eighteen elements.

Solution:

- Maximum number of electrons that can be accommodated in a shell is given by the formula: $2n^2$, where $n= 1, 2, 3\dots$
- Maximum number of electrons in different shells are:

K shell – $n=1$; $2n^2 = 2(1)^2 = 2$

L shell – $n=2$; $2n^2 = 2(2)^2 = 8$

M shell – $n=3$; $2n^2 = 2(3)^2 = 18$

N shell- $n=4$; $2n^2 = 2(4)^2 = 32$

- The outermost orbit can be accommodated with 8 electrons at the maximum.

NCERT Solution for Class 9 Science - Chapter 4 Structure Of The Atom

- The electrons are not taken in unless the inner shells are filled which are filled step-wise, hence the highest element has K-2; L-8 ; M-8 distribution of electrons.

7. Define valency by taking examples of silicon and oxygen.

Solution:

The definite combining capacity of the atoms of each element, wherein electrons are lost, gained or shared to make the octet of electrons present in the outermost shell is defined as valency. To measure valency, we can figure out the number of electrons that are required to complete the shell in which it is contained or losing excess electrons if present, once the filling is complete.

Example : To find the valency of silicon:

The atomic number of silicon is 14

Number of electrons is equal to the number of protons in silicon i.e., 14

The distribution of electrons in silicon atom is K - 2, L - 8, M - 4

Hence, from the distribution of silicon it is clearly evident that to fill the M shell 4 electrons are required. Therefore its valency is $8-4=4$.

To find the valency of oxygen:

The atomic number of oxygen is 8

Number of electrons is equal to the number of protons in oxygen i.e., 8

The distribution of electrons in oxygen atom is K - 2, L - 6

Hence, from the distribution of oxygen it is clearly evident that to fill the M shell 6 more electrons are required. Therefore its valency is $8-6=2$.

8. Explain with examples

(i) Atomic number,

(ii) Mass number,

(iii) Isotopes and

(iv) Isobars.

Give any two uses of isotopes.

Solution:

NCERT Solution for Class 9 Science - Chapter 4 Structure Of The Atom

(i) The number of positively charged protons present in the nucleus of an atom is defined as the atomic number and is denoted by Z . Example: Hydrogen has one proton in its nucleus, hence its atomic number is one.

(ii) The total number of protons and neutrons present in the nucleus of an atom is known as the mass number. It is denoted by A . ${}_{20}\text{Ca}^{40}$. Mass number is 40. Atomic number is 20.

(iii) The atoms which have the same number of protons but different number of neutrons are referred to as isotopes. Hence the mass number varies.

Example: The most simple example is the Carbon molecule which exists as ${}_{6}\text{C}^{12}$ and ${}_{6}\text{C}^{14}$

(d) Isobars: Isobars are atoms which have the same mass number but differ in the atomic number.

Examples are, ${}_{20}\text{Ca}^{40}$ and ${}_{18}\text{Ar}^{40}$

Uses of isotopes:

- The isotope of Iodine atom is used to treat goitre and iodine deficient disease.
- In the treatment of cancer, an isotope of cobalt is used.
- Fuel for nuclear reactors is derived from the isotopes of the Uranium atom.

9. Na^+ has completely filled K and L shells. Explain.

Solution:

The atomic number of sodium is 11. It has 11 electrons in its orbitals wherein the number of protons is equal to the number of electrons. Hence, its electronic configuration is K-2 ; L-8 ; M-1 ; The one electron in the M shell is lost and it obtains a positive charge since it has one more proton than electrons, and obtains a positive charge, Na^+ . The new electronic configuration is K-1 ; L-8 which is the filled state. Hence it is very difficult to eliminate the electron from a filled state as it is very stable.

10. If bromine atom is available in the form of, say, two isotopes ${}_{35}\text{Br}^{79}$ (49.7%) and ${}_{35}\text{Br}^{81}$ (50.3%), calculate the average atomic mass of Bromine atom.

Solution:

The atomic masses of two isotopic atoms are 79 (49.7%) and 81 (50.3%).

Thus, total mass = $(79 * 49.7 / 100) + (81 * 50.3 / 100) = 39.263 + 40.743 = 80.006 \text{ u}$

11. The average atomic mass of a sample of an element X is 16.2 u. What are the percentages of isotopes ${}_{8}\text{X}^{16}$ and ${}_{8}\text{X}^{18}$ in the sample?

Solution:

Let the percentage of ${}_{8}\text{X}^{16}$ be 'a' and that of ${}_{8}\text{X}^{18}$ be '100-a'.

As per given data,

$$16.2u = 16 a / 100 + 18 (100-a) / 100$$

$$1620 = 16a + 1800 - 18a$$

$$1620 = 1800 - 2a$$

$$a = 90\%$$

Hence, the percentage of isotope in the sample ${}_8\text{X}^{16}$ is 90% and that of

$${}_8\text{X}^{18} = 100 - a = 100 - 90 = 10\%$$

12. If $Z=3$, what would be the valency of the element? Also, name the element.

Solution:

Given: Atomic number, $Z = 3$

The electronic configuration of the element = K-2; L-1, hence its valency = 1

The element with atomic number 3 is Lithium.

13. Composition of the nuclei of two atomic species X and Y are given as under

	X	Y
Protons =	6	6
Neutrons =	6	8

Give the mass numbers of X and Y. What is the relation between the two species?

Solution:

Mass number of X: Protons + neutrons = $6+6 = 12$

Mass number of Y: Protons + neutrons = $6+8 = 14$

They are the same element as their atomic numbers are the same.

They are isotopes as they differ in the number of neutrons and hence their mass numbers.

14. For the following statements, write T for true and F for false.

- (a) J.J. Thomson proposed that the nucleus of an atom contains only nucleons.
- (b) A neutron is formed by an electron and a proton combining together. Therefore it is neutral.
- (c) The mass of an electron is about $1/2000$ times that of proton.

NCERT Solution for Class 9 Science - Chapter 4 Structure Of The Atom

(d) An isotope of iodine is used for making tincture iodine, which is used as a medicine.

Solution:

- (a) Statement is False
- (b) Statement is False
- (c) Statement is True
- (d) Statement is False

Put a tick(✓) against correct choice and cross(x) against wrong choice in questions 15, 16 and 17.

15. Rutherford's alpha – particle scattering experiment was responsible for the discovery of

- (a) Atomic nucleus
- (b) Electron
- (c) Proton
- (d) Neutron

Solution:

- (a) Atomic nucleus

16. Isotopes of an element have

- (a) The same physical properties
- (b) Different chemical properties
- (c) Different number of neutrons
- (d) Different atomic numbers.

Solution:

- (c) Different number of neutrons

17. Number of valence electrons in Cl⁻ ion are:

- (a) 16
- (b) 8

(c) 17

(d) 18

Solution:

(b) 8

Electronic distribution of Cl is K-2, L-8, M-7. Valence electrons are 7, hence chlorine gains one electron for the formation of Cl^- . Therefore, its valency is 8.

18. Which one of the following is a correct electronic configuration of Sodium?

(a) 2, 8

(b) 8, 2, 1

(c) 2, 1, 8

(d) 2, 8, 1

Solution:

(d) 2, 8, 1

19. Complete the following table.

Atomic number	Mass number	Number of neutrons	Number of Protons	Number of electrons	Name of the atomic species
9	—	10	—	—	—
16	32	—	—	—	Sulphur
—	24	—	12	—	—
—	2	—	1	—	—
—	1	0	1	0	—

NCERT Solution for Class 9 Science - Chapter 4 Structure Of The Atom

Solution:

The following table depicts the missing data:

Atomic number(Z) = Number of protons

Mass number = Number of neutrons + atomic number

(or)

Mass number(A) = Number of neutrons + number of neutrons

Atomic number	Mass number	Number of neutrons	Number of Protons	Number of electrons	Name of the atomic species
9	19	10	9	9	Fluorine
16	32	16	16	16	Sulphur
12	24	12	12	12	Magnesium
1	2	1	1	1	Deuterium
					Hydrogen
1	1	0	1	0	