

Exercise Solutions

Physics Class 11

Question 1: Find the dimensions of

- (i) Linear Momentum
- (ii) Frequency and
- (iii) Pressure

Solution:

(i)

Linear momentum can be written as “mv”

Dimensions of Linear momentum: $mv = [MLT^{-1}]$

(ii) Frequency can be written as “1/T”, where T is time.

Dimensions of Frequency: $1/T = [M^0L^0T^{-1}]$

(iii) All units of pressure represent some ratio of force to area

So, dimensions of pressure: $\text{Force/Area} = \frac{[MLT^{-2}]}{[L^2]} = [ML^{-1}T^{-2}]$

Question 2: Find the dimensions of

- (a) Angular speed, ω
- (b) Angular acceleration, α
- (c) Torque, τ and
- (d) Moment of Inertia, I

$$\omega = \frac{\theta_2 - \theta_1}{t_2 - t_1}, \quad \alpha = \frac{\omega_2 - \omega_1}{t_2 - t_1}, \quad \Gamma = F.r \text{ and } I = mr^2.$$

Some of the equations involving these quantities are.

Solution:

(a) Dimensions of Angular speed, ω

We know, $\omega = \theta/t$

Dimensions are $[M^0L^0T^{-1}]$

(b) Dimensions of Angular acceleration, α

We know, $\alpha = \omega/t$

So required dimensions are $= \frac{[M^0L^0T^{-1}]}{[T]} = [M^0L^0T^{-2}]$

[using (a) result)]

(c) Dimensions of Torque, τ and

We know, $\tau = Fr$

So, $\tau = [MLT^{-2}][L] = [ML^2T^{-2}]$

(d) Dimensions of Moment of Inertia, I

Here $I = Mr^2 = [M][L^2] = [ML^2T^0]$

Question 3: Find the dimensions of

(a) Electric Field E

(b) Magnetic field B and

(c) Magnetic permeability μ_0

The relevant equations are

$$F = qE, F = qvB, \text{ and } B = \frac{\mu_0 I}{2\pi a};$$

where F is force, q is charge, v is speed, I is current, and a is distance.

Solution:

(a) Dimensions of Electric Field $E = F/q = \frac{[MLT^{-2}]}{[TI]} = [MLT^{-3}I^{-1}]$

(b) Dimensions of Magnetic field $B = F/qv = \frac{[MLT^{-2}]}{[TI][LT^{-1}]} = [MT^{-2}I^{-1}]$

(c) Dimensions of Magnetic permeability $\mu_0 = (B \times 2 \pi a) / I = \frac{[MT^{-2}I^{-1}][L]}{[I]} = [MLT^{-2}I^{-2}]$

Question 4: Find the dimensions of

(a) Electric dipole moment p and

(b) Magnetic dipole moment M .

The defining equations are $p = qd$ and $M = IA$

Where d is distance, A is area, q is charge and I is current.

Solution:

a) Dimensions of Electric dipole moment $p = qd = [IT][L] = [LTI]$

(b) Dimensions of Magnetic dipole moment $M = IA = [I][L^2] = [L^2I]$

Question 5: Find the dimensions of Planck's constant h from the equation $E = hv$ where E is the energy and v is the frequency.

Solution:

Planck's constant can be written as, $h = E/v$

Where $E =$ energy and $v =$ frequency

$$\Rightarrow h = E/v = \frac{[ML^2T^{-2}]}{[T^{-1}]} = [ML^2 T^{-1}]$$

Question 6: Find the dimensions of

(a) the specific heat capacity c ,

(b) the coefficient of linear expansion α and

(c) the gas constant R .

Some of the equations involving these quantities are

$$Q = mc(T_2 - T_1), l_t = l_0[1 + \alpha(T_2 - T_1)] \text{ and } PV = nRT.$$

Solution:

(a) Dimensions of specific heat capacity,

$$c = Q/m\Delta T = \frac{[ML^2T^{-2}]}{[M][K]} = [L^2 T^{-2}K^{-1}]$$

(b) Dimensions of coefficient of linear expansion,

$$\alpha = \frac{L_1 - L_2}{L_0 \Delta T} = \frac{[L]}{[L][R]} = K^{-1}$$

(c) Dimensions of gas constant,

$$R = PV/nT = \frac{[ML^{-1}T^{-2}][L^3]}{[mol][K]} = [ML^2 T^{-2}K^{-1} (\text{mol})^{-1}]$$

Question 7: Taking force, length and time to be the fundamental quantities find the dimensions of

- (a) Density
- (b) Pressure
- (c) Momentum and
- (d) Energy

Solution:

As per given instruction, considering force, length and time to be the fundamental quantities

(a) Density = mass/volume ... (1)

and, mass = Force/acceleration = (Force × time²)/displacement.

(1) => Density = {(Force × time²)/displacement}/Volume

Dimensions of Density = [FL⁻⁴T²]

(b)

Pressure = F/A

Dimensions of A = [L²]

Dimensions of Pressure = [FL⁻²]

(c)

Momentum = mv = (force/acceleration) × velocity

= [F/LT²] × [LT⁻¹] = [FT]

Dimensions of Momentum [FT]

(d)

$$\text{Energy} = \frac{1}{2} mv^2 = \text{Force/acceleration} \times (\text{velocity})^2$$

$$= \frac{[F]}{[LT^{-2}]} [LT^{-1}]^2 = [FL]$$

Dimensions of Energy = [FL]

Question 8: Suppose the acceleration due to gravity at a place is 10 m/s^2 . Find its value in $\text{cm}/(\text{minute})^2$.

Solution:

Given: acceleration due to gravity at a place is 10 m/s^2
Convert units into cm/min^2

$$\text{Here, } g = 10 \text{ m/sec}^2 = 36 \times 10^5 \text{ cm/min}^2$$

Question 9: The average speed of a snail is 0.020 miles/hour and that of a leopard is 70 miles/hour . Convert these speeds in SI units.

Solution:

Average speed of a snail = 0.020 miles/hour (Given)

Average speed of a leopard = 70 miles/hour (Given)

In SI Units:

$$0.020 \text{ miles/hour} = (0.02 \times 1.6 \times 1000) / 3600 = 0.0089 \text{ m/s}$$

[Using $1 \text{ mile} = 1.6 \text{ km} = 1600 \text{ m}$]

And,

$$70 \text{ miles/hr} = (70 \times 1.6 \times 1000) / 3600 = 31 \text{ m/s}$$

Question 10: The height of mercury column in a barometer in a Calcutta laboratory was recorded to be 75 cm . Calculate this pressure in SI and CGS units using the following data.

Specific gravity of mercury = 13.6

Density of water = 10^3 kg/m^3 ,

gravity, $g = 9.8 \text{ m/s}^2$ at Calcutta.
Pressure = $h\rho g$ in usual symbols.

Solution:

The height of mercury column in a barometer in a Calcutta laboratory was recorded to be 75 cm. (Given)

Say, $h = 75 \text{ cm}$

Calculate pressure in SI and CGS units.

Pressure = $h\rho g = 10 \times 10^4 \text{ N/m}^2$ approx

In C.G.S. units, Pressure = $10 \times 10^5 \text{ dyne/cm}^2$

Question 11: Express the power of a 100 watt bulb in CGS unit.

Solution:

Write power 100 watt in CGS units.

In S.I. units: 100 watt = 100 Joule/sec

In C.G.S. unit = 10^9 erg/sec

Question 12: The normal duration of I.Sc. Physics practical period in Indian colleges is 100 minutes. Express this period in microcenturies. 1 microcentury = $10^{-6} \times 100$ years. How many microcenturies did you sleep yesterday?

Solution:

Given: 1 microcentury = $10^{-6} \times 100$ years.

1 year = $365 \times 24 \times 60 \text{ min}$

Or 1 microcentury = $10^{-4} \times 365 \times 24 \times 60 \text{ min}$

So, 100 min = $10^5 / 52560 = 1.9$ microcentury

Question 13: The surface tension of water is 72 dyne/cm. Convert it in SI units.

Solution:

Given: surface tension of water is 72 dyne/cm

In S.I units: 72 dyne/cm = 0.072 N/m

Question 14: The kinetic energy K of a rotating body depends on its moment of inertia I and its angular speed ω . Assuming the relation to be $K = kI^a \omega^b$ where k is a dimensionless constant, find a and b . Moment of inertia of a sphere about its diameter is $\frac{2}{5} Mr^2$.

Solution:

$K = kI^a \omega^b$; where k is a dimensionless constant, K = kinetic energy and ω = angular speed

To find: a and b

Now, $K = [ML^2T^{-2}]$

$$I^a = [ML^2]^a \text{ and } \omega^b = [T^{-1}]^b$$

$$\Rightarrow [ML^2T^{-2}] = [ML^2]^a [T^{-1}]^b$$

[using principle of homogeneity of dimension]

Equating the dimensions, we get

$$2a = 2 \Rightarrow a = 1$$

$$-b = -2 \Rightarrow b = 2$$

Question 15: Theory of relativity reveals that mass can be converted into energy. The energy E so obtained is proportional to certain powers of mass m and the speed c of light. Guess a relation among the quantities using the method of dimensions.

Solution:

The relationship between energy, mass and speed of light is,

$$E \propto M^a C^b$$

Where M = mass and C = speed of light

$$\text{or } E = K M^a C^b \dots\dots(1)$$

where K = constant of proportionality

Find the dimensions of (1)

$$[ML^2T^{-2}] = [M]^a [LT^{-1}]^b$$

By comparing values, we have

$a = 1$ and $b = 2$

So, we have required relation is $E = KMC^2$

Question 16: Let I = current through a conductor, R = its resistance and V = potential difference across its ends. According to Ohm's law, product of two of these quantities equals the third. Obtain Ohm's law from dimensional analysis. Dimensional formulae for R and V are $[ML^2I^{-2}T^{-3}]$ and $[ML^2T^{-3}I^{-1}]$ respectively.

Solution:

Given:

Dimensional formulae for $R = [ML^2I^{-2}T^{-3}]$ and

Dimensional formulae for $V = [ML^2T^{-3}I^{-1}]$

Therefore,

$$[ML^2T^{-3}I^{-1}] = [ML^2I^{-2}T^{-3}] [I]$$

$$\Rightarrow V = IR$$

Question 17: The frequency of vibration of a string depends on the length L between the nodes, the tension F in the string and its mass per unit length m . Guess the expression for its frequency from dimensional analysis.

Solution:

L = length, M = mass and F = Force

$$\text{Here, } f = KL^aF^bM^c \dots(1)$$

Dimension of frequency, $f = [T^{-1}]$ or $[M^0L^0T^{-1}]$

Dimension of length, $L = [L]$

Dimension of mass, $M = [ML^{-1}]$

Dimension of force, $F = [MLT^{-2}]$

$$(1) \Rightarrow [M^0L^0T^{-1}] = K [L]^a [MLT^{-2}]^b [ML^{-1}]^c$$

Equating both sides, we get

$$b + c = 0$$

$$-c + a + b = 0$$

$$-2b = -1$$

Solving above three equations, we have

$$a = -1, b = \frac{1}{2} \text{ and } c = -1/2$$

Therefore, frequency is

$$f = KL^{-1}F^{1/2}M^{-1/2}$$

$$f = KL^{-1}F^{1/2}M^{-1/2} = \frac{K}{L} \sqrt{\frac{F}{M}}$$

Question 18: Test if the following equations are dimensionally correct:

$$(a) h = \frac{2S \cos\theta}{\rho g},$$

$$(b) v = \sqrt{\frac{P}{\rho}},$$

$$(c) V = \frac{\pi P r^4 t}{8 \eta l},$$

$$(d) v = \frac{1}{2\pi} \sqrt{\frac{mgl}{I}};$$

where h = height, S = surface tension, ρ = density, P = pressure, V = volume, η = coefficient of viscosity, v = frequency and I = moment of inertia

Solution:

(a) Dimension of $h = [L]$

Dimension of $S = F/l = MLT^{-2}/L = [MT^{-2}]$

Dimension of density = $M/V = [ML^{-3}T^0]$

Dimension of radius = $[L]$

Dimension of gravity = $[LT^{-2}]$

Now,

$$\frac{2S \cos\theta}{\rho g} = \frac{[MT^{-2}]}{[ML^{-3}T^0][L][LT^{-2}]} = [M^0L^1T^0] = [L]$$

Relation is correct.

(b) Let velocity = $v = \sqrt{p/\rho}$ (1)

Dimension of $v = [LT^{-1}]$

Dimension of $p = F/A = [ML^{-1}T^{-2}]$

Dimension of $\rho = m/v = [ML^{-3}]$

Substituting dimensions in (1), we have

$$\sqrt{\frac{\rho}{\rho}} = \sqrt{\frac{[ML^{-1}T^{-2}]}{[ML^{-3}]}} = [L^2T^{-2}]^{1/2} = [LT^{-1}]$$

Therefore, relation is correct.

(c)

Dimension of $V = [L^3]$

Dimension of $\rho = [ML^{-1}T^{-2}]$

Dimension of $r^4 = [L^4]$

Dimension of $t = [T]$

Dimension of $\eta = [ML^{-1}T^{-1}]$

$$\frac{\pi \rho r^4 t}{8 \eta l} = \frac{[ML^{-1}T^{-2}][L^4][T]}{[ML^{-1}T^{-1}][L]}$$

Therefore, relation is correct.

(d)

Dimension of $v = [T^{-1}]$

Dimension of $m = [M]$

Dimension of $g = [LT^{-2}]$

Dimension of $l = [L]$

Dimension of inertia = $[ML^2]$

$$\sqrt{(mgl/l)} = \sqrt{\frac{[M][LT^{-2}][L]}{[ML^2]}} = [T^{-1}]$$

Therefore, relation is correct.

Question 19: Let x and a stand for distance. Is below equation dimensionally correct?

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \frac{1}{a} \sin^{-1} \frac{a}{x}$$

Solution:

Dimensions of $a = [L]$

Dimensions of $x = [L]$

LHS

$$\int \frac{dx}{\sqrt{(a^2 - x^2)}} = \int \frac{L}{\sqrt{(L^2 - L^2)}} = [L^0]$$

RHS

$$\frac{1}{a} \sin^{-1}\left(\frac{a}{x}\right) = [L^{-1}]$$

$$\int \frac{dx}{\sqrt{(a^2 - x^2)}} \neq \frac{1}{a} \sin^{-1}\left(\frac{a}{x}\right)$$

