

Exercise 6.1

Page: 122

1. Fill in the blanks using correct v	word given in the brackets:-
--	------------------------------

(i) All circles are _____. (congruent, similar) Answer: Similar

(ii) All squares are _____. (similar, congruent) Answer: Similar

(iii) All ______ triangles are similar. (isosceles, equilateral) Answer: Equilateral


(iv) Two polygons of the same number of sides are similar, if (a) their corresponding angles are ______ and (b) their corresponding sides are ______. (equal, proportional)


Answer: (a) Equal (b) Proportional

2. Give two different examples of pair of(i) Similar figures(ii) Non-similar figures

Solution:


(i)Example of two similar figure;







Two Rectangle

(ii) Example of two Non-similar figure;








Rectangle

Trapezium



3. State whether the following quadrilaterals are similar or not:



Solution: From the given two figures, we can see their corresponding angles are different or unequal. Therefore they are not similar.






Exercise 6.2

Page: 128

1. In figure. (i) and (ii), DE || BC. Find EC in (i) and AD in (ii).



Solution:


(i) Given, in \triangle ABC, DE||BC \therefore AD/DB = AE/EC [Using Basic proportionality theorem] $\Rightarrow 1.5/3 = 1/EC$ $\Rightarrow EC = 3/1.5$ EC = $3 \times 10/15 = 2$ cm Hence, EC = 2 cm.

(ii) Given, in \triangle ABC, DE||BC \therefore AD/DB = AE/EC [Using Basic proportionality theorem] \Rightarrow AD/7.2 = 1.8 / 5.4 \Rightarrow AD = 1.8 ×7.2/5.4 = (18/10)×(72/10)×(10/54) = 24/10 \Rightarrow AD = 2.4 Hence, AD = 2.4 cm.

2. E and F are points on the sides PQ and PR respectively of a \triangle PQR. For each of the following cases, state whether EF || QR. (i) PE = 3.9 cm, EQ = 3 cm, PF = 3.6 cm and FR = 2.4 cm (ii) PE = 4 cm, QE = 4.5 cm, PF = 8 cm and RF = 9 cm (iii) PQ = 1.28 cm, PR = 2.56 cm, PE = 0.18 cm and PF = 0.63 cm Solution:

Given, in $\triangle PQR$, E and F are two points on side PQ and PR respectively. See the figure below;






(i) Given, PE = 3.9 cm, EQ = 3 cm, PF = 3.6 cm and FR = 2,4 cm Therefore, by using Basic proportionality theorem, we get, PE/EQ = 3.9/3 = 39/30 = 13/10 = 1.3And PF/FR = 3.6/2.4 = 36/24 = 3/2 = 1.5So, we get, PE/EQ \neq PF/FR Hence, EF is not parallel to QR.

(ii) Given, PE = 4 cm, QE = 4.5 cm, PF = 8cm and RF = 9cm Therefore, by using Basic proportionality theorem, we get, PE/QE = 4/4.5 = 40/45 = 8/9And, PF/RF = 8/9So, we get here, PE/QE = PF/RFHence, EF is parallel to QR.

(iii) Given, PQ = 1.28 cm, PR = 2.56 cm, PE = 0.18 cm and PF = 0.36 cm From the figure, EQ = PQ - PE = 1.28 - 0.18 = 1.10 cm And, FR = PR - PF = 2.56 - 0.36 = 2.20 cm So, PE/EQ = 0.18/1.10 = 18/110 = 9/55.....(i) And, PE/FR = 0.36/2.20 = 36/220 = 9/55.....(ii) So, we get here, PE/EQ = PF/FRHence, EF is parallel to QR.

3. In the figure, if LM || CB and LN || CD, prove that AM/MB = AN/AD B



Solution:



In the given figure, we can see, $LM \parallel CB$, By using basic proportionality theorem, we get, AM/MB = AL/LC.....(i)


Similarly, given, LN || CD and using basic proportionality theorem,

 \therefore AN/AD = AL/LC.....(ii)

From equation (i) and (ii), we get, AM/MB = AN/AD

Hence, proved.

4. In the figure, DE||AC and DF||AE. Prove that BF/FE = BE/EC



Solution:

In \triangle ABC, given as, DE || AC Thus, by using Basic Proportionality Theorem, we get,

 \therefore BD/DA = BE/EC(i)


In $\triangle ABC$, given as, DF || AE Thus, by using Basic Proportionality Theorem, we get,

 \therefore BD/DA = BF/FE(ii)

From equation (i) and (ii), we get BE/EC = BF/FE Hence, proved.

5. In the figure, DE||OQ and DF||OR, show that EF||QR.






Solution:

Given, In \triangle PQO, DE || OQ So by using Basic Proportionality Theorem, PD/DO = PE/EQ(i)

Again given, in \triangle PQO, DE || OQ , So by using Basic Proportionality Theorem, PD/DO = PF/FR(ii)

From equation (i) and (ii), we get, PE/EQ = PF/FRTherefore, by converse of Basic Proportionality Theorem, $EF \parallel QR$, in ΔPQR .

6. In the figure, A, B and C are points on OP, OQ and OR respectively such that AB || PQ and AC || PR. Show that BC || QR.



Solution:

Given here, In $\triangle OPQ$, AB || PQ By using Basic Proportionality Theorem,


B BYJU'S The Learning App NCERT Solutions for Class 10 Maths Chapter 6 – Triangles

OA/AP = OB/BQ....(i)Also given, In $\triangle OPR$, $AC \parallel PR$ By using Basic Proportionality Theorem $\therefore OA/AP = OC/CR...(ii)$

From equation (i) and (ii), we get, OB/BQ = OC/CR

Therefore, by converse of Basic Proportionality Theorem, In ΔOQR , BC || QR.

7. Using Basic proportionality theorem, prove that a line drawn through the mid-points of one side of a triangle parallel to another side bisects the third side. (Recall that you have proved it in Class IX).



Solution:

Given, in $\triangle ABC$, D is the midpoint of AB such that AD=DB. A line parallel to BC intersects AC at E as shown in above figure such that DE || BC.

We have to prove that E is the mid point of AC.

Since, D is the mid-point of AB. \therefore AD=DB \Rightarrow AD/DB = 1(i) In \triangle ABC, DE || BC, By using Basic Proportionality Theorem, Therefore, AD/DB = AE/EC From equation (i), we can write, \Rightarrow 1 = AE/EC

 $\Rightarrow 1 = AE/EC$ $\therefore AE = EC$


Hence, proved, E is the midpoint of AC.

8. Using Converse of basic proportionality theorem, prove that the line joining the mid-points of any two sides of a triangle is parallel to the third side. (Recall that you have done it in Class IX).



Solution:

Given, in $\triangle ABC$, D and E are the mid points of AB and AC respectively, such that, AD=BD and AE=EC.




9. ABCD is a trapezium in which AB || DC and its diagonals intersect each other at the point O. Show that AO/BO = CO/DO.

Solution:

Given, ABCD is a trapezium where AB || DC and diagonals AC and BD intersect each other at O.





We have to prove, AO/BO = CO/DO

From the point O, draw a line EO touching AD at E, in such a way that, EO \parallel DC \parallel AB

In \triangle ADC, we have OE || DC Therefore, By using Basic Proportionality Theorem AE/ED = AO/CO(i)

Now, In $\triangle ABD$, OE || AB Therefore, By using Basic Proportionality Theorem

DE/EA = DO/BO.....(ii)


From equation (i) and (ii), we get, AO/CO = BO/DO $\Rightarrow AO/BO = CO/DO$ Hence, proved.

10. The diagonals of a quadrilateral ABCD intersect each other at the point O such that AO/BO = CO/DO. Show that ABCD is a trapezium.

Solution:

Given, Quadrilateral ABCD where AC and BD intersects each other at O such that, AO/BO = CO/DO.






We have to prove here, ABCD is a trapezium

From the point O, draw a line EO touching AD at E, in such a way that, EO \parallel DC \parallel AB

In ΔDAB , EO || AB Therefore, By using Basic Proportionality Theorem DE/EA = DO/OB(i) Also, given, AO/BO = CO/DO

 $\Rightarrow AO/CO = BO/DO$ $\Rightarrow CO/AO = DO/BO$ $\Rightarrow DO/OB = CO/AO$ (ii) From equation (i) and (ii), we get DE/EA = CO/AO

Therefore, By using converse of Basic Proportionality Theorem, EO || DC also EO || AB \Rightarrow AB || DC. Hence, quadrilateral ABCD is a trapezium with AB || CD.






Exercise 6.3

Page: 138

1. State which pairs of triangles in Figure, are similar. Write the similarity criterion used by you for answering the question and also write the pairs of similar triangles in the symbolic form:



Solution:

(i) Given, in $\triangle ABC$ and $\triangle PQR$, $\angle A = \angle P = 60^{\circ}$ $\angle B = \angle Q = 80^{\circ}$ $\angle C = \angle R = 40^{\circ}$ Therefore by AAA similarity criterion, $\therefore \triangle ABC \sim \triangle PQR$

(ii) Given, in $\triangle ABC$ and $\triangle PQR$,



AB/QR = BC/RP = CA/PQ

By SSS similarity criterion, $\Delta ABC \sim \Delta QRP$

(iii) Given, in Δ LMP and Δ DEF,

LM = 2.7, MP = 2, LP = 3, EF = 5, DE = 4, DF = 6 MP/DE = 2/4 = 1/2PL/DF = 3/6 = 1/2LM/EF = 2.7/5 = 27/50Here , MP/DE = PL/DF \neq LM/EF

Therefore, Δ LMP and Δ DEF are not similar.

(iv) In Δ MNL and Δ QPR, it is given, MN/QP = ML/QR = 1/2 \angle M = \angle Q = 70° Therefore, by SAS similarity criterion $\therefore \Delta$ MNL $\sim \Delta$ QPR

(v) In $\triangle ABC$ and $\triangle DEF$, given that, AB = 2.5, BC = 3, $\angle A = 80^{\circ}$, EF = 6, DF = 5, $\angle F = 80^{\circ}$ Here, AB/DF = 2.5/5 = 1/2And, BC/EF = 3/6 = 1/2 $\Rightarrow \angle B \neq \angle F$ Hence, $\triangle ABC$ and $\triangle DEF$ are not similar.


(vi) In $\triangle DEF$, by sum of angles of triangles, we know that, $\angle D + \angle E + \angle F = 180^{\circ}$ $\Rightarrow 70^{\circ} + 80^{\circ} + \angle F = 180^{\circ}$ $\Rightarrow \angle F = 180^{\circ} - 70^{\circ} - 80^{\circ}$ $\Rightarrow \angle F = 30^{\circ}$

Similarly, In $\triangle PQR$, $\angle P + \angle Q + \angle R = 180$ (Sum of angles of \triangle) $\Rightarrow \angle P + 80^{\circ} + 30^{\circ} = 180^{\circ}$ $\Rightarrow \angle P = 180^{\circ} - 80^{\circ} - 30^{\circ}$ $\Rightarrow \angle P = 70^{\circ}$

Now, comparing both the triangles, ΔDEF and ΔPQR , we have $\angle D = \angle P = 70^{\circ}$ $\angle F = \angle Q = 80^{\circ}$ $\angle F = \angle R = 30^{\circ}$ Therefore, by AAA similarity criterion, Hence, $\Delta DEF \sim \Delta PQR$



2. In the figure, $\triangle ODC \propto \frac{1}{4} \triangle OBA$, $\angle BOC = 125^{\circ}$ and $\angle CDO = 70^{\circ}$. Find $\angle DOC$, $\angle DCO$ and $\angle OAB$.



Solution:


As we can see from the figure, DOB is a straight line. Therefore, $\angle DOC + \angle COB = 180^{\circ}$ $\Rightarrow \angle DOC = 180^{\circ} - 125^{\circ}$ (Given, $\angle BOC = 125^{\circ}$) $= 55^{\circ}$

In $\triangle DOC$, Sum of the measures of the angles of a triangle is 180° Therefore, $\angle DCO + \angle CDO + \angle DOC = 180^{\circ}$ $\Rightarrow \angle DCO + 70^{\circ} + 55^{\circ} = 180^{\circ}$ (Given, $\angle CDO = 70^{\circ}$) $\Rightarrow \angle DCO = 55^{\circ}$

It is given that, $\triangle ODC \propto \frac{1}{4} \triangle OBA$, Therefore, $\triangle ODC \sim \triangle OBA$. Hence, Corresponding angles are equal in similar triangles $\angle OAB = \angle OCD$ $\Rightarrow \angle OAB = 55^{\circ}$ $\angle OAB = \angle OCD$ $\Rightarrow \angle OAB = 55^{\circ}$

3. Diagonals AC and BD of a trapezium ABCD with AB || DC intersect each other at the point O. Using a similarity criterion for two triangles, show that AO/OC = OB/OD

Solution:



In $\triangle DOC$ and $\triangle BOA$, AB || CD, thus alternate interior angles will be equal,



 $\therefore \angle CDO = \angle ABO$ Similarly, $\angle DCO = \angle BAO$ Also, for the two triangles $\triangle DOC$ and $\triangle BOA$, vertically opposite angles will be equal; $\therefore \angle DOC = \angle BOA$

Hence, by AAA similarity criterion, $\Delta DOC \sim \Delta BOA$ Thus, the corresponding sides are proportional. DO/BO = OC/OA $\Rightarrow OA/OC = OB/OD$

Hence, proved.

4. In the fig.6.36, QR/QS = QT/PR and $\angle 1 = \angle 2$. Show that $\triangle PQS \sim \triangle TQR$.

Q 1 2 R

Solution:

In $\triangle PQR$, $\angle PQR = \angle PRQ$ $\therefore PQ = PR$ (i) Given, QR/QS = QT/PRUsing equation (i), we get QR/QS = QT/QP....(ii)


In $\triangle PQS$ and $\triangle TQR$, by equation (ii), QR/QS = QT/QP $\angle Q = \angle Q$ $\therefore \triangle PQS \sim \triangle TQR$ [By SAS similarity criterion]

5. S and T are point on sides PR and QR of \triangle PQR such that \angle P = \angle RTS. Show that \triangle RPQ ~ \triangle RTS.

Solution:




Given, S and T are point on sides PR and QR of \triangle PQR And \angle P = \angle RTS.



In $\triangle RPQ$ and $\triangle RTS$, $\angle RTS = \angle QPS$ (Given) $\angle R = \angle R$ (Common angle) $\therefore \triangle RPQ \sim \triangle RTS$ (AA similarity criterion)



6. In the figure, if $\triangle ABE \cong \triangle ACD$, show that $\triangle ADE \sim \triangle ABC$.



Solution:


Given, $\triangle ABE \cong \triangle ACD$. $\therefore AB = AC [By CPCT]$(i) And, AD = AE [By CPCT].....(ii)

In $\triangle ADE$ and $\triangle ABC$, dividing eq.(ii) by eq(i), AD/AB = AE/AC $\angle A = \angle A$ [Common angle]

 $\therefore \Delta ADE \sim \Delta ABC \text{ [SAS similarity criterion]}$

7. In the figure, altitudes AD and CE of ΔABC intersect each other at the point P. Show that:

B BYJU'S The Learning App NCERT Solutions for Class 10 Maths Chapter 6 – Triangles



(i) $\triangle AEP \sim \triangle CDP$ (ii) $\triangle ABD \sim \triangle CBE$ (iii) $\triangle AEP \sim \triangle ADB$ (iv) $\triangle PDC \sim \triangle BEC$

Solution:

Given, altitudes AD and CE of \triangle ABC intersect each other at the point P.

(i) In $\triangle AEP$ and $\triangle CDP$, $\angle AEP = \angle CDP$ (90° each) $\angle APE = \angle CPD$ (Vertically opposite angles) Hence, by AA similarity criterion, $\triangle AEP \sim \triangle CDP$

(ii) In $\triangle ABD$ and $\triangle CBE$, $\angle ADB = \angle CEB$ (90° each) $\angle ABD = \angle CBE$ (Common Angles) Hence, by AA similarity criterion, $\triangle ABD \sim \triangle CBE$


(iii) In $\triangle AEP$ and $\triangle ADB$, $\angle AEP = \angle ADB$ (90° each) $\angle PAE = \angle DAB$ (Common Angles) Hence, by AA similarity criterion, $\triangle AEP \sim \triangle ADB$ (iv) In $\triangle PDC$ and $\triangle BEC$, $\angle PDC = \angle BEC$ (90° each) $\angle PCD = \angle BCE$ (Common angles) Hence, by AA similarity criterion, $\triangle PDC \sim \triangle BEC$

8. E is a point on the side AD produced of a parallelogram ABCD and BE intersects CD at F. Show that $\triangle ABE \sim \triangle CFB$.

Solution:




Given, E is a point on the side AD produced of a parallelogram ABCD and BE intersects CD at F. Consider the figure below,



In $\triangle ABE$ and $\triangle CFB$, $\angle A = \angle C$ (Opposite angles of a parallelogram) $\angle AEB = \angle CBF$ (Alternate interior angles as $AE \parallel BC$) $\therefore \triangle ABE \sim \triangle CFB$ (AA similarity criterion)

9. In the figure, ABC and AMP are two right triangles, right angled at B and M respectively, prove that:



(i) $\triangle ABC \sim \triangle AMP$ (ii) CA/PA = BC/MP

Solution:

Given, ABC and AMP are two right triangles, right angled at B and M respectively.

(i) In $\triangle ABC$ and $\triangle AMP$, we have, $\angle CAB = \angle MAP$ (common angles) $\angle ABC = \angle AMP = 90^{\circ}$ (each 90°) $\therefore \triangle ABC \sim \triangle AMP$ (AA similarity criterion)

(ii) As, $\triangle ABC \sim \triangle AMP$ (AA similarity criterion) If two triangles are similar then the corresponding sides are always equal, Hence, CA/PA = BC/MP


B BYJU'S

NCERT Solutions for Class 10 Maths Chapter 6 – Triangles

10. CD and GH are respectively the bisectors of ∠ACB and ∠EGF such that D and H lie on sides AB and FE of ΔABC and ΔEFG respectively. If ΔABC ~ ΔFEG, Show that: (i) CD/GH = AC/FG


(ii) ΔDCB ~ ΔHGE(iii) ΔDCA ~ ΔHGF

Solution: Given, CD and GH are respectively the bisectors of $\angle ACB$ and $\angle EGF$ such that D and H lie on sides AB and FE of $\triangle ABC$ and $\triangle EFG$ respectively.




11. In the following figure, E is a point on side CB produced of an isosceles triangle ABC with AB = AC. If AD \perp BC and EF \perp AC, prove that \triangle ABD ~ \triangle ECF.

BYJU'S The Learning App NCERT Solutions for Class 10 Maths Chapter 6 – Triangles



Solution: Given, ABC is an isosceles triangle. \therefore AB = AC $\Rightarrow \angle$ ABD = \angle ECF In \triangle ABD and \triangle ECF, \angle ADB = \angle EFC (Each 90°) \angle BAD = \angle CEF (Already proved) $\therefore \triangle$ ABD ~ \triangle ECF (using AA similarity criterion)

12. Sides AB and BC and median AD of a triangle ABC are respectively proportional to sides PQ and QR and median PM of Δ PQR (see Fig 6.41). Show that Δ ABC ~ Δ PQR.



Solution: Given, $\triangle ABC$ and $\triangle PQR$, AB, BC and median AD of $\triangle ABC$ are proportional to sides PQ, QR and median PM of $\triangle PQR$

i.e. AB/PQ = BC/QR = AD/PM

We have to prove: $\triangle ABC \sim \triangle PQR$

As we know here, AB/PQ = BC/QR = AD/PM

 $\frac{AB}{PQ} = \frac{\frac{1}{2}BC}{\frac{1}{2}QR} = \frac{AD}{PM}.$ (i)


 $\Rightarrow AB/PQ = BC/QR = AD/PM \text{ (D is the midpoint of BC. M is the midpoint of QR)}$ $\Rightarrow \Delta ABD \sim \Delta PQM \text{ [SSS similarity criterion]}$ $\therefore \angle ABD = \angle PQM \text{ [Corresponding angles of two similar triangles are equal]}$ $\Rightarrow \angle ABC = \angle PQR$



In $\triangle ABC$ and $\triangle PQR$ AB/PQ = BC/QR(i) $\angle ABC = \angle PQR$ (ii) From equation (i) and (ii), we get, $\triangle ABC \sim \triangle PQR$ [SAS similarity criterion]

13. D is a point on the side BC of a triangle ABC such that $\angle ADC = \angle BAC$. Show that $CA^2 = CB.CD$

Solution: Given, D is a point on the side BC of a triangle ABC such that $\angle ADC = \angle BAC$.




14. Sides AB and AC and median AD of a triangle ABC are respectively proportional to sides PQ and PR and median PM of another triangle PQR. Show that \triangle ABC ~ \triangle PQR.

Solution: Given: Two triangles $\triangle ABC$ and $\triangle PQR$ in which AD and PM are medians such that; AB/PQ = AC/PR = AD/PM

We have to prove, $\triangle ABC \sim \triangle PQR$

Let us construct first: Produce AD to E so that AD = DE. Join CE, Similarly produce PM to N such that PM = MN, also Join RN.





In $\triangle ABD$ and $\triangle CDE$, we have AD = DE [By Construction.] BD = DC [Since, AP is the median] and, $\angle ADB = \angle CDE$ [Vertically opposite angles] $\therefore \triangle ABD \cong \triangle CDE$ [SAS criterion of congruence] $\Rightarrow AB = CE$ [By CPCT](i)

Also, in ΔPQM and ΔMNR , PM = MN [By Construction.] QM = MR [Since, PM is the median] and, $\angle PMQ = \angle NMR$ [Vertically opposite angles] $\therefore \Delta PQM = \Delta MNR$ [SAS criterion of congruence] $\Rightarrow PQ = RN$ [CPCT](ii)

Now, AB/PQ = AC/PR = AD/PM From equation (i) and (ii), \Rightarrow CE/RN = AC/PR = AD/PM \Rightarrow CE/RN = AC/PR = 2AD/2PM \Rightarrow CE/RN = AC/PR = AE/PN [Since 2AD = AE and 2PM = PN] $\therefore \Delta ACE \sim \Delta PRN$ [SSS similarity criterion] Therefore, $\angle 2 = \angle 4$ Similarly, $\angle 1 = \angle 3$ $\therefore \angle 1 + \angle 2 = \angle 3 + \angle 4$ $\Rightarrow \angle A = \angle P$(iii)


Now, In $\triangle ABC$ and $\triangle PQR$, we have AB/PQ = AC/PR (Already given) From equation (iii), $\angle A = \angle P$ $\therefore \triangle ABC \sim \triangle PQR$ [SAS similarity criterion]



15. A vertical pole of a length 6 m casts a shadow 4m long on the ground and at the same time a tower casts a shadow 28 m long. Find the height of the tower.

Solution:


Given, Length of the vertical pole = 6mShadow of the pole = 4mLet Height of tower = hmLength of shadow of the tower = 28m



In $\triangle ABC$ and $\triangle DEF$, $\angle C = \angle E$ (angular elevation of sum) $\angle B = \angle F = 90^{\circ}$ $\therefore \triangle ABC \sim \triangle DEF$ (AA similarity criterion) $\therefore AB/DF = BC/EF$ (If two triangles are similar corresponding sides are proportional)

 $\therefore 6/h = 4/28$ $\Rightarrow h = (6 \times 28)/4$ $\Rightarrow h = 6 \times 7$ $\Rightarrow h = 42 \text{ m}$ Hence, the height of the tower is 42 m.

16. If AD and PM are medians of triangles ABC and PQR, respectively where $\triangle ABC \sim \triangle PQR$ prove that AB/PQ = AD/PM.



BYJU'S

NCERT Solutions for Class 10 Maths Chapter 6 – Triangles

We know that the corresponding sides of similar triangles are in proportion.

 $\therefore AB/PQ = AC/PR = BC/QR....(i)$ $Also, <math>\angle A = \angle P, \angle B = \angle Q, \angle C = \angle R$ (ii) Since AD and PM are medians, they will divide their opposite sides. $\therefore BD = BC/2 \text{ and } QM = QR/2 \dots (iii)$ From equations (i) and (iii), we get $AB/PQ = BD/QM \dots (iv)$

In $\triangle ABD$ and $\triangle PQM$, From equation (ii), we have $\angle B = \angle Q$ From equation (iv), we have, AB/PQ = BD/QM

 $\therefore \Delta ABD \sim \Delta PQM \text{ (SAS similarity criterion)}$ $\Rightarrow AB/PQ = BD/QM = AD/PM$



Exercise 6.4

Page: 143

1. Let $\triangle ABC \sim \triangle DEF$ and their areas be, respectively, 64 cm² and 121 cm². If EF = 15.4 cm, find BC.

Solution: Given, $\triangle ABC \sim \triangle DEF$, Area of $\triangle ABC = 64 \text{ cm}^2$ Area of $\triangle DEF = 121 \text{ cm}^2$ EF = 15.4 cm


 $\therefore \frac{Area \ of \ \Delta ABC}{Area \ of \ \Delta DEF} = \frac{AB^2}{DE^2}$

As we know, if two triangles are similar, ratio of their areas are equal to the square of the ratio of their corresponding sides, = $AC^2/DF^2 = BC^2/EF^2$

 $\therefore 64/121 = BC^2/EF^2$ $\Rightarrow (8/11)^2 = (BC/15.4)^2$ $\Rightarrow 8/11 = BC/15.4$ $\Rightarrow BC = 8 \times 15.4/11$ $\Rightarrow BC = 8 \times 1.4$ $\Rightarrow BC = 11.2 \text{ cm}$

2. Diagonals of a trapezium ABCD with AB || DC intersect each other at the point O. If AB = 2CD, find the ratio of the areas of triangles AOB and COD.

Solution: Given, ABCD is a trapezium with AB || DC. Diagonals AC and BD intersect each other at point O.



In $\triangle AOB$ and $\triangle COD$, we have $\angle 1 = \angle 2$ (Alternate angles) $\angle 3 = \angle 4$ (Alternate angles) $\angle 5 = \angle 6$ (Vertically opposite angle) $\therefore \triangle AOB \sim \triangle COD$ [AAA similarity criterion]


B BYJU'S

NCERT Solutions for Class 10 Maths Chapter 6 – Triangles

As we know, If two triangles are similar then the ratio of their areas are equal to the square of the ratio of their corresponding sides. Therefore, Area of $(\Delta AOB)/Area$ of $(\Delta COD) = AB^2/CD^2$ = $(2CD)^2/CD^2$ [$\therefore AB = CD$] \therefore Area of $(\Delta AOB)/Area$ of (ΔCOD) = $4CD^2/CD = 4/1$

Hence, the required ratio of the area of $\triangle AOB$ and $\triangle COD = 4:1$


3. In the figure, ABC and DBC are two triangles on the same base BC. If AD intersects BC at O, show that area (ΔABC)/area (ΔDBC) = AO/DO.



Solution: Given, ABC and DBC are two triangles on the same base BC. ADintersects BC at O.

We have to prove: Area (ΔABC)/Area (ΔDBC) = AO/DO

Let us draw two perpendiculars AP and DM on line BC.



We know that area of a triangle = $1/2 \times Base \times Height$

$$\therefore \frac{\operatorname{ar}(\Delta ABC)}{\operatorname{ar}(\Delta DEF)} = \frac{\frac{1}{2} BC \times AP}{\frac{1}{2} BC \times DM} = \frac{AP}{DM}$$


In $\triangle APO$ and $\triangle DMO$, $\angle APO = \angle DMO$ (Each 90°)



 $\angle AOP = \angle DOM$ (Vertically opposite angles) $\therefore \Delta APO \sim \Delta DMO$ (AA similarity criterion) $\therefore AP/DM = AO/DO$ \Rightarrow Area ($\triangle ABC$)/Area ($\triangle DBC$) = AO/DO.


4. If the areas of two similar triangles are equal, prove that they are congruent.

Solution: Say \triangle ABC and \triangle PQR are two similar triangles and equal in area



5. D, E and F are respectively the mid-points of sides AB, BC and CA of \triangle ABC. Find the ratio of the area of \triangle DEF and \triangle ABC.

Solution: Given, D, E and F are respectively the mid-points of sides AB, BC and CA of \triangle ABC.



In \triangle ABC, F is the mid point of AB (Already given) E is the mid-point of AC (Already given)




So, by the mid-point theorem, we have, FE || BC and FE = 1/2BC \Rightarrow FE || BC and FE || BD [BD = 1/2BC] Since, opposite sides of parallelogram are equal and parallel \therefore BDEF is parallelogram. Similarly in \triangle FBD and \triangle DEF, we have FB = DE (Opposite sides of parallelogram BDEF) FD = FD (Common sides) BD = FE (Opposite sides of parallelogram BDEF) $\therefore \Delta FBD \cong \Delta DEF$ Similarly, we can prove that $\Delta AFE \cong \Delta DEF$ $\Delta EDC \cong \Delta DEF$ As we know, if triangles are congruent, then they are equal in area. So, $Area(\Delta FBD) = Area(\Delta DEF)$ (i) Area(ΔAFE) = Area(ΔDEF)(ii) and, $Area(\Delta EDC) = Area(\Delta DEF)$ (iii) Now, $Area(\Delta ABC) = Area(\Delta FBD) + Area(\Delta DEF) + Area(\Delta AFE) + Area(\Delta EDC) \dots (iv)$ $Area(\Delta ABC) = Area(\Delta DEF) + Area(\Delta DEF) + Area(\Delta DEF) + Area(\Delta DEF)$ From equation (i), (ii) and (iii), \Rightarrow Area(\triangle DEF) = (1/4)Area(\triangle ABC) \Rightarrow Area(Δ DEF)/Area(Δ ABC) = 1/4

Hence, Area(ΔDEF):Area(ΔABC) = 1:4

6. Prove that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding medians.

Solution: Given: AM and DN are the medians of triangles ABC and DEF respectively and $\triangle ABC \sim \triangle DEF$.






We have to prove: Area(ΔABC)/Area(ΔDEF) = AM²/DN²

Since, $\triangle ABC \sim \triangle DEF$ (Given) $\therefore \text{Area}(\Delta ABC)/\text{Area}(\Delta DEF) = (AB^2/DE^2) \dots (i)$ and, AB/DE = BC/EF = CA/FD(ii) $\Rightarrow \frac{AB}{DE} = \frac{\frac{1}{2}BC}{\frac{1}{2}EF} =$ $\frac{\text{CD}}{\text{FD}}$ In $\triangle ABM$ and $\triangle DEN$, Since $\triangle ABC \sim \triangle DEF$ $\therefore \angle B = \angle E$ AB/DE = BM/EN [Already Proved in equation (i)] $\therefore \Delta ABC \sim \Delta DEF$ [SAS similarity criterion] \Rightarrow AB/DE = AM/DN(iii) $\therefore \Delta ABM \sim \Delta DEN$ As the areas of two similar triangles are proportional to the squares of the corresponding sides. \therefore area($\triangle ABC$)/area($\triangle DEF$) = $AB^2/DE^2 = AM^2/DN^2$ Hence, proved.

7. Prove that the area of an equilateral triangle described on one side of a square is equal to half the area of the equilateral triangle described on one of its diagonals.

Solution: Given, ABCD is a square whose one diagonal is AC. \triangle APC and \triangle BQC are two equilateral



triangles described on the diagonals AC and side BC of the square ABCD.

Area(Δ BQC) = $\frac{1}{2}$ Area(Δ APC)



Since, $\triangle APC$ and $\triangle BQC$ are both equilateral triangles, as per given, $\therefore \triangle APC \sim \triangle BQC$ [AAA similarity criterion] $\therefore area(\triangle APC)/area(\triangle BQC) = (AC^2/BC^2) = AC^2/BC^2$ Since, Diagonal = $\sqrt{2}$ side = $\sqrt{2}$ BC

$$(\frac{\sqrt{2}BC}{BC})^2 = 2$$

 $\Rightarrow \operatorname{area}(\Delta APC) = 2 \times \operatorname{area}(\Delta BQC)$ $\Rightarrow \operatorname{area}(\Delta BQC) = 1/2\operatorname{area}(\Delta APC)$ Hence, proved.


Tick the correct answer and justify:

8. ABC and BDE are two equilateral triangles such that D is the mid-point of BC. Ratio of the area of triangles ABC and BDE is

(A) 2:1
(B) 1:2
(C) 4:1

(D) 1 : 4

Solution: Given, $\triangle ABC$ and $\triangle BDE$ are two equilateral triangle. D is the midpoint of BC.




 $\therefore BD = DC = 1/2BC$ Let each side of triangle is 2*a*. As, $\triangle ABC \sim \triangle BDE$ $\therefore Area(\triangle ABC)/Area(\triangle BDE) = AB^2/BD^2 = (2a)^2/(a)^2 = 4a^2/a^2 = 4/1 = 4:1$ Hence, the correct answer is (C).

9. Sides of two similar triangles are in the ratio 4 : 9. Areas of these triangles are in the ratio (A) 2 : 3
(B) 4 : 9



(C) 81 : 16 (D) 16 : 81

Solution: Given, Sides of two similar triangles are in the ratio 4 : 9.



Let ABC and DEF are two similar triangles, such that,

 $\Delta ABC \sim \Delta DEF$

And AB/DE = AC/DF = BC/EF = 4/9

As, the ratio of the areas of these triangles will be equal to the square of the ratio of the corresponding sides,

 $\therefore \text{Area}(\Delta \text{ABC})/\text{Area}(\Delta \text{DEF}) = \text{AB}^2/\text{DE}^2$

: Area(ΔABC)/Area(ΔDEF) = (4/9)² = 16/81 = 16:81

Hence, the correct answer is (D).



Exercise 6.5

Page: 150

1. Sides of triangles are given below. Determine which of them are right triangles? In case of a right triangle, write the length of its hypotenuse.

(i) 7 cm, 24 cm, 25 cm
(ii) 3 cm, 8 cm, 6 cm
(iii) 50 cm, 80 cm, 100 cm

(iv) 13 cm, 12 cm, 5 cm

Solution:

(i) Given, sides of the triangle are 7 cm, 24 cm, and 25 cm.

Squaring the lengths of the sides of the, we will get 49, 576, and 625.

49 + 576 = 625

 $(7)^2 + (24)^2 = (25)^2$

Therefore, the above equation satisfies, Pythagoras theorem. Hence, it is right angled triangle. Length of Hypotenuse = 25 cm

(ii) Given, sides of the triangle are 3 cm, 8 cm, and 6 cm.

Squaring the lengths of these sides, we will get 9, 64, and 36.

Clearly, $9 + 36 \neq 64$

Or, $3^2 + 6^2 \neq 8^2$

Therefore, the sum of the squares of the lengths of two sides is not equal to the square of the length of the hypotenuse.

Hence, the given triangle does not satisfies Pythagoras theorem.

(iii) Given, sides of triangle's are 50 cm, 80 cm, and 100 cm. Squaring the lengths of these sides, we will get 2500, 6400, and 10000. However, $2500 + 6400 \neq 10000$ Or, $50^2 + 80^2 \neq 100^2$ As you can see, the sum of the squares of the lengths of two sides is not equal to the square of the length of the third side.

Therefore, the given triangle does not satisfies Pythagoras theorem. Hence, it is not a right triangle.


fichce, it is not a right triangle.

(iv) Given, sides are 13 cm, 12 cm, and 5 cm. Squaring the lengths of these sides, we will get 169, 144, and 25. Thus, 144 + 25 = 169Or, $12^2 + 5^2 = 13^2$ The sides of the given triangle are satisfying Pythagoras theorem. Therefore, it is a right triangle. Hence, length of the hypotenuse of this triangle is 13 cm.



2. PQR is a triangle right angled at P and M is a point on QR such that PM \perp QR. Show that PM² = QM × MR.

Solution: Given, $\triangle PQR$ is right angled at P is a point on QR such that PM $\perp QR$




We have to prove, $PM^2 = QM \times MR$

In $\triangle PQM$, by Pythagoras theorem $PQ^2 = PM^2 + QM^2$ Or, $PM^2 = PQ^2 - QM^2$ (i)

In $\triangle PMR$, by Pythagoras theorem $PR^2 = PM^2 + MR^2$ Or, $PM^2 = PR^2 - MR^2$ (ii)

Adding equation, (i) and (ii), we get, $2PM^2 = (PQ^2 + PM^2) - (QM^2 + MR^2)$ $= QR^2 - QM^2 - MR^2 \quad [: QR^2 = PQ^2 + PR^2]$ $= (QM + MR)^2 - QM^2 - MR^2$ $= 2QM \times MR$ $\therefore PM^2 = QM \times MR$

3. In Figure, ABD is a triangle right angled at A and AC ⊥ BD. Show that
(i) AB² = BC × BD
(ii) AC² = BC × DC
(iii) AD² = BD × CD



Solution:



(i) In $\triangle ADB$ and $\triangle CAB$, $\angle DAB = \angle ACB$ (Each 90°) $\angle ABD = \angle CBA$ (Common angles) $\therefore \triangle ADB \sim \triangle CAB$ [AA similarity criterion] $\Rightarrow AB/CB = BD/AB$ $\Rightarrow AB^2 = CB \times BD$

(ii) Let $\angle CAB = x$ In $\triangle CBA$, $\angle CBA = 180^{\circ} - 90^{\circ} - x$ $\angle CBA = 90^{\circ} - x$ Similarly, in $\triangle CAD$ $\angle CAD = 90^{\circ} - \angle CBA$ $= 90^{\circ} - x$ $\angle CDA = 180^{\circ} - 90^{\circ} - (90^{\circ} - x))$ $\angle CDA = x$




In \triangle CBA and \triangle CAD, we have \angle CBA = \angle CAD \angle CAB = \angle CDA \angle ACB = \angle DCA (Each 90°) $\therefore \triangle$ CBA ~ \triangle CAD [AAA similarity criterion] \Rightarrow AC/DC = BC/AC \Rightarrow AC² = DC × BC

(iii) In \triangle DCA and \triangle DAB, \angle DCA = \angle DAB (Each 90°) \angle CDA = \angle ADB (common angles) $\therefore \triangle$ DCA ~ \triangle DAB [AA similarity criterion] \Rightarrow DC/DA = DA/DA \Rightarrow AD² = BD × CD

4. ABC is an isosceles triangle right angled at C. Prove that $AB^2 = 2AC^2$.

Solution: Given, \triangle ABC is an isosceles triangle right angled at C.




In $\triangle ACB$, $\angle C = 90^{\circ}$ AC = BC (By isosceles triangle property) AB² = AC² + BC² [By Pythagoras theorem]



 $= AC^{2} + AC^{2}$ [Since, AC = BC] AB² = 2AC²

5. ABC is an isosceles triangle with AC = BC. If $AB^2 = 2AC^2$, prove that ABC is a right triangle.


Solution: Given, $\triangle ABC$ is an isosceles triangle having AC = BC and $AB^2 = 2AC^2$



In $\triangle ACB$, AC = BC $AB^2 = 2AC^2$ $AB^2 = AC^2 + AC^2$ $= AC^2 + BC^2$ [Since, AC = BC] Hence, by Pythagoras theorem $\triangle ABC$ is right angle triangle.

6. ABC is an equilateral triangle of side 2a. Find each of its altitudes.

Solution: Given, ABC is an equilateral triangle of side 2a.



Draw, $AD \perp BC$ In $\triangle ADB$ and $\triangle ADC$, AB = ACAD = AD $\angle ADB = \angle ADC$ [Both are 90°]

Therefore, $\triangle ADB \cong \triangle ADC$ by RHS congruence. Hence, BD = DC [by CPCT]


In right angled $\triangle ADB$, $AB^2 = AD^2 + BD^2$



 $(2a)^{2} = AD^{2} + a^{2}$ $\Rightarrow AD^{2} = 4a^{2} - a^{2}$ $\Rightarrow AD^{2} = 3a^{2}$ $\Rightarrow AD = \sqrt{3}a$

7. Prove that the sum of the squares of the sides of rhombus is equal to the sum of the squares of its diagonals.

Solution: Given, ABCD is a rhombus whose diagonals AC and BD intersect at O.



We have to prove, as per the question, $AB^2 + BC^2 + CD^2 + AD^2 = AC^2 + BD^2$

Since, the diagonals of a rhombus bisect each other at right angles. Therefore, AO = CO and BO = DOIn $\triangle AOB$, $\angle AOB = 90^{\circ}$ $AB^2 = AO^2 + BO^2$(i) [By Pythagoras theorem]


Similarly, $AD^2 = AO^2 + DO^2$(ii) $DC^2 = DO^2 + CO^2$(iii) $BC^2 = CO^2 + BO^2$(iv)

Adding equations (i) + (ii) + (iii) + (iv), we get, $AB^{2} + AD^{2} + DC^{2} + BC^{2} = 2(AO^{2} + BO^{2} + DO^{2} + CO^{2})$ $= 4AO^{2} + 4BO^{2}$ [Since, AO = CO and BO =DO] $= (2AO)^{2} + (2BO)^{2} = AC^{2} + BD^{2}$

 $AB^2 + AD^2 + DC^2 + BC^2 = AC^2 + BD^2$ Hence, proved.

8. In Fig. 6.54, O is a point in the interior of a triangle.






ABC, OD \perp BC, OE \perp AC and OF \perp AB. Show that: (i) OA² + OB² + OC² - OD² - OE² - OF² = AF² + BD² + CE², (ii) AF² + BD² + CE² = AE² + CD² + BF².

Solution: Given, in $\triangle ABC$, O is a point in the interior of a triangle. And OD \perp BC, OE \perp AC and OF \perp AB.

Join OA, OB and OC



(i) By Pythagoras theorem in $\triangle AOF$, we have $OA^2 = OF^2 + AF^2$ Similarly, in $\triangle BOD$ $OB^2 = OD^2 + BD^2$

Similarly, in $\triangle COE$ $OC^2 = OE^2 + EC^2$


Adding these equations, $OA^2 + OB^2 + OC^2 = OF^2 + AF^2 + OD^2 + BD^2 + OE^2 + EC^2$ $OA^2 + OB^2 + OC^2 - OD^2 - OE^2 - OF^2 = AF^2 + BD^2 + CE^2$.

(ii) $AF^2 + BD^2 + EC^2 = (OA^2 - OE^2) + (OC^2 - OD^2) + (OB^2 - OF^2)$ $\therefore AF^2 + BD^2 + CE^2 = AE^2 + CD^2 + BF^2.$

9. A ladder 10 m long reaches a window 8 m above the ground. Find the distance of the foot of the ladder from base of the wall.



Solution: Given, a ladder 10 m long reaches a window 8 m above the ground.




Let BA be the wall and AC be the ladder, Therefore, by Pythagoras theorem, $AC^2 = AB^2 + BC^2$ $10^2 = 8^2 + BC^2$ $BC^2 = 100 - 64$ $BC^2 = 36$ BC = 6m

Therefore, the distance of the foot of the ladder from the base of the wall is 6 m.

10. A guy wire attached to a vertical pole of height 18 m is 24 m long and has a stake attached to the other end. How far from the base of the pole should the stake be driven so that the wire will be taut?

Solution: Given, a guy wire attached to a vertical pole of height 18 m is 24 m long and has a stake attached to the other end.



Let AB be the pole and AC be the wire. By Pythagoras theorem,

 $AC^{2} = AB^{2} + BC^{2}$ $24^{2} = 18^{2} + BC^{2}$ $BC^{2} = 576 - 324$ $BC^{2} = 252$ $BC = 6\sqrt{7}m$ Therefore, the distance from the base is $6\sqrt{7}m$.



11. An aeroplane leaves an airport and flies due north at a speed of 1,000 km per hour. At the same time, another aeroplane leaves the same airport and flies due west at a speed of 1,200 km per

 $1\frac{1}{2}$ hours?


Solution: Given,

Speed of first aeroplane = 1000 km/hr

hour. How far apart will be the two planes after

Distance covered by first aeroplane flying due north in $1\frac{1}{2}$ hours (OA) = $100 \times 3/2$ km = 1500 km Speed of second aeroplane = 1200 km/hr

Distance covered by second aeroplane flying due west in $1\frac{1}{2}$ hours (OB) = $1200 \times 3/2$ km = 1800 km




In right angle $\triangle AOB$, by Pythagoras Theorem, $AB^2 = AO^2 + OB^2$ $\Rightarrow AB^2 = (1500)^2 + (1800)^2$ $\Rightarrow AB = \sqrt{(2250000 + 3240000)}$ $= \sqrt{5490000}$ $\Rightarrow AB = 300\sqrt{61}$ km

Hence, the distance between two aeroplanes will be $300\sqrt{61}$ km.

12. Two poles of heights 6 m and 11 m stand on a plane ground. If the distance between the feet of the poles is 12 m, find the distance between their tops.

Solution: Given, Two poles of heights 6 m and 11 m stand on a plane ground. And distance between the feet of the poles is 12 m.






Let AB and CD be the poles of height 6m and 11m. Therefore, CP = 11 - 6 = 5m

From the figure, it can be observed that AP = 12mBy Pythagoras theorem for $\triangle APC$, we get, $AP^2 = PC^2 + AC^2$ $(12m)^2 + (5m)^2 = (AC)^2$ $AC^2 = (144+25)m^2 = 169 m^2$ AC = 13mTherefore, the distance between their tops is 13 m.



13. D and E are points on the sides CA and CB respectively of a triangle ABC right angled at C. Prove that $AE^2 + BD^2 = AB^2 + DE^2$.

Solution: Given, D and E are points on the sides CA and CB respectively of a triangle ABC right angled at C.




By Pythagoras theorem in $\triangle ACE$, we get $AC^2 + CE^2 = AE^2$(i) In $\triangle BCD$, by Pythagoras theorem, we get $BC^2 + CD^2 = BD^2$(ii)

From equations (i) and (ii), we get, $AC^2 + CE^2 + BC^2 + CD^2 = AE^2 + BD^2$ (iii) In $\triangle CDE$, by Pythagoras theorem, we get $DE^2 = CD^2 + CE^2$ In $\triangle ABC$, by Pythagoras theorem, we get



 $AB^2 = AC^2 + CB^2$ Putting the above two values in equation (iii), we get $DE^2 + AB^2 = AE^2 + BD^2$.

14. The perpendicular from A on side BC of a \triangle ABC intersects BC at D such that DB = 3CD (see Figure). Prove that $2AB^2 = 2AC^2 + BC^2$.




Solution: Given, the perpendicular from A on side BC of a \triangle ABC intersects BC at D such that; DB = 3CD.

In \triangle ABC, AD \perp BC and BD = 3CD In right angle triangle, ADB and ADC, by Pythagoras theorem, AB² = AD² + BD²(i) AC² = AD² + DC²(ii) Subtracting equation (ii) from equation (i), we get AB² - AC² = BD² - DC² = 9CD² - CD² [Since, BD = 3CD] = 9CD² = 8(BC/4)² [Since, BC = DB + CD = 3CD + CD = 4CD] Therefore, AB² - AC² = BC²/2 $\Rightarrow 2(AB² - AC²) = BC²$ $\Rightarrow 2AB² - 2AC² = BC²$.

15. In an equilateral triangle ABC, D is a point on side BC such that BD = 1/3BC. Prove that $9AD^2 = 7AB^2$.


Solution: Given, ABC is an equilateral triangle. And D is a point on side BC such that BD = 1/3BC





16. In an equilateral triangle, prove that three times the square of one side is equal to four times the square of one of its altitudes.

Solution: Given, an equilateral triangle say ABC,



Let the sides of the equilateral triangle be of length a, and AE be the altitude of $\triangle ABC$. $\therefore BE = EC = BC/2 = a/2$ In $\triangle ABE$, by Pythagoras Theorem, we get $AB^2 = AE^2 + BE^2$



$$a^{2} = AE^{2} + \left(\frac{a}{2}\right)^{2}$$

$$AE^{2} = a^{2} - \frac{a^{2}}{4}$$

$$AE^{2} = \frac{3a^{2}}{4}$$

$$4AE^{2} = 3a^{2}$$

$$\Rightarrow 4 \times (\text{Square of altitude}) = 3 \times (\text{Square of one side})$$
Hence, proved.


17. Tick the correct answer and justify: In $\triangle ABC$, $AB = 6\sqrt{3}$ cm, AC = 12 cm and BC = 6 cm. The angle B is: (A) 120°

(B) 60°

(C) 90°

(D) 45°

Solution: Given, in $\triangle ABC$, $AB = 6\sqrt{3}$ cm, AC = 12 cm and BC = 6 cm.



We can observe that, $AB^2 = 108$ $AC^2 = 144$ And, $BC^2 = 36$ $AB^2 + BC^2 = AC^2$ The given triangle, ΔABC , is satisfying Pythagoras theorem. Therefore, the triangle is a right triangle, right-angled at B. $\therefore \angle B = 90^{\circ}$ Hence, the correct answer is (C).



Exercise 6.6

Page: 152


1. In Figure, PS is the bisector of \angle QPR of \triangle PQR. Prove that QS/PQ = SR/PR

R

Solution:

Let us draw a line segment RT parallel to SP which intersects extended line segment QP at point T.

Given, PS is the angle bisector of $\angle QPR$. Therefore, $\angle QPS = \angle SPR$(i)




As per the constructed figure, $\angle SPR = \angle PRT(Since, PS||TR).....(ii)$ $\angle QPS = \angle QRT(Since, PS||TR)(iii)$ From the above equations, we get, $\angle PRT = \angle QTR$ Therefore, PT = PR

In \triangle QTR, by basic proportionality theorem, QS/SR = QP/PT Since, PT=TR Therefore, QS/SR = PQ/PR Hence, proved.



2. In Fig. 6.57, D is a point on hypotenuse AC of △ABC, such that BD ⊥AC, DM ⊥ BC and DN ⊥ AB. Prove




that: (i) $DM^2 = DN \cdot MC$ (ii) $DN^2 = DM \cdot AN$.

Solution:

Let us join Point D and B.

(i) Given,

BD \perp AC, DM \perp BC and DN \perp AB Now from the figure we have, DN || CB, DM || AB and \angle B = 90 ° Therefore, DMBN is a rectangle. So, DN = MB and DM = NB



The given condition which we have to prove, is when D is the foot of the perpendicular drawn from B to AC.

 $\therefore \angle CDB = 90^{\circ} \Rightarrow \angle 2 + \angle 3 = 90^{\circ} \dots (i)$ In $\triangle CDM$, $\angle 1 + \angle 2 + \angle DMC = 180^{\circ}$ $\Rightarrow \angle 1 + \angle 2 = 90^{\circ} \dots (ii)$ In $\triangle DMB$, $\angle 3 + \angle DMB + \angle 4 = 180^{\circ}$ $\Rightarrow \angle 3 + \angle 4 = 90^{\circ} \dots (iii)$ From equation (i) and (ii), we get $\angle 1 = \angle 3$ From equation (i) and (iii), we get $\angle 2 = \angle 4$


In Δ DCM and Δ BDM, $\angle 1 = \angle 3$ (Already Proved) $\angle 2 = \angle 4$ (Already Proved) $\therefore \Delta$ DCM ~ Δ BDM (AA similarity criterion) BM/DM = DM/MC \Rightarrow DN/DM = DM/MC (BM = DN) \Rightarrow DM² = DN × MC

Hence, proved.




From equation (iv) and (vi), we get, $\angle 6 = \angle 7$ From equation (v) and (vi), we get, $\angle 8 = \angle 5$ In \triangle DNA and \triangle BND, $\angle 6 = \angle 7$ (Already proved) $\angle 8 = \angle 5$ (Already proved) $\therefore \triangle$ DNA ~ \triangle BND (AA similarity criterion) AN/DN = DN/NB \Rightarrow DN² = AN × NB \Rightarrow DN² = AN × DM (Since, NB = DM) Hence, proved.

3. In Figure, ABC is a triangle in which $\angle ABC > 90^{\circ}$ and $AD \perp CB$ produced. Prove that $AC^2 = AB^2 + BC^2 + 2$ BC.BD.



Solution: By applying Pythagoras Theorem in $\triangle ADB$, we get, $AB^2 = AD^2 + DB^2$ (i) Again, by applying Pythagoras Theorem in $\triangle ACD$, we get, $AC^2 = AD^2 + DC^2$ $AC^2 = AD^2 + (DB + BC)^2$ $AC^2 = AD^2 + DB^2 + BC^2 + 2DB \times BC$ From equation (i), we can write, $AC^2 = AB^2 + BC^2 + 2DB \times BC$ Hence, proved.


4. In Figure, ABC is a triangle in which \angle ABC < 90° and AD \perp BC. Prove that





Solution: By applying Pythagoras Theorem in $\triangle ADB$, we get, $AB^2 = AD^2 + DB^2$ We can write it as; $\Rightarrow AD^2 = AB^2 - DB^2$ (i) By applying Pythagoras Theorem in $\triangle ADC$, we get, $AD^2 + DC^2 = AC^2$ From equation (i), $AB^2 - BD^2 + DC^2 = AC^2$ $AB^2 - BD^2 + (BC - BD)^2 = AC^2$ $AC^2 = AB^2 - BD^2 + BC^2 + BD^2 - 2BC \times BD$ $AC^2 = AB^2 + BC^2 - 2BC \times BD$ Hence, proved.

5. In Figure, AD is a median of a triangle ABC and AM ⊥ BC. Prove that :
(i) AC² = AD² + BC.DM + 2 (BC/2)²
(ii) AB² = AD² - BC.DM + 2 (BC/2)²
(iii) AC² + AB² = 2 AD² + ¹/₂ BC²



Solution:

(i) By applying Pythagoras Theorem in $\triangle AMD$, we get, $AM^2 + MD^2 = AD^2$ (i)

Again, by applying Pythagoras Theorem in $\triangle AMC$, we get, $AM^2 + MC^2 = AC^2$ $AM^2 + (MD + DC)^2 = AC^2$ $(AM^2 + MD^2) + DC^2 + 2MD.DC = AC^2$

From equation(i), we get, $AD^2 + DC^2 + 2MD.DC = AC^2$ Since, DC=BC/2, thus, we get, $AD^2 + (BC/2)^2 + 2MD.(BC/2)^2 = AC^2$ $AD^2 + (BC/2)^2 + 2MD \times BC = AC^2$ Hence, proved.

(ii) By applying Pythagoras Theorem in $\triangle ABM$, we get; $AB^2 = AM^2 + MB^2$




=> $(AD^2 - DM^2) + MB^2$ => $(AD^2 - DM^2) + (BD - MD)^2$ => $AD^2 - DM^2 + BD^2 + MD^2 - 2BD \times MD$ => $AD^2 + BD^2 - 2BD \times MD$ => $AD^2 + (BC/2)^2 - 2(BC/2) \times MD$ => $AD^2 + (BC/2)^2 - BC \times MD$ Hence, proved.

6. Prove that the sum of the squares of the diagonals of parallelogram is equal to the sum of the squares of its sides.

Solution:

Let us consider, ABCD be a parallelogram. Now, draw perpendicular DE on extended side of AB, and draw a perpendicular AF meeting DC at point F.



By applying Pythagoras Theorem in $\triangle DEA$, we get, $DE^2 + EA^2 = DA^2$ (i) By applying Pythagoras Theorem in $\triangle DEB$, we get, $DE^2 + EB^2 = DB^2$ $DE^2 + (EA + AB)^2 = DB^2$

 $(DE² + EA²) + AB² + 2EA \times AB = DB²$



 $DA^2 + AB^2 + 2EA \times AB = DB^2$ (ii) By applying Pythagoras Theorem in $\triangle ADF$, we get, $AD^2 = AF^2 + FD^2$ Again, applying Pythagoras theorem in $\triangle AFC$, we get, $AC^{2} = AF^{2} + FC^{2} = AF^{2} + (DC - FD)^{2}$ $= AF^2 + DC^2 + FD^2 - 2DC \times FD$ $= (AF^2 + FD^2) + DC^2 - 2DC \times FD AC^2$ $AC^2 = AD^2 + DC^2 - 2DC \times FD$ (iii) Since ABCD is a parallelogram, AB = CD(iv) And BC = AD(v) In $\triangle DEA$ and $\triangle ADF$, $\angle DEA = \angle AFD$ (Each 90°) $\angle EAD = \angle ADF (EA \parallel DF)$ AD = AD (Common Angles) $\therefore \Delta EAD \cong \Delta FDA$ (AAS congruence criterion) \Rightarrow EA = DF (vi) Adding equations (i) and (iii), we get, $DA^2 + AB^2 + 2EA \times AB + AD^2 + DC^2 - 2DC \times FD = DB^2 + AC^2$ $DA^{2} + AB^{2} + AD^{2} + DC^{2} + 2EA \times AB - 2DC \times FD = DB^{2} + AC^{2}$ From equation (iv) and (vi), $BC^2 + AB^2 + AD^2 + DC^2 + 2EA \times AB - 2AB \times EA = DB^2 + AC^2$ $AB^2 + BC^2 + CD^2 + DA^2 = AC^2 + BD^2$

7. In Figure, two chords AB and CD intersect each other at the point P. Prove that :
(i) △APC ~ △ DPB
(ii) AP . PB = CP . DP


Solution: Firstly, let us join CB, in the given figure.

(i) In \triangle APC and \triangle DPB, \angle APC = \angle DPB (Vertically opposite angles) \angle CAP = \angle BDP (Angles in the same segment for chord CB) Therefore, \triangle APC ~ \triangle DPB (AA similarity criterion)

(ii) In the above, we have proved that $\triangle APC \sim \triangle DPB$

We know that the corresponding sides of similar triangles are proportional.

 $\therefore AP/DP = PC/PB = CA/BD$ $\Rightarrow AP/DP = PC/PB$ $\therefore AP. PB = PC. DP$ Hence, proved.




B BYJU'S

NCERT Solutions for Class 10 Maths Chapter 6 – Triangles

8. In Fig. 6.62, two chords AB and CD of a circle intersect each other at the point P (when produced) outside the circle. Prove that:

(i) \triangle PAC ~ \triangle PDB (ii) PA . PB = PC . PD.



Solution:

(i) In \triangle PAC and \triangle PDB, $\angle P = \angle P$ (Common Angles) As we know, exterior angle of a cyclic quadrilateral is \angle PCA and \angle PBD is opposite interior angle, which are both equal. $\angle PAC = \angle PDB$ Thus, $\triangle PAC \sim \triangle PDB$ (AA similarity criterion)


(ii) We have already proved above, $\Delta APC \sim \Delta DPB$ We know that the corresponding sides of similar triangles are proportional. Therefore, AP/DP = PC/PB = CA/BD AP/DP = PC/PB $\therefore AP. PB = PC. DP$

9. In Figure, D is a point on side BC of \triangle ABC such that BD/CD = AB/AC. Prove that AD is the bisector of \angle BAC.


R

Solutions: In the given figure, let us extend BA to P such that; AP = AC. Now join PC.





10. Nazima is fly fishing in a stream. The tip of her fishing rod is 1.8 m above the surface of the water and the fly at the end of the string rests on the water 3.6 m away and 2.4 m from a point directly under the tip of the rod. Assuming that her string (from the tip of her rod to the fly) is taut, how much string does she have out (see Figure)? If she pulls in the string at the rate of 5 cm per second, what will be the horizontal distance of the fly from her after 12 seconds?




Solution:

Let us consider, AB is the height of the tip of the fishing rod from the water surface and BC is the



horizontal distance of the fly from the tip of the fishing rod. Therefore, AC is now the length of the string.




To find AC, we have to use Pythagoras theorem in $\triangle ABC$, is such way; $AC^2 = AB^2 + BC^2$ $AB^2 = (1.8 \text{ m})^2 + (2.4 \text{ m})^2$ $AB^2 = (3.24 + 5.76) \text{ m}^2$ $AB^2 = 9.00 \text{ m}^2$ $\Rightarrow AB = \sqrt{9} \text{ m} = 3\text{m}$

Thus, the length of the string out is 3 m.

As its given, she pulls the string at the rate of 5 cm per second.

Therefore, string pulled in 12 seconds = $12 \times 5 = 60$ cm = 0.6 m



Let us say now, the fly is at point D after 12 seconds. Length of string out after 12 seconds is AD. AD = AC - String pulled by Nazima in 12 seconds = (3.00 - 0.6) m = 2.4 m In $\triangle ADB$, by Pythagoras Theorem, $AB^2 + BD^2 = AD^2$ $(1.8 \text{ m})^2 + BD^2 = (2.4 \text{ m})^2$ $BD^2 = (5.76 - 3.24)$ m² = 2.52 m² BD = 1.587 m Horizontal distance of fly = BD + 1.2 m = (1.587 + 1.2) m = 2.787 m = 2.79 m