

# CBSE Class 12 Maths Question Paper 2020

## Set 2

### General Instructions:

Read the following instructions very carefully and strictly follow them:

- (i) This question paper comprises **four** Sections A, B, C and D. This question paper carries **36** questions. **All** questions are compulsory.
- (ii) **Section A** – Questions no. **1 to 20** comprises of **20** questions of **1** mark each.
- (iii) **Section B** – Questions no. **21 to 26** comprises of **6** questions of **2** mark each.
- (iv) **Section C** – Questions no. **27 to 32** comprises of **6** questions of **4** mark each.
- (v) **Section D** – Questions no. **33 to 36** comprises of **4** questions of **6** mark each.
- (vi) There is no overall choice in the question paper. However, an internal choice has been provided in 3 questions of one mark, 2 questions of two marks, 2 questions of four marks and 2 questions of six marks. Only one of the choices in such questions have to be attempted.
- (vii) In addition to this, separate instructions are given with each section and question, wherever necessary.
- (viii) Use of calculators is **not** permitted.

### SECTION - A

**Question numbers 1 to 20 carry 1 mark each.**

**Question numbers 1 to 10 are multiple choice type questions. Select the correct option.**

1. If  $f$  and  $g$  are two functions from  $R$  to  $R$  defined as  $f(x) = |x| + x$  and  $g(x) = |x| - x$ , then  $f \circ g(x)$  for  $x < 0$  is
  - (a)  $4x$
  - (b)  $2x$
  - (c)  $0$
  - (d)  $-4x$
2. The principal value of  $\cot^{-1}(-\sqrt{3})$  is
  - (a)  $-\frac{\pi}{6}$
  - (b)  $\frac{\pi}{6}$
  - (c)  $\frac{2\pi}{3}$
  - (d)  $\frac{5\pi}{6}$
3. If  $A = \begin{bmatrix} -2 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -2 \end{bmatrix}$ , then the value of  $|\text{adj } A|$  is
  - (a) 64
  - (b) 16
  - (c) 0
  - (d) -8
4. The maximum value of slope of the curve  $y = -x^3 + 3x^2 + 12x - 5$  is
  - (a) 15
  - (b) 12
  - (c) 9
  - (d) 0
5.  $\int \frac{e^x(1+x)}{\cos^2(xe^x)} dx$  is equal to

DATE:

SUBJECT

CLASS:

CENTRE:

TOPIC

(a)  $\tan(xe^x) + c$       (b)  $\cot(xe^x) + c$       (c)  $\cot(e^x) + c$       (d)  $\tan[e^x(1+x)] + c$

6. The degree of the differential equation  $x^2 \frac{d^2 y}{dx^2} = \left(x \frac{dy}{dx} - y\right)^3$  is

- (a) 1                      (b) 2                      (c) 3                      (d) 6

7. The value of  $p$  for which  $p(\hat{i} + \hat{j} + \hat{k})$  is a unit vector is

- (a) 0                      (b)  $\frac{1}{\sqrt{3}}$                       (c) 1                      (d)  $\sqrt{3}$

8. The coordinates of the foot of the perpendicular drawn from the point  $(-2, 8, 7)$  on the ZX-plane is


- (a)  $(-2, -8, 7)$       (b)  $(2, 8, -7)$       (c)  $(-2, 0, 7)$       (d)  $(0, 8, 0)$

9. The vector equation of XY-plane is

- (a)  $\vec{r} \cdot \hat{k} = 0$       (b)  $\vec{r} \cdot \hat{j} = 0$       (c)  $\vec{r} \cdot \hat{i} = 0$       (d)  $\vec{r} \cdot \vec{n} = 1$

10. The feasible region for an LPP is shown below:

Let  $z = 3x - 4y$  be the objective function. Minimum of  $z$  occurs at



- (a)  $(0, 0)$                       (b)  $(0, 8)$                       (c)  $(5, 0)$                       (d)  $(4, 10)$

Fill in the blanks in question numbers 11 to 15.

11. If  $y = \tan^{-1} x + \cot^{-1} x$ ,  $x \in R$ , then  $\frac{dy}{dx}$  is equal to \_\_\_\_\_.

(OR)

If  $\cos(xy) = k$ , where  $k$  is a constant and  $xy \neq n\pi$ ,  $n \in Z$ , then  $\frac{dy}{dx}$  is equal to \_\_\_\_\_.

DATE:

SUBJECT

CLASS:

CENTRE:

TOPIC

12. The value of  $\lambda$  so that the function  $f$  defined by  $f(x) = \begin{cases} \lambda x, & \text{if } x \leq \pi \\ \cos x, & \text{if } x > \pi \end{cases}$  is continuous at  $x = \pi$  is \_\_\_\_\_.

13. The equation of the tangent to the curve  $y = \sec x$  at the point  $(0, 1)$  is \_\_\_\_\_.

14. The area of the parallelogram whose diagonals are  $2\hat{i}$  and  $-3\hat{k}$  is \_\_\_\_\_ square units.

(OR)

The value of  $\lambda$  for which the vectors  $2\hat{i} - \lambda\hat{j} + \hat{k}$  and  $i + 2\hat{j} - \hat{k}$  are orthogonal is \_\_\_\_\_.

15. A bag contains 3 black, 4 red and 2 green balls. If three balls are drawn simultaneously at random, then the probability that the balls are of different colours is \_\_\_\_\_.

**Question numbers 16 to 20 are very short answer type questions.**

16. Construct a  $2 \times 2$  matrix  $A = [a_{ij}]$  whose elements are given by  $a_{ij} = |(i)^2 - j|$ .

17. Differentiate  $\sin^2(\sqrt{x})$  with respect to  $x$ .

18. Find the interval in which the function  $f$  given by  $f(x) = 7 - 4x - x^2$  is strictly increasing.

19. Evaluate:  $\int_{-2}^2 |x| dx$

(OR)

Find:  $\int \frac{dx}{3 + 4x^2}$

20. An unbiased coin is tossed 4 times. Find the probability of getting at least one head.

### SECTION - B

**Question numbers 21 to 26 carry 2 marks each.**

21. Solve for  $x$ :

$$\sin^{-1} 4x + \sin^{-1} 3x = -\frac{\pi}{2}$$

(OR)

Express  $\tan^{-1}\left(\frac{\cos x}{1 - \sin x}\right)$ ,  $-\frac{3\pi}{2} < x < \frac{\pi}{2}$  in the simplest form.

22. Express  $A = \begin{bmatrix} 4 & -3 \\ 2 & -1 \end{bmatrix}$  as a sum of a symmetric and a skew symmetric matrix.

DATE:

SUBJECT

CLASS:

CENTRE:

TOPIC

23. If  $y^2 \cos\left(\frac{1}{x}\right) = a^2$ , then find  $\frac{dy}{dx}$ .

24. Show that for any two non-zero vectors  $\vec{a}$  and  $\vec{b}$ ,  $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$  if  $\vec{a}$  and  $\vec{b}$  are perpendicular vectors.

(OR)

Show that the vectors  $2\hat{i} - \hat{j} + \hat{k}$ ,  $3\hat{i} + 7\hat{j} + \hat{k}$  and  $5\hat{i} + 6\hat{j} + 1\hat{k}$  form the sides of a right-angled triangle.

25. Find the coordinates of the point where the line through  $(-1, 1, -8)$  and  $(5, -2, 10)$  crosses the ZX-plane.

26. If A and B are two events such that  $P(A) = 0.4$ ,  $P(B) = 0.3$  and  $P(A \cup B) = 0.6$ , then find  $P(B' \cap A)$ .

### SECTION - C

Question numbers 27 to 32 carry 4 marks each.

27. Show that the function  $f : (-\infty, 0) \rightarrow (-1, 0)$  defined by  $f(x) = \frac{x}{1+|x|}$ ,  $x \in (-\infty, 0)$  is one-one and onto.

(OR)

Show that the relation R in the set  $A = \{1, 2, 3, 4, 5, 6\}$  given by  $R = \{(a, b) : |a - b| \text{ is divisible by } 2\}$  is an equivalence relation.

28. If  $y = x^3 (\cos x)^x + \sin^{-1} \sqrt{x}$ , find  $\frac{dy}{dx}$ .

29. Evaluate:  $\int_{-1}^5 (|x| + |x+1| + |x-5|) dx$

30. Find the general solution of the differential equation  $x^2 y dx - (x^3 + y^3) dy = 0$ .

31. Solve the following LPP graphically:

Minimize  $z = 5x + 7y$

subject to the constraints

$$2x + y \geq 8$$

$$x + 2y \geq 10$$

$$x, y \geq 0$$

32. A bag contains two coins, one biased and the other unbiased. When tossed, the biased coin has a 60% chance of showing heads. One of the coin is selected at random and on tossing it shows tails. What is the probability it was an unbiased coin?

DATE:

SUBJECT

CLASS:

CENTRE:

TOPIC

(OR)

The probability distribution of a random variable X, where  $k$  is a constant is given below:

$$P(X = x) = \begin{cases} 0.1 & \text{if } x = 0 \\ kx^2, & \text{if } x = 1 \\ kx, & \text{if } x = 2 \text{ or } 3 \\ 0, & \text{otherwise} \end{cases}$$

Determine

- (a) the value of  $k$
- (b)  $P(X \leq 2)$
- (c) Mean of the distribution

#### SECTION - D

**Question numbers 33 to 36 carry 6 marks each.**

33. Solve the following system of equations by matrix method:

$$x - y + 2z = 7$$

$$2x - y + 3z = 12$$

$$3x + 2y - z = 5$$

(OR)

Obtain the inverse of the following matrix using elementary operations:

$$A = \begin{bmatrix} 2 & 1 & -3 \\ -1 & -1 & 4 \\ 3 & 0 & 2 \end{bmatrix}$$

34. Find the points on the curve  $9y^2 = x^3$ , where the normal to the curve makes equal intercepts with both the axes. Also find the equation of the normals.

35. Find the area of the following region using integration:  $\{(x, y) : y \leq |x| + 2, y \geq x^2\}$

(OR)

Using integration, find the area of a triangle whose vertices are  $(1,0)$ ,  $(2,2)$  and  $(3,1)$ .

36. Show that the lines  $\frac{x-2}{1} = \frac{y-2}{3} = \frac{z-3}{1}$  and  $\frac{x-2}{1} = \frac{y-3}{4} = \frac{z-4}{2}$  intersect. Also, find the coordinates of the point of intersection. Find the equation of the plane containing the two lines.