

**CBSE Class 12 Chemistry Question Paper
Solution 2020 Set 56/2/1**



Marking scheme – 2020

CHEMISTRY (043)/ CLASS XII


56/2/1

Q.No	Expected Answer / Value Points	Marks
SECTION A		
1	Simple organic compounds containing both carboxyl ($-\text{COOH}$) and amino ($-\text{NH}_2$) group. / Building blocks of protein. / Amino substituted carboxylic acid / structure	1
2	They contain both acid ($-\text{COOH}$) and basic(NH_2) group / Zwitter ion form or structure / It can react both acids and bases.	1
3	Acidic aminoacids- will have more number of carboxyl groups than amino groups. / Basic amino acids- will have more number of amino groups than carboxyl groups.	1
4	Those which are not synthesized in our body or must be supplied through diet.	1
5	Peptide linkage/amide linkage/ $\begin{array}{c} \text{O} \\ \parallel \\ \text{---C---NH---} \end{array}$	1

6	Leaching	1	
7	Zinc, Mercury or any other suitable metal	1	
8	Linkage	1	
9	Desorption	1	
10	2	1	
11	(D)	1	
12	(A)	1	
13	1 Mark will be given if attempted / if written none of the answer is right / 4	1	
14	(A)	1	
15	(C)	1	
16	(iii)	1	
17	(i)	1	
18	(iii)	1	
19	(iii)	1	
20	(i)	1	
SECTION B			
21	<p>a) Tranquilizers: Used to treat mild to severe mental diseases.</p> <p>b) Antiseptics- are applied to living tissues.</p> <p style="text-align: center;">OR</p> <p>Cationic detergent-are quaternary ammonium salts of amines with acetates, chlorides or bromides as anions. e.g. cetyltrimethyl ammonium bromide.</p>	<p>Analgesics: Used to reduce pain</p> <p>Disinfectants – are applied to inanimate objects or non living objects (or any other correct difference)</p> <p>Anionic detergent – are sodium salts of sulphonated long chain alcohols or hydrocarbons. e.g. Sodium lauryl sulphate / sodium dodecyl benzene sulphonate. (or any other suitable example)</p>	<p>1</p> <p>1</p> <p>1</p> <p>½ + ½</p>
22	<p>(a) Due to lone pair -lone pair repulsion on oxygen.</p> <p>(b) Due to resonance in phenol and not in methanol / sp² hybridised carbon atom in phenol / sp³ hybridised carbon atom in methanol.</p>	<p>1</p> <p>1</p>	
23	<p>(a) $2\text{MnO}_4^- + \text{H}_2\text{O} + \text{I}^- \rightarrow 2\text{MnO}_2 + 2\text{OH}^- + \text{IO}_3^-$</p> <p>(b) $2\text{MnO}_4^- + 10\text{I}^- + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 5\text{I}_2$ (deduct half mark if balancing is incorrect)</p>	<p>1</p> <p>1</p>	
24	<p>The curve obtained by plotting the amount of gas adsorbed by the adsorbent with pressure at constant temperature.</p> <p>$x/m = kp^{1/n}$</p> <p style="text-align: center;">OR</p>	<p>1</p> <p>1</p>	
24	<p>The catalytic reaction that depends upon the pore structure of the catalyst and the size of the reactant and product molecules.</p> <p>Shape selective catalysis / Catalytic dehydration .</p>	<p>1</p> <p>1</p>	
25	<p>Rate = k[A][B]</p> <p>Average rate - Rate of a reaction for a particular period or interval of time.</p> <p>Instantaneous rate - Rate of a reaction at a particular instant of time.</p> <p style="text-align: right;">(or any other suitable difference)</p>	<p>1</p> <p>½</p> <p>½</p>	

26.	$\text{Mg}_{(s)} \text{Mg}^{2+}_{(aq)} \text{Ag}^{+}_{(aq)} \text{Ag}_{(s)}$ $E_{\text{cell}} = E^{\circ}_{\text{cell}} - \frac{0.059}{2} \log \frac{[\text{Mg}^{2+}]}{[\text{Ag}^{+}]^2}$	1 1
27	(a) Dissociated (b) Associated	1 1
SECTION C		
28	(a) $\text{CF}_2=\text{CF}_2$ (b)  and $\text{CH}_2\text{OH}-\text{CH}_2\text{OH}$ (c)  / $\text{NH}_2-(\text{CH}_2)_5-\text{COOH}$	1 $\frac{1}{2} + \frac{1}{2}$ 1
28	OR (i) Hexamethylene diamine and Adipic acid (ii) 1,3-Butadiene and Styrene (iii) Chloroprene (or IUPAC names of the monomers)	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ 1
29	(a) the lone pair of nitrogen in aniline is in resonance or conjugation with the benzene ring. (b) Aniline forms salt with anhydrous AlCl_3 . (c) As only alkyl halides undergo nucleophilic substitution.	1 1 1
30	(a) 1-bromobutane < 2-bromobutane < 2-bromo-2-methylpropane. Tertiary carbo cation is more stable than secondary than primary. (b) 2-bromo-2-methyl propane < 2-bromobutane < 1-bromobutane. due to decrease in steric hindrance.	1 $\frac{1}{2}$ 1 $\frac{1}{2}$
31	(a) Potassiumhexacyanomanganate(II) / Potassiumhexacyanomanganate(II) $t_{2g}^5 e_g^0$ (b) Increased stability of the complex due to presence of chelating or didentate or polydentate ligands. e.g. $[\text{Cr}(\text{en})_3]^{3+}$	1 1 $\frac{1}{2}$ $\frac{1}{2}$
31	OR (i) d^2sp^3 , diamagnetic (ii) sp^3d^2 , paramagnetic (iii) sp^3 , diamagnetic	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
32	$\text{Al}_2\text{O}_3 + 2\text{NaOH} + 3\text{H}_2\text{O} \rightarrow 2\text{Na}[\text{Al}(\text{OH})_4]$ $2\text{Na}[\text{Al}(\text{OH})_4] + \text{CO}_2 \rightarrow \text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O} + 2\text{NaHCO}_3$ $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O} \xrightarrow{\Delta} \text{Al}_2\text{O}_3 + x\text{H}_2\text{O}$	1 1 1


33	$\Lambda_m = k \times 1000 / M$ $= 8 \times 10^{-5} \times 1000 / 2 \times 10^{-3}$ $= 40 \text{ Scm}^2 \text{ mol}^{-1}$ $\alpha = \Lambda_m / \Lambda_m^0$ $= 40 / 404$ $= 0.099.$ <p style="text-align: right;">(1/2 mark to be deducted if no or incorrect unit)</p>	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p>
34	$\Delta T_f = K_f m$ $= 1.86 \times 31 \times 1000 / 62 \times 600$ $= 1.55 \text{ }^\circ\text{C or K}$ $\Delta T_f = T_f^0 - T_f$ $T_f = -1.55^\circ\text{C OR } T_f = 271.45\text{K OR } 271.6\text{K}$ <p style="text-align: right;">(1/2 mark to be deducted if no or incorrect unit)</p>	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p>
SECTION D		
35	<p>(a) (i) zero order (ii) slope = -k (iii) $\text{molL}^{-1}\text{s}^{-1}$</p> <p>(b) $k = 2.303/t \log [A_0]/[A]$ $= 2.303/25 \log 100/75$ $= 2.303/25 \log 4/3$ $= 2.303/25(0.6021 - 0.4771)$ $= 0.0115 \text{ min}^{-1}$ $t_{1/2} = 0.693/k = 0.693/0.0115$ $= 60.26 \text{ min or } 60.2 \text{ min}$</p> <p style="text-align: right;">(or by any other suitable method) (deduct half mark for no or incorrect unit)</p>	<p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p>
OR		
35	<p>(a) $t = 2.303/k \log [A_0]/[A]$ $= 2.303/60 \log 1/1/16$ $= 0.046 \text{ s}$</p> <p style="text-align: right;">(or by any other suitable method) (deduct half mark for no or incorrect unit)</p> <p>(b) Concentration of reactants, Temperature, Pressure, surface area and catalyst (any two factors)</p> <p>(c) Proper orientation and energy of colliding particles should be equal to or greater than threshold energy.</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
36	<p>a) A= S₈ B= SO₂, C= SO₃, D= H₂S₂O₇, E= H₂SO₄, F=CuSO₄ b) $\text{Cu} + 2\text{H}_2\text{SO}_4 \rightarrow \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$ c) dehydrating agent, oxidising agent, electrolyte and catalyst (any two)</p> <p style="text-align: center;">OR</p> <p>(a) (i) readily accept one electron to attain noble gas configuration (ii) due to weak dispersion forces / van der Waal forces. (iii) due to smaller size of oxygen as compared to chlorine / due to higher electron density on O than on Cl / due to larger size of chlorine as compared to oxygen.</p> <p>(b) (i) $2\text{NaOH} + \text{Cl}_2 \rightarrow \text{NaCl} + \text{NaOCl} + \text{H}_2\text{O}$ (cold, dil.)</p>	<p>1/2 x 6</p> <p>1</p> <p>1/2 x 2</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>

	(ii) $2\text{I}^-_{(\text{aq.})} + \text{H}_2\text{O}_{(\text{l})} + \text{O}_3_{(\text{g})} \rightarrow \text{I}_{2(\text{s})} + \text{O}_{2(\text{g})} + 2\text{OH}^-_{(\text{aq.})}$	1
37	<p>a) A = $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$ / pentan-3-one</p> $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3 \xrightarrow{\text{Zn-Hg, HCl(conc.)}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ <p>b) i) $\text{CH}_3\text{CH}_2\text{COOH} + \text{Br}_2 \xrightarrow{\text{Red P}} \text{CH}_3\text{CHBrCOOH}$ ii) </p> <p style="text-align: center;">Benzoyl chloride Benzaldehyde</p> <p>c) On heating with NaOH / I_2, acetaldehyde will give yellow ppt of CHI_3, while benzaldehyde doesn't. (OR any other suitable chemical test)</p> <p style="text-align: center;">OR</p>	1 1 1 1
37	<p>a) i) A = $(\text{CH}_3)_2\text{CH}(\text{OH})\text{CH}_2\text{COCH}_3$, B = $(\text{CH}_3)_2\text{CH}=\text{CHCOCH}_3$, C and D = CHI_3 and $(\text{CH}_3)_2\text{CH}=\text{CHCOONa}$</p> <p>ii) 4-Hydroxy-4-methylpentan-2-one</p> <p>b) i) Propanone will give yellow coloured solution with 2,4-DNP but ethanol doesn't. (or any other suitable chemical test) ii) benzoic acid will give brisk effervescence with NaHCO_3 but phenol doesn't. (or any other suitable chemical test)</p>	$\frac{1}{2} \times 4$ 1 1 1


Marking scheme – 2020

CHEMISTRY (043)/ CLASS XII

56/2/2

Q.No	Expected Answer / Value Points	Marks
SECTION A		
1	Acidic aminoacids- will have more number of carboxyl groups than amino groups. / Basic amino acids- will have more number of amino groups than carboxyl groups.	1
2	Those which are not synthesized in our body or must be supplied through diet.	1
3	They contain both acid (-COOH) and basic(NH ₂) group / Zwitter ion form or structure / It can react both acids and bases.	1
4	Peptide linkage/amide linkage/ 	1
5	Simple organic compounds containing both carboxyl (-COOH) and amino (-NH ₂) group. / Building blocks of protein. / Amino substituted carboxylic acid / structure	1
6	Vapour phase refining	1
7	Desorption	1
8	Zinc, Mercury or any other suitable metal	1
9	Linkage	1
10	Order =3	1
11	(C)	1
12	(D)	1
13	(A)	1
14	(D)	1
15	(A)	1
16	(i)	1
17	(iii)	1
18	(iii)	1
19	(i)	1
20	(iii)	1
SECTION B		
21	(a) Due to lone pair -lone pair repulsion on oxygen. (b) Due to resonance in phenol and not in methanol / sp ² hybridised carbon atom in phenol / sp ³ hybridised carbon atom in methanol.	1 1
22	<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>a) Tranquilizers:used to treat mild to severe mental diseases.</p> <p>b) Antiseptics- are applied to living tissues.</p> </div> <div style="width: 45%;"> <p>Analgesics: Used to reduce pain</p> <p>Disinfectants – are applied to inanimate objects or non living objects (or any other correct difference)</p> </div> </div> <p style="text-align: center;">OR</p> <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>Cationic detergent-are quaternary ammonium</p> </div> <div style="width: 45%;"> <p>Anionic detergent – are sodium salts</p> </div> </div>	1 1 1

	salts of amines with acetates, chlorides or bromides as anions. e.g. cetyltrimethyl ammonium bromide.	of sulphonated long chain alcohols or hydrocarbons. e.g. Sodium lauryl sulphate / sodium dodecyl benzene sulphonate.	½ + ½
23	$8 \text{MnO}_4^- + 3\text{S}_2\text{O}_3^{2-} + \text{H}_2\text{O} \rightarrow 8\text{MnO}_2 + 6\text{SO}_4^{2-} + 2\text{OH}^-$ $2\text{MnO}_4^- + 5\text{C}_2\text{O}_4^{2-} + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 10\text{CO}_2$ (deduct 1/2 mark if not balanced)		1 1
24	Rate = k[A][B] Average rate - Rate of a reaction for a particular period or interval of time. Instantaneous rate - Rate of a reaction at a particular instant of time. (or any other suitable difference)		1 ½ ½
25	The curve obtained by plotting the amount of gas adsorbed by the adsorbent with pressure at constant temperature. $x/m = kp^{1/n}$		1 1
25	OR The catalytic reaction that depends upon the pore structure of the catalyst and the size of the reactant and product molecules. Shape selective catalysis / Catalytic dehydration.		1 1
26	$\text{Ni (s) Ni}^{2+} \text{ (aq) Cu}^{2+} \text{ (aq) Cu (s)}$ $E_{\text{cell}} = E_{\text{cell}}^0 - 0.059/2 \log [\text{Ni}^{2+}] / [\text{Cu}^{2+}]$		1 1
27	a) Associated b) Dissociated		1 1
SECTION C			
28	a) $\text{CH}_2 = \text{C}(\text{Cl}) - \text{CH} = \text{CH}_2$ b) $\text{HOOC} - (\text{CH}_2)_4 - \text{COOH}$ and $\text{NH}_2 - (\text{CH}_2)_6 - \text{NH}_2$ c) $\text{CH}_2\text{OH} - \text{CH}_2\text{OH}$ and $\text{HOOC} - \text{C}_6\text{H}_4 - \text{COOH}$ OR		1 x 3
28	i) Amino caproic acid / Caprolactum / 6 -Aminohexanoic acid ii) Vinyl cyanide / Acrylonitrile iii) Melamine and formaldehyde.		1 1 1
29	(a) Potassiumhexacyanidomanganate(II) / Potassiumhexacyanomanganate(II) $t_{2g}^5 e_g^0$ (b) Increased stability of the complex due to presence of chelating or didentate or polydentate ligands. e.g. $[\text{Cr}(\text{en})_3]^{3+}$ OR	(or any other suitable example.)	1 1 ½ ½
29	(i) $d^2 sp^3$, diamagnetic (ii) $sp^3 d^2$, paramagnetic (iii) sp^3 , diamagnetic		½ + ½ ½ + ½ ½ + ½
30	a) $\text{C}_6\text{H}_5\text{CH}_2\text{Br} < \text{C}_6\text{H}_5\text{CH}(\text{Br})\text{CH}_3 < \text{C}_6\text{H}_5\text{C}(\text{Br})(\text{CH}_3)_2$ due to increasing stability of carbocation b) $\text{C}_6\text{H}_5\text{C}(\text{Br})(\text{CH}_3)_2 < \text{C}_6\text{H}_5\text{CH}(\text{Br})\text{CH}_3 < \text{C}_6\text{H}_5\text{CH}_2\text{Br}$ due to decreasing steric hindrance.		1 ½ 1 ½
31	(a) The lone pair of nitrogen in aniline is in resonance or conjugation with the benzene ring. (b) Aniline forms salt with anhydrous AlCl_3 . (c) As only alkyl halides undergo nucleophilic substitution.		1 1 1


32	$4 \text{ Au} + 8 \text{ CN}^- + 2 \text{ H}_2\text{O} + \text{ O}_2 \rightarrow 4[\text{Au}(\text{CN})_2]^- + 4 \text{ OH}^-$ $2 [\text{Au}(\text{CN})_2]^- + \text{ Zn} \rightarrow [\text{Zn}(\text{CN})_4]^{2-} + 2 \text{ Au}$ <p>Zn acts as a reducing agent.</p>	1 1 1
33	<p>Limiting molar conductivity of an electrolyte can be represented as the sum of the individual contributions of the anion and cation of the electrolyte.</p> <p>Molar conductivity of $\text{Sr}(\text{NO}_3)_2 = \lambda^\circ_{\text{Sr}^{2+}} + 2 \lambda^\circ_{\text{NO}_3^-}$ $= 119 + (2 \times 72)$ $= 263 \text{ Scm}^2/\text{mol}$ (deduct ½ mark if no or incorrect unit)</p>	1 ½ ½ 1
34	$\Delta T_f = K_f m$ $= 1.86 \times 31 \times 1000 / 62 \times 600$ $= 1.55^\circ\text{C or K}$ $\Delta T_f = T_f^\circ - T_f$ $T_f = -1.55^\circ\text{C OR } T_f = 271.45\text{K OR } 271.6\text{K}$ <p>(1/2 mark to be deducted if no or incorrect unit)</p>	½ ½ 1 1
Section D		
35	<p>a) A = $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$ / pentan-3-one</p> $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3 \xrightarrow{\text{Zn-Hg, HCl(conc.)}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ <p>b) i) $\text{CH}_3\text{CH}_2\text{COOH} + \text{Br}_2 \xrightarrow{\text{Red P}} \text{CH}_3\text{CHBrCOOH}$</p> <p>ii)</p> <div style="text-align: center;">  <p style="text-align: center;">Benzoyl chloride Benzaldehyde</p> </div> <p>c) On heating with NaOH / I_2, acetaldehyde will give yellow ppt of CHI_3, while benzaldehyde doesn't. (OR any other suitable chemical test)</p> <p style="text-align: center;">OR</p>	1 1 1 1 1
35	<p>a) i) A = $(\text{CH}_3)_2\text{CH}(\text{OH})\text{CH}_2\text{COCH}_3$, B = $(\text{CH}_3)_2\text{CH}=\text{CHCOCH}_3$, C and D = CHI_3 and $(\text{CH}_3)_2\text{CH}=\text{CHCOONa}$ ii) 4-Hydroxy-4-methylpentan-2-one</p> <p>b) i) Propanone will give yellow coloured solution with 2,4-DNP but ethanol doesn't. (or any other suitable chemical test) ii) benzoic acid will give brisk effervescence with NaHCO_3 but phenol doesn't. (or any other suitable chemical test)</p>	½ x 4 1 1 1

36	<p>(a) (i) zero order (ii) slope = - k (iii) molL⁻¹s⁻¹</p> <p>(b) $k = 2.303/t \log [A_0]/[A]$ $= 2.303/25 \log 100/75$ $= 2.303/25 \log 4/3$ $= 2.303/25(0.6021 - 0.4771)$ $= 0.0115 \text{ min}^{-1}$ $t_{1/2} = 0.693/k = 0.693/0.0115$ $= 60.26 \text{ min or } 60.2 \text{ min}$</p> <p style="text-align: right;">(or by any other suitable method) (deduct half mark for no or incorrect unit)</p> <p style="text-align: center;">OR</p> <p>(a) $t = 2.303/k \log [A_0]/[A]$ $= 2.303/ 60 \log 1/1/16$ $= 0.046 \text{ s}$</p> <p style="text-align: right;">(or by any other suitable method) (deduct half mark for no or incorrect unit)</p> <p>(b) Concentration of reactants, Temperature, Pressure, surface area and catalyst (any two factors)</p> <p>(c) Proper orientation and energy should be equal to or greater than threshold energy.</p>	<p>1 1 1</p> <p>½ ½</p> <p>1</p> <p>1 1 1</p> <p>1 1 1</p>
37	<p>(a) A= S₈ B= SO₂, C= SO₃, D= H₂S₂O₇, E= H₂SO₄, F=CuSO₄</p> <p>(b) $\text{Cu} + 2\text{H}_2\text{SO}_4 \rightarrow \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$</p> <p>(c) Dehydrating agent, oxidising agent, electrolyte in automobile batteries and catalyst (any two)</p> <p style="text-align: center;">OR</p> <p>(a) (i) Readily accept one electron to attain noble gas configuration (ii) Due to weak dispersion forces / van der Waal forces. (iii) Due to smaller size of oxygen as compared to chlorine / due to higher electron density on O than on Cl / due to larger size of chlorine as compared to oxygen.</p> <p>(b) (i) $2\text{NaOH} + \text{Cl}_2 \rightarrow \text{NaCl} + \text{NaOCl} + \text{H}_2\text{O}$ (cold, dil.)</p> <p>(ii) $2\text{I}^-_{(\text{aq.})} + \text{H}_2\text{O}_{(\text{l})} + \text{O}_{3(\text{g})} \rightarrow \text{I}_{2(\text{s})} + \text{O}_{2(\text{g})} + 2\text{OH}^-_{(\text{aq.})}$</p>	<p>½ x 6 1 ½ x 2</p> <p>1 1 1</p> <p>1</p> <p>1</p>

Marking scheme – 2020

CHEMISTRY (043)/ CLASS XII

56/2/3

Q.No	Expected Answer / Value Points	Marks
SECTION A		
1	Peptide linkage/amide linkage/ 	1
2	Those which are not synthesized in our body or must be supplied through diet.	1
3	Simple organic compounds containing both carboxyl (—COOH) and amino (—NH ₂) group. / Building blocks of protein. / Amino substituted carboxylic acid / structure	1
4	They contain both acid (-COOH) and basic(NH ₂) group / Zwitter ion form or structure / It can react both acids and bases.	1
5	Acidic aminoacids- will have more number of carboxyl groups than amino groups. / Basic amino acids- will have more number of amino groups than carboxyl groups.	1
6	Hydraulic washing / gravity separation	1
7	Linkage	1
8	Desorption	1
9	Zinc / Mercury or any other suitable metal	1
10	1	1
11	(A)	1
12	(C)	1
13	(D)	1
14	(C)	1
15	(A)	1
16	(i)	1
17	(iii)	1
18	(iii)	1
19	(i)	1
20	(iii)	1
SECTION B		
21	(a) Associated (b) Associated	1 1
22	Pt I _(aq.) I _{2(s)} F _{2(g)} F _(aq.) Pt (Pt may be ignored) $E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.059}{2} \log \frac{[F^-]^2}{[I]^-2}$	1 1
23	a) $5\text{Fe}^{2+} + \text{MnO}_4^- + 8\text{H}^+ \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O} + 5\text{Fe}^{3+}$ b) $2\text{MnO}_4^- + 3\text{Mn}^{2+} + 2\text{H}_2\text{O} \rightarrow 5\text{MnO}_2 + 4\text{H}^+$	1 1

24	<p>a) Tranquilizers: Used to treat mild to severe mental diseases.</p> <p>b) Antiseptics- are applied to living tissues.</p> <p style="text-align: center;">OR</p> <p>Cationic detergent-are quaternary ammonium salts of amines with acetates, chlorides or bromides as anions. e.g. cetyltrimethyl ammonium bromide.</p>	<p>Analgesics: Used to reduce pain</p> <p>Disinfectants – are applied to inanimate objects or non living objects (or any other correct difference)</p> <p>Anionic detergent – are sodium salts of sulphonated long chain alcohols or hydrocarbons. e.g. Sodium lauryl sulphate / sodium dodecyl benzene sulphonate. (or any other suitable example)</p>	<p>1</p> <p>1</p> <p>1</p> <p>½ + ½</p>
25	<p>(a) Due to lone pair -lone pair repulsion on oxygen.</p> <p>(b) Due to resonance in phenol and not in methanol / sp^2 hybridised carbon atom in phenol / sp^3 hybridised carbon atom in methanol.</p>		<p>1</p> <p>1</p>
26	<p>The curve obtained by plotting the amount of gas adsorbed by the adsorbent with pressure at constant temperature. $x/m = kp^{1/n}$</p> <p style="text-align: center;">OR</p>		<p>1</p> <p>1</p>
26	<p>The catalytic reaction that depends upon the pore structure of the catalyst and the size of the reactant and product molecules. Shape selective catalysis / Catalytic dehydration .</p>		<p>1</p> <p>1</p>
27	<p>Rate = $k[A][B]$ Average rate - Rate of a reaction for a particular period or interval of time. Instantaneous rate - Rate of a reaction at a particular instant of time. (or any other suitable difference)</p>		<p>1</p> <p>½</p> <p>½</p>
SECTION C			
28	<p>a) $CH_2 = C(CH_3) - CH = CH_2$ b) $CH_2 = CH - CH = CH_2$ and $C_6H_5 - CH = CH_2$ c) $C_6H_5 - OH$ and $HCHO$</p> <p style="text-align: center;">OR</p>		<p>1</p> <p>1</p> <p>1</p>
28	<p>i) Ethylene glycol and Terephthalic acid ii) Tetrafluoro ethene iii) β-Hydroxy butyric acid and β-Hydroxy valeric acid / 3-Hydroxybutanoic acid and 3-Hydroxypentanoic acid</p>		<p>1</p> <p>1</p> <p>1</p>
29	<p>$4 Ag + 8CN^- + 2H_2O + O_2 \rightarrow 4[Ag(CN)_2]^- + 4OH^-$</p> <p>$2 [Ag(CN)_2]^- + Zn \rightarrow [Zn(CN)_4]^{2-} + 2 Ag$</p> <p>Zn acts as a reducing agent.</p>		<p>1</p> <p>1</p> <p>1</p>
30	<p>Limiting molar conductivity of an electrolyte can be represented as the sum of the individual contributions of the anion and cation of the electrolyte. Molar conductivity of $Ba(OH)_2 = \lambda^{\circ} Ba^{2+} + 2 \lambda^{\circ} OH^-$ $= 127 + (2 \times 199)$ $= 525 Scm^2/mol$ (deduct ½ mark if no or incorrect unit)</p>		<p>1</p> <p>½</p> <p>½</p> <p>1</p>

31	$\Delta T_f = K_f m$ $= 1.86 \times 31 \times 1000 / 62 \times 600$ $= 1.55^\circ\text{C or K}$ $\Delta T_f = T_f^\circ - T_f$ $T_f = -1.55^\circ\text{C or } T_f = 271.45\text{K or } 271.6\text{K}$	$\frac{1}{2}$ $\frac{1}{2}$ 1 1
	(deduct $\frac{1}{2}$ mark if no or incorrect unit)	
32	(a) the lone pair of nitrogen in aniline is in resonance or conjugation with the benzene ring. (b) Aniline forms salt with anhydrous AlCl_3 . (c) As only alkyl halides undergo nucleophilic substitution.	1 1 1
33	(a) Potassiumhexacyanomanganate(II) / Potassiumhexacyanomanganate(II) $t_{2g}^5 e_g^0$ (b) Increased stability of the complex due to presence of chelating or didentate or polydentate ligands. e.g. $[\text{Cr}(\text{en})_3]^{3+}$	1 1 $\frac{1}{2}$
	(or any other suitable example.)	$\frac{1}{2}$
	OR	
33	(i) $d^2 sp^3$, diamagnetic (ii) $sp^3 d^2$, paramagnetic (iii) sp^3 , diamagnetic	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
34	a) ethyl chloride < isopropyl chloride < tertiarybutyl chloride Due to increasing stability of carbocation	1 $\frac{1}{2}$
	b) tertiarybutylchloride < isopropylchloride < ethylchloride Due to decreasing steric hindrance.	1 $\frac{1}{2}$
	SECTION D	
35	a) A= S_8 B= SO_2 , C= SO_3 , D= $\text{H}_2\text{S}_2\text{O}_7$, E= H_2SO_4 , F= CuSO_4 b) $\text{Cu} + 2\text{H}_2\text{SO}_4 \rightarrow \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$ c) dehydrating agent, oxidising agent, electrolyte and catalyst	$\frac{1}{2} \times 6$ 1 $\frac{1}{2} \times 2$
	OR	
35	(a) (i) readily accept one electron to attain noble gas configuration (ii) due to weak dispersion forces / vander Waal forces. (iii) due to smaller size of oxygen as compared to chlorine / due to higher electron density on O than on Cl / due to larger size of chlorine as compared to oxygen.	1 1 1
	(b) (i) $2\text{NaOH} + \text{Cl}_2 \rightarrow \text{NaCl} + \text{NaOCl} + \text{H}_2\text{O}$ (cold, dil.)	1
	(ii) $2\text{I}^-_{(\text{aq.})} + \text{H}_2\text{O}_{(\text{l})} + \text{O}_3_{(\text{g})} \rightarrow \text{I}_2_{(\text{s})} + \text{O}_2_{(\text{g})} + 2\text{OH}^-_{(\text{aq.})}$	1
36	a) A= $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$ / pentan-3-one $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3 \xrightarrow{\text{Zn-Hg, HCl(conc.)}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$	1 1
	b) i) $\text{CH}_3\text{CH}_2\text{COOH} + \text{Br}_2 \xrightarrow{\text{Red P}} \text{CH}_3\text{CHBrCOOH}$	1

