

**CBSE Class 12 Chemistry Question Paper
Solution 2020 Set 56/3/1**

Set – 1 (56/3/1)

MARKING SCHEME

SR. SECONDARY SCHOOL EXAMINATION, 2020





Subject: CHEMISTRY




Q.No	Expected Answer / Value Points	Distribution of Marks
SECTION - A		
1.	Zn , Cd and Hg have completely filled d^{10} configuration in their ground state as well as in their oxidized state.	1
2.	Scandium / Sc	1
3.	Because of multiple oxidation states / ability to form complexes / having large surface area.	1
4.	Involvement of (n-1)d and ns electrons in inter atomic metallic bonding / strong metal-metal bonding.	1
5.	Presence of unpaired e^- showing d-d transition in Cu^{2+} , while in Zn^{2+} there is no unpaired electron.	1
6.	$H_2 - O_2$ Fuel cell	1
7.	$2 \times 96500 C / 193000 C$	1
8.	Slope = -k	1
9.	Maltose	1
10.	Anionic detergent	1
11.	(C)	1
12.	(B)	1
13.	(B)	1
14.	(A)	1
15.	(C)	1
16.	(iii)	1
17.	(i)	1
18.	(iv)	1
19.	(i)	1
20.	(iv)	1

SECTION – B





21.	$\text{Rate} = -\frac{\Delta[A]}{\Delta t} = -\frac{1}{3} \frac{\Delta[B]}{\Delta t} = \frac{1}{2} \frac{\Delta[C]}{\Delta t}$ <p>(i) $\text{Rate} = \frac{1}{2} \frac{\Delta[C]}{\Delta t}$ $2 \times \text{Rate} = \frac{\Delta[C]}{\Delta t} = 2.5 \times 10^{-4}$ $\text{Rate} = \frac{2.5 \times 10^{-4}}{2} = 1.25 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$</p> <p>ii) $\text{Rate} = -\frac{1}{3} \frac{\Delta[B]}{\Delta t}$ $1.25 \times 10^{-4} = -\frac{1}{3} \frac{\Delta[B]}{\Delta t}$ $-\frac{\Delta[B]}{\Delta t} = 3 \times 1.25 \times 10^{-4} = 3.75 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$</p>	<p style="text-align: right;">½</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p>						
22.	<p>i) Acts as solvent / lowers the melting point of the mixture / Increases the conductivity.</p> <p>ii) It forms a volatile compound Ni(CO)₄, which decomposes at higher temperature to give pure Nickel.</p> <p style="text-align: center;">OR</p> $\text{Al}_2\text{O}_3 + 2 \text{NaOH} + 3\text{H}_2\text{O} \rightarrow 2 \text{Na} [\text{Al}(\text{OH})_4]$ $2\text{Na} [\text{Al}(\text{OH})_4] + \text{CO}_2 \rightarrow \text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O} + 2 \text{NaHCO}_3$ $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O} \xrightarrow{\Delta} \text{Al}_2\text{O}_3 + x\text{H}_2\text{O}$	<p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p>						
23.	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #f08080;"> <th style="padding: 5px;">Physisorption</th> <th style="padding: 5px;">Chemisorption</th> </tr> </thead> <tbody> <tr> <td style="padding: 5px;">1. It arises because of van der Waals' forces.</td> <td style="padding: 5px;">1. It is caused by chemical bond formation.</td> </tr> <tr> <td style="padding: 5px;">2. It is not specific in nature.</td> <td style="padding: 5px;">2. It is highly specific in nature.</td> </tr> </tbody> </table> <p style="text-align: center;">(Or any other two correct differences)</p> <p style="text-align: center;">OR</p>	Physisorption	Chemisorption	1. It arises because of van der Waals' forces.	1. It is caused by chemical bond formation.	2. It is not specific in nature.	2. It is highly specific in nature.	<p style="text-align: right;">1</p> <p style="text-align: right;">1</p>
Physisorption	Chemisorption							
1. It arises because of van der Waals' forces.	1. It is caused by chemical bond formation.							
2. It is not specific in nature.	2. It is highly specific in nature.							
23.	<p>i) There are some substances which at low concentration behave as normal strong electrolytes, but at higher concentration exhibit colloidal behavior due to the formation of aggregates. Example: Micelles / soap</p> <p>ii) Oil is dispersed phase and water is dispersion medium. Example : Milk (or any other correct example)</p>	<p style="text-align: right;">½ + ½</p> <p style="text-align: right;">½ + ½</p>						
24.	<p>a) Hexafluoridocobaltate(III) sp^3d^2</p> <p>b) Isomerism – Geometrical / optical cis isomer is optically active</p>	<p style="text-align: right;">½ + ½</p> <p style="text-align: right;">½ + ½</p>						
25.	<p>(i) To prevent rapid drying.</p> <p>(ii) Antacids will make the stomach alkaline and trigger the production of more acid.</p>	<p style="text-align: right;">1</p> <p style="text-align: right;">1</p>						
26.	<p>i) Carbohydrates that yield two to ten monosaccharides units on hydrolysis.</p> <p>ii) Hydrolysis of sucrose brings about a change in the sign of rotation, from dextro (+) to laevo (-) and the product is named as invert sugar.</p>	<p style="text-align: right;">1</p> <p style="text-align: right;">1</p>						
27.	$6\text{XeF}_4 + 12 \text{H}_2\text{O} \rightarrow 4\text{Xe} + 2\text{XeO}_3 + 24\text{HF} + 3\text{O}_2$ <p>Yes Xe^{+4} changes to Xe^0 and Xe^{+6} / Xe gets oxidized as well as reduced in the same reaction.</p>	<p style="text-align: right;">1</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p>						

SECTION – C

28.	$\Delta G^\circ = -nF E^\circ_{\text{cell}}$ $= -2 \times 96500 \times \{0.80 - (-0.25)\}$ $= -202650 \text{ J/ mol}$ <p>Maximum work = 202650 J/ mol</p> $E^\circ_{\text{cell}} = \frac{0.059}{2} \log Kc$ $\log Kc = \frac{2 \times 1.15}{0.059}$ $= 35.6$	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p>
29.	$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$ $= \frac{2.303}{80} \log \frac{100}{60}$ $= \frac{2.303}{80} \times (1 - 0.7782)$ $= 0.0064 \text{ min}^{-1}$ $t = \frac{2.303}{k} \log \frac{[A]_0}{[A]}$ $= \frac{2.303}{0.0064} \log \frac{100}{10}$ $= 360 \text{ min}$ <p>(deduct 1/2 mark for no or incorrect unit)</p>	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p>
30.	<p>(i)</p> <p>$\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$, </p> <p>1, 3-Butadiene , Styrene</p> <p>ii)</p> <p>$\text{HOOC}(\text{CH}_2)_4\text{COOH}$, $\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2$</p> <p>Adipic acid and Hexamethylene diamine</p> <p>iii)</p> <p> + HCHO , Phenol and formaldehyde</p>	<p>1/2 x 6</p>
31.	<p>i) </p> <p>ii) </p> <p>iii) $\text{H}_3\text{C} - \text{C}(\text{CH}_3) = \text{CH}_2$</p>	<p>1</p> <p>1</p> <p>1</p>

32.	<p>a.</p> $\text{CH}_3\text{-CH}_2\text{-}\ddot{\text{O}}\text{-H} + \text{H}^+ \rightarrow \text{CH}_3\text{-CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-H}$ $\text{CH}_3\text{CH}_2\text{-}\ddot{\text{O}}\text{:} + \text{CH}_3\text{-CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-H} \rightarrow \text{CH}_3\text{CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-CH}_2\text{CH}_3 + \text{H}_2\text{O}$ $\text{CH}_3\text{CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-CH}_2\text{CH}_3 \rightarrow \text{CH}_3\text{CH}_2\text{-O-CH}_2\text{CH}_3 + \text{H}^+$ <p>b.</p>  <p style="text-align: center;">OR</p>	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p>
32.	<p>i)</p>  <p>ii) $\text{CH}_3\text{COCH}_3 \xrightarrow{\text{LiAlH}_4} \text{CH}_3\text{CH(OH)CH}_3 \xrightarrow[443\text{ K}]{\text{H}_2\text{SO}_4(\text{conc})} \text{CH}_3\text{CH=CH}_2$</p> <p>iii)</p>  <p style="text-align: right;">(or any other correct method)</p>	<p>1</p> <p>1</p> <p>1</p>
33.	<p>i) $(\text{CH}_3)_3\text{CCH(OH)CH}_2\text{COCH}_3$ ii) $(\text{CH}_3)_3\text{CCH(OH)CN}$ iii) $(\text{CH}_3)_3\text{CCOONa} + (\text{CH}_3)_3\text{CCH}_2\text{OH}$</p>	<p>1</p> <p>1</p> <p>1/2 + 1/2</p>
34.	<p>Lyophobic sol is liquid or water repelling. Example: Fe(OH)_3 sol / gold sol (or any other suitable example) Lyophilic sol is liquid or water attracting. Example: starch sol / rubber sol (or any other suitable example)</p> <p>Due to the charge and solvation of the colloidal particles.</p> <p style="text-align: center;">OR</p>	<p>1/2 + 1/2</p> <p>1/2 + 1/2</p> <p>1/2 + 1/2</p>
34.	<p>i) The catalytic reaction that depends upon the pore structure of the catalyst and the size of the reactant and product molecules.</p> <p>ii) The formation of micelles takes place only above a particular temperature is called Kraft temperature (T_k)</p> <p>iii) Process of converting a precipitate into colloidal sol by shaking it with dispersion medium in the presence of a small amount of electrolyte.</p>	<p>1</p> <p>1</p> <p>1</p>


SECTION – D


<p>35.</p>	<p>a) i) Due to high ionization enthalpy ii) Because of higher oxidation state of Cl in HClO₄ than in HOCl / ClO₄⁻ is more stable than ClO⁻ iii) Because oxygen can form pπ- pπ multiple bond effectively but sulphur can not. b) i) $\text{Cu} + 2 \text{H}_2\text{SO}_4(\text{conc.}) \rightarrow \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$ ii) $\text{C}_{12}\text{H}_{22}\text{O}_{11} \xrightarrow{\text{H}_2\text{SO}_4} 12\text{C} + 11\text{H}_2\text{O}$ (or any other suitable reaction in both above cases) OR a) i) Because of smaller size of F⁻ ion than Cl⁻ ion. ii) Because sulphur is more stable in +6 state and Tellurium is more stable in +4 state b) $2\text{F}_2 + 2\text{H}_2\text{O} \longrightarrow 4\text{HF} + \text{O}_2$ Because I₂ is a weak oxidizing agent. c)</p> 	<p>1 1 1 1 1 1 1 1 1 1</p>
<p>36.</p>	<p>a) i) Because aniline gets protonated to give anilinium ion which is deactivating in nature and is meta directing. ii) Because of combined factors of solvation and inductive effects. iii) Because it gives a mixture of amines which is difficult to separate. b) i) On heating with CHCl₃ and KOH (alcoholic) CH₃CH₂NH₂ gives a foul smelling isocyanide while (CH₃CH₂)₂NH doesn't. ii) On adding benzenediazonium chloride, aniline gives a yellow coloured dye while CH₃NH₂ doesn't. (or any other suitable chemical test) OR</p> <p>a) i) A =  B =  ii) A = CH₃CONH₂ B = CH₃NH₂</p> <p>b)  N-Methylbenzamide</p> <p>c) (C₂H₅)₂NH < C₂H₅NH₂ < NH₃ < C₆H₅NH₂</p>	<p>1 1 1 1 1 1/2 + 1/2 1/2 + 1/2 1 1 1</p>
<p>37.</p>	<p>a) $\pi = i \text{ CRT}$ $4.75 = i \times \frac{5.85}{58.5} \times \frac{1}{1} \times 0.082 \times 300$ $i = 1.93$ $\alpha = \frac{i-1}{n-1} = \frac{1.93-1}{2-1} = 0.93$ or 93% b) Partial pressure of gas in liquid is directly proportional to its solubility or</p>	<p>1/2 1/2 1 1 1</p>




	mole fraction. To prevent 'Bends'	
	OR	
37.	a) $\Delta T_f = i K_f m$ $1 = i \times 1.86 \times \frac{19.5}{78} \times \frac{1000}{500}$ $i = 1.075$ $\alpha = \frac{i-1}{n-1} = \frac{1.075-1}{2-1} = 0.075$ or 7.5%	1 1/2 1/2 1
	b) i) Due to dissociation of KCl / number of particles in 0.1 M KCl is more.	1
	ii) Due to osmosis bacteria loses its water and dies which causes preservation.	1










Set – (56/3/2)
MARKING SCHEME
SR. SECONDARY SCHOOL EXAMINATION, 2020
Subject: CHEMISTRY

Q.No	Expected Answer / Value Points	Distribution of Marks
SECTION - A		
1.	Zn , Cd and Hg have completely filled d^{10} configuration in their ground state as well as in their oxidized state.	1
2.	Scandium / Sc	1
3.	Because of multiple oxidation states / ability to form complexes / having large surface area.	1
4.	Involvement of (n-1)d and ns electrons in inter atomic metallic bonding / strong metal-metal bonding.	1
5.	Presence of unpaired e^- showing d-d transition in Cu^{2+} , while in Zn^{2+} there is no unpaired electron.	1
6.	p-dichlorobenzene	1
7.	Benzene / C_6H_6 / 	1
8.	$H_{2(g)}$ / Hydrogen gas	1
9.	Lactose	1
10.	Tranquilizers	1
11.	(D)	1
12.	(B)	1
13.	(C)	1
14.	(D)	1
15.	(C)	1
16.	(iv)	1
17.	(iii)	1
18.	(iv)	1
19.	(i)	1
20.	(i)	1
SECTION – B		
21.	i) Acts as solvent / lowers the melting point of the mixture / Increases the conductivity. ii) It forms a volatile compound $Ni(CO)_4$, which decomposes at higher temperature to give pure Nickel. OR $Al_2O_3 + 2 NaOH + 3H_2O \rightarrow 2 Na [Al(OH)_4]$ $2Na [Al(OH)_4] + CO_2 \rightarrow Al_2O_3 \cdot xH_2O + 2 NaHCO_3$ $Al_2O_3 \cdot xH_2O \xrightarrow{\Delta} Al_2O_3 + xH_2O$	1 1 1 ½ ½

22.	$\text{Rate} = -\frac{\Delta[A]}{\Delta t} = -\frac{1}{3} \frac{\Delta[B]}{\Delta t} = \frac{1}{2} \frac{\Delta[C]}{\Delta t}$ <p>(i) $\text{Rate} = \frac{1}{2} \frac{\Delta[C]}{\Delta t}$ $2 \times \text{Rate} = \frac{\Delta[C]}{\Delta t} = 2.5 \times 10^{-4}$ $\text{Rate} = \frac{2.5 \times 10^{-4}}{2} = 1.25 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$</p> <p>ii) $\text{Rate} = -\frac{1}{3} \frac{\Delta[B]}{\Delta t}$ $1.25 \times 10^{-4} = -\frac{1}{3} \frac{\Delta[B]}{\Delta t}$ $-\frac{\Delta[B]}{\Delta t} = 3 \times 1.25 \times 10^{-4} = 3.75 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$</p>	<p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>						
23.	<p>a) Hexafluoridocobaltate(III) sp^3d^2</p> <p>b) Isomerism – Geometrical / optical Cis isomer is optically active</p>	<p>1/2 + 1/2</p> <p>1/2 + 1/2</p>						
24.	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #f8d7da;"> <th style="padding: 5px;">Physisorption</th> <th style="padding: 5px;">Chemisorption</th> </tr> </thead> <tbody> <tr> <td style="padding: 5px;">1. It arises because of van der Waals' forces.</td> <td style="padding: 5px;">1. It is caused by chemical bond formation.</td> </tr> <tr> <td style="padding: 5px;">2. It is not specific in nature.</td> <td style="padding: 5px;">2. It is highly specific in nature.</td> </tr> </tbody> </table> <p style="text-align: center;">(or any other two correct differences)</p> <p style="text-align: center;">OR</p> <p>24. iii) There are some substances which at low concentration behave as normal strong electrolytes, but at higher concentration exhibit colloidal behavior due to the formation of aggregates. Example: Micelles / soap</p> <p>iv) Oil is dispersed phase and water is dispersion medium. Example : Milk</p> <p style="text-align: right;">(or any other correct example)</p>	Physisorption	Chemisorption	1. It arises because of van der Waals' forces.	1. It is caused by chemical bond formation.	2. It is not specific in nature.	2. It is highly specific in nature.	<p>1</p> <p>1</p> <p>1/2 + 1/2</p> <p>1/2 + 1/2</p>
Physisorption	Chemisorption							
1. It arises because of van der Waals' forces.	1. It is caused by chemical bond formation.							
2. It is not specific in nature.	2. It is highly specific in nature.							
25.	$\text{XeF}_6 + 2\text{H}_2\text{O} \longrightarrow \text{XeO}_2\text{F}_2 + 4\text{HF}$ <p>No.</p> 	<p>1</p> <p>1/2</p> <p>1/2</p>						
26.	<p>i) A linkage between two monosaccharide units through oxygen atom.</p> <p>ii) Protein having a unique three-dimensional structure and biological activity.</p>	<p>1</p> <p>1</p>						
27.	<p>i) Because these are biodegradable</p> <p>ii) Because it is unstable at cooking temperature</p>	<p>1</p> <p>1</p>						
28.	$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$ $= \frac{2.303}{80} \log \frac{100}{60}$ $= \frac{2.303}{80} \times (1 - 0.7782)$ $= 0.0064 \text{ min}^{-1}$ $t = \frac{2.303}{k} \log \frac{[A]_0}{[A]}$ $= \frac{2.303}{0.0064} \log \frac{100}{10}$ $= 360 \text{ min}$	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p>						

29.	$\Delta G^{\circ} = -nF E^{\circ}_{\text{cell}}$ $= -2 \times 96500 \times \{0.80 - (-0.25)\}$ $= -202650 \text{ J/mol}$ <p>Maximum work = 202650 J/mol</p> $E^{\circ}_{\text{cell}} = \frac{0.059}{2} \log Kc$ $\log Kc = \frac{2 \times 1.15}{0.059}$ $= 35.6$	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p>
30.	<p>i) </p> <p>ii) </p> <p>iii) $\text{H}_3\text{C}-\underset{\text{CH}_3}{\text{C}}=\text{CH}_2$</p>	<p>1</p> <p>1</p> <p>1</p>
31.	<p>i) $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2, \text{CH}_2=\underset{\text{CN}}{\text{CH}}$ 1,3-Butadiene Acrylonitrile</p> <p>ii) $\text{H}_2\text{C}=\underset{\text{CH}_3}{\text{C}}-\text{CH}=\text{CH}_2$ Isoprene</p> <p>iii)  Caprolactam or $\text{NH}_2(\text{CH}_2)_5\text{COOH}$ and Aminocaproic acid</p>	<p>1/2 + 1/2</p> <p>1/2 + 1/2</p> <p>1/2 + 1/2</p>
32.	<p>Lyophobic sol is liquid or water repelling. Example: $\text{Fe}(\text{OH})_3$ sol / gold sol (or any other suitable example)</p> <p>Lyophilic sol is liquid or water attracting. Example: starch sol / rubber sol (or any other suitable example)</p> <p>Due to the charge and solvation of the colloidal particles.</p> <p style="text-align: center;">OR</p> <p>i) The catalytic reaction that depends upon the pore structure of the catalyst and the size of the reactant and product molecules.</p> <p>ii) The formation of micelles takes place only above a particular temperature is called Kraft temperature (T_k)</p> <p>iii) Process of converting a precipitate into colloidal sol by shaking it with dispersion medium in the presence of a small amount of electrolyte.</p>	<p>1/2 + 1/2</p> <p>1/2 + 1/2</p> <p>1/2 + 1/2</p> <p>1</p> <p>1</p> <p>1</p>






33.	<p>a.</p> $\text{CH}_3\text{-CH}_2\text{-}\ddot{\text{O}}\text{-H} + \text{H}^+ \rightarrow \text{CH}_3\text{-CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-H}$ $\text{CH}_3\text{CH}_2\text{-}\ddot{\text{O}}\text{:} + \text{CH}_3\text{-CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-H} \rightarrow \text{CH}_3\text{CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-CH}_2\text{CH}_3 + \text{H}_2\text{O}$ $\text{CH}_3\text{CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-CH}_2\text{CH}_3 \rightarrow \text{CH}_3\text{CH}_2\text{-O-CH}_2\text{CH}_3 + \text{H}^+$ <p>b.</p>  <p style="text-align: center;">OR</p> <p>i)</p>  <p>ii) $\text{CH}_3\text{COCH}_3 \xrightarrow{\text{LiAlH}_4} \text{CH}_3\text{CHOHCH}_3 \xrightarrow[443\text{ K}]{\text{H}_2\text{SO}_4(\text{conc})} \text{CH}_3\text{CH}=\text{CH}_2$</p> <p>iii)</p> 	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1</p>
34.	<p>i) $(\text{CH}_3)_3\text{CCH}(\text{OH})\text{CH}_2\text{COCH}_3$ ii) $(\text{CH}_3)_3\text{CCH}(\text{OH})\text{CN}$ iii) $(\text{CH}_3)_3\text{CCOONa} + (\text{CH}_3)_3\text{CCH}_2\text{OH}$</p>	<p>1</p> <p>1</p> <p>1/2 + 1/2</p>
SECTION - D		
35.	<p>a) $\pi = i \text{ CRT}$ $4.75 = i \times \frac{5.85}{58.5} \times \frac{1}{1} \times 0.082 \times 300$ $i = 1.93$ $\alpha = \frac{i-1}{n-1} = \frac{1.93-1}{2-1} = 0.93$ or 93%</p> <p>b) Partial pressure of gas in liquid is directly proportional to its solubility or mole fraction. To prevent 'Bends'</p> <p style="text-align: center;">OR</p> <p>a) $\Delta T_f = i K_f m$ $1 = i \times 1.86 \times \frac{19.5}{78} \times \frac{1000}{500}$ $i = 1.075$ $\alpha = \frac{i-1}{n-1} = \frac{1.075-1}{2-1} = 0.075$ or 7.5%</p> <p>b) i) Due to dissociation of KCl / number of particles in 0.1 M KCl is more. ii) Due to osmosis bacteria loses its water and dies which causes preservation.</p>	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>

36.	<p>a) i) Due to high ionization enthalpy ii) Because of higher oxidation state of Cl in HClO_4 than in $\text{HOCl} / \text{ClO}_4^-$ is more stable than ClO^- iii) Because oxygen can form $p\pi-p\pi$ multiple bond effectively but sulphur can not.</p> <p>b) i) $\text{Cu} + 2 \text{H}_2\text{SO}_4(\text{conc.}) \rightarrow \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$ ii) $\text{C}_{12}\text{H}_{22}\text{O}_{11} \xrightarrow{\text{H}_2\text{SO}_4} 12\text{C} + 11\text{H}_2\text{O}$ (or any other suitable reaction in both above cases)</p> <p style="text-align: center;">OR</p> <p>a) i) Because of smaller size of F^- ion than Cl^- ion. ii) Because sulphur is more stable in +6 state and Tellurium is more stable in +4 state b) $2\text{F}_2 + 2\text{H}_2\text{O} \longrightarrow 4\text{HF} + \text{O}_2$ Because I_2 is a weak oxidizing agent. c)</p> <div style="text-align: center;">  </div>	<p>1 1 1 1 1 1 1 1 1 1</p>
37.	<p>a) i) Because aniline gets protonated to give anilinium ion which is deactivating in nature and is meta directing. ii) Because of combined factors of solvation and inductive effects. iii) Because it gives a mixture of amines which is difficult to separate.</p> <p>b) i) On heating with CHCl_3 and KOH (alcoholic) $\text{CH}_3\text{CH}_2\text{NH}_2$ gives a foul smelling isocyanide while $(\text{CH}_3\text{CH}_2)_2\text{NH}$ doesn't. ii) On adding benzenediazonium chloride, aniline gives a yellow coloured dye while CH_3NH_2 doesn't. (or any other suitable chemical test)</p> <p style="text-align: center;">OR</p> <p>a) i) A =  B = </p> <p>ii) A = CH_3CONH_2 B = CH_3NH_2</p> <p>b)  N-Methylbenzamide</p> <p>c) $(\text{C}_2\text{H}_5)_2\text{NH} < \text{C}_2\text{H}_5\text{NH}_2 < \text{NH}_3 < \text{C}_6\text{H}_5\text{NH}_2$</p>	<p>1 1 1 1 1 1/2 + 1/2 1/2 + 1/2 1 1 1</p>

Set – (56/3/3)
MARKING SCHEME
SR. SECONDARY SCHOOL EXAMINATION, 2020
Subject: CHEMISTRY

Q.No	Expected Answer / Value Points	Distribution of Marks						
SECTION - A								
1.	Zn , Cd and Hg have completely filled d ¹⁰ configuration in their ground state as well as in their oxidized state.	1						
2.	Scandium / Sc	1						
3.	Because of multiple oxidation states / ability to form complexes / having large surface area.	1						
4.	Involvement of greater no. of electrons in inter atomic metallic bonding / strong metal-metal bonding.	1						
5.	Presence of unpaired e ⁻ showing d-d transition in Cu ²⁺ , while in Zn ²⁺ there is no unpaired electron / has d ¹⁰ configuration.	1						
6.	Iodobenzene	1						
7.	N-Phenylethanamide	1						
8.	Zn	1						
9.	B ₆ / Pyridoxine	1						
10.	Bithional	1						
11.	(C)	1						
12.	(C)	1						
13.	(B)	1						
14.	(D)	1						
15.	(A)	1						
16.	(ii)	1						
17.	(i)	1						
18.	(iii)	1						
19.	(iv)	1						
20.	(i)	1						
SECTION – B								
21.	i) Acts as solvent / lowers the melting point of the mixture / Increases the conductivity. ii) It forms a volatile compound Ni(CO) ₄ , which decomposes at higher temperature to give pure Nickel. <p style="text-align: center;">OR</p> $\text{Al}_2\text{O}_3 + 2 \text{NaOH} + 3\text{H}_2\text{O} \rightarrow 2 \text{Na} [\text{Al}(\text{OH})_4]$ $2\text{Na} [\text{Al}(\text{OH})_4] + \text{CO}_2 \rightarrow \text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O} + 2 \text{NaHCO}_3$ $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O} \xrightarrow{\Delta} \text{Al}_2\text{O}_3 + x\text{H}_2\text{O}$	1 1 1 ½ ½						
22.	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="background-color: #f8d7da; text-align: center;">Physisorption</th> <th style="background-color: #f8d7da; text-align: center;">Chemisorption</th> </tr> </thead> <tbody> <tr> <td style="padding: 5px;">1. It arises because of van der Waals' forces.</td> <td style="padding: 5px;">1. It is caused by chemical bond formation.</td> </tr> <tr> <td style="padding: 5px;">2. It is not specific in nature.</td> <td style="padding: 5px;">2. It is highly specific in nature.</td> </tr> </tbody> </table> <p style="text-align: center;">(or any other two correct differences)</p> <p style="text-align: center;">OR</p>	Physisorption	Chemisorption	1. It arises because of van der Waals' forces.	1. It is caused by chemical bond formation.	2. It is not specific in nature.	2. It is highly specific in nature.	1 1
Physisorption	Chemisorption							
1. It arises because of van der Waals' forces.	1. It is caused by chemical bond formation.							
2. It is not specific in nature.	2. It is highly specific in nature.							

22.	<p>i) There are some substances which at low concentration behave as normal strong electrolytes, but at higher concentration exhibit colloidal behavior due to the formation of aggregates. Example: Micelles / soap</p> <p>ii) Oil is dispersed phase and water is dispersion medium. Example : Milk (or any other correct example)</p>	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
23.	<p>Rate = $-\frac{\Delta[A]}{\Delta t} = -\frac{1}{3} \frac{\Delta[B]}{\Delta t} = \frac{1}{2} \frac{\Delta[C]}{\Delta t}$</p> <p>(i) Rate = $\frac{1}{2} \frac{\Delta[C]}{\Delta t}$ $2 \times \text{Rate} = \frac{\Delta[C]}{\Delta t} = 2.5 \times 10^{-4}$ Rate = $\frac{2.5 \times 10^{-4}}{2} = 1.25 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$</p> <p>(ii) Rate = $-\frac{1}{3} \frac{\Delta[B]}{\Delta t}$ $1.25 \times 10^{-4} = -\frac{1}{3} \frac{\Delta[B]}{\Delta t}$ $-\frac{\Delta[B]}{\Delta t} = 3 \times 1.25 \times 10^{-4} = 3.75 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
24.	<p>i) Chemical compounds which reduce or abolish pain Example: aspirin / paracetamol. (or any other suitable example)</p> <p>ii) Quaternary ammonium salts with acetate / chloride / bromide Example: cetyltrimethylammonium bromide. (or any other suitable example)</p>	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
25.	<p>(a) On adding AgNO_3, $[\text{Co}(\text{NH}_3)_5(\text{SO}_4)]\text{Cl}$ will give white ppt of AgCl but $[\text{Co}(\text{NH}_3)_5\text{Cl}] \text{SO}_4$ doesn't / On adding BaCl_2, $[\text{Co}(\text{NH}_3)_5\text{Cl}] \text{SO}_4$ will give white ppt of BaSO_4 but $[\text{Co}(\text{NH}_3)_5(\text{SO}_4)]\text{Cl}$ doesn't.</p> <p>(b) $\text{K}_3[\text{Cr}(\text{C}_2\text{O}_4)_3] / \text{K}_3[\text{Cr}(\text{ox})_3]$</p>	<p>1</p> <p>1</p>
26.	<p>a) The pentaacetate of glucose does not react with hydroxylamine / HCN / Schiff's reagent indicating the absence of free $-\text{CHO}$ group.</p> <p>b) Adenine, Guanine, Uracil and Cytosine Uracil</p>	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
27.	<p>$6\text{XeF}_4 + 12 \text{H}_2\text{O} \rightarrow 4\text{Xe} + 2\text{XeO}_3 + 24\text{HF} + 3\text{O}_2$</p> <p>Yes Xe^{+4} changes to Xe^0 and Xe^{+6} / Xe gets oxidized as well as reduced in the same reaction.</p>	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
28.	<p>a.</p> <p>$\text{CH}_3\text{-CH}_2\text{-}\ddot{\text{O}}\text{-H} + \text{H}^+ \rightarrow \text{CH}_3\text{-CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-H}$</p> <p>$\text{CH}_3\text{CH}_2\text{-}\ddot{\text{O}}\text{:} + \text{CH}_3\text{-CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-H} \rightarrow \text{CH}_3\text{CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-CH}_2\text{CH}_3 + \text{H}_2\text{O}$</p> <p>$\text{CH}_3\text{CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-CH}_2\text{CH}_3 \rightarrow \text{CH}_3\text{CH}_2\text{-O-CH}_2\text{CH}_3 + \text{H}^+$</p>	<p>$\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$</p>

28.	<p>b.</p>  <p>Cumene $\xrightarrow{O_2}$ $\xrightarrow[H_2O]{H^+}$ Phenol</p> <p>OR</p> <p>i)</p>  <p>ii) $CH_3COCH_3 \xrightarrow{LiAlH_4} CH_3CHOHCH_3 \xrightarrow[443\text{ K}]{H_2SO_4(\text{conc})} CH_3CH=CH_2$</p> <p>iii)</p> 	1 1 1
29.	<p>i)</p>  <p>ii)</p>  <p>iii) $H_3C-C(CH_3)=CH_2$</p>	1 1 1
30.	$\Delta G^\circ = -nF E^\circ_{\text{cell}}$ $= -2 \times 96500 \times \{0.80 - (-0.25)\}$ $= -202650 \text{ J/mol}$ <p>Maximum work = 202650 J/mol</p> $E^\circ_{\text{cell}} = \frac{0.059}{2} \log Kc$ $\log Kc = \frac{2 \times 1.15}{0.059}$ $= 35.6$	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1
31.	$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$ $= \frac{2.303}{80} \log \frac{100}{60}$ $= \frac{2.303}{80} \times (1 - 0.7782)$ $= 0.0064 \text{ min}^{-1}$ $t = \frac{2.303}{k} \log \frac{[A]_0}{[A]}$ $= \frac{2.303}{0.0064} \log \frac{100}{10}$ $= 360 \text{ min}$	$\frac{1}{2}$ $\frac{1}{2}$ 1 1

