

JEE 2009
Paper I
Part 1 Chemistry
Section -1
Single Correct Choice Type

This section contains 8 multiple choice questions. Each question has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONLY ONE** is correct.

1. Given that the abundance of isotopes ^{54}Fe , ^{56}Fe and ^{57}Fe are 5%, 90% and 5% respectively, the atomic mass of Fe is

- (A) 55.85
- (B) 55.95
- (C) 55.75
- (D) 56.05

Answer:

- ☐ (A) ☒ (B) ☐ (C) ☐ (D)

2. The term that corrects for the attractive forces present in a real gas in the van der Waals equation is

- (A) nb
- (B) an^2/V^2
- (C) $-an^2/V^2$
- (D) $-nb$


Answer:

- ☐ (A) ☒ (B) ☐ (C) ☐ (D)

3. Among the electrolytes Na_2SO_4 , CaCl_2 , $\text{Al}_2(\text{SO}_4)_3$ and NH_4Cl , the most effective coagulating agent for Sb_2S_3 sol is

- (A) Na_2SO_4
- (B) CaCl_2
- (C) $\text{Al}_2(\text{SO}_4)_3$
- (D) NH_4Cl


Answer:



4. The Henry's law constant for the solubility of N_2 gas in water at 298 K is 1.0×10^5 atm. The mole fraction of N_2 in air is 0.8. the number of moles of N_2 from air dissolved in 10 moles of water at 298 K and 5 atm pressure is

- (A) 4.0×10^{-4}
(B) 4.0×10^{-5}
(C) 5.0×10^{-4}
(D) 4.0×10^{-6}


Answer:




5. The reaction of P_4 with X leads selectively to P_4O_6 . The X is

- (A) Dry O_2
(B) A mixture of O_2 and N_2
(C) Moist O_2
(D) O_2 in the presence of aqueous NaOH

Answer:




6. The correct acidity of the following is



- (A) (III) > (IV) > (II) > (I)

- (B) (IV) > (III) > (I) > (II)
 (C) (III) > (II) > (I) > (IV)
 (D) (II) > (III) > (IV) > (I)

Answer:




7. Among cellulose, poly (vinyl chloride), nylon and natural rubber, the polymer in which the intermolecular force of attraction is weakest is

- (A) Nylon
 (B) Poly vinyl chloride
 (C) Cellulose
 (D) Natural Rubber

Answer:




8. The IUPAC name of the following compound is



- (A) 4- Bromo -3-cyanophenol
 (B) 2- Bromo-5-hydroxybenzonitrile
 (C) 2-Cyano-4-hydroxybromobenzene
 (D) 6- Bromo-3-hydroxybenzonitrile

Answer:



Section - II
Multiple Correct Choice Type

This section contains 4 multiple choice questions. Each questions has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONE OR MORE** is / are correct.

9. The correct statement(s) regarding defects in solids is (are)

- (A) Frenkel defect is usually favoured by a very small difference in the sizes of cation and anion.
- (B) Frenkel defect is a dislocation defect
- (C) Trapping of an electron in the lattice leads to the formation of F- Center
- (D) Schottky defects have no effect on the physical properties of solids.


Answer:



10. The compound(s) that exhibit(s) geometrical isomerism is (are)

- (A) $[\text{Pt}(\text{en})\text{Cl}_2]$
- (B) $[\text{Pt}(\text{en})_2]\text{Cl}_2$
- (C) $[\text{Pt}(\text{en})_2\text{Cl}_2]\text{Cl}_2$
- (D) $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$

Answer:



11. The compound(s) formed upon combustion of sodium metal in excess air is(are)

- (A) Na_2O_2
- (B) Na_2O
- (C) NaO_2
- (D) NaOH


Answer:



12. The correct statement(s) about the compound $\text{H}_3\text{C}(\text{HO})\text{HC}-\text{CH}=\text{CH}-\text{CH}(\text{OH})\text{CH}_3$ (X) is(are)

- (A) The total number of stereoisomers possible for X is 6
- (B) The total number of diastereomers possible for X is 3
- (C) If the stereochemistry about the double bond in X is trans, the number of enantiomers possible for X is 4
- (D) If the stereochemistry about the double bond in X is cis, the number of enantiomers possible for X is 2.

Answer:



SECTION-III

Comprehension Type

This section contains 2 groups of questions. Each group has 3 multiple choice questions based on a paragraph. Each question has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONLY ONE** is correct.

Paragraph for Question Nos.13 to 15

p-Amino-N, N-dimethylaniline is added to a strongly acidic solution of X. The resulting solution is treated with a few drops of aqueous solution of Y to yield blue coloration due to the formation of methylene blue. Treatment of the aqueous solution of Y with the reagent potassium hexacyanoferrate(II) leads to the formation of an intense blue precipitate. The precipitate dissolves on excess addition of the reagent. Similarly, treatment of the solution of Y with the solution of potassium hexacyanoferrate(III) leads to a brown colouration due to the formation of Z.

13. The compound X is

- (A) NaNO_3
- (B) NaCl
- (C) Na_2SO_4
- (D) Na_2S

Answer:

- ☐ (A) ☐ (B) ☐ (C) ☒ (D)

14. The compound Y is

- (A) MgCl_2
(B) FeCl_2
(C) FeCl_3
(D) ZnCl_2

Answer:

- ☐ (A) ☐ (B) ☒ (C) ☐ (D)

15. The compound Z is


- (A) $\text{Mg}_2[\text{Fe}(\text{CN})_6]$
(B) $\text{Fe}[\text{Fe}(\text{CN})_6]$
(C) $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$
(D) $\text{K}_2\text{Zn}_3[\text{Fe}(\text{CN})_6]_2$

Answer

- ☐ (A) ☒ (B) ☐ (C) ☐ (D)

Paragraph for Question Nos. 16 to 18

A carbonyl compound **P**, which gives positive iodoform test, undergoes reaction with MeMgBr followed by dehydration to give an olefin **Q**. Ozonolysis of **Q** leads to a dicarbonyl compound **R**, which undergoes intramolecular aldol reaction to give predominantly **S**.




16. The structure of the carbonyl compound **P** is

- (A)
- (B)
- (C)
- (D)


Answer:

- ☐ (A)
 ☒ (B)
 ☐ (C)
 ☐ (D)


17. The structure of the products **Q** and **R**, respectively are





Answer:



18. The structure of the product **S** is





Answer:



(A) (B) (C) (D)

Section -IV
Matrix- Match Type

This section contains 2 questions. Each question contains statements given in two columns, which have to be matched. The statements in **Column I** are labelled A, B, C and D, while the statements in **Column II** are labelled p, q, r, s and t. Any given statement in **Column I** can have correct matching with **ONE OR MORE** statement(s) in **Column II**. The appropriate bubbles corresponding to the Answer:s to these questions have to be darkened as illustrated in the following example.

If the correct matches are A-p, s and t; B-q and r; C-p and q; and D-s and t;

Then the correct darkening of bubbles will look like the following.

	p	q	r	s	t
A	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>
B	<input type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>
C	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
D	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>

19. Match each of the diatomic molecules in **Column I** with its property/properties in **Column II**.

Column I	Column II
(A) B_2	(p) Paramagnetic
(B) N_2	(q) Undergoes oxidation
(C) O_2^-	(r) Undergoes reduction
(D) O_2	(s) Bond order ≥ 2
	(t) Mixing of 's' and 'p' orbitals

Answer:

	p	q	r	s	t
A	<input checked="" type="radio"/> p	<input checked="" type="radio"/> q	<input checked="" type="radio"/> r	<input type="radio"/> s	<input checked="" type="radio"/> t
B	<input type="radio"/> p	<input checked="" type="radio"/> q	<input checked="" type="radio"/> r	<input checked="" type="radio"/> s	<input checked="" type="radio"/> t
C	<input checked="" type="radio"/> p	<input checked="" type="radio"/> q	<input checked="" type="radio"/> r	<input type="radio"/> s	<input type="radio"/> t
D	<input checked="" type="radio"/> p	<input checked="" type="radio"/> q	<input checked="" type="radio"/> r	<input checked="" type="radio"/> s	<input type="radio"/> t

20. Match each of the compounds in **Column I** with its characteristic reaction(s) in **Column II**.

Column I	Column II
(A) $CH_3CH_2CH_2CN$	(p) Reduction with $Pd-C/H_2$
(B) $CH_3CH_2OCOCH_3$	(q) Reduction with $SnCl_2/HCl$
(C) $CH_3-CH=CH-CH_2OH$	(r) Development of foul smell on treatment with chloroform and alcoholic KOH
(D) $CH_3CH_2CH_2CH_2NH_2$	(s) Reduction with diisobutylaluminium hydride (DIBAL-H)
	(t) Alkaline hydrolysis

Answer:

	p	q	r	s	t
A	<input checked="" type="radio"/> p	<input checked="" type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s	<input checked="" type="radio"/> t
B	<input type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s	<input checked="" type="radio"/> t
C	<input checked="" type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input type="radio"/> s	<input type="radio"/> t
D	<input type="radio"/> p	<input type="radio"/> q	<input checked="" type="radio"/> r	<input type="radio"/> s	<input type="radio"/> t

PART II - MATHEMATICS

Section -I

Single Correct Choice Type

This section contains 8 multiple choice questions. Each question has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONLY ONE** is correct.

21. Let P(3,2,6) be a point in space and Q be a point on the line

$$\vec{r} = (\hat{i} - \hat{j} + 2\hat{k}) + \mu(-3\hat{i} + \hat{j} + 5\hat{k})$$


Then the value of μ for which the vector PQ is parallel to the plane $x-4y+3z = 1$ is

- (A) $\frac{1}{4}$
- (B) $-\frac{1}{4}$
- (C) $\frac{1}{8}$
- (D) $-\frac{1}{8}$

Answer:




(A)



(B)



(C)




(D)

22. Tangents drawn from the point $P(1,8)$ to the circle $x^2+y^2-6x-4y-11=0$ touch the circle at points A and B. The equation of the circumcircle of the triangle PAB is

- (A) $x^2+y^2+4x-6y+19=0$
 (B) $x^2+y^2-4x-10y+19=0$
 (C) $x^2+y^2-2x+6y-29=0$
 (D) $x^2+y^2-6x-4y+19=0$

Answer:




23. Let f be a non-negative function defined on the interval $[0,1]$. If $\int_0^x \sqrt{1-(f'(t))^2} dt = \int_0^x f(t) dt$,

$0 \leq x \leq 1$, and $f(0) = 0$, then

- (A) $f(1/2) < 1/2$ and $f(1/3) > 1/3$
 (B) $f(1/2) > 1/2$ and $f(1/3) > 1/3$
 (C) $f(1/2) < 1/2$ and $f(1/3) < 1/3$
 (D) $f(1/2) > 1/2$ and $f(1/3) < 1/3$

Answer:




24. Let $z = x+iy$ be a complex number where x and y are integers. Then the area of the rectangle whose vertices are the roots of the equation

$$z\bar{z}^3 + \bar{z}z^3 = 350 \text{ is}$$

- (A) 48
 (B) 32
 (C) 40
 (D) 80


Answer:



25. The line passing through the extremity A of the major axis and extremity B of the minor axis of the ellipse $x^2 + 9y^2 = 9$ meets its auxiliary circle at the point M. Then the area of the triangle with vertices at A, M and the origin O is

- (A) 31/10
- (B) 29/10
- (C) 21/10
- (D) 27/10


Answer:



26. If \vec{a} , \vec{b} , \vec{c} and \vec{d} are unit vectors such that $(\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d}) = 1$ and $\vec{a} \cdot \vec{c} = \frac{1}{2}$, then

- (A) \vec{a} , \vec{b} , \vec{c} are non-coplanar
- (B) \vec{b} , \vec{c} and \vec{d} are non - coplanar
- (C) \vec{b} , \vec{d} are non - parallel
- (D) \vec{a} , \vec{d} are parallel and \vec{b} , \vec{c} are parallel


Answer:



27. Let $z = \cos \theta + i \sin \theta$. Then the value of $\sum_{m=1}^{15} \text{Im}(z^{2m-1})$ at $\theta = 2^\circ$ is

- (A) $1/\sin 2^\circ$
- (B) $1/3 \sin 2^\circ$
- (C) $1/2 \sin 2^\circ$
- (D) $1/4 \sin 2^\circ$


Answer:



28. The number of seven digit integers with sum of the digits equal to 10 and formed by using 1,2 and 3 only, is

- (A) 55
(B) 66
(C) 77
(D) 88

Answer:




Section II
Multiple Correct Choice Type

This section contains 4 multiple choice questions. Each question has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONE OR MORE** is/are correct.

29. Area of the region bounded by the curve $y = e^x$ and lines $x = 0$ and $y = e$ is
(A) $e-1$
(B) $\int_1^e \ln(e+1-y) dy$
(C) $e - \int_0^1 e^x dx$
(D) $\int_1^e \ln y dy$

Answer:



30. Let


$$L = \lim_{x \rightarrow 0} \frac{a - \sqrt{a^2 - x^2} - \frac{x^2}{4}}{x^4}$$

, $a > 0$.

If L is finite, then

- (A) $a = 2$
 (B) $a = 1$
 (C) $L = 1/64$
 (D) $L = 1/32$

Answer:




31. In a triangle ABC with fixed base BC, the vertex A moves such that $\cos B + \cos C = 4 \sin^2 A/2$.

If a, b and c denote the lengths of the sides of the triangle opposite to the angles A, B and C, respectively, then

- (A) $b+c=4a$
 (B) $b+c=2a$
 (C) locus of point A is an ellipse
 (D) locus of point A is a pair of straight lines


Answer:



32. If $\sin^4 x/2 + \cos^4 x/3 = 1/5$, then

- (A) $\tan^2 x = 2/3$
 (B) $\frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} = \frac{1}{125}$
 (C) $\tan^2 x = 1/3$
 (D) $\frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} = \frac{2}{125}$

Answer:



Section III
Comprehension Type

This section contains 2 groups of questions. Each group has 3 multiple choice questions based on a paragraph. Each question has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONLY ONE** is correct.

Paragraph for Question Nos. 33 to 35

Let A be the set of all 3×3 symmetric matrices all of whose entries are either 0 or 1. Five of these entries are 1 and four of them are 0.

33. The number of matrices in A is

- (A) 12
- (B) 6
- (C) 9
- (D) 3

Answer:



(A)



(B)



(C)



(D)

34. The number of matrices A in A for which the system of linear equations

$$A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

has a unique solution, is

- (A) less than 4
- (B) at least 4 but less than 7
- (C) at least 7 but less than 10
- (D) at least 10


Answer:



(A)



(B)



(C)



(D)


35. The number of matrices A in A for which the system of linear equations

$$A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

is inconsistent, is

- (A) 0
- (B) more than 2
- (C) 2
- (D) 1

Answer:



(A)



(B)



(C)



(D)

Paragraph for Question No. 36 to 38

A fair die is tossed repeatedly until a six is obtained. Let X denote the number of tosses required.

36. The probability that $X = 3$ equals

- (A) $25/216$
- (B) $25/36$
- (C) $5/36$
- (D) $125/216$

Answer:



(A)



(B)



(C)




(D)

37. The probability that $X \geq 23$ equals

- (A) $125/216$
- (B) $25/36$
- (C) $5/36$

(D) 25/216


Answer:



38. The conditional probability that $X \geq 6$ given $X > 3$ equals

- (A) 125/216
(B) 25/216
(C) 5/36
(D) 25/36

Answer:



Section -IV
Matrix Match Type

This section contains 2 questions. Each question contains statements given in two columns, which have to be matched. The statements in **Column I** are labelled A, B, C and D, while the statements in **Column II** are labelled p, q, r, s and t. Any given statement in **Column I** can have correct matching with **ONE OR MORE** statement(s) in **Column II**. The appropriate bubbles corresponding to the Answer:s to these questions have to be darkened as illustrated in the following example:

If the correct matches are A p, s and t; B-q and r; C-p and q; and D-s and t; then the correct darkening of bubbles will look like the following.

	p	q	r	s	t
A	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
B	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
C	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
D	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>

39. Match the statements/expressions in Column I with the open intervals in Column II.

Column I	Column II
(A) Interval definition of non-zero solutions of the differential equation $(x-3)^2 y' + y = 0$	(p) $(-\pi/2, \pi/2)$
(B) Interval containing the value of the integral $\int (x-1)(x-2)(x+3)(x-4)(x-5) dx$	(q) $(0, \pi/2)$
(C) Interval in which at least one of the points of local maximum of $\cos^2 x + \sin x$ lies	(r) $(\pi/8, 5\pi/4)$
(D) Interval in which $\tan^{-1}(\sin x + \cos x)$ is increasing	(s) $(0, \pi/8)$
	(t) $(-\pi, \pi)$

Answer:

	p	q	r	s	t
A	<input checked="" type="radio"/> p	<input checked="" type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s	<input type="radio"/> t
B	<input checked="" type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input type="radio"/> s	<input checked="" type="radio"/> t
C	<input checked="" type="radio"/> p	<input checked="" type="radio"/> q	<input checked="" type="radio"/> r	<input type="radio"/> s	<input checked="" type="radio"/> t
D	<input type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s	<input type="radio"/> t

40. Match the conics in Column I with the statements/expressions in Column II.

Column I	Column II
(A) Circle	(p) The locus of the point (h, k) for which the line $hx + ky = 1$ touches the circle $x^2 + y^2 = 4$
(B) Parabola	(q) Points z in the complex plane satisfying $ z+2 - z-2 = \pm 3$
(C) Ellipse	(r) Points of the conic have parametric representation $x = \sqrt{3}(1-t^2)/(1+t^2)$, $y = 2t/(1+t^2)$
(D) Hyperbola	(s) The eccentricity of the conic lies in the interval $1 \leq e < \infty$
	(t) Points z in the complex plane satisfying $\operatorname{Re}(z+1)^2 = z^2 + 1$

Answer:

	p	q	r	s	t
A	<input checked="" type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input type="radio"/> s	<input type="radio"/> t
B	<input type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s	<input checked="" type="radio"/> t
C	<input type="radio"/> p	<input type="radio"/> q	<input checked="" type="radio"/> r	<input type="radio"/> s	<input type="radio"/> t
D	<input type="radio"/> p	<input checked="" type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s	<input type="radio"/> t


PART III : PHYSICS
SECTION - I
Single Correct Choice Type

This section contains 8 multiple choice questions. Each question has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONLY ONE** is correct.

41. Three concentric metallic spherical shells of radii R , $2R$, $3R$, are given charges Q_1 , Q_2 , Q_3 , respectively. It is found that the surface charge densities on the outer surfaces of the shells are equal. Then, the ratio of the charges given to the shells, $Q_1:Q_2:Q_3$ is

- (A) 1:2:3
(B) 1:3:5
(C) 1:4:9
(D) 1:8:18

Answer:




42. A block of base 10 cm x 10 cm and height 15 cm is kept on an inclined plane. The coefficient of

friction between them is $\sqrt{3}$. The inclination θ of this inclined plane from the horizontal plane is gradually increased from 0° . Then

- (A) at $\theta = 30^\circ$, the block will start sliding down the plane
- (B) the block will remain at rest on the plane up to certain θ and then it will topple
- (C) at $\theta = 60^\circ$, the block will start sliding down the plane and continue to do so at higher angles
- (D) at $\theta = 60^\circ$, the block will start sliding down the plane and on further increasing θ , it will topple at certain θ


Answer:




43. A ball is dropped from a height of 20 m above the surface of water in a lake. The refractive index of water is $4/3$. A fish inside the lake, in the line of fall of the ball, is looking at the ball. At an instant, when the ball is 12.8 m above the water surface, the fish sees the speed of ball as [Take $g = 10 \text{ m/s}^2$.]

- (A) 9 m/s
- (B) 12m/s
- (C) 16 m/s
- (D) 21.33.m/s

Answer:




44. Look at the drawing given in the figure which has been drawn with ink of uniform line-thickness. The mass of ink used to draw each of the two inner circles, and each of the two line segments is m . The mass of the ink used to draw the outer circle is $6m$. The coordinates of the center of the different parts are: outer circle $(0,0)$, left inner circle $(-a, a)$, right inner circle (a, a) , vertical line $(0,0)$ and horizontal line $(0,-a)$. The y -coordinate of the centre of mass of the ink in this drawing is




- (A) $a/10$
- (B) $a/8$
- (C) $a/12$
- (D) $a/3$

Answer:




45. Two small particles of equal masses start moving in opposite directions from a point A in a horizontal circular orbit. Their tangential velocities are v and $2v$, respectively, as shown in the figure. Between collisions, the particles move with constant speed. After making how many elastic collisions, other than that at A, these two particles will again reach the point A?




- (A) 4
- (B) 3
- (C) 2
- (D) 1

Answer:




46. The figure shows certain wire segments joined together to form a coplanar loop. The loop is placed in a perpendicular magnetic field in the direction going into the plane of the figure. The magnitude of the field increases with time. I_1 , and I_2 are the currents in the segments ab and cd. Then,




- (A) $I_1 > I_2$
 (B) $I_1 < I_2$
 (C) I_1 is in the direction ba and I_2 is in the direction cd
 (D) I_1 is in the direction ab and I_2 is in the direction dc

Answer:



47. A disk of radius $a/4$ having a uniformly distributed charge $6C$ is placed in the x - y plane with its centre at $(-a/2, 0, 0)$. A rod of length a carrying a uniformly distributed charge $8C$ is placed on the x -axis from $x = a/4$ to $x = 5a/4$. Two point charges $-7C$ and $3C$ are placed at $(a/4, -a/4, 0)$ and $(-3a/4, 3a/4, 0)$, respectively. Consider a cubical surface formed by six surfaces $x = \pm a/2$, $y = \pm a/2$, $z = \pm a/2$. The electric flux through this cubical surface is




- (A) $-2C/\epsilon_0$
 (B) $2C/\epsilon_0$

(C) $10C/\epsilon_0$

(D) $12C/\epsilon_0$

Answer:




(A)

(B)

(C)

(D)

48. The x - t graph of a particle undergoing simple harmonic motion is shown below. The acceleration of the particle at $t = 4/3$ s is




(A) $(\sqrt{3}/32)\pi^2 \text{ cm/s}^2$

(B) $-\pi^2/32 \text{ cm/s}^2$

(C) $\pi^2/32 \text{ cm/s}^2$

(D) $-(\sqrt{3}/32)\pi^2 \text{ cm/s}^2$

Answer:



(A)

(B)

(C)

(D)


SECTION - II
Multiple Correct Choice Type

This section contains 4 multiple choice questions. Each question has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONE OR MORE** is /are correct.

49. If the resultant of all the external forces acting on a system of particles is zero, then from an inertial frame, one can surely say that

- (A) linear momentum of the system does not change in time
- (B) kinetic energy of the system does not change in time
- (C) angular momentum of the system does not change in time
- (D) potential energy of the system does not change in time


Answer:




50. A student performed the experiment of determination of focal length of a concave mirror by u-v method using an optical bench of length 1.5 meter. The focal length of the mirror used is 24 cm. The maximum error in the location of the image can be 0.2 cm. The 5 sets of (u, v) values recorded by the student in cm are (42, 56), (48, 48), (60, 40), (66, 33), (78, 39). The data set(s) that **cannot** come from experiment and is (are) incorrectly recorded, is (are)

- (A) (42, 56)
- (B) (48, 48)
- (C) (66, 33)
- (D) (78, 39)

Answer:




51. For the circuit shown in the figure



- (A) the current I through the battery is 7.5 mA
- (B) the potential difference across R_L is 18 V
- (C) ratio of powers dissipated in R_1 and R_2 is 3
- (D) if R_1 and R_2 are interchanged, magnitude of the power dissipated in R_L will decrease by a factor of 9


Answer:



52. C_v and C_p Denote the molar specific heat capacities of a gas at constant volume and constant pressure, respectively. Then

- (A) $C_p - C_v$ is larger for a diatomic ideal gas than for a monoatomic ideal gas
- (B) $C_p - C_v$ is larger for a diatomic ideal gas than for a monoatomic ideal gas
- (C) C_p / C_v is larger for a diatomic ideal gas than for a monoatomic ideal gas
- (D) $C_p \cdot C_v$ is larger for a diatomic ideal gas than for a monoatomic ideal gas

Answer:




SECTION -III
Comprehension Type

This section contains 2 groups of questions. Each group has 3 multiple choice questions based on a paragraph. Each question has 4 choices (A), (B), (C) and (D) for its Answer:, out of which **ONLY ONE** is correct.

Paragraph for Question No. 53 to 55

Scientists are working hard to develop nuclear fusion reactor. Nuclei of heavy hydrogen, ${}_1^2\text{H}$, known as deuteron and denoted by D, can be thought of as a candidate for fusion reactor. The D-D reaction is



In the core of fusion reactor, a gas of heavy hydrogen is fully ionized into deuteron nuclei and electrons. This collection of H nuclei and electrons is known as plasma. The nuclei move randomly in the reactor core and occasionally come close enough for nuclear fusion to take place. Usually, the temperatures in the reactor core are too high and no material wall can be used to confine the plasma. Special techniques are used which confine the plasma for a time to before the particles fly away from the core. If n is the density (number/volume) of deuterons, the product nt_0 , is called Lawson number. In one of the criteria, a reactor is termed successful if Lawson number is greater than $5 \times 10^{14} \text{ s/cm}^3$.

It may be helpful to use the following: Boltzmann constant $k = 8.6 \times 10^{-5} \text{ eV/K}$; $e^2/4\pi\epsilon_0 = 1.44 \times 10^{-9} \text{ eVm}$.

53. In the core of nuclear fusion reactor, the gas becomes plasma because of

- (A) strong nuclear force acting between the deuterons
- (B) Coulomb force acting between the deuterons
- (C) Coulomb force acting between deuteron-electron pairs
- (D) the high temperature maintained inside the reactor core

Answer:

- ☐ (A) ☐ (B) ☐ (C) ☒ (D)

54. Assume that two deuteron nuclei in the core of fusion reactor at temperature T are moving towards each other, each with kinetic energy 1.5 kT , when the separation between them is large enough to neglect Coulomb potential energy. Also neglect any interaction from other particles in the core. The minimum temperature T required for them to reach a separation of $4 \times 10^{-15} \text{ m}$ is in the range

- (A) $1.0 \times 10^9 \text{ K} < T < 2.0 \times 10^9 \text{ K}$
- (B) $2.0 \times 10^9 \text{ K} < T < 3.0 \times 10^9 \text{ K}$
- (C) $3.0 \times 10^9 \text{ K} < T < 4.0 \times 10^9 \text{ K}$
- (D) $4.0 \times 10^9 \text{ K} < T < 5.0 \times 10^9 \text{ K}$

Answer:

- ☒ (A)
 ☐ (B)
 ☐ (C)
 ☐ (D)

55. Results of calculations for four different designs of a fusion reactor using D-D reaction are given below. Which of these is most promising based on Lawson criterion?

- (A) deuteron density = 2.0×10^2 cm, confinement time = 5.0×10^{-3} s
 (B) deuteron density = 8.0×10^{14} cm, confinement time = 9.0×10^{-1}
 (C) deuteron density = 4.0×10^{23} cm, confinement time = 1.0×10^{-11} s
 (D) deuteron density = 1.0×10^{24} cm, confinement time = 4.0×10^{-12} s

Answer:

- ☐ (A)
 ☒ (B)
 ☐ (C)
 ☐ (D)

Paragraph for Question No. 56 to 58


When a particle is restricted to move along x-axis between $x = 0$ and $x = a$, where a is of nanometer dimension, its energy can take only certain specific values. The allowed energies of the particle moving in such a restricted region, correspond to the formation of standing waves with nodes at its ends $x = 0$ and $x = a$. The wavelength of this standing wave is related to the linear momentum p of the particle according to the de Broglie relation. The energy of the particle of mass m is related to its linear momentum as $E = p^2/2m$. Thus, the energy of the particle can be denoted by a quantum number 'n' taking values 1, 2, 3, . ($n = 1$, called the ground state) corresponding to the number of loops in the standing wave.

Use the model described above to Answer: the following three questions for a particle moving in the line $x = 0$ to $x = a$. Take $h = 6.6 \times 10^{-34}$ Js and $e = 1.6 \times 10^{-19}$ C.

56. The allowed energy for the particle for a particular value of n is proportional to

- (A) a^{-2}
 (B) $a^{-3/2}$
 (C) a^{-1}
 (D) a^2


Answer:



57. If the mass of the particle is $m = 1.0 \times 10^{-30}$ kg and $a = 6.6$ nm, the energy of the particle in its ground state is closest to

- (A) 0.8 meV
- (B) 8 meV
- (C) 80 meV
- (D) 800 meV


Answer:







58. The speed of the particle, that can take discrete values, is proportional to


- (A) $n^{-3/2}$
- (B) n^{-1}
- (C) $n^{-1/2}$
- (D) n

Answer:























59. Six point charges, each of the same magnitude q , are arranged in different manners as shown in Column II. In each case, a point M and a line PQ passing through M are shown. Let E be the electric field and V be the electric potential at M (potential at infinity is zero) due to the given charge distribution when it is at rest. Now, the whole system is set into rotation with a constant angular velocity about the line PQ. Let B be the magnetic field at M and u be the magnetic moment of the system in this condition. Assume each rotating charge to be equivalent to a steady current.




Column I	Column II
(A) $E = 0$	<p>(p)</p>  <p>Charges are at the corners of a regular hexagon. M is at the centre of the hexagon. PQ is perpendicular to the plane of the hexagon.</p>
(B) $V \neq 0$	<p>(q)</p>  <p>Charges are on a line perpendicular to PQ at equal intervals. M is the midpoint between the two innermost charges.</p>
(C) $B = 0$	<p>(r)</p>  <p>Charges are placed on two coplanar insulating rings at equal intervals. M is the common centre of the rings. PQ is perpendicular to the plane of the rings.</p>
(D) $\mu \neq 0$	<p>(s)</p>  <p>Charges are placed at the corners of a rectangle of sides a and 2a and at the mid points of the longer</p>



	sides. M is at the centre of the rectangle. PQ is parallel to the longer sides.
	<p>(t)</p>  <p>Charges are placed on two coplanar, identical insulating rings at equal intervals. M is the mid-point between the centres of the rings. PQ is perpendicular to the line joining the centre and coplanar to the rings.</p>

Answer:

	p	q	r	s	t
A					
B					
C					
D					

60. Column II shows five systems in which two objects are labelled as X and Y. Also in each case a point P is shown. Column I gives some statements about X and/or Y. Match these statements to the appropriate system(s) from Column II.

Column I	Column II
(A) The force exerted by X on Y has a magnitude Mg .	<p>(p)</p>  <p>Block Y of mass M is on a fixed inclined plane X, slides on it with a constant velocity.</p>
(B) The gravitational potential energy of X is continuously increasing	<p>(q)</p>  <p>Two ring magnets Y and Z, each of mass M, are kept in frictionless vertical plastic stand so that they repel each other. Y rests on the base X and Z hangs in air in equilibrium. P is the topmost point of the stand on the common axis of the two rings. The whole system is in a lift that is going up with a constant velocity.</p>
(C) Mechanical energy of the system X+Y is continuously decreasing	<p>(r)</p>  <p>A pulley Y of mass m_0 is fixed to a table through a clamp X. A block of mass M hangs from a string that goes over the pulley and is fixed at point P of the table. The whole system is kept in a lift that is going down with a constant velocity.</p>
(D) The torque of the weight of Y about point P	<p>(s)</p>

Is zero.	 <p>A sphere Y of mass M is put in non viscous liquid X kept in a container at rest. The sphere is released and it moves down in the liquid.</p>
	<p>(t)</p>  <p>A sphere Y of mass M is falling with its terminal velocity in a viscous liquid X kept in a container.</p>

Answer:

	p	q	r	s	t
A	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>
B	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>
C	<input checked="" type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>
D	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>