

DATE:

# JEE ADVANCED 2019 MATHS - PAPER 2



CLASS:

ALL CENTRE

## SECTION - 1

1. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be given by  $f(x) = (x - 1)(x - 2)(x - 5)$ . Define  $F(x) = \int_0^x f(t) dt, x > 0$ . Then which of the

following options is/are correct?

(a)  $F$  has a local minimum at  $x = 1$

(b)  $F$  has a local maximum at  $x = 2$

(c)  $F(x) \neq 0$  for all  $x \in (0, 5)$


(d)  $F$  has two local maxima and one local minimum in  $(0, \infty)$

### Solution:

$$f(x) = (x - 1)(x - 2)(x - 5)$$

$$\text{Given } F(x) = \int_0^x f(t) dt$$

$$F'(x) = (x - 1)(x - 2)(x - 5)$$



At  $x = 1$  and  $x = 5$ ,  $F'(x)$  changes from  $-$  to  $+$

$\therefore F(x)$  has two local minima points at  $x = 1$  and  $x = 5$

$F(x)$  has one local maxima point at  $x = 2$ .

2. For a  $\in \mathbb{R}$ ,  $|a| > 1$ , let  $\lim_{n \rightarrow \infty} \left( \frac{1 + \sqrt[3]{2} + \dots + \sqrt[3]{n}}{n^{7/3} \left( \frac{1}{(an+1)^2} + \frac{1}{(an+2)^2} + \dots + \frac{1}{(an+n)^2} \right)} \right) = 54$ . Then the possible value(s) of a

is/are:

(a) 8

(b) -9

(c) -6

(d) 7

**Solution:**

$$\lim_{n \rightarrow \infty} \frac{\sqrt[3]{1} + \sqrt[3]{2} + \dots + \sqrt[3]{n}}{n^{7/3} \left[ \frac{1}{(an+1)^2} + \frac{1}{(an+2)^2} + \dots + \frac{1}{(an+n)^2} \right]} = 54$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{\frac{1}{n} \sum_{r=1}^n \left( \frac{r}{n} \right)^{1/3}}{\frac{1}{n} \left[ \frac{n^2}{(an+1)^2} + \frac{n^2}{(an+2)^2} + \dots + \frac{n^2}{(an+n)^2} \right]} = 54$$

$$\Rightarrow \frac{\int_0^1 x^{1/3} dx}{\int_0^1 \frac{dx}{(a+x)^2}} = 54 \quad \Rightarrow \frac{\left[ \frac{3}{4} x^{4/3} \right]_0^1}{\left[ \frac{-1}{a+x} \right]_0^1} = \frac{3/4}{\frac{1}{a} - \frac{1}{a+1}} = 54$$

$$\Rightarrow \frac{(a+1) - a}{a(a+1)} = \frac{3}{4} \times \frac{1}{54} \quad \Rightarrow \frac{1}{a(a+1)} = \frac{1}{72} \quad \Rightarrow a(a+1) = 72$$

$$\Rightarrow a = 8 \text{ or } a = -9$$

3. Three lines

$$L_1 : \vec{r} = \lambda \hat{i}, \lambda \in \mathbb{R},$$

$$L_2 : \vec{r} = \vec{k} + \mu \hat{j}, \mu \in \mathbb{R} \text{ and}$$

$$L_3 : \vec{r} = \hat{i} + \hat{j} + \nu \hat{k}, \nu \in \mathbb{R}$$

are given. For which point(s) Q and  $L_2$  can we find a point P on  $L_1$  and a point R on  $L_3$  so that P, Q and R are collinear?

(a)  $\hat{k} + \hat{j}$

(b)  $\hat{k}$

(c)  $\hat{k} + \frac{1}{2} \hat{j}$

(d)  $\hat{k} - \frac{1}{2} \hat{j}$

**Solution:**

$$P(\lambda, 0, 0), Q(0, \mu, 1), R(1, 1, \nu)$$

$$\text{Given } \overrightarrow{PQ} = k \cdot \overrightarrow{PR} \Rightarrow \frac{\lambda}{\lambda-1} = \frac{-\mu}{-1} = \frac{-1}{-\nu}$$

$\therefore \mu$  cannot take the values 0 and 1

4. Let  $F: \mathbb{R} \rightarrow \mathbb{R}$  be a function. We say that  $f$  has

PROPERTY 1 if  $\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{\sqrt{|h|}}$  exists and is finite and

PROPERTY 2 if  $\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h^2}$  exists and is finite

Then which of the following options is/are correct?

- (a)  $f(x) = x|x|$  has PROPERTY 2  
 (b)  $F(x) = x^{2/3}$  has PROPERTY 1  
 (c)  $f(x) = \sin x$  has PROPERTY 2  
 (d)  $f(x) = |x|$  has PROPERTY 1

**Solution:**

(a)  $f(x) = x|x|$

$$\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h^2} = \lim_{h \rightarrow 0} \frac{h|h| - 0}{h^2} \text{ which does not exist.}$$

(b)  $\lim_{h \rightarrow 0} \frac{h^{2/3} - 0}{\sqrt{|h|}} = 0$

(c)  $\lim_{h \rightarrow 0} \frac{\sin h - 0}{h^2}$  does not exist

(d)  $\lim_{h \rightarrow 0} \frac{|h| - 0}{\sqrt{|h|}} = 0$

5. For non-negative integers  $n$ , let

$$f(n) = \frac{\sum_{k=0}^n \sin\left(\frac{k+1}{n+2}\pi\right) \sin\left(\frac{k+2}{n+2}\pi\right)}{\sum_{k=0}^n \sin^2\left(\frac{k+1}{n+2}\pi\right)}$$

Assuming  $\cos^{-1} x$  takes value in  $[0, \pi]$ , which of the following options is/are correct?

- (a)  $\sin(7 \cos^{-1} f(5)) = 0$   
 (b)  $f(4) = \frac{\sqrt{3}}{2}$   
 (c)  $\lim_{n \rightarrow \infty} f(n) = \frac{1}{2}$   
 (d) If  $\alpha = \tan(\cos^{-1} f(6))$ , then  $\alpha^2 + 2\alpha - 1 = 0$

**Solution:**

$$\begin{aligned}
 f(n) &= \frac{\sum_{k=0}^n \sin\left(\frac{k+1}{n+2}\pi\right) \cdot \sin\left(\frac{k+2}{n+2}\pi\right)}{\sum_{k=0}^n 2\sin^2\left(\frac{k+1}{n+2}\pi\right)} \\
 &= \frac{\sum_{k=0}^n \cos\frac{\pi}{n+2} - \cos\left(\frac{2k+3}{n+2}\right)\pi}{\sum_{k=0}^n 2\sin^2\left(\frac{k+1}{n+2}\right)\pi} \\
 &= \frac{(n+1)\cos\frac{\pi}{n+2} - \frac{\cos\left(\frac{n+3}{n+2}\right)\pi \cdot \sin\left(\frac{n+1}{n+2}\right)\pi}{\sin\frac{\pi}{n+2}}}{(n+1) - \frac{\cos\pi \cdot \sin\left(\frac{n+1}{n+2}\right)\pi}{\sin\left(\frac{\pi}{n+2}\right)}} \\
 &= \frac{(n+1)\cos\left(\frac{\pi}{n+2}\right) + \cos\left(\frac{n+3}{n+2}\right)\pi}{(n+1)+1} \\
 &= \cos\left(\frac{\pi}{n+2}\right)
 \end{aligned}$$

$$(A) \alpha = \tan\left(\cos^{-1} f(6)\right) = \tan\cos^{-1}\left(\cos\frac{\pi}{8}\right) = \tan\frac{\pi}{8}$$

$$\alpha^2 + 2\alpha - 1 = \tan^2\frac{\pi}{8} + 2\tan\frac{\pi}{8} - 1$$

$$\tan 2\left(\frac{\pi}{8}\right) = \frac{2\tan\frac{\pi}{8}}{1 - \tan^2\frac{\pi}{8}}$$

$$\Rightarrow 1 = \frac{2\alpha}{1 - \alpha^2} \Rightarrow \alpha^2 + 2\alpha - 1 = 0$$

\(\therefore\) option (A) is correct.

$$(B) \lim_{n \rightarrow \infty} f(x) = \lim_{n \rightarrow \infty} \cos\left(\frac{\pi}{n+2}\right) = \lim_{\frac{1}{n} \rightarrow 0} \cos\left(\frac{\pi/n}{1 + 2/n}\right) = 1$$

Option (B) correct.

$$(C) f(4) = \cos\left(\frac{\pi}{4+2}\right) = \cos\frac{\pi}{6} = \frac{\sqrt{3}}{2}$$

Option (C) wrong

$$(D) \sin\left[7 \cos^{-1} f(5)\right] = \sin\left[7 \cos^{-1}\left(\cos\frac{\pi}{7}\right)\right] = \sin\left[7 \times \frac{\pi}{7}\right] = 0$$

$$6. \text{ Let } P_1 = I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, P_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, P_3 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, P_4 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, P_5 = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix},$$

$$P_6 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \text{ and } X = \sum_{k=1}^6 P_k \begin{bmatrix} 2 & 1 & 3 \\ 1 & 0 & 2 \\ 3 & 2 & 1 \end{bmatrix} P_k^T$$

Where  $P_k^T$  denotes the transpose of the matrix  $P_k$ . Then which of the following options is/are correct?

(a)  $X - 30I$  is an invertible matrix

(b) The sum of diagonal entries of  $X$  is 18

(c) If  $X \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \alpha \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ , then  $\alpha = 30$

(d)  $X$  is a symmetric matrix

### Solution:

From the given data it is clear that

$$P_1 = P_1^T = P_1^{-1}$$

$$P_2 = P_2^T = P_2^{-1}$$

$$P_6 = P_6^T = P_6^{-1}$$

$$\text{And Let } A = \begin{bmatrix} 2 & 1 & 3 \\ 1 & 0 & 2 \\ 3 & 2 & 1 \end{bmatrix}$$

Here  $A^T = A \rightarrow A$  is symmetric matrix

$$X^T = \left(P_1 A P_1^T + \dots + P_6 A P_6^T\right)^T$$

$$= P_1 A^T P_1^T + \dots + P_6 A^T P_6^T$$

$$= X$$

$\therefore X$  is symmetric

$$\text{Let } B = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$XB = P_1 A P_1^T G + P_2 A P_2^T B + \dots + P_6 A P_6^T B$$

$$= P_1 AB + P_2 AB + \dots + P_6 AB$$

$$= (P_1 + P_2 + P_3 + \dots + P_6) \begin{bmatrix} 6 \\ 3 \\ 6 \end{bmatrix}$$

$$= \begin{bmatrix} 30 \\ 30 \\ 30 \end{bmatrix} = 30B \quad \Rightarrow \alpha = 30$$

$$\text{Since } X \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = 30 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$\Rightarrow (X - 30I)B = 0 \text{ has a nontrivial solution } B = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$\Rightarrow (X - 30I) = 0$$

$$X = P_1 A P_1^T + \dots + P_6 A P_6^T$$

$$\begin{aligned}\text{Trace}(X) &= \text{tr}(P_1AP_1^T) + \dots + \text{Tr}(P_6AP_6^T) \\ &= (2 + 0 + 1) + \dots + (2 + 0 + 1) = 3 + 3 + \dots \text{ (6 times)} = 18\end{aligned}$$

7. Let  $x \in \mathbb{R}$  and let  $P = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix}$ ,  $Q = \begin{bmatrix} 2 & x & x \\ 0 & 4 & 0 \\ x & x & 6 \end{bmatrix}$  and  $R = PQP^{-1}$

Then which of the following options is/are correct?

(a) For  $x = 1$ , there exists a unit vector  $\alpha\hat{i} + \beta\hat{j} + \gamma\hat{k}$  for which  $R \begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

(b) There exists a real number  $x$  such that  $PQ = QP$

(c)  $\det R = \det \begin{bmatrix} 2 & x & x \\ 0 & 4 & 0 \\ x & x & 5 \end{bmatrix} + 8$ , for all  $x \in \mathbb{R}$

(d) for  $x = 0$ , if  $R \begin{bmatrix} 1 \\ a \\ b \end{bmatrix} = 6 \begin{bmatrix} 1 \\ a \\ b \end{bmatrix}$ , then  $a + b = 5$

**Solution:**

$$R = PQP^{-1}$$

$$|R| = |P||Q||P^{-1}|$$

$$\Rightarrow \det Q = 2(24) - x(0) + x(-4x) = 48 - 4x^2$$

$$P = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix}, Q(X=0) = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 6 \end{bmatrix}$$

$$R = PQR^{-1}$$

$$= \begin{bmatrix} 2 & 4 & 6 \\ 0 & 8 & 12 \\ 0 & 0 & 18 \end{bmatrix} \cdot \frac{1}{6} \begin{bmatrix} 6 & -3 & 0 \\ 0 & 3 & -2 \\ 0 & 0 & 2 \end{bmatrix}$$

$$= \frac{1}{6} \begin{bmatrix} 12 & 6 & 4 \\ 0 & 24 & 8 \\ 0 & 0 & 36 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 2/3 \\ 0 & 4 & 4/3 \\ 0 & 0 & 6 \end{bmatrix}$$

$$(R-6I) \begin{pmatrix} 1 \\ a \\ b \end{pmatrix} = \begin{pmatrix} -4 & 1 & 2/3 \\ 0 & -2 & 4/3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ a \\ b \end{pmatrix} = \begin{bmatrix} -4 & +a & +\frac{2b}{3} \\ 0 & -2a & +4b/3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$-4 + a + \frac{2b}{3} = 0 \text{ and } -2a + \frac{4b}{3} = 0 \Rightarrow a = 2 \text{ \& } b = 3$$

$$\therefore a + b = 5$$

$$PQ = QP \Rightarrow x + 4 + x = 2 + 2x + 0 \Rightarrow \text{No value exist}$$

8. Let  $f(x) = \frac{\sin \pi x}{x^2}, x > 0$

Let  $x_1 < x_2 < x_3 < \dots < x_n < \dots$  be all the points of local maximum of  $f$   
and  $y_1 < y_2 < y_3 < \dots < y_n < \dots$  be all the points of local minimum of  $f$ .

Then which of the following options is/are correct?

(a)  $|x_n - y_n| > 1$  for every  $n$

(b)  $x_1 < y_1$

(c)  $x_n \in \left(2n, 2n + \frac{1}{2}\right)$  for every  $n$

(d)  $x_{n+1} - x_n > 2$  for every  $n$

**Solution:**

$$f(x) = \frac{\sin \pi x}{x^2} \Rightarrow f'(x) = \frac{x^2 \cdot (\cos \pi x) \cdot (\pi) - \sin \pi x \cdot (2x)}{x^4}$$

$$\Rightarrow f'(x) = \frac{2x \cos \pi x \left(\frac{\pi x}{2} - \tan \pi x\right)}{x^4}$$

By using graph we can say that option (1) (3) (4) are correct.



**SECTION – 2**

1. The value of  $\sec^{-1}\left(\frac{1}{4}\sum_{k=0}^{10}\sec\left(\frac{7\pi}{12}+\frac{k\pi}{2}\right)\sec\left(\frac{7\pi}{12}+\frac{(k+1)\pi}{2}\right)\right)$  in the interval  $\left[-\frac{\pi}{4}, \frac{3\pi}{4}\right]$  equals

**Solution:**

$$\begin{aligned} & \sec^{-1}\left(\frac{1}{4}\sum_{k=0}^{10}\sec\left(\frac{7\pi}{12}+\frac{k\pi}{2}\right)\sec\left(\frac{7\pi}{12}+\frac{(k+1)\pi}{2}\right)\right) \\ &= \sec^{-1}\left(\frac{-1}{4}\sum_{k=0}^{10}\sec\left(\frac{7\pi}{12}+\frac{k\pi}{2}\right)\operatorname{cosec}\left(\frac{7\pi}{12}+\frac{k\pi}{2}\right)\right) \\ &= \sec^{-1}\left(\frac{-1}{4}\sum_{k=0}^{10}\frac{2}{\sin\left(\frac{7\pi}{6}+k\pi\right)}\right) \\ &= \sec^{-1}\left(\frac{-1}{2}\sum_{k=0}^{10}\frac{1}{(-1)^{k+1}\sin\frac{\pi}{6}}\right) \\ &= \sec^{-1}\left(-\sum_{k=0}^{10}\frac{1}{(-1)^{k+1}}\right) = \sec^{-1}(1) = 0 \end{aligned}$$

2. Let  $|X|$  denote the number of elements in set  $X$ . Let  $S = \{1,2,3,4,5,6\}$  be a sample space, where each element is equally likely to occur. If  $A$  and  $B$  are independent events associated with  $S$ , then the number of ordered pairs  $(A,B)$  such that  $1 \leq |B| < |A|$ , equals.

**Solution:**

The number of ordered pairs of  $(A, B)$  are

$$\begin{aligned} & 6c_1(6c_2 + 6c_3 + \dots + 6c_6) + 6c_2(6c_2(6c_3 + 6c_4 + \dots + 6c_6) + 6c_3(6c_4 + 6c_5 + 6c_6) + 6c_4(6c_5 + 6c_6) + 6c_5 \cdot 6c_6) \\ &= (6c_1 \cdot 6c_2 + 6c_1 \cdot 6c_3 + \dots + 6c_1 6c_6) + (6c_2 \cdot 6c_3 + 6c_2 \cdot 6c_4 + \dots + 6c_2 \cdot 6c_6) + (6c_3 \cdot 6c_4 + 6c_3 \cdot 6c_5 + 6c_3 \cdot 6c_6) \end{aligned}$$

DATE:

MATHS

CLASS:

JEE ADVANCED 2019

ALL CENTRE

PAPER 2

$$\begin{aligned}
 &+ 6C_4 \cdot 6C_5 + 6C_4 \cdot 6C_6 + 6C_5 \cdot 6C_6. \\
 &= (12C_5 - 6C_1) + (12C_4 - 6C_2) + (12C_3 - 6C_3) + (12C_2 - 6C_4) + (12C_1 - 6C_5) \\
 &= (12C_1 + 12C_2 + 12C_3 + 12C_4 + 12C_5) - (6C_1 + 6C_2 + \dots + 6C_5) \\
 &= 1585 - 62 = 1523.
 \end{aligned}$$

3. Five person A, B, C, D and E are seated in a circular arrangement. If each of them is given a hat of one of the three colours red, blue and green, then the number of ways of distributing the hats such that the persons seated in adjacent seats get different coloured hats is

**Solution:**

Maximum number of hats used of same colour are 2.

They cannot be 3 otherwise atleast 2 hats of same colour are consecutive.

Now the hats used are consider as B B G G B

Which can be selected in 3 ways.


It can be R G G B B or R R G B B

The number of ways of distributing blue hat (single one) in 5 persons equal to 5

Now either position B and D are filled by green hats and C and E are filled by Red hats or B & D are filled by Red hats and C & E are filled by Green hats.

→ 2 ways are possible.

Hence number of ways =  $3 \times 5 \times 2 = 30$  ways.



4. Suppose

$$\det \begin{bmatrix} \sum_{k=0}^n k & \sum_{k=0}^n {}^n C_k k^2 \\ \sum_{k=0}^n {}^n C_k k & \sum_{k=0}^n {}^n C_k 3^k \end{bmatrix} = 0, \text{ holds for some positive integer } n. \text{ Then } \sum_{k=0}^n \frac{{}^n C_k}{k+1} \text{ equals}$$

**Solution:**

$$\left| \begin{array}{cc} \sum_{k=0}^n k & \sum_{k=0}^n {}^n C_k \cdot k^2 \\ \sum_{k=0}^n {}^n C_k \cdot k & \sum_{k=0}^n {}^n C_k \cdot 3^k \end{array} \right| = 0$$

$$\left| \begin{array}{cc} \frac{n(n+1)}{2} & n \cdot 2^{n-1} + n(n-1) \cdot 2^{n-2} \\ n \cdot 2^{n-1} & 4^n \end{array} \right| = 0$$

$$\Rightarrow \frac{n(n+1)}{2} \cdot 4^n - n \cdot 2^{2n-1} (n \cdot 2^{n-1} + n(n-1) \cdot 2^{n-2}) = 0$$

$$\Rightarrow \frac{n(n+1)}{2} \cdot 4^n - n^2 \cdot 2^{2n-2} \cdot n(n-1) \cdot 2^{2n-3} = 0$$

$$\Rightarrow \frac{n(n+1)}{2} - \frac{n^2}{4} - \frac{n^2(n-1)}{8} = 0 \Rightarrow \frac{n}{2} \left[ n+1 - \frac{n}{2} - \frac{n(n-1)}{4} \right] = 0$$

$$\Rightarrow n = 0 \text{ or } 4(n+1) - 2n - 1(n-1) = 0 \Rightarrow n = 0 \text{ or } n = 4$$

$$\sum_{r=0}^4 \frac{4c\pi}{r+1} = \sum_{r=0}^4 \frac{5cr+1}{5} = \frac{2^5-1}{5} = \frac{31}{5} = 6.20$$

5. The value of the integral  $\int_0^{\pi/2} \frac{3\sqrt{\cos \theta}}{(\sqrt{\cos \theta} + \sqrt{\sin \theta})^5} d\theta$  equals

**Solution:**

$$I = \int_0^{\pi/2} \frac{3\sqrt{\cos \theta}}{(\sqrt{\sin \theta} + \sqrt{\cos \theta})^5} \cdot d\theta$$

$$I = 3 \int_0^{\pi/2} \frac{\sqrt{\cos \theta}}{(\sqrt{\sin \theta} + \sqrt{\cos \theta})^5} \rightarrow 1$$

$$I = 3 \int_0^{\pi/2} \frac{\sqrt{\sin \theta}}{(\sqrt{\cos \theta} + \sqrt{\sin \theta})^5} \rightarrow 2 \left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) \cdot dx \right]$$

$$2I = 3 \int_0^{\pi/2} \frac{\sqrt{\cos \theta} \sqrt{\sin \theta}}{(\sqrt{\cos \theta} + \sqrt{\sin \theta})^5} \cdot d\theta = 3 \int_0^{\pi/2} \frac{d\theta}{(\sqrt{\cos \theta} + \sqrt{\sin \theta})^4}$$

$$\frac{2I}{3} = \int_0^{\pi/2} \frac{\sec 2\theta \cdot d\theta}{(\sqrt{\tan \theta} + 1)^4}$$

Let  $\tan \theta = t^2 \Rightarrow \sec 2\theta \cdot d\theta = 2t dt$

$$\frac{2I}{3} = \int_0^{\infty} \frac{2t dt}{(t+1)^4}$$

$$\frac{I}{3} = \int_0^{\infty} \left[ \frac{1}{(t+1)^3} - \frac{1}{(t+1)^4} \right] dt$$

$$I = \left[ \frac{-3}{2(t+1)^2} + \frac{1}{(t+1)^3} \right]_0^{\infty}$$

$$= \frac{3}{2} - 1 = \frac{1}{2}$$

6. Let  $\vec{a} = 2\hat{i} + \hat{j} - \hat{k}$  and  $\vec{b} = \hat{i} + 2\hat{j} + \hat{k}$  be two vectors. Consider a vector  $\vec{c} = \alpha\vec{a} + \beta\vec{b} + \alpha, \beta \in \mathbb{R}$ . If the projection of  $\vec{c}$  on the vector  $(\vec{a} + \vec{b})$  is  $3\sqrt{2}$ , then the minimum value of  $(\vec{c} - (\vec{a} \times \vec{b})) \cdot \vec{c}$  equals

**Solution:**

$$\vec{a} = 2i + j - k$$

$$\vec{b} = i + 2j + k$$

DATE:

MATHS

CLASS:

JEE ADVANCED 2019

ALL CENTRE

PAPER 2

$$\begin{aligned}\vec{c} &= \alpha\vec{a} + \beta\vec{b} = \alpha(2\vec{i} + \vec{j} - \vec{k}) + \beta(\vec{i} + 2\vec{j} + \vec{k}) \\ &= (2\alpha + \beta)\vec{i} + (\alpha + 2\beta)\vec{j} + (\beta - \alpha)\vec{k}\end{aligned}$$

$$\text{Given } \frac{\vec{c} \cdot (\vec{a} + \vec{b})}{|\vec{a} + \vec{b}|} = 3\sqrt{2}$$

$$\Rightarrow 9(\alpha + \beta) = 18 \Rightarrow \alpha + \beta = 2$$

$$\begin{aligned}(\vec{c} - \alpha \times \vec{b}) \cdot \vec{c} &= (\alpha\vec{a} + \beta\vec{b} + \vec{a} \times \vec{b}) \cdot (\alpha\vec{a} + \beta\vec{b}) \\ &= 6\alpha^2 + 6\alpha\beta + 6\beta^2 = 6[\alpha^2 + \alpha(2 - \alpha) + (2 - \alpha)^2] \\ &= 6(\alpha^2 - 2\alpha + 4)\end{aligned}$$

Minimum value = 18

### SECTION - 3

1. Answer the following by appropriately matching the lists based on the information given in the paragraph  
Let  $f(x) = \sin(\pi \cos x)$  and  $g(x) = \cos(2\pi \sin x)$  be two functions defined for  $x > 0$ . Define the following sets whose elements are written in the increasing order:

$$X = \{x : f(x) = 0\}, \quad Y = \{x : f'(x) = 0\}$$

$$Z = \{x : g(x) = 0\}, \quad W = \{x : g'(x) = 0\}$$

List - I contains the sets X, Y, Z and W. List - II contains some information regarding these sets.

#### List I

(I) X

(II) Y

(III) Z

(IV) W

#### List - II

$$(P) \supseteq \left\{ \frac{\pi}{2}, \frac{3\pi}{2}, 4\pi, 7\pi \right\}$$

(Q) an arithmetic progression

(R) Not an arithmetic progression

$$(S) \supseteq \left\{ \frac{\pi}{6}, \frac{7\pi}{6}, \frac{13\pi}{6} \right\}$$

$$(T) \supseteq \left\{ \frac{\pi}{3}, \frac{2\pi}{3}, \pi \right\}$$

$$(U) \supseteq \left\{ \frac{\pi}{6}, \frac{3\pi}{4} \right\}$$

Which of the following is the only correct combination?

- (a) (II), (R), (S)                      (b) (I), (P), (R)                      (c) (II), (Q), (T)                      (d) (I), (Q), (U)

2. Answer the following by appropriately matching the lists based on the information given in the paragraph  
Let  $f(x) = \sin(\pi \cos x)$  and  $g(x) = \cos(2\pi \sin x)$  be two functions defined for  $x > 0$ . Define the following sets whose element are written in the increasing order:

$$X = \{x : f(x) = 0\}, \quad Y = \{x : f'(x) = 0\}$$

$$Z = \{x : g(x) = 0\}, \quad W = \{x : g'(x) = 0\}$$

List –I contains the sets X,Y,Z and W. List – II contains some information regarding these sets.

**List I**

(I) X

(II) Y

(III) Z

(IV) W

**List – II**

$$(P) \supseteq \left\{ \frac{\pi}{2}, \frac{3\pi}{2}, 4\pi, 7\pi \right\}$$

(Q) an arithmetic progression

(R) Not an arithmetic progression

$$(S) \supseteq \left\{ \frac{\pi}{6}, \frac{7\pi}{6}, \frac{13\pi}{6} \right\}$$

$$(T) \supseteq \left\{ \frac{\pi}{3}, \frac{2\pi}{3}, \pi \right\}$$

$$(U) \supseteq \left\{ \frac{\pi}{6}, \frac{3\pi}{4} \right\}$$

Which of the following is the only correct combination?

- (a) (IV), (Q), (T)                      (b) (IV), (P), (R), (S)                      (c) (III), (R), (U)                      (d) (III), (P), (Q), (U)

**Solution:**

$$f(x) = 0 \Rightarrow \sin(\pi \cos x) = 0$$

$$\Rightarrow \pi \cos x = n\pi \Rightarrow \cos x = n \Rightarrow \cos x = -1, 0, 1$$

$$x = \left\{ n\pi, (2n\pi) \frac{\pi}{2} \right\}$$

$$x = \left\{ \frac{n\pi}{2}, n \in \mathbb{I} \right\}$$

DATE:

MATHS

CLASS:

JEE ADVANCED 2019 PAPER 2

ALL CENTRE

$$f'(x) = 0 \Rightarrow \cos(\pi \cos x)(-\pi \sin x) = 0$$

$$\Rightarrow \pi \cos x = (2n+1)\frac{\pi}{2} \text{ or } x = n\pi$$

$$\Rightarrow \cos x = n + \frac{1}{2} \text{ or } x = n\pi$$

$$\Rightarrow \cos x = \pm \frac{1}{2} \text{ or } x = n\pi$$

$$\therefore y = \left\{ 2n\pi \pm \frac{\pi}{3}, 2n\pi \pm \frac{2\pi}{3}, n\pi \right\}$$

$$g(x) = 0 \Rightarrow \cos(2\pi \sin x) = 0$$

$$\Rightarrow 2\pi \sin x = (2n+1)\frac{\pi}{2}$$

$$\Rightarrow \sin x = \frac{2n+1}{4} = \pm \frac{1}{4}, \pm \frac{3}{4}$$

$$z = \left\{ n\pi \pm \sin^{-1} \frac{1}{4}, n\pi \pm \sin^{-1} \frac{3}{4}, n \in I \right\}$$

$$g'(x) = 0 \Rightarrow -\sin(2\pi \sin x)(2\pi \cos x) = 0$$

$$\Rightarrow 2\pi \sin x = n\pi \text{ or } x = (2n+1)\frac{\pi}{2}$$

$$\Rightarrow \sin x = \frac{n}{2} = 0, \pm \frac{1}{2}, \pm 1 \text{ or } x = (2n+1)\frac{\pi}{2}$$

$$\Rightarrow W = \left\{ n\pi, (2n+1)\frac{\pi}{2}, n\pi \pm \frac{\pi}{6}, n \in I \right\}$$

(1) Option – 3

(2) Option – 2

3. Answer the following by appropriately matching the lists based on the information given in the paragraph

DATE:

MATHS

CLASS:

JEE ADVANCED 2019 PAPER 2

ALL CENTRE

Let the circles  $C_1 : x^2 + y^2 = 9$  and  $C_2 : (x - 3)^2 + (y - 4)^2 = 16$ , intersect at the points X and Y. Suppose that another circle  $C_3 : (x - h)^2 + (y - k)^2 = r^2$  satisfies the following conditions:

- (i) centre of  $C_3$  is collinear with the centres of  $C_1$  and  $C_2$
- (ii)  $C_1$  and  $C_2$  both lie inside  $C_3$ , and
- (iii)  $C_3$  touches  $C_1$  at M and  $C_2$  at N

Let the line through X and Y intersect  $C_3$  at Z and W, and let a common tangent of  $C_1$  and  $C_3$  be a tangent to the parabola  $x^2 = 8\alpha y$ .

There are some expression given in the List – I whose values are given in List – II below:

- | List – I  | List – II          |
|---|--------------------|
| (I) $2h + k$  | (P) 6              |
| (II) $\frac{\text{Length of } ZW}{\text{Length of } XY}$                  | (Q) $\sqrt{6}$     |
| (III) $\frac{\text{Area of triangle } MZN}{\text{Area of triangle } ZMW}$ | (R) $\frac{5}{4}$  |
| (IV) $\alpha$   | (S) $\frac{21}{5}$ |
|   | (T) $2\sqrt{6}$    |
|   | (U) $\frac{10}{3}$ |

Which of the following is the only INCORRECT combination?

- (a) (IV), (S)                      (b) (IV), (U)                      (c) (III), (R)                      (d) (I), (P)

**Solution:**

(ii) Equation of line zw

$$C_1 = C_2$$

$$\Rightarrow 3x + 4y = 9$$

$\Rightarrow$  Distance of zw from (0, 0)

$$\left| \frac{-9}{\sqrt{3^2 + 4^2}} \right| = \frac{9}{5}$$



DATE:

MATHS

CLASS:

JEE ADVANCED 2019 PAPER 2

ALL CENTRE

$$\text{Length of } xy = 2\sqrt{9 - \left(\frac{9}{5}\right)^2} = \frac{24}{5}$$

Distance of  $zw$  from  $c$

$$\frac{\left| \frac{3 \times 9}{5} + 4 \times \frac{12}{5} - 9 \right|}{\sqrt{3^2 + 4^2}} = \frac{6}{5}$$

$$\text{Length of } zw = 2\sqrt{6^2 - \frac{6^2}{5^2}} = \frac{24\sqrt{6}}{5}$$

$$\frac{\text{length of } zw}{\text{length of } xy} = \sqrt{6}$$

$$\text{(iii) Area of } \Delta mzn = \frac{1}{2} \cdot Nm \cdot \left(\frac{1}{2} zw\right) = \frac{72\sqrt{6}}{5}$$

$$\text{Area of } \Delta zmw = \frac{1}{2} \cdot zw \cdot (om + op) = \frac{1}{2} \cdot \frac{24\sqrt{6}}{5} \cdot \left(3 + \frac{9}{5}\right) = \frac{288\sqrt{6}}{25}$$

$$\therefore \frac{\text{Area of } \Delta mzn}{\text{Area of } \Delta zmw} = \frac{5}{4}$$

$$\text{(iv) Slope of tangent to } C_1 \text{ at } m = \frac{-1}{\frac{4}{3}} = -\frac{3}{4}$$

$$\text{Equation of Tangent } y = mx - 2\sqrt{1+m^2}$$

$$y = \frac{-3x}{4} - 3\sqrt{1 + \frac{9}{16}}$$

$$y = \frac{-3x - 15}{4}$$

DATE:

MATHS

CLASS:

JEE ADVANCED 2019 PAPER 2

ALL CENTRE

$$\Rightarrow x = \frac{-4y}{3} - 5 \quad \rightarrow 1$$

$$\text{Tangent to } x^2 = 4(2d)y \text{ is } x = m'y + \frac{2d}{m'} \quad \rightarrow 2$$

Compare 1 and 2

$$m' = \frac{-4}{3} \text{ and } \frac{2\infty}{m'} = -5 \quad \Rightarrow \infty = \frac{10}{3}$$

4. Answer the following by appropriately matching the lists based on the information given in the paragraph  
Let the circles  $C_1 : x^2 + y^2 = 9$  and  $C_2 : (x - 3)^2 + (y - 4)^2 = 16$ , intersect at the points X and Y. Suppose that another circle  $C_3 : (x - h)^2 + (y - k)^2 = r^2$  satisfies the following conditions:

- (i) centre of  $C_3$  is collinear with the centres of  $C_1$  and  $C_2$
- (ii)  $C_1$  and  $C_2$  both lie inside  $C_3$ , and
- (iii)  $C_3$  touches  $C_1$  at M and  $C_2$  at N

Let the line through X and Y intersect  $C_3$  at Z and W, and let a common tangent of  $C_1$  and  $C_3$  be a tangent to the parabola  $x^2 = 8\alpha y$ .

There are some expression given in the List – I whose values are given in List – II below:

- | List – I  | List – II          |
|---|--------------------|
| (I) $2h + k$  | (P) 6              |
| (II) $\frac{\text{Length of } ZW}{\text{Length of } XY}$                  | (Q) $\sqrt{6}$     |
| (III) $\frac{\text{Area of triangle } MZN}{\text{Area of triangle } ZMW}$ | (R) $\frac{5}{4}$  |
| (IV) $\alpha$   | (S) $\frac{21}{5}$ |
|   | (T) $2\sqrt{6}$    |
|   | (U) $\frac{10}{3}$ |

Which of the following is the only INCORRECT combination?

- (a) (II), (T)                      (b) (I), (S)                      (c) (I), (U)                      (d) (II), (Q)

**Solution:**

DATE:

MATHS

CLASS:

JEE ADVANCED 2019

ALL CENTRE

PAPER 2

$$2r = MN = 3 + \sqrt{3^2 + 4^2} + 4 = 12$$

$$\Rightarrow r = 6$$

Centre  $c$  of circle  $c_3$  lies on  $y = \frac{4}{3}x$

$$\text{Let } c \left( h, \frac{4}{3}h \right)$$

$$OC = MC - OM = \frac{12}{2} - 3 = 3$$

$$\sqrt{h^2 + \frac{16}{9}h^2} = 3 \Rightarrow h = \frac{9}{5}$$

$$k = \frac{4}{3}h = \frac{12}{5} \Rightarrow 2h + k = 6$$

