

Complex Numbers & Quadratic Equations

EXERCISE 5A

PAGE: 153

Q. 1. Evaluate:

(i) i^{19}

(ii) i^{62}

(ii) i^{373} .

Solution: We all know that $i = \sqrt{-1}$.

And $i^{4n} = 1$

$i^{4n+1} = i$ (where n is any positive integer)

$i^{4n+2} = -1$

$i^{4n+3} = -i$

So,

(i) L.H.S = i^{19}

= $i^{4 \times 4 + 3}$

= i^{4n+3}

Since it is of the form i^{4n+3} so the solution would be simply $-i$

Hence the value of i^{19} is $-i$.

(ii) L. H. S = i^{62}

$\Rightarrow i^{4 \times 15 + 2}$

$\Rightarrow i^{4n+2} \Rightarrow i^2 = -1$

so it is of the form i^{4n+2} so its solution would be -1

(iii) L.H.S. = i^{373}

$$\Rightarrow i^{4 \times 93 + 1}$$

$$\Rightarrow i^{4n+1}$$

$$\Rightarrow i$$

So, it is of the form of i^{4n+1} so the solution would be i .

Q. 2. Evaluate:

(i) $(\sqrt{-1})^{192}$

(ii) $(\sqrt{-1})^{93}$

(iii) $(\sqrt{-1})^{30}$.

Solution: Since $i = \sqrt{-1}$ so

(i) L.H.S. = $(\sqrt{-1})^{192}$

$$\Rightarrow i^{192}$$

$$\Rightarrow i^{4 \times 48} = 1$$

Since it is of the form $i^{4n} = 1$ so the solution would be 1

(ii) L.H.S. = $(\sqrt{-1})^{93}$

$$\Rightarrow i^{4 \times 23 + 1}$$

$$\Rightarrow i^{4n+1}$$

$$\Rightarrow i^1 = i$$

Since it is of the form of $i^{4n+1} = i$ so the solution would be simply i .

$$\text{(iii) L.H.S} = (\sqrt{-1})^{30}$$

$$\Rightarrow i^{4 \times 7 + 2}$$

$$\Rightarrow i^{4n+2}$$

$$\Rightarrow i^2 = -1$$

Since it is of the form i^{4n+2} so the solution would be -1

Q. 3. Evaluate:

(i) i^{-50}

(ii) i^{-9}

(ii) i^{-131} .

Solution: (i) L.H.S. = i^{-50}

$$\Rightarrow i^{-4 \times 13 + 2}$$

$$\Rightarrow i^{4n+2}$$

$$\Rightarrow -1$$

Since it is of the form i^{4n+2} so the solution would be -1

(ii) L.H.S. = i^{-9}

$$\Rightarrow i^{-4 \times 3 + 3}$$

$$\Rightarrow i^{4n+3}$$

$$\Rightarrow i^3 = -i$$

Since it is of the form of i^{4n+3} so the solution would be simply $-i$.

(iii) L.H.S. = i^{-131}

$$\Rightarrow i^{-4 \times 33 + 1}$$

$$\Rightarrow i^{4n+1}$$

$$\Rightarrow i^1 = i$$

Since it is of the form i^{4n+1} , so the solution would be i

Q. 4. Evaluate:

(i) $\left(i^{41} + \frac{1}{i^{71}} \right)$

(ii) $\left(i^{53} + \frac{1}{i^{53}} \right)$

Solution:

(i) $\left(i^{41} + \frac{1}{i^{71}} \right) = i^{41} + i^{-71}$

$$\Rightarrow i^{4 \times 10 + 1} + i^{-4 \times 18 + 1} \quad (\text{Since } i^{4n+1} = i)$$

$$\Rightarrow i^1 + i^1$$

$$\Rightarrow 2i$$

Hence, $\left(i^{41} + \frac{1}{i^{71}} \right) = 2i$

(ii) $\left(i^{53} + \frac{1}{i^{53}} \right)$

$$\Rightarrow i^{53} + i^{-53}$$

$$\Rightarrow i^{4 \times 13 + 1} + i^{-4 \times 14 + 3} \quad (\text{Since } i^{4n+1} = i$$

$$\Rightarrow i^1 + i^3 \quad (\text{As } i^3 = -1)$$

$$\Rightarrow 0$$

Hence, $\left(i^{53} + \frac{1}{i^{53}} \right) = 0$

Q. 5. Prove that $1 + i^2 + i^4 + i^6 = 0$

Solution: L.H.S. = $1 + i^2 + i^4 + i^6$

To Prove: $1 + i^2 + i^4 + i^6 = 0$

$$\Rightarrow 1 + (-1) + 1 + i^2$$

Since, $i^{4n} = 1$

(Where n is any positive integer)

$$\Rightarrow i^{4n+2}$$

$$\Rightarrow i^2 = -1$$

$$\Rightarrow 1 + -1 + 1 + -1 = 0$$

\Rightarrow L.H.S = R.H.S

Hence proved.

Q. 6. Prove that $6i^{50} + 5i^{33} - 2i^{15} + 6i^{48} = 7i$.

Solution: Given: $6i^{50} + 5i^{33} - 2i^{15} + 6i^{48}$

To prove: $6i^{50} + 5i^{33} - 2i^{15} + 6i^{48} = 7i$

$$\Rightarrow 6i^{4 \times 12 + 2} + 5i^{4 \times 8 + 1} - 2i^{4 \times 3 + 3} + 6i^{4 \times 12}$$

$$\Rightarrow 6i^2 + 5i^1 - 2i^3 + 6i^0$$

$$\Rightarrow -6 + 5i + 2i + 6$$

$$\Rightarrow 7i$$

$$\Rightarrow \text{L.H.S} = \text{R.H.S}$$

Hence proved.

Q. 7. Prove that $\frac{1}{i} - \frac{1}{i^2} + \frac{1}{i^3} - \frac{1}{i^4} = 0.$

Solution:

Given: $\frac{1}{i} - \frac{1}{i^2} + \frac{1}{i^3} - \frac{1}{i^4}$

To prove: $\frac{1}{i} - \frac{1}{i^2} + \frac{1}{i^3} - \frac{1}{i^4} = 0.$

$$\Rightarrow \text{L.H.S.} = i^{-1} - i^{-2} + i^{-3} - i^{-4}$$

$$\Rightarrow i^{-4 \times 1 + 3} - i^{-4 \times 1 + 2} + i^{-4 \times 1 + 3} - i^{-4 \times 1}$$

Since $i^{4n} = 1$

$$\Rightarrow i^{4n+1} = i$$

$$\Rightarrow i^{4n+2} = -1$$

$$\Rightarrow i^{4n+3} = -i$$

So,

$$\Rightarrow i^1 - i^2 + i^3 - 1$$

$$\Rightarrow i+1-i-1$$

$$\Rightarrow 0$$

$$\Rightarrow \text{L.H.S} = \text{R.H.S}$$

Hence Proved

Q. 8. Prove that $(1 + i^{10} + i^{20} + i^{30})$ is a real number.

$$\text{Solution: L.H.S} = (1 + i^{10} + i^{20} + i^{30})$$

$$= (1 + i^{4 \times 2 + 2} + i^{4 \times 5} + i^{4 \times 7 + 2})$$

$$\text{Since } \Rightarrow i^{4n} = 1$$

$$\Rightarrow i^{4n+1} = i$$

$$\Rightarrow i^{4n+2} = -1$$

$$\Rightarrow i^{4n+3} = -i$$

$$= 1 + i^2 + 1 + i^2$$

$$= 1 + -1 + 1 + -1$$

$$= 0, \text{ which is a real no.}$$

Hence, $(1 + i^{10} + i^{20} + i^{30})$ is a real number.

$$\text{Q. 9. Prove that } \left\{ i^{21} - \left(\frac{1}{i} \right)^{46} \right\}^2 = 2i.$$

$$\text{Solution: L.H.S.} = \left\{ i^{21} - \left(\frac{1}{i} \right)^{46} \right\}^2$$

$$= \left\{ i^{4 \times 5 + 1} - i^{-4 \times 12 + 2} \right\}^2$$

Since $i^{4n} = 1$

$$i^{4n+1} = i$$

$$i^{4n+2} = i^2 = -1$$

$$i^{4n+3} = i^3 = -i$$

$$= \left\{ i^1 - i^2 \right\}^2$$

$$= \left\{ i + 1 \right\}^2$$

Now, applying the formula $(a + b)^2 = a^2 + b^2 + 2ab$

$$= i^2 + 1 + 2i.$$

$$= -1 + 1 + 2i$$

$$= 2i$$

L.H.S = R.H.S

Hence proved.

Q. 10. $\left\{ i^{18} + \frac{1}{i^{25}} \right\}^3 = 2(1 - i).$

Solution: L.H.S = $\left\{ i^{18} + \frac{1}{i^{25}} \right\}^3$

$$\Rightarrow \left\{ i^{4 \times 4 + 2} + i^{-4 \times 7 + 3} \right\}^3$$

Since $i^{4n} = 1$

$$i^{4n+1} = i$$

$$i^{4n+2} = -1$$

$$i^{4n+3} = -i$$

$$= \left\{ i^2 + i^3 \right\}^3$$

$$= (-1 - i)^3$$

Applying the formula $(a + b)^3 = a^3 + b^3 + 3ab(a + b)$

We have,

$$+ 3i^2 + 3i + 1)$$

$$i + 3 - 3i - 1$$

$$= 2(1-i)$$

$$\text{L.H.S} = \text{R.H.S}$$

Hence proved.

Q. 11. Prove that $(1 - i)^n \left(1 - \frac{1}{i}\right)^n = 2^n$ for all values of $n \in \mathbb{N}$

$$\text{Solution: L.H.S} = (1-i)^n \left(1 - \frac{1}{i}\right)^n$$

$$= (1-i)^n (1-i^{-4*1+3})^n$$

$$= (1-i)^n (1-i^3)^n$$

Since, $i^{4n+3} = -i$

$$= (1-i)^n (1+i)^n$$

Applying $a^n b^n = (ab)^n$

$$= ((1-i)(1+i))^n$$

$$= (1-i^2)^n$$

$$= 2^n$$

L.H.S = R.H.S

Q. 12. Prove that $\sqrt{-16} + 3\sqrt{-25} + \sqrt{-36} - \sqrt{-625} = 0$.

Solution: L.H.S = $\sqrt{-16} + 3\sqrt{-25} + \sqrt{-36} - \sqrt{-625}$

Since we know that $i = \sqrt{-1}$,

So,

$$= \sqrt{16} i + 3\sqrt{25} i + \sqrt{36} i - \sqrt{625} i$$

$$=4i + 15i + 6i - 25i$$

$$= 0$$

$$\text{L.H.S} = \text{R.H.S}$$

Hence proved.

Q. 13. Prove that $(1 + i^2 + i^4 + i^6 + i^8 + \dots + i^{20}) = 1$.

$$\text{Solution: L.H.S} = (1 + i^2 + i^4 + i^6 + i^8 + \dots + i^{20})$$

$$= \sum_{n=0}^{n=20} i^n$$

$$= 1 + -1 + 1 + -1 + \dots + 1$$

As there are 11 times 1 and 6 times it is with positive sign as $i^0 = 1$ as this is the extra term and there are 5 times 1 with negative sign -

So, these 5 cancel out the positive one leaving one positive value i.e. 1

$$= \sum_{n=0}^{20} i^n = 1$$

$$\text{L.H.S} = \text{R.H.S}$$

Hence proved.

Q. 14. Prove that $i^{53} + i^{72} + i^{93} + i^{102} = 2i$.

$$\text{Solution: L.H.S} = i^{53} + i^{72} + i^{93} + i^{102}$$

$$= i^{4 \times 13 + 1} + i^{4 \times 18} + i^{4 \times 23 + 1} + i^{4 \times 25 + 2}$$

$$\text{Since } i^{4n} = 1$$

$$\Rightarrow i^{4n+1} = i \text{ (where } n \text{ is any positive integer)}$$

$$\Rightarrow i^{4n+2} = -1$$

$$\Rightarrow i^{4n+3} = -i$$

$$= i + 1 + i + i^2$$

$$= i + 1 + i - 1$$

$$= 2i$$

$$\text{L.H.S} = \text{R.H.S}$$

Hence proved.

Q. 15. Prove that $\sum_{n=1}^{13} (i^n + i^{n+1}) = (-1 + i),$ **n N.**

Solution: L.H.S $= \sum_{n=1}^{13} (i^n + i^{n+1})$

$$= i^1 + i^2 + i^3 + i^4 + i^5 + i^6 + \dots + i^{13} + i^{14}$$

Since $i^{4n} = 1$

$$\Rightarrow i^{4n+1} = i$$

$$\Rightarrow i^{4n+2} = -1$$

$$\Rightarrow i^{4n+3} = -i$$

$$= i - 1 - i + 1 + i - 1 \dots \dots + i - 1$$

As, all terms will get cancel out consecutively except the first two terms. So that will get remained will be the answer.

$$= i - 1$$

$$\text{L.H.S} = \text{R.H.S}$$

Hence proved.

EXERCISE 5B

PAGE: 171

Q. 1. A. Simplify each of the following and express it in the form $a + ib$:

$$2(3 + 4i) + i(5 - 6i)$$

Solution: Given: $2(3 + 4i) + i(5 - 6i)$

Firstly, we open the brackets

$$2 \times 3 + 2 \times 4i + i \times 5 - i \times 6i$$

$$= 6 + 8i + 5i - 6i^2$$

$$= 6 + 13i - 6(-1) [\because i^2 = -1]$$

$$= 6 + 13i + 6$$

$$= 12 + 13i$$

$\underbrace{\hspace{1.5cm}}$ Real part	$\underbrace{\hspace{1.5cm}}$ Imaginary part
---	--

Q. 1. B. Simplify each of the following and express it in the form $a + ib$:

$$(3 + \sqrt{-16}) - (4 - \sqrt{-9})$$

Solution: Given: $(3 + \sqrt{-16}) - (4 - \sqrt{-9})$

We re - write the above equation

$$(3 + \sqrt{(-1) \times 16}) - (4 - \sqrt{(-1) \times 9})$$

$$= (3 + \sqrt{16i^2}) - (4 - \sqrt{9i^2}) [\because i^2 = -1]$$

$$= (3 + 4i) - (4 - 3i)$$

Now, we open the brackets, we get

$$3 + 4i - 4 + 3i$$

$$= -1 + 7i$$

$$\begin{array}{cc} \underbrace{\quad} & \underbrace{\quad} \\ \text{Real} & \text{Imaginary} \\ \text{part} & \text{part} \end{array}$$

Q. 1. C. Simplify each of the following and express it in the form $a + ib$:

$$(-5 + 6i) - (-2 + i)$$

Solution: Given: $(-5 + 6i) - (-2 + i)$

Firstly, we open the brackets

$$-5 + 6i + 2 - i$$

$$= -3 + 5i$$

$$\begin{array}{cc} \underbrace{\quad} & \underbrace{\quad} \\ \text{Real} & \text{Imaginary} \\ \text{part} & \text{part} \end{array}$$

Q. 1. D. Simplify each of the following and express it in the form $a + ib$:

$$(8 - 4i) - (-3 + 5i)$$

Solution: Given: $(8 - 4i) - (-3 + 5i)$

Firstly, we open the brackets

$$8 - 4i + 3 - 5i$$

$$= 11 - 9i$$

$$\begin{array}{cc} \underbrace{\quad} & \underbrace{\quad} \\ \text{Real} & \text{Imaginary} \\ \text{part} & \text{part} \end{array}$$

Q. 1. E. Simplify each of the following and express it in the form $a + ib$:

$$(1 - i)^2(1 + i) - (3 - 4i)^2$$

Solution: Given: $(1 - i)^2(1 + i) - (3 - 4i)^2$

$$= (1 + i^2 - 2i)(1 + i) - (9 + 16i^2 - 24i)$$

$$[\because (a - b)^2 = a^2 + b^2 - 2ab]$$

$$= (1 - 1 - 2i)(1 + i) - (9 - 16 - 24i) [\because i^2 = -1]$$

$$= (-2i)(1 + i) - (-7 - 24i)$$

Now, we open the brackets

$$= -2i \times 1 - 2i \times i + 7 + 24i$$

$$= -2i - 2i^2 + 7 + 24i$$

$$= -2(-1) + 7 + 22i [\because i^2 = -1]$$

$$= 2 + 7 + 22i$$

$$= 9 + 22i$$

$\underbrace{\quad}$ $\underbrace{\quad}$
 Real Imaginary
 part part

Q. 1. F. Simplify each of the following and express it in the form $a + ib$:

$$(5 + \sqrt{-3})(5 - \sqrt{-3})$$

Solution: Given: $(5 + \sqrt{-3})(5 - \sqrt{-3})$

We re - write the above equation

$$(5 + \sqrt{(-1) \times 3})(5 - \sqrt{(-1) \times 3})$$

$$= (5 + \sqrt{3i^2})(5 - \sqrt{3i^2}) [\because i^2 = -1]$$

$$= (5 + i\sqrt{3})(5 - i\sqrt{3})$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

Here, $a = 5$ and $b = i\sqrt{3}$

$$= (5)^2 - (i\sqrt{3})^2$$

$$= 25 - (3i^2)$$

$$= 25 - [3 \times (-1)]$$

$$= 25 + 3$$

$$= 28 + 0$$

$$= 28 + 0i$$

$\underbrace{\hspace{1.5cm}}_{\text{Real part}} \quad \underbrace{\hspace{1.5cm}}_{\text{Imaginary part}}$

Q. 1. G. Simplify each of the following and express it in the form $a + ib$:

$$(3 + 4i)(2 - 3i)$$

Solution: Given: $(3 + 4i)(2 - 3i)$

Firstly, we open the brackets

$$3 \times 2 + 3 \times (-3i) + 4i \times 2 - 4i \times 3i$$

$$= 6 - 9i + 8i - 12i^2$$

$$= 6 - i - 12(-1) [\because i^2 = -1]$$

$$= 6 - i + 12$$

$$= 18 - i$$

$\underbrace{\quad}$ $\underbrace{\quad}$
 Real part Imaginary part

Q. 1. H. Simplify each of the following and express it in the form $a + ib$:

$$(-2 + \sqrt{-3})(-3 + 2\sqrt{-3})$$

Solution: Given: $(-2 + \sqrt{-3})(-3 + 2\sqrt{-3})$

We re – write the above equation

$$(-2 + \sqrt{(-1) \times 3})(-3 + 2\sqrt{(-1) \times 3})$$

$$= (-2 + \sqrt{3i^2})(-3 + 2\sqrt{3i^2}) \quad [\because i^2 = -1]$$

$$= (-2 + i\sqrt{3})(-3 + 2i\sqrt{3})$$

Now, open the brackets,

$$= -2 \times (-3) + (-2) \times 2i\sqrt{3} + i\sqrt{3} \times (-3) + i\sqrt{3} \times 2i\sqrt{3}$$

$$= 6 - 4i\sqrt{3} - 3i\sqrt{3} + 6i^2$$

$$= 6 - 7i\sqrt{3} + [6 \times (-1)] \quad [\because i^2 = -1]$$

$$= 6 - 7i\sqrt{3} - 6$$

$$= 0 - 7i\sqrt{3}$$

$\underbrace{\quad}$ $\underbrace{\quad}$
 Real part Imaginary part

Q. 2. A. Simplify each of the following and express it in the form $(a + ib)$:

$$(2 + \sqrt{-3})^2$$

Solution: Given: $(2 - \sqrt{-3})^2$

We know that,

$$(a - b)^2 = a^2 + b^2 - 2ab \dots(i)$$

So, on replacing a by 2 and b by $\sqrt{-3}$ in eq. (i), we get

$$(2)^2 + (\sqrt{-3})^2 - 2(2)(\sqrt{-3})$$

$$= 4 + (-3) - 4\sqrt{-3}$$

$$= 4 - 3 - 4\sqrt{-3}$$

$$= 1 - 4\sqrt{3}i^2 [\because i^2 = -1]$$

$$= 1 - 4i\sqrt{3}$$

$\underbrace{\quad}$ $\underbrace{\quad}$
 Real Imaginary
 part part

Q. 2. B. Simplify each of the following and express it in the form (a + ib) :

$$(5 - 2i)^2$$

Solution: Given: $(5 - 2i)^2$

We know that,

$$(a - b)^2 = a^2 + b^2 - 2ab \dots(i)$$

So, on replacing a by 5 and b by 2i in eq. (i), we get

$$(5)^2 + (2i)^2 - 2(5)(2i)$$

$$= 25 + 4i^2 - 20i$$

$$= 25 - 4 - 20i [\because i^2 = -1]$$

$$= 21 - 20i$$

$\underbrace{\hspace{1cm}}$ $\underbrace{\hspace{1cm}}$
 Real part Imaginary part

Q. 2. C. Simplify each of the following and express it in the form (a + ib) :

$$(-3 + 5i)^3$$

Solution: Given: $(-3 + 5i)^3$

We know that,

$$(-a + b)^3 = -a^3 + 3a^2b - 3ab^2 + b^3 \dots(i)$$

So, on replacing a by 3 and b by 5i in eq. (i), we get

$$\begin{aligned} &-(3)^3 + 3(3)^2(5i) - 3(3)(5i)^2 + (5i)^3 \\ &= -27 + 3(9)(5i) - 3(3)(25i^2) + 125i^3 \\ &= -27 + 135i - 225i^2 + 125i^3 \\ &= -27 + 135i - 225 \times (-1) + 125i \times i^2 \\ &= -27 + 135i + 225 - 125i [\because i^2 = -1] \end{aligned}$$

$$= 198 + 10i$$

$\underbrace{\hspace{1cm}}$ $\underbrace{\hspace{1cm}}$
 Real part Imaginary part

Q. 2. D. Simplify each of the following and express it in the form (a + ib) :

$$\left(-2 - \frac{1}{3}i\right)^3$$

Solution: Given: $\left(-2 - \frac{1}{3}i\right)^3$

We know that,

$$(-a - b)^3 = -a^3 - 3a^2b - 3ab^2 - b^3 \dots (i)$$

So, on replacing a by 2 and b by $\frac{1}{3}i$ in eq. (i), we get

$$-(2)^3 - 3(2)^2\left(\frac{1}{3}i\right) - 3(2)\left(\frac{1}{3}i\right)^2 - \left(\frac{1}{3}i\right)^3$$

$$= -8 - 4i - 6\left(\frac{1}{9}i^2\right) - \left(\frac{1}{27}i^3\right)$$

$$= -8 - 4i - \frac{2}{3}i^2 - \frac{1}{27}i(i^2)$$

$$= -8 - 4i - \frac{2}{3}(-1) - \frac{1}{27}i(-1) \quad [\because i^2 = -1]$$

$$= -8 - 4i + \frac{2}{3} + \frac{1}{27}i$$

$$= \left(-8 + \frac{2}{3}\right) + \left(-4i + \frac{1}{27}i\right)$$

$$= \left(\frac{-24 + 2}{3}\right) + \left(\frac{-108i + i}{27}\right)$$

$$= -\frac{22}{3} + \left(-\frac{107}{27}i\right)$$

$$= -\frac{22}{3} - \frac{107}{27}i$$

$\underbrace{\hspace{1.5cm}}$
 $\underbrace{\hspace{1.5cm}}$

Real part Imaginary part

Q. 2. E. Simplify each of the following and express it in the form (a + ib) :

$$(4 - 3i)^{-1}$$

Solution: Given: $(4 - 3i)^{-1}$

We can re- write the above equation as

$$= \frac{1}{4 - 3i}$$

Now, rationalizing

$$= \frac{1}{4 - 3i} \times \frac{4 + 3i}{4 + 3i}$$

$$= \frac{4 + 3i}{(4 - 3i)(4 + 3i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{4 + 3i}{(4)^2 - (3i)^2}$$

$$= \frac{4 + 3i}{16 - 9i^2}$$

$$= \frac{4 + 3i}{16 - 9(-1)} [\because i^2 = -1]$$

$$= \frac{4 + 3i}{16 + 9}$$

$$= \frac{4 + 3i}{25}$$

$$= \frac{4}{25} + \frac{3}{25}i$$

$$\underbrace{\quad}_{\text{Real part}} \quad \underbrace{\quad}_{\text{Imaginary part}}$$

Real part Imaginary part

Q. 2. F. Simplify each of the following and express it in the form (a + ib) :

$$\left(-2 + \sqrt{-3}\right)^{-1}$$

Solution: Given: $(-2 + \sqrt{-3})^{-1}$

We can re- write the above equation as

$$= \frac{1}{-2 + \sqrt{-3}}$$

$$= \frac{1}{-2 + \sqrt{3i^2}} \quad [\because i^2 = -1]$$

$$= \frac{1}{-2 + i\sqrt{3}}$$

Now, rationalizing

$$= \frac{1}{-2 + i\sqrt{3}} \times \frac{-2 - i\sqrt{3}}{-2 - i\sqrt{3}}$$

$$= \frac{-2 - i\sqrt{3}}{(-2 + i\sqrt{3})(-2 - i\sqrt{3})} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{-2 - i\sqrt{3}}{(-2)^2 - (i\sqrt{3})^2}$$

$$= \frac{-2 - i\sqrt{3}}{4 - (3i^2)}$$

$$= \frac{-2 - i\sqrt{3}}{4 - 3(-1)} \quad [\because i^2 = -1]$$

$$= \frac{-2 - i\sqrt{3}}{4 + 3}$$

$$= \frac{-2 - i\sqrt{3}}{7}$$

$$= -\frac{2 + i\sqrt{3}}{7}$$

$$= -\frac{2}{7} - \frac{\sqrt{3}}{7}i$$

$$\underbrace{\quad\quad}_{\text{Real part}} \quad \underbrace{\quad\quad}_{\text{Imaginary part}}$$

Real part Imaginary part

Q. 2. G. Simplify each of the following and express it in the form (a + ib) :

$$(2 + i)^{-2}$$

Solution: Given: $(2 + i)^{-2}$

Above equation can be re – written as

$$= \frac{1}{(2 + i)^2}$$

Now, rationalizing

$$= \frac{1}{(2 + i)^2} \times \frac{(2 - i)^2}{(2 - i)^2}$$

$$= \frac{(2 - i)^2}{(2 + i)^2(2 - i)^2}$$

$$= \frac{4 + i^2 - 4i}{(4 + i^2 + 4i)(4 + i^2 - 4i)} \quad [\because (a - b)^2 = a^2 + b^2 - 2ab]$$

$$= \frac{4 - 1 - 4i}{(4 - 1 + 4i)(4 - 1 - 4i)} \quad [\because i^2 = -1]$$

$$= \frac{3 - 4i}{(3 + 4i)(3 - 4i)} \quad \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{3 - 4i}{(3)^2 - (4i)^2}$$

$$= \frac{3 - 4i}{9 - 16i^2}$$

$$= \frac{3 - 4i}{9 - 16(-1)}$$

$$= \frac{3 - 4i}{25}$$

$$= \frac{3}{25} - \frac{4}{25}i$$

$$\underbrace{\quad}_{\text{Real part}} \quad \underbrace{\quad}_{\text{Imaginary part}}$$

Real part Imaginary part

Q. 2. H. Simplify each of the following and express it in the form (a + ib) :

$$(1 + 2i)^{-3}$$

Solution: Given: $(1 + 2i)^{-3}$

Above equation can be re – written as

$$= \frac{1}{(1 + 2i)^3}$$

Now, rationalizing

$$= \frac{1}{(1 + 2i)^3} \times \frac{(1 - 2i)^3}{(1 - 2i)^3}$$

$$= \frac{(1 - 2i)^3}{(1 + 2i)^3(1 - 2i)^3}$$

We know that,

$$(a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$$

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

$$= \frac{(1)^3 - 3(1)^2(2i) + 3(1)(2i)^2 - (2i)^3}{[(1)^3 + 3(1)^2(2i) + 3(1)(2i)^2 + (2i)^3][(1)^3 - 3(1)^2(2i) + 3(1)(2i)^2 - (2i)^3]}$$

$$= \frac{1 - 6i + 6i^2 - 8i^3}{[1 + 6i + 6i^2 + 8i^3][1 - 6i + 6i^2 - 8i^3]}$$

$$= \frac{1 - 6i + 6(-1) - 8i(-1)}{[1 + 6i + 6(-1) + 8i(-1)][1 - 6i + 6(-1) - 8i(-1)]} [\because i^2 = -1]$$

$$= \frac{1 - 6i - 6 + 8i}{[1 + 6i - 6 - 8i][1 - 6i - 6 + 8i]}$$

$$= \frac{-5 + 2i}{[-5 - 2i][-5 + 2i]}$$


$$= \frac{-5 + 2i}{-5(-5) - 5(2i) - 2i(-5) - 2i(2i)}$$

$$= \frac{-5 + 2i}{25 - 10i + 10i - 4i^2}$$

$$= \frac{-5 + 2i}{25 - 4(-1)} [\because i^2 = -1]$$

$$= \frac{-5 + 2i}{29}$$

$$= -\frac{5}{29} + \frac{2}{29}i$$



 Real part Imaginary part

Q. 2. I. Simplify each of the following and express it in the form (a + ib) :

$$(1 + i)^3 - (1 - i)^3$$

Solution: Given: $(1 + i)^3 - (1 - i)^3 \dots(i)$

We know that,

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

$$(a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$$

By applying the formulas in eq. (i), we get

$$(1)^3 + 3(1)^2(i) + 3(1)(i)^2 + (i)^3 - [(1)^3 - 3(1)^2(i) + 3(1)(i)^2 - (i)^3]$$

$$= 1 + 3i + 3i^2 + i^3 - [1 - 3i + 3i^2 - i^3]$$

$$= 1 + 3i + 3i^2 + i^3 - 1 + 3i - 3i^2 + i^3$$

$$= 6i + 2i^3$$

$$= 6i + 2i(i^2)$$

$$= 6i + 2i(-1) [\because i^2 = -1]$$

$$= 6i - 2i$$

$$= 4i$$

$$= 0 + 4i$$

$\underbrace{\quad}$ $\underbrace{\quad}$
 Real Imaginary
 part part

Q. 3. A. Express each of the following in the form (a + ib):

$$\frac{1}{(4 + 3i)}$$

Solution: Given: $\frac{1}{4+3i}$

Now, rationalizing

$$= \frac{1}{4+3i} \times \frac{4-3i}{4-3i}$$

$$= \frac{4-3i}{(4+3i)(4-3i)} \dots(i)$$

Now, we know that,

$$(a+b)(a-b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{4-3i}{(4)^2 - (3i)^2}$$

$$= \frac{4-3i}{16-9i^2}$$

$$= \frac{4-3i}{16-9(-1)} [\because i^2 = -1]$$

$$= \frac{4-3i}{16+9}$$

$$= \frac{4-3i}{25}$$

$$= \frac{4}{25} - \frac{3}{25}i$$

$$\underbrace{\quad} \quad \underbrace{\quad}$$

Real part Imaginary part

Q. 3. B. Express each of the following in the form (a + ib):

$$\frac{(3+4i)}{(4+5i)}$$

Solution: Given: $\frac{3+4i}{4+5i}$

Now, rationalizing

$$= \frac{3 + 4i}{4 + 5i} \times \frac{4 - 5i}{4 - 5i}$$

$$= \frac{(3+4i)(4-5i)}{(4+5i)(4-5i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{(3 + 4i)(4 - 5i)}{(4)^2 - (5i)^2}$$

$$= \frac{3(4) + 3(-5i) + 4i(4) + 4i(-5i)}{16 - 25i^2}$$

$$= \frac{12 - 15i + 16i - 20i^2}{16 - 25(-1)} \quad [\because i^2 = -1]$$

$$= \frac{12 + i - 20(-1)}{16 + 25}$$

$$= \frac{12 + i + 20}{41}$$

$$= \frac{32 + i}{41}$$

$$= \frac{32}{41} + \frac{1}{41}i$$

$\underbrace{\hspace{1.5cm}}$ $\underbrace{\hspace{1.5cm}}$
 Real Imaginary
 part part

Q. 3. C. Express each of the following in the form (a + ib):

$$\frac{(5 + \sqrt{2}i)}{(1 - \sqrt{2}i)}$$

$$(1 - \sqrt{2}i)$$

Solution: Given: $\frac{5+\sqrt{2}i}{1-\sqrt{2}i}$

Now, rationalizing

$$= \frac{5 + \sqrt{2}i}{1 - \sqrt{2}i} \times \frac{1 + \sqrt{2}i}{1 + \sqrt{2}i}$$

$$= \frac{(5+\sqrt{2}i)(1+\sqrt{2}i)}{(1-\sqrt{2}i)(1+\sqrt{2}i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{(5 + \sqrt{2}i)(1 + \sqrt{2}i)}{(1)^2 - (\sqrt{2}i)^2}$$

$$= \frac{5(1) + 5(\sqrt{2}i) + \sqrt{2}i(1) + \sqrt{2}i(\sqrt{2}i)}{1 - 2i^2}$$

$$= \frac{5+5\sqrt{2}i+\sqrt{2}i+2i^2}{1-2(-1)} \quad [\because i^2 = -1]$$

$$= \frac{5 + 6i\sqrt{2} + 2(-1)}{1 + 2}$$

$$= \frac{3 + 6i\sqrt{2}}{3}$$

$$= \frac{3(1 + 2i\sqrt{2})}{3}$$

$$= 1 + 2i\sqrt{2}$$

$\underbrace{\hspace{1.5cm}}$
 $\underbrace{\hspace{1.5cm}}$

Real part Imaginary part

Q. 3. D. Express each of the following in the form (a + ib):

$$\frac{(-2 + 5i)}{(3 - 5i)}$$

Solution: Given: $\frac{-2+5i}{3-5i}$

Now, rationalizing

$$= \frac{-2 + 5i}{3 - 5i} \times \frac{3 + 5i}{3 + 5i}$$

$$= \frac{(-2+5i)(3+5i)}{(3-5i)(3+5i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{(-2 + 5i)(3 + 5i)}{(3)^2 - (5i)^2}$$

$$= \frac{-2(3) + (-2)(5i) + 5i(3) + 5i(5i)}{9 - 25i^2}$$

$$= \frac{-6 - 10i + 15i + 25i^2}{9 - 25(-1)} \quad [\because i^2 = -1]$$

$$= \frac{-6 + 5i + 25(-1)}{9 + 25}$$

$$= \frac{-31 + 5i}{34}$$

$$= -\frac{31}{34} + \frac{5}{34}i$$

$\underbrace{\hspace{1.5cm}}$ $\underbrace{\hspace{1.5cm}}$
 Real part Imaginary part

Q. 3. E. Express each of the following in the form (a + ib):

$$\frac{(3 - 4i)}{(4 - 2i)(1 + i)}$$

Solution: Given: $\frac{3-4i}{(4-2i)(1+i)}$

Solving the denominator, we get

$$\frac{3 - 4i}{(4 - 2i)(1 + i)} = \frac{3 - 4i}{4(1) + 4(i) - 2i(1) - 2i(i)}$$

$$= \frac{3 - 4i}{4 + 4i - 2i - 2i^2}$$

$$= \frac{3 - 4i}{4 + 2i - 2(-1)}$$

$$= \frac{3 - 4i}{6 + 2i}$$

Now, we rationalize the above by multiplying and divide by the conjugate of $6 + 2i$

$$= \frac{3 - 4i}{6 + 2i} \times \frac{6 - 2i}{6 - 2i}$$

$$= \frac{(3-4i)(6-2i)}{(6+2i)(6-2i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{(3 - 4i)(6 - 2i)}{(6)^2 - (2i)^2}$$

$$= \frac{3(6) + 3(-2i) + (-4i)(6) + (-4i)(-2i)}{36 - 4i^2}$$

$$= \frac{18 - 6i - 24i + 8i^2}{36 - 4(-1)} \quad [\because i^2 = -1]$$

$$= \frac{18 - 30i + 8(-1)}{36 + 4}$$

$$= \frac{18 - 30i - 8}{40}$$

$$= \frac{10 - 30i}{40}$$

$$= \frac{10(1 - 3i)}{40}$$

$$= \frac{1 - 3i}{4}$$

$$= \frac{1}{4} - \frac{3}{4}i$$

$$\underbrace{\quad}_{\text{Real part}} \quad \underbrace{\quad}_{\text{Imaginary part}}$$

Q. 3. F. Express each of the following in the form $(a + ib)$:

$$\frac{(3 - 2i)(2 + 3i)}{(1 + 2i)(2 - i)}$$

$$\text{Solution: Given: } \frac{(3 - 2i)(2 + 3i)}{(1 + 2i)(2 - i)}$$

Firstly, we solve the given equation

$$= \frac{3(2) + 3(3i) - 2i(2) + (-2i)(3i)}{(1)(2) + 1(-i) + 2i(2) + 2i(-i)}$$

$$= \frac{6 + 9i - 4i - 6i^2}{2 - i + 4i - 2i^2}$$

$$= \frac{6 + 5i - 6(-1)}{2 + 3i - 2(-1)}$$

$$= \frac{6 + 6 + 5i}{2 + 3i + 2}$$

$$= \frac{12 + 5i}{4 + 3i}$$

Now, we rationalize the above by multiplying and divide by the conjugate of $4 + 3i$

$$= \frac{12 + 5i}{4 + 3i} \times \frac{4 - 3i}{4 - 3i}$$

$$= \frac{(12+5i)(4-3i)}{(4+3i)(4-3i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{(12 + 5i)(4 - 3i)}{(4)^2 - (3i)^2}$$

$$= \frac{12(4) + 12(-3i) + 5i(4) + 5i(-3i)}{16 - 9i^2}$$

$$= \frac{48 - 36i + 20i - 15i^2}{16 - 9(-1)} \quad [\because i^2 = -1]$$

$$= \frac{48 - 16i - 15(-1)}{16 + 9} \quad [\because i^2 = -1]$$

$$= \frac{48 - 16i + 15}{25}$$

$$= \frac{63 - 16i}{25}$$

$$= \frac{63}{25} - \frac{16}{25}i$$

$\underbrace{\hspace{1.5cm}}$ $\underbrace{\hspace{1.5cm}}$
 Real part Imaginary part

Q. 3. G. Express each of the following in the form $(a + ib)$:

$$\frac{(2 + 3i)^2}{(2 - i)}$$

Solution: Given: $\frac{(2+3i)^2}{(2-i)}$

Now, we rationalize the above equation by multiply and divide by the conjugate of $(2 - i)$

$$= \frac{(2 + 3i)^2}{(2 - i)} \times \frac{(2 + i)}{(2 + i)}$$

$$= \frac{(2 + 3i)^2(2 + i)}{(2 - i)(2 + i)}$$

$$= \frac{(4 + 9i^2 + 12i)(2 + i)}{(2)^2 - (i)^2}$$

$$[\because (a + b)(a - b) = (a^2 - b^2)]$$

$$= \frac{[4 + 9(-1) + 12i](2 + i)}{4 - i^2} \quad [\because i^2 = -1]$$

$$= \frac{[4 - 9 + 12i](2 + i)}{4 - (-1)}$$

$$= \frac{(-5 + 12i)(2 + i)}{5}$$

$$= \frac{-10 - 5i + 24i + 12i^2}{5}$$

$$= \frac{-10 + 19i + 12(-1)}{5}$$

$$= \frac{-10 - 12 + 19i}{5}$$

$$= \frac{-22 + 19i}{5}$$

$$= -\frac{22}{5} + \frac{19}{5}i$$

$$\underbrace{\quad}_{\text{Real part}} \quad \underbrace{\quad}_{\text{Imaginary part}}$$

Q. 3. H. Express each of the following in the form $(a + ib)$:

$$\frac{(1-i)^3}{(1-i^3)}$$

Solution: Given: $\frac{(1-i)^3}{(1-i^3)}$

The above equation can be re-written as

$$= \frac{(1)^3 - (i)^3 - 3(1)^2(i) + 3(1)(i)^2}{(1 - i \times i^2)}$$

$$[\because (a - b)^3 = a^3 - b^3 - 3a^2b + 3ab^2]$$

$$= \frac{1 - i^3 - 3i + 3i^2}{[1 - i(-1)]} \quad [\because i^2 = -1]$$

$$= \frac{1 - i \times i^2 - 3i + 3(-1)}{(1 + i)}$$

$$= \frac{1 - i(-1) - 3i - 3}{1 + i}$$

$$= \frac{-2 + i - 3i}{1 + i}$$

$$= \frac{-2 - 2i}{1 + i}$$

$$= \frac{-2(1 + i)}{1 + i}$$

$$= -2 + 0i$$

$$\underbrace{\quad}_{\text{Real part}} \quad \underbrace{\quad}_{\text{Imaginary part}}$$

Q. 3. I. Express each of the following in the form (a + ib):

$$\frac{(1 + 2i)^3}{(1 + i)(2 - i)}$$

Solution: Given: $\frac{(1+2i)^3}{(1+i)(2-i)}$

We solve the above equation by using the formula

$$(a + b)^3 = a^3 + b^3 + 3a^2b + 3ab^2$$

$$= \frac{(1)^3 + (2i)^3 + 3(1)^2(2i) + 3(1)(2i)^2}{1(2) + 1(-i) + i(2) + i(-i)}$$

$$= \frac{1 + 8i^3 + 6i + 12i^2}{2 - i + 2i - i^2}$$

$$= \frac{1 + 8i \times i^2 + 6i + 12(-1)}{2 + i - (-1)} \quad [\because i^2 = -1]$$

$$= \frac{1 + 8i(-1) + 6i - 12}{2 + i + 1}$$

$$= \frac{1 - 8i + 6i - 12}{3 + i}$$

$$= \frac{-11 - 2i}{3 + i}$$

Now, we rationalize the above by multiplying and divide by the conjugate of $3 + i$

$$= \frac{-11 - 2i}{3 + i} \times \frac{3 - i}{3 - i}$$

$$= \frac{(-11-2i)(3-i)}{(3+i)(3-i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{(-11 - 2i)(3 - i)}{(3)^2 - (i)^2}$$

$$= \frac{-11(3) + (-11)(-i) + (-2i)(3) + (-2i)(-i)}{9 - i^2}$$

$$= \frac{-33+11i-6i+2i^2}{9-(-1)} \quad [\because i^2 = -1]$$

$$= \frac{-33+5i+2(-1)}{9+1} \quad [\because i^2 = -1]$$

$$= \frac{-33 + 5i - 2}{10}$$

$$= \frac{-35 + 5i}{10}$$

$$= \frac{5(-7 + i)}{10}$$

$$= \frac{-7 + i}{2}$$

$$= \underbrace{\frac{-7}{2}}_{\text{Real part}} + \underbrace{\frac{1}{2}i}_{\text{Imaginary part}}$$

Q. 4. Simplify each of the following and express it in the form (a + ib):

$$(i) \left(\frac{5}{-3+2i} + \frac{2}{1-i} \right) \left(\frac{4-5i}{3+2i} \right)$$

$$(ii) \left(\frac{1}{1+4i} - \frac{2}{1+i} \right) \left(\frac{1-i}{5+3i} \right)$$

Solution: Given:

$$\left(\frac{5}{-3+2i} + \frac{2}{1-i} \right) \left(\frac{4-5i}{3+2i} \right)$$

$$= \left[\frac{5(1-i)+2(-3+2i)}{(-3+2i)(1-i)} \right] \left(\frac{4-5i}{3+2i} \right) \quad [\text{Taking the LCM}]$$

$$= \left[\frac{5-5i-6+4i}{(-3)(1-i)+2i(1-i)} \right] \left(\frac{4-5i}{3+2i} \right)$$

$$= \left[\frac{-1-i}{-3+3i+2i-2i^2} \right] \left(\frac{4-5i}{3+2i} \right)$$

$$= \left[\frac{-(1+i)}{-3+5i-2(-1)} \right] \left(\frac{4-5i}{3+2i} \right)$$

$$= \left(\frac{-(1+i)}{-1+5i} \right) \left(\frac{4-5i}{3+2i} \right)$$

$$= \frac{-1(4-5i) - i(4-5i)}{-1(3+2i) + 5i(3+2i)}$$

$$= \frac{-4+5i-4i+5i^2}{-3-2i+15i+10i^2}$$

$$= \frac{-4+i+5(-1)}{-3+13i+10(-1)} \text{ [Putting } i^2 = -1]$$

$$= \frac{-9+i}{-13+13i}$$

$$= \frac{-(9-i)}{-(13-13i)}$$

$$= \frac{9-i}{13-13i}$$

Now, rationalizing by multiply and divide by the conjugate of $(13 - 13i)$

$$= \frac{9-i}{13-13i} \times \frac{13+13i}{13+13i}$$

$$= \frac{(9-i)(13+13i)}{(13-13i)(13+13i)}$$

$$= \frac{117+117i-13i-13i^2}{(13)^2-(13i)^2} \text{ [}\because (a-b)(a+b) = (a^2 - b^2)\text{]}$$

$$= \frac{117+104i-13(-1)}{169-169i^2} \text{ [}\because i^2 = -1\text{]}$$

$$= \frac{130+104i}{169(1-i^2)}$$

$$= \frac{13(10+8i)}{169[1-(-1)]} \text{ [Taking 13 common]}$$

$$= \frac{10+8i}{13 \times 2}$$

$$= \frac{5+4i}{13}$$

$$= \frac{5}{13} + \frac{4}{13}i$$

(ii) Given:

$$\begin{aligned}
 & \left(\frac{1}{1+4i} - \frac{2}{1+i} \right) \left(\frac{1-i}{5+3i} \right) \\
 &= \left[\frac{1(1+i) - 2(1+4i)}{(1+4i)(1+i)} \right] \left(\frac{1-i}{5+3i} \right) \quad [\text{Taking the LCM}] \\
 &= \left[\frac{1+i-2-8i}{(1)(1+i)+4i(1+i)} \right] \left(\frac{1-i}{5+3i} \right) \\
 &= \left[\frac{-1-7i}{1+i+4i+4i^2} \right] \left(\frac{1-i}{5+3i} \right) \\
 &= \left[\frac{-1-7i}{1+5i+4(-1)} \right] \left(\frac{1-i}{5+3i} \right) \\
 &= \left(\frac{-1-7i}{-3+5i} \right) \left(\frac{1-i}{5+3i} \right) \\
 &= \frac{-1(1-i) - 7i(1-i)}{-3(5+3i) + 5i(5+3i)} \\
 &= \frac{-1+i-7i+7i^2}{-15-9i+25i+15i^2} \\
 &= \frac{-1-6i+7(-1)}{-15+16i+15(-1)} \\
 &= \frac{-6i-8}{16i-30} \\
 &= \frac{-2(4+3i)}{-2(15-8i)} \\
 &= \frac{4+3i}{15-8i}
 \end{aligned}$$

Now, rationalizing by multiply and divide by the conjugate of $(15 + 8i)$

$$\begin{aligned}
 &= \frac{4 + 3i}{15 - 8i} \times \frac{15 + 8i}{15 + 8i} \\
 &= \frac{(4+3i)(15+8i)}{(15)^2 - (8i)^2} \quad [\because (a-b)(a+b) = (a^2 - b^2)] \\
 &= \frac{4(15 + 8i) + 3i(15 + 8i)}{225 - 64i^2} \\
 &= \frac{60 + 32i + 45i + 24i^2}{225 - 64(-1)} \quad [\because i^2 = -1] \\
 &= \frac{60 + 77i + 24(-1)}{225 + 64} \\
 &= \frac{36 + 77i}{289} \\
 &= \frac{36}{289} + \frac{77}{289}i
 \end{aligned}$$

Q. 5. Show that

- (i) $\left\{ \frac{(3+2i)}{(2-3i)} + \frac{(3-2i)}{(2+3i)} \right\}$ is purely real,
- (ii) $\left\{ \frac{(\sqrt{7} + i\sqrt{3})}{(\sqrt{7} - i\sqrt{3})} + \frac{(\sqrt{7} - i\sqrt{3})}{(\sqrt{7} + i\sqrt{3})} \right\}$ is purely real.

Solution: Given: $\frac{3+2i}{2-3i} + \frac{3-2i}{2+3i}$

Taking the L.C.M, we get

$$= \frac{(3 + 2i)(2 + 3i) + (3 - 2i)(2 - 3i)}{(2 - 3i)(2 + 3i)}$$

$$= \frac{3(2) + 3(3i) + 2i(2) + 2i(3i) + 3(2) + 3(-3i) - 2i(2) + (-2i)(-3i)}{(2)^2 - (3i)^2}$$

$$[\because (a + b)(a - b) = (a^2 - b^2)]$$

$$= \frac{6 + 9i + 4i + 6i^2 + 6 - 9i - 4i + 6i^2}{4 - 9i^2}$$

$$= \frac{12 + 12i^2}{4 - 9i^2}$$

Putting $i^2 = -1$

$$= \frac{12 + 12(-1)}{4 - 9(-1)}$$

$$= \frac{12 - 12}{4 + 9}$$

$$= 0 + 0i$$

Hence, the given equation is purely real as there is no imaginary part.

(ii) Given: $\frac{\sqrt{7}+i\sqrt{3}}{\sqrt{7}-i\sqrt{3}} + \frac{\sqrt{7}-i\sqrt{3}}{\sqrt{7}+i\sqrt{3}}$

Taking the L.C.M, we get

$$= \frac{(\sqrt{7} + i\sqrt{3})(\sqrt{7} + i\sqrt{3}) + (\sqrt{7} - i\sqrt{3})(\sqrt{7} - i\sqrt{3})}{(\sqrt{7} - i\sqrt{3})(\sqrt{7} + i\sqrt{3})}$$

$$= \frac{(\sqrt{7}+i\sqrt{3})^2 + (\sqrt{7}-i\sqrt{3})^2}{(\sqrt{7})^2 - (i\sqrt{3})^2} \dots(i)$$

$$[\because (a + b)(a - b) = (a^2 - b^2)]$$

Now, we know that,

$$(a + b)^2 + (a - b)^2 = 2(a^2 + b^2)$$

So, by applying the formula in eq. (i), we get

$$= \frac{2[(\sqrt{7})^2 + (i\sqrt{3})^2]}{7 - 3i^2}$$

$$= \frac{2[7 + 3i^2]}{7 - 3(-1)}$$

Putting $i^2 = -1$

$$= \frac{2[7 + 3(-1)]}{7 + 3}$$

$$= \frac{2[7 - 3]}{10}$$

$$= \frac{8}{10} + 0i$$

$$= \frac{4}{5} + 0i$$

Hence, the given equation is purely real as there is no imaginary part.

Q. 6. Find the real values of θ for which $\frac{1+i \cos \theta}{1-2i \cos \theta}$ is purely real.

Solution: Since $\frac{1+i \cos \theta}{1-2i \cos \theta}$ is purely real

Firstly, we need to solve the given equation and then take the imaginary part as 0

$$\frac{1+i \cos \theta}{1-2i \cos \theta}$$

We rationalize the above by multiply and divide by the conjugate of $(1 - 2i \cos \theta)$

$$= \frac{1+i \cos \theta}{1-2i \cos \theta} \times \frac{1+2i \cos \theta}{1+2i \cos \theta}$$

$$= \frac{(1+i \cos \theta)(1+2i \cos \theta)}{(1-2i \cos \theta)(1+2i \cos \theta)}$$

We know that,

$$(a - b)(a + b) = (a^2 - b^2)$$

$$= \frac{1(1) + 1(2i \cos \theta) + i \cos \theta(1) + i \cos \theta(2i \cos \theta)}{(1)^2 - (2i \cos \theta)^2}$$

$$= \frac{1 + 2i \cos \theta + i \cos \theta + 2i^2 \cos^2 \theta}{1 - 4i^2 \cos^2 \theta}$$

$$= \frac{1 + 3i \cos \theta + 2(-1) \cos^2 \theta}{1 - 4(-1) \cos^2 \theta} \quad [\because i^2 = -1]$$

$$= \frac{1 + 3i \cos \theta - 2 \cos^2 \theta}{1 + 4 \cos^2 \theta}$$

$$= \frac{1 - 2 \cos^2 \theta}{1 + 4 \cos^2 \theta} + i \frac{3 \cos \theta}{1 + 4 \cos^2 \theta}$$

Since $\frac{1+i \cos \theta}{1-2i \cos \theta}$ is purely real [given]

Hence, imaginary part is equal to 0

$$\text{i.e. } \frac{3 \cos \theta}{1 + 4 \cos^2 \theta} = 0$$

$$\Rightarrow 3 \cos \theta = 0 \times (1 + 4 \cos^2 \theta)$$

$$\Rightarrow 3 \cos \theta = 0$$

$$\Rightarrow \cos \theta = 0$$

$$\Rightarrow \cos \theta = \cos 0$$

Since, $\cos \theta = \cos y$

Then $\theta = (2n + 1) \frac{\pi}{2} \pm y$ where $n \in \mathbb{Z}$

Putting $y = 0$

$$\theta = (2n + 1)\frac{\pi}{2} \pm 0$$

$$\theta = (2n + 1)\frac{\pi}{2} \text{ where } n \in \mathbb{Z}$$

Hence, for $\theta = (2n + 1)\frac{\pi}{2}$, where $n \in \mathbb{Z}$ $\frac{1+i\cos\theta}{1-2i\cos\theta}$ is purely real.

Q. 7. If $|z + i| = |z - i|$, prove that z is real.

Solution: Let $z = x + iy$

Consider, $|z + i| = |z - i|$

$$\Rightarrow |x + iy + i| = |x + iy - i|$$

$$\Rightarrow |x + i(y + 1)| = |x + i(y - 1)|$$

$$\Rightarrow \sqrt{(x)^2 + (y + 1)^2} = \sqrt{(x)^2 + (y - 1)^2}$$

$$[\because |z| = \text{modulus} = \sqrt{a^2 + b^2}]$$

$$\Rightarrow \sqrt{x^2 + y^2 + 1 + 2y} = \sqrt{x^2 + y^2 + 1 - 2y}$$

Squaring both the sides, we get

$$\Rightarrow x^2 + y^2 + 1 + 2y = x^2 + y^2 + 1 - 2y$$

$$\Rightarrow x^2 + y^2 + 1 + 2y - x^2 - y^2 - 1 + 2y = 0$$

$$\Rightarrow 2y + 2y = 0$$

$$\Rightarrow 4y = 0$$

$$\Rightarrow y = 0$$

Putting the value of y in eq. (i), we get

$$z = x + i(0)$$

$$\Rightarrow z = x$$

Hence, z is purely real.

Q. 8. Give an example of two complex numbers z_1 and z_2 such that $z_1 \neq z_2$ and $|z_1| = |z_2|$.

Solution: Let $z_1 = 3 - 4i$ and $z_2 = 4 -$

$3i$ Here, $z_1 \neq z_2$

Now, calculating the modulus, we get,

$$|z_1| = \sqrt{3^2 + (4)^2} = \sqrt{25} = 5$$

$$|z_2| = \sqrt{4^2 + (3)^2} = \sqrt{25} = 5$$

Q. 9. A. Find the conjugate of each of the following:

(-5 - 2i)

Solution: Given: $z = (-5 - 2i)$

Here, we have to find the conjugate of $(-5 - 2i)$

So, the conjugate of $(-5 - 2i)$ is $(-5 + 2i)$

Q. 9. B. Find the conjugate of each of the following:

$$\frac{1}{(4 + 3i)}$$

Solution: Given: $\frac{1}{4+3i}$

First, we calculate $\frac{1}{4+3i}$ and then find its conjugate

Now, rationalizing

$$= \frac{1}{4 + 3i} \times \frac{4 - 3i}{4 - 3i}$$

$$= \frac{4 - 3i}{(4 + 3i)(4 - 3i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{4 - 3i}{(4)^2 - (3i)^2}$$

$$= \frac{4 - 3i}{16 - 9i^2}$$

$$= \frac{4 - 3i}{16 - 9(-1)} [\because i^2 = -1]$$

$$= \frac{4 - 3i}{16 + 9}$$

$$= \frac{4 - 3i}{25}$$

$$= \frac{4}{25} - \frac{3}{25}i$$

Hence, $\frac{1}{4+3i} = \frac{4}{25} - \frac{3}{25}i$

So, a conjugate of $\frac{1}{4+3i}$ is $\frac{4}{25} + \frac{3}{25}i$

Q. 9. C. Find the conjugate of each of the following:

$$\frac{(1+i)^2}{(3-i)}$$

Solution: Given: $\frac{(1+i)^2}{(3-i)}$

Firstly, we calculate $\frac{(1+i)^2}{(3-i)}$ and then find its conjugate

$$\frac{(1+i)^2}{(3-i)} = \frac{1+i^2+2i}{(3-i)} \quad [\because (a + b)^2 = a^2 + b^2 + 2ab]$$

$$= \frac{1+(-1)+2i}{3-i} \quad [\because i^2 = -1]$$

$$= \frac{2i}{3-i}$$

Now, we rationalize the above by multiplying and divide by the conjugate of $3 - i$

$$= \frac{2i}{3-i} \times \frac{3+i}{3+i}$$

$$= \frac{(2i)(3+i)}{(3+i)(3-i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{(2i)(3+i)}{(3)^2 - (i)^2}$$

$$= \frac{2i(3) + 2i(i)}{9 - i^2}$$

$$= \frac{6i+2i^2}{9-(-1)} \quad [\because i^2 = -1]$$

$$= \frac{6i+2(-1)}{9+1} \quad [\because i^2 = -1]$$

$$= \frac{6i - 2}{10}$$

$$= \frac{2(3i - 1)}{10}$$

$$= \frac{(-1 + 3i)}{5}$$

$$= -\frac{1}{5} + \frac{3}{5}i$$

Hence, $\frac{(1+i)^2}{(3-i)} = -\frac{1}{5} + \frac{3}{5}i$

So, the conjugate of $\frac{(1+i)^2}{(3-i)}$ is $-\frac{1}{5} - \frac{3}{5}i$

Q. 9. D. Find the conjugate of each of the following:

$$\frac{(1+i)(2+i)}{(3+i)}$$

Solution: Given: $\frac{(1+i)(2+i)}{(3+i)}$

Firstly, we calculate $\frac{(1+i)(2+i)}{(3+i)}$ and then find its conjugate

$$\frac{(1+i)(2+i)}{(3+i)} = \frac{1(2) + 1(i) + i(2) + i(i)}{(3+i)}$$

$$= \frac{2 + i + 2i + i^2}{3+i}$$

$$= \frac{2+3i-1}{3+i} \quad [\because i^2 = -1]$$

$$= \frac{1+3i}{3+i}$$

Now, we rationalize the above by multiplying and divide by the conjugate of $3+i$

$$= \frac{1+3i}{3+i} \times \frac{3-i}{3-i}$$

$$= \frac{(1+3i)(3-i)}{(3+i)(3-i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{(1 + 3i)(3 - i)}{(3)^2 - (i)^2}$$

$$= \frac{1(3) + 1(-i) + 3i(3) + 3i(-i)}{9 - i^2}$$

$$= \frac{3 - i + 9i - 3i^2}{9 - (-1)} \quad [\because i^2 = -1]$$

$$= \frac{3 + 8i - 3(-1)}{9 + 1} \quad [\because i^2 = -1]$$

$$= \frac{3 + 8i + 3}{10}$$

$$= \frac{6 + 8i}{10}$$

$$= \frac{2(3 + 4i)}{10}$$

$$= \frac{3 + 4i}{5}$$

$$= \frac{3}{5} + \frac{4}{5}i$$

Hence, $\frac{(1+i)(2+i)}{(3+i)} = \frac{3}{5} + \frac{4}{5}i$

So, the conjugate of $\frac{(1+i)^2}{(3-i)}$ is $\frac{3}{5} - \frac{4}{5}i$

Q. 9. E. Find the conjugate of each of the following:

$$\sqrt{-3}$$

Solution: Given: $z = \sqrt{-3}$

The above can be re – written as

$$z = \sqrt{(-1) \times 3}$$

$$z = \sqrt{3i^2} \quad [\because i^2 = -1]$$

$$z = 0 + i\sqrt{3}$$

So, the conjugate of $z = 0 + i\sqrt{3}$ is

$$\bar{z} = 0 - i\sqrt{3}$$

$$\text{Or } \bar{z} = -i\sqrt{3} = -\sqrt{-3}$$

Q. 9. F. Find the conjugate of each of the following:

$$\sqrt{2}$$

Solution: Given: $z = \sqrt{2}$

The above can be re – written as

$$z = \sqrt{2} + 0i$$

Here, the imaginary part is zero

So, the conjugate of $z = \sqrt{2} + 0i$ is

$$\bar{z} = \sqrt{2} - 0i$$

$$\text{Or } \bar{z} = \sqrt{2}$$

Q. 9. G. Find the conjugate of each of the following:

$$-\sqrt{-1}$$

Solution: Given: $z = -\sqrt{-1}$

The above can be re – written as

$$z = -\sqrt{i^2} \quad [\because i^2 = -1]$$

$$z = 0 - i$$

So, the conjugate of $z = (0 - i)$ is

$$\bar{z} = 0 + i$$

Or $\bar{z} = i$

Q. 9. H. Find the conjugate of each of the following:

$$(2 - 5i)^2$$

Solution: Given: $z = (2 - 5i)^2$

First we calculate $(2 - 5i)^2$ and then we find the conjugate

$$(2 - 5i)^2 = (2)^2 + (5i)^2 - 2(2)(5i)$$

$$= 4 + 25i^2 - 20i$$

$$= 4 + 25(-1) - 20i \quad [\because i^2 = -1]$$

$$= 4 - 25 - 20i$$

$$= -21 - 20i$$

Now, we have to find the conjugate of $(-21 - 20i)$

So, the conjugate of $(-21 - 20i)$ is $(-21 + 20i)$

Q. 10. A. Find the modulus of each of the following:

$$(3 + \sqrt{-5})$$

Solution: Given: $z = (3 + \sqrt{-5})$

The above can be re-written as

$$z = 3 + \sqrt{(-1) \times 5}$$

$$z = 3 + i\sqrt{5} \quad [\because i^2 = -1]$$

Now, we have to find the modulus of $(3 + i\sqrt{5})$

$$\text{So, } |z| = |3 + i\sqrt{5}| = \sqrt{(3)^2 + (\sqrt{5})^2} = \sqrt{9 + 5} = \sqrt{14}$$

Hence, the modulus of $(3 + i\sqrt{5})$ is $\sqrt{14}$

Q. 10. B. Find the modulus of each of the following:

$(-3 - 4i)$

Solution: Given: $z = (-3 - 4i)$

Now, we have to find the modulus of $(-3 - 4i)$

$$\text{So, } |z| = |-3 - 4i| = \sqrt{(-3)^2 + (-4)^2} = \sqrt{9 + 16} = \sqrt{25} = 5$$

Hence, the modulus of $(-3 - 4i)$ is 5

Q. 10. C. Find the modulus of each of the following:

$(7 + 24i)$

Solution: Given: $z = (7 + 24i)$

Now, we have to find the modulus of $(7 + 24i)$

$$\text{So, } |z| = |7 + 24i| = \sqrt{(7)^2 + (24)^2} = \sqrt{49 + 576} = \sqrt{625} = 25$$

Hence, the modulus of $(7 + 24i)$ is 25

Q. 10. D. Find the modulus of each of the following:

$3i$

Solution: Given: $z = 3i$

The above equation can be re – written as

$$z = 0 + 3i$$

Now, we have to find the modulus of $(0 + 3i)$

$$\text{So, } |z| = |0 + 3i| = \sqrt{(0)^2 + (3)^2} = \sqrt{9} = 3$$

Hence, the modulus of $(3i)$ is 3

Q. 10. E. Find the modulus of each of the following:

$$\frac{(3 + 2i)^2}{(4 - 3i)}$$

Solution: Given: $\frac{(3+2i)^2}{(4-3i)}$

Firstly, we calculate $\frac{(3+2i)^2}{(4-3i)}$ and then find its modulus

$$\begin{aligned} \frac{(3+2i)^2}{(4-3i)} &= \frac{9+4i^2+12i}{(4-3i)} \quad [\because (a+b)^2 = a^2 + b^2 + 2ab] \\ &= \frac{9+4(-1)+12i}{4-3i} \quad [\because i^2 = -1] \end{aligned}$$

$$= \frac{5 + 12i}{4 - 3i}$$

Now, we rationalize the above by multiplying and divide by the conjugate of $4 + 3i$

$$\begin{aligned} &= \frac{5 + 12i}{4 - 3i} \times \frac{4 + 3i}{4 + 3i} \\ &= \frac{(5+12i)(4+3i)}{(4-3i)(4+3i)} \dots(i) \end{aligned}$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{5(4) + (5)(3i) + 12i(4) + 12i(3i)}{(4)^2 - (3i)^2}$$

$$= \frac{20 + 15i + 48i + 36i^2}{16 - 9i^2}$$

$$= \frac{20+63i+36(-1)}{16-9(-1)} \quad [\because i^2 = -1]$$

$$= \frac{20-36+63i}{16+9} \quad [\because i^2 = -1]$$

$$= \frac{-16 + 63i}{25}$$

$$= -\frac{16}{25} + \frac{63}{25}i$$

Now, we have to find the modulus of $\left(-\frac{16}{25} + \frac{63}{25}i\right)$

$$\text{So, } |z| = \left| -\frac{16}{25} + \frac{63}{25}i \right| = \sqrt{\left(-\frac{16}{25}\right)^2 + \left(\frac{63}{25}\right)^2}$$

$$= \sqrt{\frac{256}{625} + \frac{3969}{625}}$$

$$= \sqrt{\frac{4225}{625}}$$

$$= \frac{65}{25}$$

$$= \frac{13}{5}$$

Hence, the modulus of $\frac{(3+2i)^2}{(4-3i)}$ is $\frac{13}{5}$

Q. 10. F. Find the modulus of each of the following:

$$\frac{(2-i)(1+i)}{(1+i)}$$

Solution: Given: $\frac{(2-i)(1+i)}{(1+i)}$

Firstly, we calculate $\frac{(2-i)(1+i)}{(1+i)}$ and then find its modulus

$$\frac{(2-i)(1+i)}{(1+i)} = \frac{2(1) + 2(i) + (-i)(1) + (-i)(i)}{(1+i)}$$

$$= \frac{2 + 2i - i - i^2}{1+i}$$

$$= \frac{2+i-(-1)}{1+i} \quad [\because i^2 = -1]$$

$$= \frac{3+i}{1+i}$$

Now, we rationalize the above by multiplying and divide by the conjugate of $1+i$

$$= \frac{3+i}{1+i} \times \frac{1-i}{1-i}$$

$$= \frac{(3+i)(1-i)}{(1+i)(1-i)} \dots (i)$$

Now, we know that,

$$(a+b)(a-b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{3(1-i) + i(1-i)}{(1)^2 - (i)^2}$$

$$= \frac{3(1) + 3(-i) + i(1) + i(-i)}{1 - i^2}$$

$$= \frac{3 - 3i + i - i^2}{1 - (-1)} \quad [\because i^2 = -1]$$

$$= \frac{3 - 2i - (-1)}{1 + 1} \quad [\because i^2 = -1]$$

$$= \frac{3 - 2i + 1}{2}$$

$$= \frac{4 - 2i}{2}$$

$$= 2 - i$$

Now, we have to find the modulus of $(2 - i)$

$$\text{So, } |z| = |2 - i| = |2 + (-1)i| = \sqrt{(2)^2 + (-1)^2} = \sqrt{4 + 1} = \sqrt{5}$$

Q. 10. G. Find the modulus of each of the following:

5

Solution: Given: $z = 5$

The above equation can be re – written as

$$z = 5 + 0i$$

Now, we have to find the modulus of $(5 + 0i)$

$$\text{So, } |z| = |5 + 0i| = \sqrt{(5)^2 + (0)^2} = 5$$

Q. 10. H. Find the modulus of each of the following:

$(1 + 2i)(i - 1)$

Solution: Given: $z = (1 + 2i)(i - 1)$

Firstly, we calculate the $(1 + 2i)(i - 1)$ and then find the modulus

So, we open the brackets,

$$\begin{aligned} & 1(i - 1) + 2i(i - 1) \\ &= 1(i) + (1)(-1) + 2i(i) + 2i(-1) \\ &= i - 1 + 2i^2 - 2i \\ &= -i - 1 + 2(-1) [\because i^2 = -1] \\ &= -i - 1 - 2 \\ &= -i - 3 \end{aligned}$$

Now, we have to find the modulus of $(-3 - i)$

$$\text{So, } |z| = |-3 - i| = |-3 + (-1)i| = \sqrt{(-3)^2 + (-1)^2} = \sqrt{9 + 1} = \sqrt{10}$$

Q. 11. A. Find the multiplicative inverse of each of the following:

$$(1 - \sqrt{3}i)$$

Solution: Given: $(1 - i\sqrt{3})$

To find: Multiplicative inverse

We know that,

Multiplicative Inverse of $z = z^{-1}$

$$= \frac{1}{z}$$

Putting $z = 1 - i\sqrt{3}$

$$\text{So, Multiplicative inverse of } 1 - i\sqrt{3} = \frac{1}{1 - i\sqrt{3}}$$

Now, rationalizing by multiply and divide by the conjugate of $(1 - i\sqrt{3})$

$$= \frac{1}{1 - i\sqrt{3}} \times \frac{1 + i\sqrt{3}}{1 + i\sqrt{3}}$$

$$= \frac{1 + i\sqrt{3}}{(1 - i\sqrt{3})(1 + i\sqrt{3})}$$

Using $(a - b)(a + b) = (a^2 - b^2)$

$$= \frac{1 + i\sqrt{3}}{(1)^2 - (i\sqrt{3})^2}$$

$$= \frac{1 + i\sqrt{3}}{1 - 3i^2}$$

$$= \frac{1 + i\sqrt{3}}{1 - 3(-1)} \quad [\because i^2 = -1]$$

$$= \frac{1 + i\sqrt{3}}{1 + 3}$$

$$= \frac{1 + i\sqrt{3}}{4}$$

$$= \frac{1}{4} + \frac{\sqrt{3}}{4}i$$

Hence, Multiplicative Inverse of $(1 - i\sqrt{3})$ is $\frac{1}{4} + \frac{\sqrt{3}}{4}i$

Q. 11. B. Find the multiplicative inverse of each of the following:

(2 + 5i)

Solution: Given: $2 + 5i$

To find: Multiplicative inverse

We know that,

Multiplicative Inverse of $z = z^{-1}$

$$= \frac{1}{z}$$

Putting $z = 2 + 5i$

$$\text{So, Multiplicative inverse of } 2 + 5i = \frac{1}{2 + 5i}$$

Now, rationalizing by multiply and divide by the conjugate of $(2+5i)$

$$= \frac{1}{2 + 5i} \times \frac{2 - 5i}{2 - 5i}$$

$$= \frac{2 - 5i}{(2 + 5i)(2 - 5i)}$$

Using $(a - b)(a + b) = (a^2 - b^2)$

$$= \frac{2 - 5i}{(2)^2 - (5i)^2}$$

$$= \frac{2 - 5i}{4 - 25i^2}$$

$$= \frac{2 - 5i}{4 - 25(-1)} \quad [\because i^2 = -1]$$

$$= \frac{2 - 5i}{4 + 25}$$

$$= \frac{2 - 5i}{29}$$

$$= \frac{2}{29} - \frac{5}{29}i$$

Hence, Multiplicative Inverse of $(2+5i)$ is $\frac{2}{29} - \frac{5}{29}i$

Q. 11. C. Find the multiplicative inverse of each of the following:

$$\frac{(2 + 3i)}{(1 + i)}$$

Solution: Given: $\frac{2+3i}{1+i}$

To find: Multiplicative inverse

We know that,

Multiplicative Inverse of $z = z^{-1}$

$$= \frac{1}{z}$$

Putting $z = \frac{2+3i}{1+i}$

So, Multiplicative inverse of $\frac{2+3i}{1+i} = \frac{1}{\frac{2+3i}{1+i}} = \frac{1+i}{2+3i}$

Now, rationalizing by multiply and divide by the conjugate of $(2+3i)$

$$= \frac{1+i}{2+3i} \times \frac{2-3i}{2-3i}$$

$$= \frac{(1+i)(2-3i)}{(2+3i)(2-3i)}$$

Using $(a-b)(a+b) = (a^2 - b^2)$

$$= \frac{1(2-3i) + i(2-3i)}{(2)^2 - (3i)^2}$$

$$= \frac{2-3i+2i-3i^2}{4-9i^2}$$

$$= \frac{2-i-3(-1)}{4-9(-1)} \quad [\because i^2 = -1]$$

$$= \frac{5-i}{4+9}$$

$$= \frac{5 - i}{13}$$

$$= \frac{5}{13} - \frac{1}{13}i$$

Hence, Multiplicative Inverse of $\frac{(2 + 3i)}{1 + i}$ is $\frac{5}{13} - \frac{1}{13}i$

Q. 11. D. Find the multiplicative inverse of each of the following:

$$\frac{(1 + i)(1 + 2i)}{(1 + 3i)}$$

Solution: Given: $\frac{(1+i)(1+2i)}{(1+3i)}$

To find: Multiplicative inverse

We know that,

Multiplicative Inverse of $z = z^{-1}$

$$= \frac{1}{z}$$

Putting $z = \frac{(1 + i)(1 + 2i)}{(1 + 3i)}$

$$\text{So, Multiplicative inverse of } \frac{(1 + i)(1 + 2i)}{(1 + 3i)} = \frac{1}{\frac{(1 + i)(1 + 2i)}{(1 + 3i)}}$$

$$= \frac{(1 + 3i)}{(1 + i)(1 + 2i)}$$

We solve the above equation

$$= \frac{1 + 3i}{1(1) + 1(2i) + i(1) + i(2i)}$$

$$= \frac{1 + 3i}{1 + 2i + i + 2i^2}$$

$$= \frac{1+3i}{1+3i+2(-1)} \quad [\because i^2 = -1]$$

$$= \frac{1 + 3i}{-1 + 3i}$$

Now, we rationalize the above by multiplying and divide by the conjugate of $(-1 + 3i)$

$$= \frac{1 + 3i}{-1 + 3i} \times \frac{-1 - 3i}{-1 - 3i}$$

$$= \frac{(1+3i)(-1-3i)}{(-1+3i)(-1-3i)} \dots(i)$$

Now, we know that,

$$(a + b)(a - b) = (a^2 - b^2)$$

So, eq. (i) become

$$= \frac{1(-1 - 3i) + 3i(-1 - 3i)}{(-1)^2 - (3i)^2}$$

$$= \frac{-1 - 3i - 3i - 9i^2}{1 - 9i^2}$$

$$= \frac{-1 - 6i - 9(-1)}{1 - 9(-1)} \quad [\because i^2 = -1]$$

$$= \frac{-1 - 6i + 9}{1 + 9}$$

$$= \frac{8 - 6i}{10}$$

$$= \frac{2(4 - 3i)}{10}$$

$$= \frac{4 - 3i}{5}$$

$$= \frac{4}{5} - \frac{3}{5}i$$

Hence, Multiplicative inverse of $\frac{(1+i)(1+2i)}{(1+3i)} = \frac{4}{5} - \frac{3}{5}i$

Q. 12. If $\left(\frac{1-i}{1+i}\right)^{100} = (a + ib)$, find the values of a and b.

Solution: Given: $a + ib = \left(\frac{1-i}{1+i}\right)^{100}$

Consider the given equation,

$$a + ib = \left(\frac{1-i}{1+i}\right)^{100}$$

Now, we rationalize

$$= \left(\frac{1-i}{1+i} \times \frac{1-i}{1-i}\right)^{100}$$

[Here, we multiply and divide by the conjugate of $1 + i$]

$$= \left(\frac{(1-i)^2}{(1+i)(1-i)}\right)^{100}$$

$$= \left(\frac{1+i^2-2i}{(1+i)(1-i)}\right)^{100}$$

Using $(a + b)(a - b) = (a^2 - b^2)$

$$= \left(\frac{1 + (-1) - 2i}{(1)^2 - (i)^2}\right)^{100}$$

$$\begin{aligned}
 &= \left(\frac{-2i}{1-i^2} \right)^{100} \\
 &= \left(\frac{-2i}{1-(-1)} \right)^{100} \quad [\because i^2 = -1] \\
 &= \left(\frac{-2i}{2} \right)^{100} \\
 &= (-i)^{100} \\
 &= [(-i)^4]^{25} \\
 &= (i^4)^{25} \\
 &= (1)^{25} \\
 &[\because i^4 = i^2 \times i^2 = -1 \times -1 = 1]
 \end{aligned}$$

$$(a + ib) = 1 + 0i$$

On comparing both the sides, we get

$$a = 1 \text{ and } b = 0$$

Hence, the value of a is 1 and b is 0

Q. 13. If $\left(\frac{1+i}{1-i} \right)^{93} - \left(\frac{1-i}{1+i} \right)^3 = x + iy$, find x and y.

Solution: Consider,

$$x + iy = \left(\frac{1+i}{1-i} \right)^{93} - \left(\frac{1-i}{1+i} \right)^3$$

Now, rationalizing

$$x + iy = \left(\frac{1+i}{1-i} \times \frac{1+i}{1+i} \right)^{93} - \left(\frac{1-i}{1+i} \times \frac{1-i}{1-i} \right)^3$$

$$= \left(\frac{(1+i)^2}{(1-i)(1+i)} \right)^{93} - \left(\frac{(1-i)^2}{(1+i)(1-i)} \right)^3$$

In denominator, we use the identity

$$(a-b)(a+b) = a^2 - b^2$$

$$= \left(\frac{1+i^2+2i}{(1)^2-(i)^2} \right)^{93} - \left(\frac{1+i^2-2i}{(1)^2-(i)^2} \right)^3$$

$$= \left(\frac{1+(-1)+2i}{1-i^2} \right)^{93} - \left(\frac{1+(-1)-2i}{1-i^2} \right)^3$$

$$= \left(\frac{2i}{1-(-1)} \right)^{93} - \left(\frac{-2i}{1-(-1)} \right)^3$$

$$= \left(\frac{2i}{2} \right)^{93} - \left(\frac{-2i}{2} \right)^3$$

$$= (i)^{93} - (-i)^3$$

$$= (i)^{92+1} - [-(i)^3]$$

$$= [(i)^{92}(i)] - [-(i^2 \times i)]$$

$$= [(i^4)^{23}(i)] - [-(-i)]$$

$$= [(1)^{23}(i)] - i$$

$$= i - i$$

$$x + iy = 0$$

$$\therefore x = 0 \text{ and } y = 0$$

Q. 14. If $x + iy = \frac{a+ib}{a-ib}$, prove that $x^2 + y^2 = 1$.

Solution: Consider the given equation,

$$x + iy = \frac{a + ib}{a - ib}$$

Now, rationalizing

$$x + iy = \frac{a + ib}{a - ib} \times \frac{a + ib}{a + ib}$$

$$= \frac{(a + ib)(a + ib)}{(a - ib)(a + ib)}$$

$$= \frac{a(a + ib) + ib(a + ib)}{(a)^2 - (ib)^2}$$

$$[(a - b)(a + b) = a^2 - b^2]$$

$$= \frac{a^2 + iab + iab + i^2 b^2}{a^2 - i^2 b^2}$$

$$= \frac{a^2 + iab + iab + (-1)b^2}{a^2 - (-1)b^2} \quad [i^2 = -1]$$

$$x + iy = \frac{a^2 + 2iab - b^2}{a^2 + b^2}$$

$$x + iy = \frac{(a^2 - b^2)}{a^2 + b^2} + i \frac{2ab}{a^2 + b^2}$$

On comparing both the sides, we get

$$x = \frac{(a^2 - b^2)}{a^2 + b^2} \quad \& \quad y = \frac{2ab}{a^2 + b^2}$$

Now, we have to prove that $x^2 + y^2 = 1$

Taking LHS,

$$x^2 + y^2$$

Putting the value of x and y, we get

$$\begin{aligned} & \left[\frac{(a^2 - b^2)}{a^2 + b^2} \right]^2 + \left[\frac{2ab}{a^2 + b^2} \right]^2 \\ &= \frac{1}{(a^2 + b^2)^2} [(a^2 - b^2)^2 + (2ab)^2] \\ &= \frac{1}{(a^2 + b^2)^2} [a^4 + b^4 - 2a^2b^2 + 4a^2b^2] \\ &= \frac{1}{(a^2 + b^2)^2} [a^4 + b^4 + 2a^2b^2] \\ &= \frac{1}{(a^2 + b^2)^2} [(a^2 + b^2)^2] \\ &= 1 \\ &= \text{RHS} \end{aligned}$$

Q. 15. If $(a + ib) = \frac{c + i}{c - i}$, where c is real, prove that $a^2 + b^2 = 1$ and $\frac{b}{a} = \frac{2c}{c^2 - 1}$.

Solution: Consider the given equation,

$$a + ib = \frac{c + i}{c - i}$$

Now, rationalizing

$$a + ib = \frac{c + i}{c - i} \times \frac{c + i}{c + i}$$

$$= \frac{(c + i)(c + i)}{(c - i)(c + i)}$$

$$= \frac{(c + i)^2}{(c)^2 - (i)^2}$$

$$[(a - b)(a + b) = a^2 - b^2]$$

$$= \frac{c^2 + 2ic + i^2}{c^2 - i^2}$$

$$a + ib = \frac{c^2 + 2ic + (-1)}{c^2 - (-1)} \quad [i^2 = -1]$$

$$a + ib = \frac{c^2 + 2ic - 1}{c^2 + 1}$$

$$a + ib = \frac{(c^2 - 1)}{c^2 + 1} + i \frac{2c}{c^2 + 1}$$

On comparing both the sides, we get

$$a = \frac{(c^2 - 1)}{c^2 + 1} \quad \& \quad b = \frac{2c}{c^2 + 1}$$

Now, we have to prove that $a^2 + b^2 = 1$

Taking LHS,

$$a^2 + b^2$$

Putting the value of a and b, we get

$$\begin{aligned} & \left[\frac{(c^2 - 1)}{c^2 + 1} \right]^2 + \left[\frac{2c}{c^2 + 1} \right]^2 \\ &= \frac{1}{(c^2 + 1)^2} [(c^2 - 1)^2 + (2c)^2] \\ &= \frac{1}{(c^2 + 1)^2} [c^4 + 1 - 2c^2 + 4c^2] \\ &= \frac{1}{(c^2 + 1)^2} [c^4 + 1 + 2c^2] \\ &= \frac{1}{(c^2 + 1)^2} [(c^2 + 1)^2] \end{aligned}$$

$$= 1$$

$$= \text{RHS}$$

Now, we have to prove $\frac{b}{a} = \frac{2c}{c^2-1}$

Taking LHS, $\frac{b}{a}$

Putting the value of a and b, we get

$$\frac{b}{a} = \frac{\frac{2c}{c^2+1}}{\frac{c^2-1}{c^2+1}} = \frac{2c}{c^2+1} \times \frac{c^2+1}{c^2-1} = \frac{2c}{c^2-1} = \text{RHS}$$

Hence Proved

Q. 16. Show that $(1-i)^n \left(1 - \frac{1}{i}\right)^n = 2^n$ **for all n N.**

Solution: To show: $(1-i)^n \left(1 - \frac{1}{i}\right)^n = 2^n$

Taking LHS,

$$(1-i)^n \left(1 - \frac{1}{i}\right)^n$$

$$= (1-i)^n \left(1 - \frac{1}{i} \times \frac{i}{i}\right)^n \quad [\text{rationalize}]$$

$$= (1-i)^n \left(1 - \frac{i}{i^2}\right)^n$$

$$= (1-i)^n \left(1 - \frac{i}{-1}\right)^n \quad [\because i^2 = -1]$$

$$= (1-i)^n (1+i)^n$$

$$= [(1-i)(1+i)]^n$$

$$= [(1)^2 - (i)^2]^n [(a + b)(a - b) = a^2 - b^2]$$

$$= (1 - i^2)^n$$

$$= [1 - (-1)]^n [\because i^2 = -1]$$

$$= (2)^n$$

$$= 2^n$$

= RHS

Hence Proved

Q. 17. Find the smallest positive integer n for which $(1 + i)^{2n} = (1 - i)^{2n}$.

Solution:

Given: $(1 + i)^{2n} = (1 - i)^{2n}$

Consider the given equation,

$$(1 + i)^{2n} = (1 - i)^{2n}$$

$$\Rightarrow \frac{(1 + i)^{2n}}{(1 - i)^{2n}} = 1$$

$$\Rightarrow \left(\frac{1 + i}{1 - i}\right)^{2n} = 1$$

Now, rationalizing by multiply and divide by the conjugate of $(1 - i)$

$$\left(\frac{1 + i}{1 - i} \times \frac{1 + i}{1 + i}\right)^{2n} = 1$$

$$\Rightarrow \left(\frac{(1 + i)^2}{(1 - i)(1 + i)}\right)^{2n} = 1$$

$$\Rightarrow \left[\frac{1 + i^2 + 2i}{(1)^2 - (i)^2}\right]^{2n} = 1$$

$$[(a + b)^2 = a^2 + b^2 + 2ab \text{ \& } (a - b)(a + b) = (a^2 - b^2)]$$

$$\Rightarrow \left[\frac{1+(-1)+2i}{1-(-1)} \right]^{2n} = 1 \quad [i^2 = -1]$$

$$\Rightarrow \left[\frac{2i}{2} \right]^{2n} = 1$$

$$\Rightarrow (i)^{2n} = 1$$

Now, $i^{2n} = 1$ is possible if $n = 2$ because $(i)^{2(2)} = i^4 = (-1)^4 = 1$

So, the smallest positive integer $n = 2$

Q. 18. Prove that $(x + 1 + i)(x + 1 - i)(x - 1 + i)(x - 1 - i) = (x^2 + 4)$.

Solution: To Prove:

$$(x + 1 + i)(x + 1 - i)(x - 1 + i)(x - 1 - i) = (x^2 + 4)$$

Taking LHS

$$\begin{aligned} & (x + 1 + i)(x + 1 - i)(x - 1 + i)(x - 1 - i) \\ &= [(x + 1) + i][(x + 1) - i][(x - 1) + i][(x - 1) - i] \end{aligned}$$

Using $(a - b)(a + b) = a^2 - b^2$

$$\begin{aligned} & \underbrace{[(x + 1) + i][(x + 1) - i]}_{a = x + 1 \text{ \& \ } b = i} \underbrace{[(x - 1) + i][(x - 1) - i]}_{a = x - 1 \text{ \& \ } b = i} \end{aligned}$$

$$\begin{aligned} &= [(x + 1)^2 - (i)^2][(x - 1)^2 - (i)^2] \\ &= [x^2 + 1 + 2x - i^2][x^2 + 1 - 2x - i^2] \\ &= [x^2 + 1 + 2x - (-1)][x^2 + 1 - 2x - (-1)] \quad [\because i^2 = -1] \\ &= [x^2 + 2 + 2x][x^2 + 2 - 2x] \end{aligned}$$

Again, using $(a - b)(a + b) = a^2 - b^2$

Now, $a = x^2 + 2$ and $b = 2x$

$$\begin{aligned}
 &= [(x^2 + 2)^2 - (2x)^2] \\
 &= [x^4 + 4 + 2(x^2)(2) - 4x^2] [\because (a + b)^2 = a^2 + b^2 + 2ab] \\
 &= [x^4 + 4 + 4x^2 - 4x^2] \\
 &= x^4 + 4 \\
 &= \text{RHS} \\
 \therefore \text{LHS} &= \text{RHS}
 \end{aligned}$$

Hence Proved

Q. 19. If $a = (\cos\theta + i \sin\theta)$, prove that $\frac{1+a}{1-a} = \left(\cot \frac{\theta}{2}\right) i$.

Solution: Given: $a = \cos\theta + i \sin\theta$

To prove: $\frac{1+a}{1-a} = \left(\cot \frac{\theta}{2}\right) i$

Taking LHS,

$$\frac{1+a}{1-a}$$

Putting the value of a , we get

$$= \frac{1 + \cos \theta + i \sin \theta}{1 - (\cos \theta + i \sin \theta)}$$

$$= \frac{1 + \cos \theta + i \sin \theta}{1 - \cos \theta - i \sin \theta}$$

We know that,

$$1 + \cos 2\theta = 2\cos^2\theta$$

Or $1 + \cos \theta = 2 \cos^2 \frac{\theta}{2}$

And $1 - \cos \theta = 2 \sin^2 \frac{\theta}{2}$

Using the above two formulas

$$= \frac{2 \cos^2 \frac{\theta}{2} + i \sin \theta}{2 \sin^2 \frac{\theta}{2} - i \sin \theta}$$

Using, $\sin \theta = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}$

$$= \frac{2 \cos^2 \frac{\theta}{2} + i 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{2 \sin^2 \frac{\theta}{2} - 2i \sin \frac{\theta}{2} \cos \frac{\theta}{2}}$$

$$= \frac{2 \cos \frac{\theta}{2} \left[\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} \right]}{2 \sin \frac{\theta}{2} \left[\sin \frac{\theta}{2} - i \cos \frac{\theta}{2} \right]}$$

$$= \cot \frac{\theta}{2} \left[\frac{\cos \frac{\theta}{2} + i \sin \frac{\theta}{2}}{\sin \frac{\theta}{2} - i \cos \frac{\theta}{2}} \right] \left[\because \frac{\cos \theta}{\sin \theta} = \cot \theta \right]$$

Rationalizing by multiply and divide by the conjugate of $\sin \frac{\theta}{2} - i \cos \frac{\theta}{2}$

$$= \left(\cot \frac{\theta}{2} \right) \left[\frac{\cos \frac{\theta}{2} + i \sin \frac{\theta}{2}}{\sin \frac{\theta}{2} - i \cos \frac{\theta}{2}} \times \frac{\sin \frac{\theta}{2} + i \cos \frac{\theta}{2}}{\sin \frac{\theta}{2} + i \cos \frac{\theta}{2}} \right]$$

$$= \left(\cot \frac{\theta}{2} \right) \frac{(\cos \frac{\theta}{2} + i \sin \frac{\theta}{2})(\sin \frac{\theta}{2} + i \cos \frac{\theta}{2})}{(\sin \frac{\theta}{2} - i \cos \frac{\theta}{2})(\sin \frac{\theta}{2} + i \cos \frac{\theta}{2})}$$

$$= \left(\cot \frac{\theta}{2} \right) \frac{(\cos \frac{\theta}{2})(\sin \frac{\theta}{2} + i \cos \frac{\theta}{2}) + i \sin \frac{\theta}{2}(\sin \frac{\theta}{2} + i \cos \frac{\theta}{2})}{(\sin \frac{\theta}{2})^2 - (i \cos \frac{\theta}{2})^2}$$

$$= \left(\cot \frac{\theta}{2} \right) \frac{\cos \frac{\theta}{2} \sin \frac{\theta}{2} + i \cos^2 \frac{\theta}{2} + i \sin^2 \frac{\theta}{2} + i^2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{\sin^2 \frac{\theta}{2} - i^2 \cos^2 \frac{\theta}{2}}$$

Putting $i^2 = -1$, we get

$$= \left(\cot \frac{\theta}{2} \right) \frac{\cos \frac{\theta}{2} \sin \frac{\theta}{2} + i \cos^2 \frac{\theta}{2} + i \sin^2 \frac{\theta}{2} + (-1) \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{\sin^2 \frac{\theta}{2} - (-1) \cos^2 \frac{\theta}{2}}$$

$$= \left(\cot \frac{\theta}{2} \right) \frac{\cos \frac{\theta}{2} \sin \frac{\theta}{2} + i (\cos^2 \frac{\theta}{2} + \sin^2 \frac{\theta}{2}) - \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{\sin^2 \frac{\theta}{2} + \cos^2 \frac{\theta}{2}}$$

We know that,

$$\cos^2 \theta + \sin^2 \theta = 1$$

$$= \left(\cot \frac{\theta}{2} \right) \left[\frac{i (\cos^2 \frac{\theta}{2} + \sin^2 \frac{\theta}{2})}{1} \right]$$

$$= \cot \frac{\theta}{2} (i)$$

= RHS

Hence Proved

Q. 20. If $z_1 = (2 - i)$ and $z_2 = (1 + i)$, find $\left| \frac{z_1 + z_2 + 1}{z_1 - z_2 + i} \right|$.

Solution:

Given: $z_1 = (2 - i)$ and $z_2 = (1 + i)$

To find: $\left| \frac{z_1 + z_2 + 1}{z_1 - z_2 + i} \right|$

Consider,

$$\left| \frac{z_1 + z_2 + 1}{z_1 - z_2 + i} \right|$$

Putting the value of z_1 and z_2 , we get

$$= \left| \frac{2 - i + 1 + i + 1}{2 - i - (1 + i) + i} \right|$$

$$= \left| \frac{4}{2 - i - 1 - i + i} \right|$$

$$= \left| \frac{4}{1 - i} \right|$$

Now, rationalizing by multiply and divide by the conjugate of $1 - i$

$$= \left| \frac{4}{1 - i} \times \frac{1 + i}{1 + i} \right|$$

$$= \left| \frac{4(1 + i)}{(1 - i)(1 + i)} \right|$$

$$= \left| \frac{4(1 + i)}{(1)^2 - (i)^2} \right| \quad [(a - b)(a + b) = a^2 - b^2]$$

$$= \left| \frac{4(1 + i)}{1 - i^2} \right| = \left| \frac{4(1 + i)}{1 - (-1)} \right| \quad [\text{Putting } i^2 = -1]$$

$$= \left| \frac{4(1 + i)}{2} \right|$$

$$= |2(1 + i)|$$

$$= |2 + 2i|$$

Now, we have to find the modulus of $(2 + 2i)$

$$\text{So, } |z| = |2 + 2i| = \sqrt{(2)^2 + (2)^2} = \sqrt{4 + 4} = \sqrt{8} = 2\sqrt{2}$$

Hence, the value of $\left| \frac{z_1 + z_2 + 1}{z_1 - z_2 + i} \right| = 2\sqrt{2}$

Q. 21. A. Find the real values of x and y for which:

$$(1 - i)x + (1 + i)y = 1 - 3i$$

Solution:

$$(1 - i)x + (1 + i)y = 1 - 3i$$

$$\Rightarrow x - ix + y + iy = 1 - 3i$$

$$\Rightarrow (x + y) - i(x - y) = 1 - 3i$$

Comparing the real parts, we get

$$x + y = 1 \dots(i)$$

Comparing the imaginary parts, we get

$$x - y = -3 \dots(ii)$$

Solving eq. (i) and (ii) to find the value of x and y

Adding eq. (i) and (ii), we get

$$x + y + x - y = 1 + (-3)$$

$$\Rightarrow 2x = 1 - 3$$

$$\Rightarrow 2x = -2$$

$$\Rightarrow x = -1$$

Putting the value of $x = -1$ in eq. (i), we get

$$(-1) + y = 1$$

$$\Rightarrow y = 1 + 1$$

$$\Rightarrow y = 2$$

Q. 21. B. Find the real values of x and y for which:

$$(x + iy)(3 - 2i) = (12 + 5i)$$

Solution: $x(3 - 2i) + iy(3 - 2i) = 12 + 5i$

$$\Rightarrow 3x - 2ix + 3iy - 2i^2y = 12 + 5i$$

$$\Rightarrow 3x + i(-2x + 3y) - 2(-1)y = 12 + 5i \quad [\because i^2 = -1]$$

$$\Rightarrow 3x + i(-2x + 3y) + 2y = 12 + 5i$$

$$\Rightarrow (3x + 2y) + i(-2x + 3y) = 12 + 5i$$

Comparing the real parts, we get

$$3x + 2y = 12 \dots(i)$$

Comparing the imaginary parts, we get

$$-2x + 3y = 5 \dots(ii)$$

Solving eq. (i) and (ii) to find the value of x and y

Multiply eq. (i) by 2 and eq. (ii) by 3, we get

$$6x + 4y = 24 \dots(iii)$$

$$-6x + 9y = 15 \dots(iv)$$

Adding eq. (iii) and (iv), we get

$$6x + 4y - 6x + 9y = 24 + 15$$

$$\Rightarrow 13y = 39$$

$$\Rightarrow y = 3$$

Putting the value of $y = 3$ in eq. (i), we get

$$3x + 2(3) = 12$$

$$\Rightarrow 3x + 6 = 12$$

$$\Rightarrow 3x = 12 - 6$$

$$\Rightarrow 3x = 6$$

$$\Rightarrow x = 2$$

Hence, the value of $x = 2$ and $y = 3$

Q. 21. A. Find the real values of x and y for which:

$$(1 - i)x + (1 + i)y = 1 - 3i$$

Solution: $(1 - i)x + (1 + i)y = 1 - 3i$

$$x - ix + y + iy = 1 - 3i$$

$$\Rightarrow (x + y) - i(x - y) = 1 - 3i$$

Comparing the real parts, we get

$$x + y = 1 \dots(i)$$

Comparing the imaginary parts, we get

$$x - y = -3 \dots(ii)$$

Solving eq. (i) and (ii) to find the value of x and y

Adding eq. (i) and (ii), we get

$$x + y + x - y = 1 + (-3)$$

$$\Rightarrow 2x = 1 - 3$$

$$\Rightarrow 2x = -2$$

$$\Rightarrow x = -1$$

Putting the value of $x = -1$ in eq. (i), we get

$$(-1) + y = 1$$

$$\Rightarrow y = 1 + 1$$

$$\Rightarrow y = 2$$

Q. 21. B. Find the real values of x and y for which:

$$(x + iy)(3 - 2i) = (12 + 5i)$$

Solution: $x(3 - 2i) + iy(3 - 2i) = 12 + 5i$

$$\Rightarrow 3x - 2ix + 3iy - 2i^2y = 12 + 5i$$

$$\Rightarrow 3x + i(-2x + 3y) - 2(-1)y = 12 + 5i \quad [\because i^2 = -1]$$

$$\Rightarrow 3x + i(-2x + 3y) + 2y = 12 + 5i$$

$$\Rightarrow (3x + 2y) + i(-2x + 3y) = 12 + 5i$$

Comparing the real parts, we get

$$3x + 2y = 12 \dots(i)$$

Comparing the imaginary parts, we get

$$-2x + 3y = 5 \dots(ii)$$

Solving eq. (i) and (ii) to find the value of x and y

Multiply eq. (i) by 2 and eq. (ii) by 3, we get

$$6x + 4y = 24 \dots(iii)$$

$$-6x + 9y = 15 \dots(iv)$$

Adding eq. (iii) and (iv), we get

$$6x + 4y - 6x + 9y = 24 + 15$$

$$\Rightarrow 13y = 39$$

$$\Rightarrow y = 3$$

Putting the value of $y = 3$ in eq. (i), we get

$$3x + 2(3) = 12$$

$$\Rightarrow 3x + 6 = 12$$

$$\Rightarrow 3x = 12 - 6$$

$$\Rightarrow 3x = 6$$

$$\Rightarrow x = 2$$

Hence, the value of $x = 2$ and $y = 3$

Q. 21. C. Find the real values of x and y for which:

$$x + 4yi = ix + y + 3$$

Solution: Given: $x + 4yi = ix + y + 3$

$$\text{or } x + 4yi = ix + (y + 3)$$

Comparing the real parts, we get

$$x = y + 3$$

$$\text{Or } x - y = 3 \dots(i)$$

Comparing the imaginary parts, we get

$$4y = x \dots(ii)$$

Putting the value of $x = 4y$ in eq. (i), we get

$$4y - y = 3$$

$$\Rightarrow 3y = 3$$

$$\Rightarrow y = 1$$

Putting the value of $y = 1$ in eq. (ii), we get

$$x = 4(1) = 4$$

Hence, the value of $x = 4$ and $y = 1$

Q. 21. D. Find the real values of x and y for which:

$$(1 + i) y^2 + (6 + i) = (2 + i)x$$

Solution: Given: $(1 + i) y^2 + (6 + i) = (2 + i)x$

Consider, $(1 + i) y^2 + (6 + i) = (2 + i)x$

$$\Rightarrow y^2 + iy^2 + 6 + i = 2x + ix$$

$$\Rightarrow (y^2 + 6) + i(y^2 + 1) = 2x + ix$$

Comparing the real parts, we get

$$y^2 + 6 = 2x$$

$$\Rightarrow 2x - y^2 - 6 = 0 \dots(i)$$

Comparing the imaginary parts, we get

$$y^2 + 1 = x$$

$$\Rightarrow x - y^2 - 1 = 0 \dots(ii)$$

Subtracting the eq. (ii) from (i), we get

$$2x - y^2 - 6 - (x - y^2 - 1) = 0$$

$$\Rightarrow 2x - y^2 - 6 - x + y^2 + 1 = 0$$

$$\Rightarrow x - 5 = 0$$

$$\Rightarrow x = 5$$

Putting the value of $x = 5$ in eq. (i), we get

$$2(5) - y^2 - 6 = 0$$

$$\Rightarrow 10 - y^2 - 6 = 0$$

$$\Rightarrow -y^2 + 4 = 0$$

$$\Rightarrow -y^2 = -4$$

$$\Rightarrow y^2 = 4$$

$$\Rightarrow y = \sqrt{4}$$

$$\Rightarrow y = \pm 2$$

Hence, the value of $x = 5$ and $y = \pm 2$

Q. 21. E. Find the real values of x and y for which:

$$\frac{(x + 3i)}{(2 + iy)} = (1 - i)$$

Solution: Given:

$$\frac{x + 3i}{2 + iy} = (1 - i)$$

$$\Rightarrow x + 3i = (1 - i)(2 + iy)$$

$$\Rightarrow x + 3i = 1(2 + iy) - i(2 + iy)$$

$$\Rightarrow x + 3i = 2 + iy - 2i - i^2y$$

$$\Rightarrow x + 3i = 2 + i(y - 2) - (-1)y [i^2 = -1]$$

$$\Rightarrow x + 3i = 2 + i(y - 2) + y$$

$$\Rightarrow x + 3i = (2 + y) + i(y - 2)$$

Comparing the real parts, we get

$$x = 2 + y$$

$$\Rightarrow x - y = 2 \dots(i)$$

Comparing the imaginary parts, we get

$$3 = y - 2$$

$$\Rightarrow y = 3 + 2$$

$$\Rightarrow y = 5$$

Putting the value of $y = 5$ in eq. (i), we get

$$x - 5 = 2$$

$$\Rightarrow x = 2 + 5$$

$$\Rightarrow x = 7$$

Hence, the value of $x = 7$ and $y = 5$

Q. 21. F. Find the real values of x and y for which:

$$\frac{(1+i)x - 2i}{(3+i)} + \frac{(2-3i)y + i}{(3-i)} = i$$

Solution: Consider,

$$\begin{aligned} \frac{(1+i)x - 2i}{3+i} + \frac{(2-3i)y + i}{3-i} &= i \\ &= \frac{x + xi - 2i}{3+i} + \frac{2y - 3iy + i}{3-i} = i \end{aligned}$$

Taking LCM

$$\begin{aligned} \Rightarrow \frac{(x + xi - 2i)(3-i) + (2y - 3iy + i)(3+i)}{(3+i)(3-i)} &= i \\ \Rightarrow \frac{3x + 3xi - 6i - xi - xi^2 + 2i^2 + 6y - 9iy + 3i + 2iy - 3i^2y + i^2}{(3)^2 - (i)^2} &= i \end{aligned}$$

Putting $i^2 = -1$

$$\begin{aligned} \Rightarrow \frac{3x + 2xi - 6i - x(-1) + 2(-1) + 6y - 7iy + 3i - 3(-1)y + (-1)}{9 - (-1)} &= i \\ \Rightarrow \frac{3x + 2xi - 6i + x - 2 + 6y - 7iy + 3i + 3y - 1}{9 + 1} &= i \\ \Rightarrow \frac{4x + 2xi - 3i - 3 + 9y - 7iy}{10} &= i \end{aligned}$$

$$\Rightarrow 4x + 2xi - 3i - 3 + 9y - 7iy = 10i$$

$$\Rightarrow (4x - 3 + 9y) + i(2x - 3 - 7y) = 10i$$

Comparing the real parts, we get

$$4x - 3 + 9y = 0$$

$$\Rightarrow 4x + 9y = 3 \dots(i)$$

Comparing the imaginary parts, we get

$$2x - 3 - 7y = 10$$

$$\Rightarrow 2x - 7y = 10 + 3$$

$$\Rightarrow 2x - 7y = 13 \dots(ii)$$

Multiply the eq. (ii) by 2, we get

$$4x - 14y = 26 \dots(iii)$$

Subtracting eq. (i) from (iii), we get

$$4x - 14y - (4x + 9y) = 26 - 3$$

$$\Rightarrow 4x - 14y - 4x - 9y = 23$$

$$\Rightarrow -23y = 23$$

$$\Rightarrow y = -1$$

Putting the value of $y = -1$ in eq. (i), we get

$$4x + 9(-1) = 3$$

$$\Rightarrow 4x - 9 = 3$$

$$\Rightarrow 4x = 12$$

$$\Rightarrow x = 3$$

Hence, the value of $x = 3$ and $y = -1$

Q. 22

Find the real values of x and y for which $(x - iy)(3 + 5i)$ is the conjugate of $(-6 - 24i)$.

Solution: Given: $(x - iy)(3 + 5i)$ is the conjugate of $(-6 - 24i)$

We know that,

$$\text{Conjugate of } -6 - 24i = -6 + 24i$$

∴ According to the given condition,

$$(x - iy)(3 + 5i) = -6 + 24i$$

$$\Rightarrow x(3 + 5i) - iy(3 + 5i) = -6 + 24i$$

$$\Rightarrow 3x + 5ix - 3iy - 5i^2y = -6 + 24i$$

$$\Rightarrow 3x + i(5x - 3y) - 5(-1)y = -6 + 24i \quad [\because i^2 = -1]$$

$$\Rightarrow 3x + i(5x - 3y) + 5y = -6 + 24i$$

$$\Rightarrow (3x + 5y) + i(5x - 3y) = -6 + 24i$$

Comparing the real parts, we get

$$3x + 5y = -6 \dots(i)$$

Comparing the imaginary parts, we get

$$5x - 3y = 24 \dots(ii)$$

Solving eq. (i) and (ii) to find the value of x and y

Multiply eq. (i) by 5 and eq. (ii) by 3, we get

$$15x + 25y = -30 \dots(iii)$$

$$15x - 9y = 72 \dots(iv)$$

Subtracting eq. (iii) from (iv), we get

$$15x - 9y - 15x - 25y = 72 - (-30)$$

$$\Rightarrow -34y = 72 + 30$$

$$\Rightarrow -34y = 102$$

$$\Rightarrow y = -3$$

Putting the value of y = -3 in eq. (i), we get

$$3x + 5(-3) = -6$$

$$\Rightarrow 3x - 15 = -6$$

$$\Rightarrow 3x = -6 + 15$$

$$\Rightarrow 3x = 9$$

$$\Rightarrow x = 3$$

Hence, the value of $x = 3$ and $y = -3$

Q. 23. Find the real values of x and y for which the complex number $(-3 + iyx^2)$ and $(x^2 + y + 4i)$ are conjugates of each other.

Solution: Let $z_1 = -3 + iyx^2$

So, the conjugate of

$$\bar{z}_1 = -3 - iyx^2$$

And $z_2 = x^2 + y + 4i$

So, the conjugate of z_2 is

$$\bar{z}_2 = x^2 + y - 4i$$

Given that: $\bar{z}_1 = z_2$ & $z_1 = \bar{z}_2$

Firstly, consider $\bar{z}_1 = z_2$

$$-3 - iyx^2 = x^2 + y + 4i$$

$$\Rightarrow x^2 + y + 4i + iyx^2 = -3$$

$$\Rightarrow x^2 + y + i(4 + yx^2) = -3 + 0i$$

Comparing the real parts, we get

$$x^2 + y = -3 \dots(i)$$

Comparing the imaginary parts, we get

$$4 + yx^2 = 0$$

$$\Rightarrow x^2y = -4 \dots(ii)$$

Now, consider $z_1 = \bar{z}_2$

$$-3 + iyx^2 = x^2 + y - 4i$$

$$\Rightarrow x^2 + y - 4i - iyx^2 = -3$$

$$\Rightarrow x^2 + y + i(-4i - yx^2) = -3 + 0i$$

Comparing the real parts, we get

$$x^2 + y = -3$$

Comparing the imaginary parts, we get

$$-4 - yx^2 = 0$$

$$\Rightarrow x^2y = -4$$

Now, we will solve the equations to find the value of x and y

From eq. (i), we get

$$x^2 = -3 - y$$

Putting the value of x^2 in eq. (ii), we get

$$(-3 - y)(y) = -4$$

$$\Rightarrow -3y - y^2 = -4$$

$$\Rightarrow y^2 + 3y = 4$$

$$\Rightarrow y^2 + 3y - 4 = 0$$

$$\Rightarrow y^2 + 4y - y - 4 = 0$$

$$\Rightarrow y(y + 4) - 1(y + 4) = 0$$

$$\Rightarrow (y - 1)(y + 4) = 0$$

$$\Rightarrow y - 1 = 0 \text{ or } y + 4 = 0$$

$$\Rightarrow y = 1 \text{ or } y = -4$$

When $y = 1$, then

$$x^2 = -3 - 1$$

$$= -4 \text{ [It is not possible]}$$

When $y = -4$, then

$$x^2 = -3 - (-4)$$

$$= -3 + 4$$

$$\Rightarrow x^2 = 1$$

$$\Rightarrow x = \sqrt{1}$$

$$\Rightarrow x = \pm 1$$

Hence, the values of $x = \pm 1$ and $y = -4$

Q. 24. If $z = (2 - 3i)$, prove that $z^2 - 4z + 13 = 0$ and hence deduce that $4z^3 - 3z^2 + 169 = 0$.

Solution: Given: $z = 2 - 3i$

To Prove: $z^2 - 4z + 13 = 0$

Taking LHS, $z^2 - 4z + 13$

Putting the value of $z = 2 - 3i$, we get

$$(2 - 3i)^2 - 4(2 - 3i) + 13$$

$$= 4 + 9i^2 - 12i - 8 + 12i + 13$$

$$= 9(-1) + 9$$

$$= -9 + 9$$

$$= 0$$

= RHS

Hence, $z^2 - 4z + 13 = 0 \dots(i)$

Now, we have to deduce $4z^3 - 3z^2 + 169$

Now, we will expand $4z^3 - 3z^2 + 169$ in this way so that we can use the above equation
i.e. $z^2 - 4z + 13$

$$= 4z^3 - 16z^2 + 13z^2 + 52z - 52z + 169$$

Re – arrange the terms,

$$= 4z^3 - 16z^2 + 52z + 13z^2 - 52z + 169$$

$$= 4z(z^2 - 4z + 13) + 13(z^2 - 4z + 13)$$

$$= 4z(0) + 13(0) \text{ [from eq. (i)]}$$

$$= 0$$

$$= \text{RHS}$$

Hence Proved

Q. 25. If $(1 + i)z = (1 - i)\bar{z}$ then prove that $z = -i\bar{z}$.

Solution: Let $z = x + iy$

Then,

$$\bar{z} = x - iy$$

Now, Given: $(1 + i)z = (1 - i)\bar{z}$

Therefore,

$$(1 + i)(x + iy) = (1 - i)(x - iy)$$

$$x + iy + xi + i^2y = x - iy - xi + i^2y$$

We know that $i^2 = -1$, therefore,

$$x + iy + ix - y = x - iy - ix - y$$

$$2xi + 2yi = 0$$

$$x = -y$$

Now, as $x = -y$

$$z = -\bar{z}$$

Hence, Proved.

Q. 26. If $\left(\frac{z-1}{z+1}\right)$ is purely an imaginary number and $z \neq -1$ then find the value of $|z|$.

Solution: Given: $\frac{z-1}{z+1}$ is purely imaginary number

$$\text{Let } z = x + iy$$

$$\text{So, } \frac{z-1}{z+1} = \frac{x+iy-1}{x+iy+1}$$

$$= \frac{(x-1) + iy}{(x+1) + iy}$$

Now, rationalizing the above by multiply and divide by the conjugate of $[(x+1) + iy]$

$$= \frac{(x-1) + iy}{(x+1) + iy} \times \frac{(x+1) - iy}{(x+1) - iy}$$

$$= \frac{[(x-1) + iy][(x+1) - iy]}{[(x+1) + iy][(x+1) - iy]}$$

Using $(a-b)(a+b) = (a^2 - b^2)$

$$= \frac{(x-1)[(x+1) - iy] + iy[(x+1) - iy]}{(x+1)^2 - (iy)^2}$$

$$= \frac{(x-1)(x+1) + (x-1)(-iy) + iy(x+1) + (iy)(-iy)}{x^2 + 1 + 2x - i^2y^2}$$

$$= \frac{x^2 - 1 - ixy + iy + ixy + iy - i^2y^2}{x^2 + 1 + 2x - i^2y^2}$$

Putting $i^2 = -1$

$$\begin{aligned}
 &= \frac{x^2 - 1 + 2iy - (-1)y^2}{x^2 + 1 + 2x - (-1)y^2} \\
 &= \frac{x^2 - 1 + 2iy + y^2}{x^2 + 1 + 2x + y^2} \\
 &= \frac{x^2 - 1 + y^2}{x^2 + 1 + 2x + y^2} + i \frac{2y}{x^2 + 1 + 2x + y^2}
 \end{aligned}$$

Since, the number is purely imaginary it means real part is 0

$$\therefore \frac{x^2 - 1 + y^2}{x^2 + 1 + 2x + y^2} = 0$$

$$\Rightarrow x^2 + y^2 - 1 = 0$$

$$\Rightarrow x^2 + y^2 = 1$$

$$\Rightarrow \sqrt{x^2 + y^2} = \sqrt{1}$$

$$\Rightarrow \sqrt{x^2 + y^2} = 1$$

$$\therefore |z| = 1$$

Q. 27. Solve the system of equations, $\operatorname{Re}(z^2) = 0$, $|z| = 2$.

Solution: Given: $\operatorname{Re}(z^2) = 0$ and $|z| = 2$

Let $z = x + iy$

$$\therefore |z| = \sqrt{x^2 + y^2}$$

$$\Rightarrow 2 = \sqrt{x^2 + y^2} \quad [\text{Given}]$$

Squaring both the sides, we get

$$x^2 + y^2 = 4 \dots (i)$$

Since, $z = x + iy$

$$\Rightarrow z^2 = (x + iy)^2$$

$$\Rightarrow z^2 = x^2 + i^2y^2 + 2ixy$$

$$\Rightarrow z^2 = x^2 + (-1)y^2 + 2ixy$$

$$\Rightarrow z^2 = x^2 - y^2 + 2ixy$$

It is given that $\text{Re}(z^2) = 0$

$$\Rightarrow x^2 - y^2 = 0 \dots(\text{ii})$$

Adding eq. (i) and (ii), we get

$$x^2 + y^2 + x^2 - y^2 = 4 + 0$$

$$\Rightarrow 2x^2 = 4$$

$$\Rightarrow x^2 = 2$$

$$\Rightarrow x = \pm\sqrt{2}$$

Putting the value of $x^2 = 2$ in eq. (i), we get

$$2 + y^2 = 4$$

$$\Rightarrow y^2 = 2$$

$$\Rightarrow y = \pm\sqrt{2}$$

Hence, $z = \sqrt{2} \pm i\sqrt{2}, -\sqrt{2} \pm i\sqrt{2}$

Q. 28. Find the complex number z for which $|z| = z + 1 + 2i$.

Solution: Given: $|z| = z + 1 + 2i$

Consider,

$$|z| = (z + 1) + 2i$$

Squaring both the sides, we get

$$|z|^2 = [(z + 1) + (2i)]^2$$

$$\Rightarrow |z|^2 = |z + 1|^2 + 4i^2 + 2(2i)(z + 1)$$

$$\Rightarrow |z|^2 = |z|^2 + 1 + 2z + 4(-1) + 4i(z + 1)$$

$$\Rightarrow 0 = 1 + 2z - 4 + 4i(z + 1)$$

$$\Rightarrow 2z - 3 + 4i(z + 1) = 0$$

Let $z = x + iy$

$$\Rightarrow 2(x + iy) - 3 + 4i(x + iy + 1) = 0$$

$$\Rightarrow 2x + 2iy - 3 + 4ix + 4i^2y + 4i = 0$$

$$\Rightarrow 2x + 2iy - 3 + 4ix + 4(-1)y + 4i = 0$$

$$\Rightarrow 2x - 3 - 4y + i(4x + 2y + 4) = 0$$

Comparing the real part, we get

$$2x - 3 - 4y = 0$$

$$\Rightarrow 2x - 4y = 3 \dots(i)$$

Comparing the imaginary part, we get

$$4x + 2y + 4 = 0$$

$$\Rightarrow 2x + y + 2 = 0$$

$$\Rightarrow 2x + y = -2 \dots(ii)$$

Subtracting eq. (ii) from (i), we get

$$2x - 4y - (2x + y) = 3 - (-2)$$

$$\Rightarrow 2x - 4y - 2x - y = 3 + 2$$

$$\Rightarrow -5y = 5$$

$$\Rightarrow y = -1$$

Putting the value of $y = -1$ in eq. (i), we get

$$2x - 4(-1) = 3$$

$$\Rightarrow 2x + 4 = 3$$

$$\Rightarrow 2x = 3 - 4$$

$$\Rightarrow 2x = -1$$

$$\Rightarrow x = -\frac{1}{2}$$

Hence, the value of $z = x + iy$

$$= -\frac{1}{2} + i(-1)$$

$$z = -\frac{1}{2} - i$$

EXERCISE 5C

PAGE: 188

Q. 1. Express each of the following in the form $(a + ib)$ and find its conjugate.

$$(i) \frac{1}{(4 + 3i)}$$

$$(ii) (2 + 3i)^2$$

$$(iii) \frac{(2 - i)}{(1 - 2i)^2}$$

$$(iv) \frac{(1 + i)(1 + 2i)}{(1 + 3i)}$$

$$(v) \left(\frac{1 + 2i}{2 + i}\right)^2$$

$$(vi) \frac{(2 + i)}{(3 - i)(1 + 2i)}$$

Solution:

$$(i) \text{ Let } z = \frac{1}{4+3i} = \frac{1}{4+3i} \times \frac{4-3i}{4-3i}$$

$$= \frac{4-3i}{16+9} = \frac{4}{25} - \frac{3}{25}i$$

$$\Rightarrow \bar{z} = \frac{4}{25} + \frac{3}{25}i$$

$$(ii) \text{ Let } z = (2+3i)^2 = (2+3i)(2+3i)$$

$$= 4 + 6i + 6i + 9i^2$$

$$= 4 + 12i + 9i^2$$

$$= 4 + 12i - 9$$

$$= -5 + 12i$$

$$\bar{z} = -5 - 12i$$

$$(iii) \text{ Let } z = \frac{(2-i)}{(1-2i)^2} = \frac{(2-i)}{1+4i^2-4i}$$

$$= \frac{(2-i)}{1-4i-4} = \frac{2-i}{-3-4i}$$

$$= \frac{2-i}{-3-4i} \times \frac{-3+4i}{-3+4i} = \frac{(2-i)(-3+4i)}{9+16}$$

$$= \frac{-6 + 11i - 4i^2}{25} = \frac{-2 + 11i}{25}$$

$$= \frac{-2}{25} + \frac{11}{25}i$$

$$\bar{z} = \frac{-2}{25} - \frac{11}{25}i$$

$$(iv) \text{ Let } z = \frac{(1+i)(1+2i)}{(1+3i)} = \frac{1+i+2i+2i^2}{(1+3i)}$$

$$= \frac{1 + 3i - 2}{1 + 3i} = \frac{-1 + 3i}{1 + 3i}$$

$$= \frac{-1 + 3i}{1 + 3i} \times \frac{1 - 3i}{1 - 3i} = \frac{-1 + 3i + 3i - 9i^2}{1 - 9i^2} = \frac{-1 + 6i + 9}{1 + 9} = \frac{8 + 6i}{10}$$

$$= \frac{8}{10} + \frac{6}{10}i$$

$$\bar{z} = \frac{8}{10} - \frac{6}{10}i$$

$$(v) \text{ Let } z = \left(\frac{1+2i}{2+i}\right)^2 = \frac{1+4i^2+2i}{4+i^2+4i} = \frac{1-4+2i}{4-1+4i} = \frac{-3+2i}{3+4i}$$

$$= \frac{-3+2i}{3+4i} \times \frac{3-4i}{3-4i}$$

$$= \frac{-9+12i+6i-8i^2}{9+16} = \frac{-9+18i+8}{25} = \frac{-1+18i}{25}$$

$$= \frac{-1}{25} + \frac{18}{25}i$$

$$\bar{z} = \frac{-1}{25} - \frac{18}{25}i$$

$$(vi) \text{ Let } z = \frac{(2+i)}{(3-i)(1+2i)} = \frac{2+i}{3+6i-1-2i^2}$$

$$= \frac{2+i}{3+6i-1+2} = \frac{2+i}{4+6i}$$

$$= \frac{2+i}{4+6i} \times \frac{4-6i}{4-6i}$$

$$= \frac{8-12i+4i-6i^2}{16+36}$$

$$= \frac{8 - 8i + 6}{52}$$

$$= \frac{14 - 8i}{52}$$

$$= \frac{14}{52} - \frac{8}{52}i$$

$$\bar{z} = \frac{14}{52} + \frac{8}{52}i$$

Q. 2. Express each of the following in the form $(a + ib)$ and find its multiplicative inverse:

(i) $\frac{1 + 2i}{1 - 3i}$

(ii) $\frac{(1 + 7i)}{(2 - i)^2}$

(iii) $\frac{-4}{(1 + i\sqrt{3})}$

Solution:

(i) Let $z = \frac{1 + 2i}{1 - 3i}$

$$= \frac{1 + 2i}{1 - 3i} \times \frac{1 + 3i}{1 + 3i} = \frac{1 + 3i + 2i + 6i^2}{1 - 9i^2}$$

$$= \frac{1 + 5i + 6i^2}{1 + 9} = \frac{-5 + 5i}{10}$$

$$z = \frac{-1}{2} + \frac{1}{2}i$$

$$\Rightarrow \bar{z} = \frac{-1}{2} - \frac{1}{2}i$$

$$\Rightarrow |z|^2 = \left(\frac{-1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 = \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

\therefore The multiplicative inverse of $\frac{1+2i}{1-3i}$

$$z^{-1} = \frac{\bar{z}}{|z|^2} = \frac{\frac{-1}{2} - \frac{1}{2}i}{\frac{1}{2}} = -1 - i$$

(ii) Let $z = \frac{1+7i}{(2-i)^2}$

$$= \frac{1+7i}{4+i^2-4i} = \frac{1+7i}{4-1-4i} = \frac{1+7i}{3-4i}$$

$$= \frac{1+7i}{3-4i} \times \frac{3+4i}{3+4i}$$

$$= \frac{3+4i+21i+28i^2}{9+16} = \frac{3+25i-28}{25} = \frac{-25+25i}{25}$$

$$z = -1 + i$$

$$\Rightarrow \bar{z} = -1 - i$$

$$\Rightarrow |z|^2 = (-1)^2 + (1)^2 = 1 + 1 = 2$$

\therefore The multiplicative inverse of $\frac{1+7i}{(2-i)^2}$

$$z^{-1} = \frac{\bar{z}}{|z|^2} = \frac{-1-i}{2} = \frac{-1}{2} - \frac{1}{2}i$$

(iii) Let $z = \frac{-4}{(1+i\sqrt{3})}$

$$= \frac{-4}{1 + i\sqrt{3}} \times \frac{1 - i\sqrt{3}}{1 - i\sqrt{3}}$$

$$= \frac{-4 + i4\sqrt{3}}{1 + 3} = \frac{-4 + i4\sqrt{3}}{4}$$

$$= -1 + i\sqrt{3}$$

$$Z = -1 + i\sqrt{3}$$

$$\Rightarrow \bar{z} = -1 + i\sqrt{3}$$

$$\Rightarrow |z|^2 = (-1)^2 + (\sqrt{3})^2 = 1 + 3 = 4$$

\(\therefore\) The multiplicative inverse of $\frac{-4}{(1 + i\sqrt{3})}$

$$z^{-1} = \frac{\bar{z}}{|z|^2} = \frac{-1 + i\sqrt{3}}{4} = \frac{-1}{4} + \frac{i\sqrt{3}}{4}$$

Q. 3. If $(x + iy)^3 = (u + iv)$ then prove that $\left(\frac{u}{x} + \frac{v}{y}\right) = 4(x^2 - y^2)$.

Solution: Given that, $(x + iy)^3 = (u + iv)$

$$\Rightarrow x^3 + (iy)^3 + 3x^2iy + 3xi^2y^2 = u + iv$$

$$\Rightarrow x^3 - iy^3 + 3x^2iy - 3xy^2 = u + iv$$

$$\Rightarrow x^3 - 3xy^2 + i(3x^2y - y^3) = u + iv$$

On equating real and imaginary parts, we get

$$U = x^3 - 3xy^2 \text{ and } v = 3x^2y - y^3$$

$$\text{Now, } \frac{u}{x} + \frac{v}{y} = \frac{x^3 - 3xy^2}{x} + \frac{3x^2y - y^3}{y}$$

$$= \frac{x(x^2 - 3y^2)}{x} + \frac{y(3x^2 - y^2)}{y}$$

$$= x^2 - 3y^2 + 3x^2 - y^2$$

$$= 4x^2 - 4y^2$$

$$= 4(x^2 - y^2)$$

Hence, $\frac{u}{x} + \frac{v}{y} = 4(x^2 - y^2)$

Q. 4. If $(x + iy)^{1/3} = (a + ib)$ then prove that $\left(\frac{x}{a} + \frac{y}{b}\right) = 4(a^2 - b^2)$.

Solution: Given that, $(x + iy)^{1/3} = (a + ib)$

$$\Rightarrow (x + iy) = (a + ib)^3$$

$$\Rightarrow (a + ib)^3 = x + iy$$

$$\Rightarrow a^3 + (ib)^3 + 3a^2ib + 3ai^2b^2 = x + iy$$

$$\Rightarrow a^3 - ib^3 + 3a^2ib - 3ab^2 = x + iy$$

$$\Rightarrow a^3 - 3ab^2 + i(3a^2b - b^3) = x + iy$$

On equating real and imaginary parts, we get

$$x = a^3 - 3ab^2 \text{ and } y = 3a^2b - b^3$$

Now, $\frac{x}{a} + \frac{y}{b} = \frac{a^3 - 3ab^2}{a} + \frac{3a^2b - b^3}{b}$

$$= \frac{a(a^2 - 3b^2)}{a} + \frac{b(3a^2 - b^2)}{b}$$

$$= a^2 - 3b^2 + 3a^2 - b^2$$

$$= 4a^2 - 4b^2$$

$$= 4(a^2 - b^2)$$

Hence, $\frac{x}{a} + \frac{y}{b} = 4(a^2 - b^2)$

Q. 5. Express $(1 - 2i)^{-3}$ in the form $(a + ib)$.

Solution: We have, $(1 - 2i)^{-3}$

$$\Rightarrow \frac{1}{(1 - 2i)^3} = \frac{1}{1 - 8i^3 - 6i + 12i^2} = \frac{1}{1 + 8i - 6i - 12} = \frac{1}{2i - 11}$$

$$\Rightarrow \frac{1}{-11 + 2i}$$

$$= \frac{1}{-11 + 2i} \times \frac{-11 - 2i}{-11 - 2i}$$

$$= \frac{-11 - 2i}{(-11)^2 - (2i)^2} = \frac{-11 - 2i}{121 + 4}$$

$$= \frac{-11 - 2i}{125}$$

$$= \frac{-11}{125} - \frac{2i}{125}$$

Q. 6. Find real values of x and y for which

$$(x^4 + 2xi) - (3x^2 + iy) = (3 - 5i) + (1 + 2iy).$$

Solution: We have, $(x^4 + 2xi) - (3x^2 + iy) = (3 - 5i) + (1 + 2iy).$

$$\Rightarrow x^4 + 2xi - 3x^2 + iy = 3 - 5i + 1 + 2iy$$

$$\Rightarrow (x^4 - 3x^2) + i(2x - y) = 4 + i(2y - 5)$$

On equating real and imaginary parts, we get

$$x^4 - 3x^2 = 4 \text{ and } 2x - y = 2y - 5$$

$$\Rightarrow x^4 - 3x^2 - 4 = 0 \text{ eq(i) and } 2x - y - 2y + 5 = 0 \text{ eq(ii)}$$

Now from eq (i), $x^4 - 3x^2 - 4 = 0$

$$\Rightarrow x^4 - 4x^2 + x^2 - 4 = 0$$

$$\Rightarrow x^2(x^2 - 4) + 1(x^2 - 4) = 0$$

$$\Rightarrow (x^2 - 4)(x^2 + 1) = 0$$

$$\Rightarrow x^2 - 4 = 0 \text{ and } x^2 + 1 = 0$$

$$\Rightarrow x = \pm 2 \text{ and } x = \sqrt{-1}$$

Real value of $x = \pm 2$

Putting $x = 2$ in eq (ii), we get

$$2x - 3y + 5 = 0$$

$$\Rightarrow 2 \times 2 - 3y + 5 = 0$$

$$\Rightarrow 4 - 3y + 5 = 0 = 9 - 3y = 0$$

$$\Rightarrow y = 3$$

Putting $x = -2$ in eq (ii), we get

$$2x - 3y + 5 = 0$$

$$\Rightarrow 2 \times -2 - 3y + 5 = 0$$

$$\Rightarrow -4 - 3y + 5 = 0 = 1 - 3y = 0$$

$$\Rightarrow y = \frac{1}{3}$$

Q. 7. If $z^2 + |z|^2 = 0$, show that z is purely imaginary.

Solution: Let $z = a + ib$

$$\Rightarrow |z| = \sqrt{a^2 + b^2}$$

$$\text{Now, } z^2 + |z|^2 = 0$$

$$\Rightarrow (a + ib)^2 + a^2 + b^2 = 0$$

$$\Rightarrow a^2 + 2abi + i^2b^2 + a^2 + b^2 = 0$$

$$\Rightarrow a^2 + 2abi - b^2 + a^2 + b^2 = 0$$

$$\Rightarrow 2a^2 + 2abi = 0$$

$$\Rightarrow 2a(a + ib) = 0$$

Either $a = 0$ or $z = 0$

Since $z \neq 0$

$a = 0 \Rightarrow z$ is purely imaginary.

$$\frac{z-1}{z+1}$$

Q. 8. If $\frac{z-1}{z+1}$ is purely imaginary, show that $|z| = 1$.

Solution: Let $z = a + ib$

$$\text{Now, } \frac{z-1}{z+1} = \frac{a+ib-1}{a+ib+1}$$

$$= \frac{(a-1) + ib}{(a+1) + ib}$$

$$\Rightarrow \frac{(a-1) + ib}{(a+1) + ib} \times \frac{(a+1) - ib}{(a+1) - ib}$$

$$= \frac{a^2 + a - iab - a - 1 + ib + iab + ib - i^2b^2}{(a+1)^2 + b^2}$$

$$= \frac{a^2 + -1 + ib + ib + b^2}{(a+1)^2 + b^2} = \frac{a^2 + b^2 - 1 + 2ib}{(a+1)^2 + b^2}$$

Given that $\frac{z-1}{z+1}$ is purely imaginary \Rightarrow real part = 0

$$\Rightarrow \frac{a^2 + b^2 - 1}{(a+1)^2 + b^2} = 0$$

$$\Rightarrow a^2 + b^2 - 1 = 0$$

$$\Rightarrow a^2 + b^2 = 1$$

$$\Rightarrow |z| = 1$$

Hence proved.

Q. 9. If z_1 is a complex number other than -1 such that $|z_1| = 1$ and $z_2 = \frac{z_1 - 1}{z_1 + 1}$ then show that z^2 is purely imaginary.

Solution: Let $z_1 = a + ib$ such that $|z_1| = \sqrt{a^2 + b^2} = 1$

$$\text{Now, } z_2 = \frac{z_1 - 1}{z_1 + 1} = \frac{a + ib - 1}{a + ib + 1} = \frac{(a-1) + ib}{(a+1) + ib}$$

$$\begin{aligned} &\Rightarrow \frac{(a-1) + ib}{(a+1) + ib} \times \frac{(a+1) - ib}{(a+1) - ib} \\ &= \frac{a^2 + a - iab - a - 1 + ib + iab + ib - i^2 b^2}{(a+1)^2 + b^2} \\ &= \frac{a^2 + -1 + ib + ib + b^2}{(a+1)^2 + b^2} = \frac{a^2 + b^2 - 1 + 2ib}{(a+1)^2 + b^2} \\ &= \frac{(a^2 + b^2) - 1 + 2ib}{(a+1)^2 + b^2} = \frac{1 - 1 + 2ib}{(a+1)^2 + b^2} [\because a^2 + b^2 = 1] \\ &= 0 + \frac{2ib}{(a+1)^2 + b^2} \end{aligned}$$

Thus, the real part of z^2 is 0 and z^2 is purely imaginary.

Q. 10. For all $z \in \mathbb{C}$, prove that

$$(i) \quad \frac{1}{2}(z + \bar{z}) = \text{Re}(z)$$

$$(ii) \quad \frac{1}{2}(z - \bar{z}) = i \text{Im}(z)$$

$$(iii) \quad z\bar{z} = |z|^2$$

$$(iv) \quad (z + \bar{z}) \text{ is real}$$

(v) $(z - \bar{z})$ is 0 or imaginary.

Solution:

Let $z = a + ib$

$$\Rightarrow \bar{z} = a - ib$$

$$\text{Now, } \frac{z + \bar{z}}{2} = \frac{(a + ib) + (a - ib)}{2} = \frac{2a}{2} = a = \text{Re}(z)$$

Hence Proved.

(ii) Let $z = a + ib$

$$\Rightarrow \bar{z} = a - ib$$

$$\begin{aligned} \text{Now, } \frac{z + \bar{z}}{2} &= \frac{(a + ib) + (a - ib)}{2} \\ &= \frac{2a}{2} = \frac{2a}{2} = \frac{a}{1} = \text{Re}(z) \end{aligned}$$

Hence, Proved.

(iii) Let $z = a + ib$

$$\Rightarrow \bar{z} = a - ib$$

$$\text{Now, } z\bar{z} = (a + ib)(a - ib) = a^2 - (ib)^2 = a^2 + b^2 = |z|^2$$

Hence Proved.

(iv) Let $z = a + ib$

$$\Rightarrow \bar{z} = a - ib$$

$$\text{Now, } z + \bar{z} = (a + ib) + (a - ib) = 2a = 2\text{Re}(z)$$

Hence, $z + \bar{z}$ is real.

(v) Case 1. Let $z = a + 0i$

$$\Rightarrow \bar{z} = a - 0i$$

$$\text{Now, } z - \bar{z} = (a + 0i) - (a - 0i) = 0$$

Case 2. Let $z = 0 + bi$

$$\Rightarrow \bar{z} = 0 - bi$$

$$\text{Now, } z - \bar{z} = (0 + ib) - (0 - ib) = 2ib = 2i\text{Im}(z) = \text{Imaginary}$$

Case 2. Let $z = a + ib$

$$\Rightarrow \bar{z} = a - ib$$

$$\text{Now, } z - \bar{z} = (a + ib) - (a - ib) = 2ib = 2i\text{Im}(z) = \text{Imaginary}$$

Thus, $(z - \bar{z})$ is 0 or imaginary.

$$\text{Q. 11. If } z_1 = (1 + i) \text{ and } z_2 = (-2 + 4i), \text{ prove that } \text{Im} \left(\frac{z_1 z_2}{z_1} \right) = 2$$

Solution: We have, $z_1 = (1 + i)$ and $z_2 = (-2 + 4i)$

$$\text{Now, } \frac{z_1 z_2}{z_1} = \frac{(1 + i)(-2 + 4i)}{(1 + i)}$$

$$= \frac{-2 + 4i - 2i + 4i^2}{(1 - i)} = \frac{-2 + 4i - 2i - 4}{(1 - i)} = \frac{-6 + 2i}{(1 - i)}$$

$$= \frac{-6 + 2i}{(1 - i)} \times \frac{(1 + i)}{(1 + i)}$$

$$= \frac{-6 - 6i + 2i + 2i^2}{1 + 1}$$

$$= \frac{-6 - 4i - 2}{2} = \frac{-8 - 4i}{2}$$

$$= -4 - 2i$$

Hence, $Im\left(\frac{z_1 z_2}{z_2}\right) = -2$

Q. 12. If a and b are real numbers such that $a^2 + b^2 = 1$ then show that a real value

of x satisfies the equation, $\frac{1-ix}{1+ix} = (a-ib)$

Solution: We have,

$$\frac{1-ix}{1+ix} = (a-ib) = \frac{a-ib}{1}$$

Applying componendo and dividendo, we get

$$\frac{(1-ix) + (1+ix)}{(1-ix) - (1+ix)} = \frac{a-ib + 1}{a-ib - 1}$$

$$\Rightarrow \frac{1-ix + 1 + ix}{1-ix - 1 + ix} = \frac{a-ib + 1}{a-ib - 1}$$

$$\Rightarrow \frac{2}{-2ix} = \frac{a-ib + 1}{-(-a + ib + 1)}$$

$$\Rightarrow ix = \frac{1-a+ib}{1+a-ib} \times \frac{1+a+ib}{1+a+ib}$$

$$= \frac{1+a+ib-a-a^2-aib+ib+aib+i^2b^2}{(1+a)^2 - i^2b^2}$$

$$\Rightarrow ix = \frac{1-a^2-b^2+2ib}{(1+a)^2 - i^2b^2} = \frac{1-a^2-b^2+2ib}{(1+a)^2 + b^2} = \frac{1-(a^2+b^2)+2ib}{1+a^2+2a+b^2}$$

$$\Rightarrow ix = \frac{1 - (a^2 + b^2) + 2ib}{1 + 2a + (a^2 + b^2)}$$

$$\Rightarrow ix = \frac{1 - 1 + 2ib}{1 + 2a + 1} [\because a^2 + b^2 = 1]$$

$$\Rightarrow ix = \frac{2ib}{2 + 2a}$$

$$\Rightarrow x = \frac{2b}{2 + 2a} = \text{Real value}$$

EXERCISE 5D

PAGE: 202

Q. 1. Find the modulus of each of the following complex numbers and hence express each of them in polar form: 4

Solution: Let $Z = 4 = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$4 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$0 = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$16 = r^2$$

Since r is always a positive no., therefore,

$$r = 4,$$

Hence its modulus is 4.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{0}{4}$$

$$\tan\theta = 0$$

Since $\cos\theta = 1$, $\sin\theta = 0$ and $\tan\theta = 0$. Therefore the θ lies in first quadrant.

$$\tan\theta = 0, \text{ therefore } \theta = 0^\circ$$

Representing the complex no. in its polar form will be

$$Z = 4(\cos 0^\circ + i\sin 0^\circ)$$

Q. 2. Find the modulus of each of the following complex numbers and hence express each of them in polar form: -2

Solution: Let $Z = -2 = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$-2 = r\cos\theta \dots\dots\dots\text{eq.1}$$

$$0 = r\sin\theta \dots\dots\dots\text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$4 = r^2$$

Since r is always a positive no, therefore,

$$r = 2,$$

Hence its modulus is 2.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{0}{-2}$$

$$\tan\theta = 0$$

Since $\cos\theta = -1$, $\sin\theta = 0$ and $\tan\theta = 0$. Therefore the θ lies in second quadrant.

$$\tan\theta = 0, \text{ therefore } \theta = \pi$$

Representing the complex no. in its polar form will be

$$Z = 2(\cos\pi + i\sin\pi)$$

Q. 3. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $-i$

Solution: Let $Z = -i = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$0 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$-1 = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$1 = r^2$$

Since r is always a positive no., therefore,

$$r = 1,$$

Hence its modulus is 1.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{-1}{0}$$

$$\tan\theta = -\infty$$

Since $\cos\theta = 0$, $\sin\theta = -1$ and $\tan\theta = -\infty$. Therefore the θ lies in fourth quadrant.

$$\tan\theta = -\infty, \text{ therefore } \theta = -\frac{\pi}{2}$$

Representing the complex no. in its polar form will be

$$Z = 1\left\{\cos\left(-\frac{\pi}{2}\right) + i\sin\left(-\frac{\pi}{2}\right)\right\}$$

Q. 4. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $2i$

Solution: Let $Z = 2i = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$0 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$2 = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$4 = r^2$$

Since r is always a positive no., therefore,

$$r = 2,$$

Hence its modulus is 2.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{2}{0}$$

$$\tan \theta = \infty$$

Since $\cos \theta = 0$, $\sin \theta = 1$ and $\tan \theta = \infty$. Therefore the θ lies in first quadrant.

$$\tan \theta = \infty, \text{ therefore } \theta = \frac{\pi}{2}$$

Representing the complex no. in its polar form will be

$$Z = 2\left\{\cos\left(\frac{\pi}{2}\right) + i\sin\left(\frac{\pi}{2}\right)\right\}$$

Q. 5. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $1 - i$

Solution: Let $Z = 1 - i = r(\cos \theta + i \sin \theta)$

Now, separating real and complex part, we get

$$1 = r \cos \theta \dots\dots\dots \text{eq.1}$$

$$-1 = r \sin \theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$2 = r^2$$

Since r is always a positive no., therefore,

$$r = \sqrt{2},$$

Hence its modulus is $\sqrt{2}$.

Now , dividing eq.2 by eq.1 , we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{-1}{1}$$

$$\tan \theta = -1$$

Since $\cos \theta = \frac{1}{\sqrt{2}}$, $\sin \theta = -\frac{1}{\sqrt{2}}$ and $\tan \theta = -1$. Therefore the θ lies in fourth quadrant.

$$\tan \theta = -1, \text{ therefore } \theta = -\frac{\pi}{4}$$

Representing the complex no. in its polar form will be

$$Z = \sqrt{2} \left\{ \cos \left(-\frac{\pi}{4} \right) + i \sin \left(-\frac{\pi}{4} \right) \right\}$$

Q. 6. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $-1 + i$

Solution: Let $Z = 1 - i = r(\cos \theta + i \sin \theta)$

Now, separating real and complex part, we get

$$-1 = r \cos \theta \dots\dots\dots \text{eq.1}$$

$$1 = r \sin \theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$2 = r^2$$

Since r is always a positive no., therefore,

$$r = \sqrt{2},$$

Hence its modulus is $\sqrt{2}$.

Now, dividing eq.2 by eq.1 , we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{1}{-1}$$

$$\tan\theta = -1$$

Since $\cos\theta = -\frac{1}{\sqrt{2}}$, $\sin\theta = \frac{1}{\sqrt{2}}$ and $\tan\theta = -1$. Therefore the θ lies in second quadrant.

$$\tan\theta = -1, \text{ therefore } \theta = \frac{3\pi}{4}$$

Representing the complex no. in its polar form will be

$$Z = \sqrt{2}\left\{\cos\left(\frac{3\pi}{4}\right) + i\sin\left(\frac{3\pi}{4}\right)\right\}$$

Q. 7. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $\sqrt{3} + i$

Solution: Let $Z = \sqrt{3} + i = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$\sqrt{3} = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$1 = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$4 = r^2$$

Since r is always a positive no., therefore,

$$r = 2,$$

Hence its modulus is 2.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{1}{\sqrt{3}}$$

$$\tan\theta = \frac{1}{\sqrt{3}}$$

Since $\cos\theta = \frac{\sqrt{3}}{2}$, $\sin\theta = \frac{1}{2}$ and $\tan\theta = \frac{1}{\sqrt{3}}$. Therefore the θ lies in first quadrant.

$$\tan\theta = \frac{1}{\sqrt{3}}, \text{ therefore } \theta = \frac{\pi}{6}$$

Representing the complex no. in its polar form will be

$$Z = 2\left\{\cos\left(\frac{\pi}{6}\right) + i\sin\left(\frac{\pi}{6}\right)\right\}$$

Q. 8. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $-1 + \sqrt{3}i$

Solution: Let $Z = \sqrt{3}i - 1 = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$-1 = r\cos\theta \dots\dots\dots\text{eq.1}$$

$$\sqrt{3} = r\sin\theta \dots\dots\dots\text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$4 = r^2$$

Since r is always a positive no., therefore,

$$r = 2,$$

Hence its modulus is 2.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{\sqrt{3}}{-1}$$

$$\tan\theta = -\frac{\sqrt{3}}{1}$$

Since $\cos\theta = -\frac{1}{2}$, $\sin\theta = \frac{\sqrt{3}}{2}$ and $\tan\theta = -\frac{\sqrt{3}}{1}$. therefore the θ lies in second quadrant.

$$\tan\theta = -\sqrt{3}, \text{ therefore } \theta = \frac{2\pi}{3}$$

Representing the complex no. in its polar form will be

$$Z = 2\left\{\cos\left(\frac{2\pi}{3}\right) + i\sin\left(\frac{2\pi}{3}\right)\right\}$$

Q. 9. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $1 - \sqrt{3}i$

Solution: Let $Z = -\sqrt{3}i + 1 = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$1 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$-\sqrt{3} = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$4 = r^2$$

Since r is always a positive no., therefore,

$$r = 2,$$

Hence its modulus is 2.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{-\sqrt{3}}{1}$$

$$\tan\theta = -\frac{\sqrt{3}}{1}$$

Since $\cos\theta = \frac{1}{2}$, $\sin\theta = -\frac{\sqrt{3}}{2}$ and $\tan\theta = -\frac{\sqrt{3}}{1}$. Therefore the θ lies in the fourth quadrant.

$$\tan\theta = -\sqrt{3}, \text{ therefore } \theta = -\frac{\pi}{3}$$

Representing the complex no. in its polar form will be

$$Z = 2\left\{\cos\left(-\frac{\pi}{3}\right) + i\sin\left(-\frac{\pi}{3}\right)\right\}$$

Q. 10. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $2 - 2i$

Solution: Let $Z = 2 - 2i = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$2 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$-2 = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$8 = r^2$$

Since r is always a positive no. therefore,

$$r = 2\sqrt{2},$$

Hence its modulus is $2\sqrt{2}$.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{-2}{2}$$

$$\tan\theta = -1$$

Since $\cos\theta = \frac{1}{\sqrt{2}}$, $\sin\theta = -\frac{1}{\sqrt{2}}$ and $\tan\theta = -1$. Therefore the θ lies in the fourth quadrant.

Tan θ = -1, therefore $\theta = -\frac{\pi}{4}$

Representing the complex no. in its polar form will be

$$Z = 2\sqrt{2}\left\{\cos\left(-\frac{\pi}{4}\right) + i\sin\left(-\frac{\pi}{4}\right)\right\}$$

Q. 11. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $-4 + 4\sqrt{3}i$

Solution: Let $Z = 4\sqrt{2}i - 4 = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$-4 = r\cos\theta \dots\dots\dots\text{eq.1}$$

$$4\sqrt{3} = r\sin\theta \dots\dots\dots\text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$64 = r^2$$

Since r is always a positive no., therefore,

$$r = 8$$

Hence its modulus is 8.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{4\sqrt{3}}{-4}$$

$$\text{Tan}\theta = -\frac{\sqrt{3}}{1}$$

Since $\cos\theta = -\frac{1}{2}$, $\sin\theta = \frac{\sqrt{3}}{2}$ and $\tan\theta = -\frac{\sqrt{3}}{1}$. Therefore the θ lies in second the quadrant.

$$\text{Tan}\theta = -\sqrt{3}, \text{ therefore } \theta = \frac{2\pi}{3}.$$

Representing the complex no. in its polar form will be

$$Z = 8\left\{\cos\left(\frac{2\pi}{3}\right) + i\sin\left(\frac{2\pi}{3}\right)\right\}$$

Q. 12. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $-3\sqrt{2} + 3\sqrt{2}i$

Solution: Let $Z = -3\sqrt{2} + 3\sqrt{2}i = r(\cos\theta + i \sin\theta)$

Now, separating real and complex part, we get

$$-3\sqrt{2} = r\cos\theta \dots\dots\dots\text{eq.1}$$

$$3\sqrt{2} = r\sin\theta \dots\dots\dots\text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$36 = r^2$$

Since r is always a positive no., therefore,

$$r = 6$$

Hence its modulus is 6.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{3\sqrt{2}}{-3\sqrt{2}}$$

$$\text{Tan}\theta = -\frac{1}{1}$$

Since $\cos\theta = -\frac{1}{\sqrt{2}}$, $\sin\theta = \frac{1}{\sqrt{2}}$ and $\tan\theta = -1$. therefore the θ lies in second quadrant.

$$\text{Tan}\theta = -1, \text{ therefore } \theta = \frac{3\pi}{4}.$$

Representing the complex no. in its polar form will be

$$Z = 6\left\{\cos\left(\frac{3\pi}{4}\right) + i\sin\left(\frac{3\pi}{4}\right)\right\}$$

Q. 13. Find the modulus of each of the following complex numbers and hence

express each of them in polar form: $\frac{1+i}{1-i}$

Solution: $\frac{1+i}{1-i} = \frac{1+i}{1-i} \times \frac{1+i}{1+i}$

$$= \frac{1+i^2+2i}{1-i^2}$$

$$= \frac{2i}{2}$$

$$= i$$

Let $Z = i = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$0 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$1 = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$1 = r^2$$

Since r is always a positive no., therefore,

$$r = 1,$$

Hence its modulus is 1.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{1}{0}$$

$$\tan\theta = \infty$$

Since $\cos\theta = 0$, $\sin\theta = 1$ and $\tan\theta = \infty$. Therefore the θ lies in first quadrant.

$$\tan\theta = \infty, \text{ therefore } \theta = \frac{\pi}{2}$$

Representing the complex no. in its polar form will be

$$Z = 1\left\{\cos\left(\frac{\pi}{2}\right) + i\sin\left(\frac{\pi}{2}\right)\right\}$$

Q. 14. Find the modulus of each of the following complex numbers and hence

express each of them in polar form: $\frac{1-i}{1+i}$

$$\text{Solution: } \frac{1-i}{1+i} = \frac{1-i}{1+i} \times \frac{1-i}{1-i}$$

$$= \frac{1+i^2-2i}{1-i^2}$$

$$= -\frac{2i}{2}$$

$$= -i$$

$$\text{Let } Z = -i = r(\cos\theta + i\sin\theta)$$

Now, separating real and complex part, we get

$$0 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$-1 = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$1 = r^2$$

Since r is always a positive no., therefore,

$$r = 1,$$

Hence its modulus is 1.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{-1}{0}$$

$$\tan \theta = -\infty$$

Since $\cos \theta = 0$, $\sin \theta = -1$ and $\tan \theta = -\infty$, therefore the θ lies in fourth quadrant.

$$\tan \theta = -\infty, \text{ therefore } \theta = -\frac{\pi}{2}$$

Representing the complex no. in its polar form will be

$$Z = 1\left\{\cos\left(-\frac{\pi}{2}\right) + i\sin\left(-\frac{\pi}{2}\right)\right\}$$

Q. 15. Find the modulus of each of the following complex numbers and hence

express each of them in polar form: $\frac{1+3i}{1-2i}$

Solution:

$$\frac{1+3i}{1-2i} = \frac{1+3i}{1-2i} \times \frac{1+2i}{1+2i}$$

$$= \frac{1+6i^2+5i}{1-4i^2}$$

$$= \frac{5i-5}{5}$$

$$= i-1$$

$$\text{Let } Z = 1 - i = r(\cos \theta + i \sin \theta)$$

Now, separating real and complex part, we get

$$-1 = r \cos \theta \dots \dots \dots \text{eq.1}$$

$$1 = r \sin \theta \dots \dots \dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$2 = r^2$$

Since r is always a positive no., therefore,

$$r = \sqrt{2},$$

Hence its modulus is $\sqrt{2}$.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{1}{-1}$$

$$\tan \theta = -1$$

Since $\cos \theta = -\frac{1}{\sqrt{2}}$, $\sin \theta = \frac{1}{\sqrt{2}}$ and $\tan \theta = -1$. Therefore the θ lies in second quadrant.

$$\tan \theta = -1, \text{ therefore } \theta = \frac{3\pi}{4}$$

Representing the complex no. in its polar form will be

$$Z = \sqrt{2} \left\{ \cos\left(\frac{3\pi}{4}\right) + i \sin\left(\frac{3\pi}{4}\right) \right\}$$

Q. 16. Find the modulus of each of the following complex numbers and hence

express each of them in polar form: $\frac{1-3i}{1+2i}$

Solution:

$$\frac{1-3i}{1+2i} = \frac{1-3i}{1+2i} \times \frac{1-2i}{1-2i}$$

$$= \frac{1 + 6i^2 - 5i}{1 - 4i^2}$$

$$= \frac{-5i - 5}{5}$$

$$= -i - 1$$

Let $Z = -1 - i = r(\cos \theta + i \sin \theta)$

Now , separating real and complex part , we get

$$-1 = r\cos\theta \dots\dots\dots\text{eq.1}$$

$$-1 = r\sin\theta \dots\dots\dots\text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$2 = r^2$$

Since r is always a positive no., therefore,

$$r = \sqrt{2},$$

Hence its modulus is $\sqrt{2}$.

Now , dividing eq.2 by eq.1 , we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{-1}{-1}$$

$$\tan\theta = 1$$

Since $\cos\theta = -\frac{1}{\sqrt{2}}$, $\sin\theta = -\frac{1}{\sqrt{2}}$ and $\tan\theta = 1$. Therefore the θ lies in third quadrant.

$$\text{Tan}\theta = 1, \text{ therefore } \theta = -\frac{3\pi}{4}$$

Representing the complex no. in its polar form will be

$$Z = \sqrt{2}\left\{\cos\left(-\frac{3\pi}{4}\right) + i\sin\left(-\frac{3\pi}{4}\right)\right\}$$

Q. 17. Find the modulus of each of the following complex numbers and hence

$$\frac{5-i}{2-3i}$$

express each of them in polar form:

Solution:

$$\frac{5-i}{2-3i} = \frac{5-i}{2-3i} \times \frac{2+3i}{2+3i}$$

$$= \frac{10 - 3i^2 + 13i}{4 - 9i^2}$$

$$= \frac{+13i + 13}{13}$$

$$= i + 1$$

Let $Z = 1 + i = r(\cos\theta + i\sin\theta)$

Now , separating real and complex part , we get

$$1 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$1 = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$2 = r^2$$

Since r is always a positive no., therefore,

$$r = \sqrt{2},$$

Hence its modulus is $\sqrt{2}$.

Now , dividing eq.2 by eq.1 , we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{1}{1}$$

$$\text{Tan}\theta = 1$$

Since $\cos\theta = \frac{1}{\sqrt{2}}$, $\sin\theta = \frac{1}{\sqrt{2}}$ and $\tan\theta = 1$. Therefore the θ lies in first quadrant.

$$\text{Tan}\theta = 1, \text{ therefore } \theta = \frac{\pi}{4}$$

Representing the complex no. in its polar form will be

$$Z = \sqrt{2}\left\{\cos\left(\frac{\pi}{4}\right) + i\sin\left(\frac{\pi}{4}\right)\right\}$$

Q. 18. Find the modulus of each of the following complex numbers and hence

express each of them in polar form: $\frac{-16}{1 + \sqrt{3}i}$

Solution:

$$\frac{-16}{1 + \sqrt{3}i} = \frac{-16}{1 + \sqrt{3}i} \times \frac{1 - \sqrt{3}i}{1 - \sqrt{3}i}$$

$$= \frac{-16 + 16\sqrt{3}i}{1 - 3i^2}$$

$$= \frac{16\sqrt{3}i - 16}{4}$$

$$= 4\sqrt{3}i - 4$$

Let $Z = 4\sqrt{3}i - 4 = r(\cos\theta + i\sin\theta)$

Now , separating real and complex part , we get

$$-4 = r\cos\theta \dots\dots\dots\text{eq.1}$$

$$4\sqrt{3} = r\sin\theta \dots\dots\dots\text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$64 = r^2$$

Since r is always a positive no., therefore,

$$r = 8,$$

Hence its modulus is 8.

Now, dividing eq.2 by eq.1 , we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{4\sqrt{3}}{-4}$$

$$\tan\theta = -\sqrt{3}$$

Since $\cos\theta = -\frac{1}{2}$, $\sin\theta = \frac{\sqrt{3}}{2}$ and $\tan\theta = -\sqrt{3}$. Therefore the θ lies in second quadrant.

$$\tan\theta = -\sqrt{3}, \text{ therefore } \theta = \frac{2\pi}{3}$$

Representing the complex no. in its polar form will be

$$Z = 8\left\{\cos\left(\frac{2\pi}{3}\right) + i\sin\left(\frac{2\pi}{3}\right)\right\}$$

Q. 19. Find the modulus of each of the following complex numbers and hence

$$\frac{2 + 6\sqrt{3}i}{5 + \sqrt{3}i}$$

express each of them in polar form:

Solution:

$$\frac{2 + 6\sqrt{3}i}{5 + \sqrt{3}i} = \frac{2 + 6\sqrt{3}i}{5 + \sqrt{3}i} \times \frac{5 - \sqrt{3}i}{5 - \sqrt{3}i}$$

$$= \frac{10 + 28\sqrt{3}i - 18i^2}{25 - 3i^2}$$

$$= \frac{28\sqrt{3}i + 28}{28}$$

$$= \sqrt{3}i + 1$$

$$\text{Let } Z = \sqrt{3}i + 1 = r(\cos\theta + i\sin\theta)$$

Now, separating real and complex part, we get

$$1 = r\cos\theta \dots\dots\dots \text{eq.1}$$

$$\sqrt{3} = r\sin\theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$4 = r^2$$

Since r is always a positive no., therefore,

$$r = 2,$$

Hence its modulus is 2.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{\sqrt{3}}{1}$$

$$\tan \theta = \sqrt{3}$$

Since $\cos \theta = \frac{1}{2}$, $\sin \theta = \frac{\sqrt{3}}{2}$ and $\tan \theta = \sqrt{3}$. therefore the θ lies in first quadrant.

$$\tan \theta = \sqrt{3}, \text{ therefore } \theta = \frac{\pi}{3}$$

Representing the complex no. in its polar form will be

$$Z = 2\left\{\cos\left(\frac{\pi}{3}\right) + i\sin\left(\frac{\pi}{3}\right)\right\}$$

Q. 20

Find the modulus of each of the following complex numbers and hence express each of them in polar form:

$$\sqrt{\frac{1+i}{1-i}}$$

Solution:

$$\sqrt{\frac{1+i}{1-i}} = \sqrt{\frac{1+i}{1-i}} \times \sqrt{\frac{1+i}{1+i}}$$

$$= \sqrt{\frac{(1+i)^2}{1-i^2}}$$

$$= \frac{1+i}{\sqrt{2}}$$

$$= \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}}$$

Let $Z = \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}} = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part , we get

$$\frac{1}{\sqrt{2}} = r\cos\theta \quad \dots\dots\dots\text{eq.1}$$

$$\frac{1}{\sqrt{2}} = r\sin\theta \quad \dots\dots\dots\text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$1 = r^2$$

Since r is always a positive no., therefore,

$$r = 1,$$

hence its modulus is 1.

now , dividing eq.2 by eq.1 , we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{\frac{i}{\sqrt{2}}}{\frac{i}{\sqrt{2}}}$$

$$\tan \theta = 1$$

Since $\cos \theta = \frac{1}{\sqrt{2}}$, $\sin \theta = \frac{1}{\sqrt{2}}$ and $\tan \theta = 1$. therefore the θ lies in first quadrant.

Tan θ = 1, therefore $\theta = \frac{\pi}{4}$

Representing the complex no. in its polar form will be

$$Z = 1\left\{\cos\left(\frac{\pi}{4}\right) + i\sin\left(\frac{\pi}{4}\right)\right\}$$

Q. 20. Find the modulus of each of the following complex numbers and hence

express each of them in polar form: $\sqrt{\frac{1+i}{1-i}}$

Solution:

$$\sqrt{\frac{1+i}{1-i}} = \sqrt{\frac{1+i}{1-i}} \times \sqrt{\frac{1+i}{1+i}}$$

$$= \sqrt{\frac{(1+i)^2}{1-i^2}}$$

$$= \frac{1+i}{\sqrt{2}}$$

$$= \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}}$$

Let $Z = \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}} = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part, we get

$$\frac{1}{\sqrt{2}} = r\cos\theta \quad \dots\dots\dots\text{eq.1}$$

$$\frac{1}{\sqrt{2}} = r \sin \theta \quad \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$1 = r^2$$

Since r is always a positive no., therefore,

$$r = 1,$$

Hence its modulus is 1.

Now , dividing eq.2 by eq.1 , we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{\frac{i}{\sqrt{2}}}{\frac{1}{\sqrt{2}}}$$

$$\tan \theta = 1$$

Since $\cos \theta = \frac{1}{\sqrt{2}}$, $\sin \theta = \frac{1}{\sqrt{2}}$ and $\tan \theta = 1$. Therefore the θ lies in first quadrant.

$$\tan \theta = 1, \text{ therefore } \theta = \frac{\pi}{4}$$

Representing the complex no. in its polar form will be

$$Z = 1 \left\{ \cos\left(\frac{\pi}{4}\right) + i \sin\left(\frac{\pi}{4}\right) \right\}$$

Q. 21. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $-\sqrt{3} - i$

Solution: Let $Z = -\sqrt{3} - i = r(\cos \theta + i \sin \theta)$

Now, separating real and complex part, we get

$$-\sqrt{3} = r \cos \theta \quad \dots\dots\dots \text{eq.1}$$

$$-1 = r \sin \theta \quad \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$4 = r^2$$

Since r is always a positive no., therefore,

$$r = 2$$

Hence its modulus is 2.

Now, dividing eq.2 by eq.1, we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{-1}{-\sqrt{3}}$$

$$\tan \theta = \frac{1}{\sqrt{3}}$$

Since $\cos \theta = -\frac{\sqrt{3}}{2}$, $\sin \theta = -\frac{1}{2}$ and $\tan \theta = \frac{1}{\sqrt{3}}$. Therefore the θ lies in third quadrant.

$$\tan \theta = \frac{1}{\sqrt{3}}, \text{ therefore } \theta = -\frac{5\pi}{6}.$$

Representing the complex no. in its polar form will be

$$Z = 2\left\{\cos\left(-\frac{5\pi}{6}\right) + i\sin\left(-\frac{5\pi}{6}\right)\right\}$$

Q. 22. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $(i^{25})^3$

Solution: $= i^{75}$

$$= i^{4n+3} \text{ where } n = 18$$

$$\text{Since } i^{4n+3} = -i$$

$$i^{75} = -i$$

$$\text{Let } Z = -i = r(\cos \theta + i \sin \theta)$$

Now, separating real and complex part, we get

$$0 = r \cos \theta \dots\dots\dots \text{eq.1}$$

$$-1 = r \sin \theta \dots\dots\dots \text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$1 = r^2$$

Since r is always a positive no., therefore,

$$r = 1,$$

Hence its modulus is 1.

Now , dividing eq.2 by eq.1 , we get,

$$\frac{r \sin \theta}{r \cos \theta} = \frac{-1}{0}$$

$$\tan \theta = -\infty$$

Since $\cos \theta = 0$, $\sin \theta = -1$ and $\tan \theta = -\infty$. therefore the θ lies in fourth quadrant.

$$\tan \theta = -\infty , \text{ therefore } \theta = -\frac{\pi}{2}$$

Representing the complex no. in its polar form will be

$$Z = 1 \left\{ \cos\left(-\frac{\pi}{2}\right) + i \sin\left(-\frac{\pi}{2}\right) \right\}$$

Q. 23. Find the modulus of each of the following complex numbers and hence

$$\frac{(1-i)}{\left(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right)}$$

express each of them in polar form:

Solution:

$$= \frac{1-i}{\frac{1}{2} + i \frac{\sqrt{3}}{2}}$$

$$= \frac{2-2i}{1+i\sqrt{3}}$$

$$\begin{aligned}
 &= \frac{2 - 2i}{1 + \sqrt{3}i} \times \frac{1 - \sqrt{3}i}{1 - \sqrt{3}i} \\
 &= \frac{2 - 2\sqrt{3}i - 2i + 2\sqrt{3}i^2}{1 - 3i^2} \\
 &= \frac{(2 - 2\sqrt{3}) + i(2\sqrt{3} + 2)}{4} \\
 &= \frac{(1 - \sqrt{3}) + i(\sqrt{3} + 1)}{2}
 \end{aligned}$$

Let $Z = \frac{(1 - \sqrt{3}) + i(\sqrt{3} + 1)}{2} = r(\cos\theta + i\sin\theta)$

Now, separating real and complex part , we get

$$\frac{1 - \sqrt{3}}{2} = r\cos\theta \quad \dots\dots\dots\text{eq.1}$$

$$\frac{1 + \sqrt{3}}{2} = r\sin\theta \quad \dots\dots\dots\text{eq.2}$$

Squaring and adding eq.1 and eq.2, we get

$$2 = r^2$$

Since r is always a positive no., therefore,

$$r = \sqrt{2},$$

Hence its modulus is $\sqrt{2}$.

Now, dividing eq.2 by eq.1 , we get,

$$\frac{r\sin\theta}{r\cos\theta} = \frac{\frac{1 + \sqrt{3}}{2}}{\frac{1 - \sqrt{3}}{2}}$$

$$\tan\theta = \frac{1 + \sqrt{3}}{1 - \sqrt{3}}$$

Since $\cos\theta = \frac{1-\sqrt{3}}{2\sqrt{2}}$, $\sin\theta = \frac{1+\sqrt{3}}{2\sqrt{2}}$ and $\tan\theta = \frac{1+\sqrt{3}}{1-\sqrt{3}}$. Therefore the θ lies in second quadrant. As

$$\tan\theta = \frac{1+\sqrt{3}}{1-\sqrt{3}}, \text{ therefore } \theta = \frac{7\pi}{12}$$

Representing the complex no. in its polar form will be

$$Z = \sqrt{2}\left\{\cos\left(\frac{7\pi}{12}\right) + i\sin\left(\frac{7\pi}{12}\right)\right\}$$

Q. 24. Find the modulus of each of the following complex numbers and hence express each of them in polar form: $(\sin 120^\circ - i \cos 120^\circ)$

Solution: $= \sin(90^\circ + 30^\circ) - i\cos(90^\circ + 30^\circ) = \cos 30^\circ + i\sin 30^\circ$

Since, $\sin(90^\circ + \alpha) = \cos\alpha$

And $\cos(90^\circ + \alpha) = -\sin\alpha$

$$= \frac{\sqrt{3}}{2} + i\frac{1}{2}$$

Hence it is of the form

$$Z = \frac{\sqrt{3}}{2} + i\frac{1}{2} = r(\cos\theta + i\sin\theta)$$

Therefore $r = 1$

Hence its modulus is 1 and argument is $\frac{\pi}{6}$.

EXERCISE 5E

Q. 1. $x^2 + 2 = 0$

Solution: This equation is a quadratic equation.

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Given:

$$\Rightarrow x^2 + 2 = 0$$

$$\Rightarrow x^2 = -2$$

$$\Rightarrow x = \pm \sqrt{-2}$$

But we know that $\sqrt{-1} = i$

$$\Rightarrow x = \pm \sqrt{2} i$$

Ans: $x = \pm \sqrt{2} i$ Q.

Q.2. $x^2 + 5 = 0$

Solution :

Given: $x^2 + 5 = 0$

$$\Rightarrow x^2 = -5$$

$$\Rightarrow x = \pm \sqrt{-5}$$

$$\Rightarrow x = \pm \sqrt{5} i$$

Ans: $x = \pm \sqrt{5} i$

Q. 3. $2x^2 + 1 = 0$

Solution: $2x^2 + 1 = 0$

$$\Rightarrow 2x^2 = -1$$

$$\Rightarrow x^2 = -\frac{1}{2}$$

$$\Rightarrow x = \pm \sqrt{-\frac{1}{2}}$$

$$\Rightarrow x = \pm \sqrt{\frac{1}{2}}i$$

$$\Rightarrow x = \pm \frac{i}{\sqrt{2}}$$

Ans: $x = \pm \frac{i}{\sqrt{2}}$

Q. 4. $x^2 + x + 1 = 0$

Solution: Given:

$$x^2 + x + 1 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{1^2 - (4 \times 1 \times 1)}}{2 \times 1}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{1-4}}{2}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{-3}}{2}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{3}i}{2}$$

$$\Rightarrow x = -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$$

Ans: $x = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$ and $x = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$

Q. 5. $x^2 - x + 2 = 0$

Solution: Given:

$$x^2 - x + 2 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-(-1) \pm \sqrt{(-1)^2 - (4 \times 1 \times 2)}}{2 \times 1}$$

$$\Rightarrow x = \frac{1 \pm \sqrt{1-8}}{2}$$

$$\Rightarrow x = \frac{1 \pm \sqrt{-7}}{2}$$

$$\Rightarrow x = \frac{1 \pm \sqrt{7}i}{2}$$

$$\Rightarrow x = \frac{1}{2} \pm \frac{\sqrt{7}}{2}i$$

Ans: $x = \frac{1}{2} + \frac{\sqrt{7}}{2}i$ and $x = \frac{1}{2} - \frac{\sqrt{7}}{2}i$

Q. 6. $x^2 + 2x + 2 = 0$

Solution: Given:

$$x^2 + 2x + 2 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-2 \pm \sqrt{(2)^2 - (4 \times 1 \times 2)}}{2 \times 1}$$

$$\Rightarrow x = \frac{-2 \pm \sqrt{4-8}}{2}$$

$$\Rightarrow x = \frac{-2 \pm \sqrt{-4}}{2}$$

$$\Rightarrow x = \frac{-2 \pm 2i}{2}$$

$$\Rightarrow x = -\frac{2}{2} \pm \frac{2}{2}i$$

$$\Rightarrow x = -1 \pm i$$

Ans: $x = -1 + i$ and $x = -1 - i$

Q. 7. $2x^2 - 4x + 3 = 0$

Solution: Given:

$$2x^2 - 4x + 3 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-(-4) \pm \sqrt{(-4)^2 - (4 \times 2 \times 3)}}{2 \times 2}$$

$$\Rightarrow x = \frac{4 \pm \sqrt{16 - 24}}{4}$$

$$\Rightarrow x = \frac{4 \pm \sqrt{-8}}{4}$$

$$\Rightarrow x = \frac{4 \pm 2\sqrt{2}i}{4}$$

$$\Rightarrow x = \frac{4}{4} \pm \frac{2\sqrt{2}}{4}i$$

$$\Rightarrow x = 1 \pm \frac{i}{\sqrt{2}}$$

Ans: $x = 1 + \frac{i}{\sqrt{2}}$ and $x = 1 - \frac{i}{\sqrt{2}}$

Q. 8. $x^2 + 3x + 5 = 0$

Solution: Given:

$$x^2 + 3x + 5 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-3 \pm \sqrt{(3)^2 - (4 \times 1 \times 5)}}{2 \times 1}$$

$$\Rightarrow x = \frac{-3 \pm \sqrt{9 - 20}}{2}$$

$$\Rightarrow x = \frac{-3 \pm \sqrt{-11}}{2}$$

$$\Rightarrow x = \frac{-3 \pm \sqrt{11}i}{2}$$

$$\Rightarrow x = -\frac{3}{2} \pm \frac{\sqrt{11}}{2}i$$

Ans: $x = -\frac{3}{2} + \frac{\sqrt{11}}{2}i$ and $x = -\frac{3}{2} - \frac{\sqrt{11}}{2}i$

Q. 9. $\sqrt{5}x^2 + x + \sqrt{5} = 0$

Solution: Given:

$$\sqrt{5}x^2 + x + \sqrt{5} = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{(1)^2 - (4 \times \sqrt{5} \times \sqrt{5})}}{2 \times \sqrt{5}}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{1-20}}{2\sqrt{5}}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{-19}}{2\sqrt{5}}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{19}i}{2\sqrt{5}}$$

$$\Rightarrow x = -\frac{1}{2\sqrt{5}} \pm \frac{\sqrt{19}}{2\sqrt{5}}i$$

$$\text{Ans: } x = -\frac{\sqrt{5}}{10} + \frac{\sqrt{\frac{19}{5}}}{2}i \text{ and } x = -\frac{\sqrt{5}}{10} - \frac{\sqrt{\frac{19}{5}}}{2}i$$

Q. 10. $25x^2 - 30x + 11 = 0$

Solution: Given:

$$25x^2 - 30x + 11 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-(-30) \pm \sqrt{(-30)^2 - (4 \times 25 \times 11)}}{2 \times 25}$$

$$\Rightarrow x = \frac{30 \pm \sqrt{900 - 1100}}{50}$$

$$\Rightarrow x = \frac{30 \pm \sqrt{-200}}{50}$$

$$\Rightarrow x = \frac{30 \pm 10\sqrt{2}i}{50}$$

$$\Rightarrow x = -\frac{30}{50} \pm \frac{10\sqrt{2}}{50}i$$

$$\text{Ans: } x = -\frac{3}{5} + \frac{\sqrt{2}}{5}i \text{ and } x = -\frac{3}{5} - \frac{\sqrt{2}}{5}i$$

Q. 11. $8x^2 + 2x + 1 = 0$

Solution: Given:

$$8x^2 + 2x + 1 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-2 \pm \sqrt{(2)^2 - (4 \times 8 \times 1)}}{2 \times 8}$$

$$\Rightarrow x = \frac{-2 \pm \sqrt{4 - 32}}{16}$$

$$\Rightarrow x = \frac{-2 \pm \sqrt{-28}}{16}$$

$$\Rightarrow x = \frac{-2 \pm 2\sqrt{7}i}{16}$$

$$\Rightarrow x = -\frac{2}{16} \pm \frac{2\sqrt{7}}{16}i$$

$$\text{Ans: } x = -\frac{1}{8} + \frac{\sqrt{7}}{8}i \text{ and } x = -\frac{1}{8} - \frac{\sqrt{7}}{8}i$$

Q. 12. $27x^2 + 10x + 1 = 0$

Solution:

Given:

$$27x^2 + 10x + 1 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-10 \pm \sqrt{(10)^2 - (4 \times 27 \times 1)}}{2 \times 27}$$

$$\Rightarrow x = \frac{-10 \pm \sqrt{100 - 108}}{54}$$

$$\Rightarrow x = \frac{-10 \pm \sqrt{-8}}{54}$$

$$\Rightarrow x = \frac{-10 \pm 2\sqrt{2}i}{54}$$

$$\Rightarrow x = -\frac{10}{54} \pm \frac{2\sqrt{2}}{54}i$$

Ans: $x = -\frac{5}{27} + \frac{\sqrt{2}}{27}i$ and $x = -\frac{5}{27} - \frac{\sqrt{2}}{27}i$

Q. 13. $2x^2 - \sqrt{3}x + 1 = 0$

Solution: Given:

$$2x^2 - \sqrt{3}x + 1 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-(-\sqrt{3}) \pm \sqrt{(-\sqrt{3})^2 - (4 \times 2 \times 1)}}{2 \times 2}$$

$$\Rightarrow x = \frac{\sqrt{3} \pm \sqrt{3-8}}{4}$$

$$\Rightarrow x = \frac{\sqrt{3} \pm \sqrt{-5}}{4}$$

$$\Rightarrow x = \frac{\sqrt{3} \pm \sqrt{5}i}{4}$$

$$\Rightarrow x = \frac{\sqrt{3}}{4} \pm \frac{\sqrt{5}}{4}i$$

Ans: $x = \frac{\sqrt{3}}{4} + \frac{\sqrt{5}}{4}i$ and $x = \frac{\sqrt{3}}{4} - \frac{\sqrt{5}}{4}i$

Q. 14. $17x^2 - 8x + 1 = 0$

Solution: Given:

$$17x^2 - 8x + 1 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-(-8) \pm \sqrt{(-8)^2 - (4 \times 17 \times 1)}}{2 \times 17}$$

$$\Rightarrow x = \frac{8 \pm \sqrt{64 - 68}}{34}$$

$$\Rightarrow x = \frac{8 \pm \sqrt{-4}}{34}$$

$$\Rightarrow x = \frac{8 \pm 2i}{34}$$

$$\Rightarrow x = \frac{8}{34} \pm \frac{2}{34}i$$

Ans: $x = \frac{4}{17} + \frac{1}{17}i$ and $x = \frac{4}{17} - \frac{1}{17}i$

Q. 15. $3x^2 + 5 = 7x$

Solution: Given:

$$3x^2 + 5 = 7x$$

$$\Rightarrow 3x^2 - 7x + 5 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-(-7) \pm \sqrt{(-7)^2 - (4 \times 3 \times 5)}}{2 \times 3}$$

$$\Rightarrow x = \frac{7 \pm \sqrt{49 - 60}}{6}$$

$$\Rightarrow x = \frac{7 \pm \sqrt{-11}}{6}$$

$$\Rightarrow x = \frac{7 \pm \sqrt{11}i}{6}$$

$$\Rightarrow x = \frac{7}{6} \pm \frac{\sqrt{11}}{6}i$$

Ans: $x = \frac{7}{6} + \frac{\sqrt{11}}{6}i$ and $x = \frac{7}{6} - \frac{\sqrt{11}}{6}i$

Q. 16. $3x^2 - 4x + \frac{20}{3} = 0$

Solution: Given:

$$3x^2 - 4x + \frac{20}{3} = 0$$

Multiplying both the sides by 3 we get,

$$9x^2 - 12x + 20 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\Rightarrow x = \frac{-(-12) \pm \sqrt{(-12)^2 - (4 \times 9 \times 20)}}{2 \times 9}$$

$$\Rightarrow x = \frac{12 \pm \sqrt{144 - 720}}{18}$$

$$\Rightarrow x = \frac{12 \pm \sqrt{-576}}{18}$$

$$\Rightarrow x = \frac{12 \pm 24i}{18}$$

$$\Rightarrow x = \frac{12}{18} \pm \frac{24}{18}i$$

$$\Rightarrow x = \frac{2}{3} \pm \frac{4}{3}i$$

Ans: $x = \frac{2}{3} + \frac{4}{3}i$ and $x = \frac{2}{3} - \frac{4}{3}i$

Q. 17. $3x^2 + 7ix + 6 = 0$

Solution: Given:

$$3x^2 + 7ix + 6 = 0$$

$$\Rightarrow 3x^2 + 9ix - 2ix + 6 = 0$$

$$\Rightarrow 3x(x + 3i) - 2i\left(x - \frac{6}{2i}\right) = 0$$

$$\Rightarrow 3x(x + 3i) - 2i\left(x - \frac{3 \times i}{i \times i}\right) = 0 \quad \dots (i^2 = -1)$$

$$\Rightarrow 3x(x + 3i) - 2i\left(x - \frac{3 \times i}{-1}\right) = 0$$

$$\Rightarrow 3x(x + 3i) - 2i(x + 3i) = 0$$

$$\Rightarrow (x + 3i)(3x - 2i) = 0$$

$$\Rightarrow x + 3i = 0 \text{ \& } 3x - 2i = 0$$

$$\Rightarrow x = 3i \text{ \& } x = \frac{2}{3}i$$

$$\text{Ans: } x = 3i \text{ and } x = \frac{2}{3}i$$

Q. 18. $21x^2 - 28x + 10 = 0$

Solution: Given:

$$21x^2 - 28x + 10 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\Rightarrow x = \frac{-(-28) \pm \sqrt{(-28)^2 - (4 \times 21 \times 10)}}{2 \times 21}$$

$$\Rightarrow x = \frac{28 \pm \sqrt{784 - 840}}{42}$$

$$\Rightarrow x = \frac{28 \pm \sqrt{-56}}{42}$$

$$\Rightarrow x = \frac{28 \pm 2\sqrt{14}i}{42}$$

$$\Rightarrow x = \frac{28}{42} \pm \frac{2\sqrt{14}}{42}i$$

$$\text{Ans: } x = \frac{2}{3} + \frac{\sqrt{14}}{21}i \text{ and } x = \frac{2}{3} - \frac{\sqrt{14}}{21}i$$

Q. 19. $x^2 + 13 = 4x$

Solution: Given:

$$x^2 + 13 = 4x$$

$$\Rightarrow x^2 - 4x + 13 = 0$$

Solution of a general quadratic equation $ax^2 + bx + c = 0$ is given by:

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$\Rightarrow x = \frac{-(-4) \pm \sqrt{(-4)^2 - (4 \times 1 \times 13)}}{2 \times 1}$$

$$\Rightarrow x = \frac{4 \pm \sqrt{16 - 52}}{2}$$

$$\Rightarrow x = \frac{4 \pm \sqrt{-36}}{2}$$

$$\Rightarrow x = \frac{4 \pm 6i}{2}$$

$$\Rightarrow x = \frac{4}{2} \pm \frac{6}{2}i$$

$$\Rightarrow x = 2 \pm 3i$$

Ans: $x = 2 + 3i$ & $x = 2 - 3i$

Q. 20. $x^2 + 3ix + 10 = 0$

Solution:

Given: $x^2 + 3ix + 10 = 0$

$$\Rightarrow x^2 + 5ix - 2ix + 10 = 0$$

$$\Rightarrow x(x + 5i) - 2i\left(x - \frac{10}{2i}\right) = 0$$

$$\Rightarrow x(x + 5i) - 2i\left(x - \frac{5 \times i}{i \times i}\right) = 0$$

$$\Rightarrow x(x + 5i) - 2i\left(x - \frac{5 \times i}{-1}\right) = 0$$

$$\Rightarrow x(x + 5i) - 2i(x + 5i) = 0$$

$$\Rightarrow (x + 5i)(x - 2i) = 0$$

$$\Rightarrow x + 5i = 0 \text{ \& } x - 2i = 0$$

$$\Rightarrow x = -5i \text{ \& } x = 2i$$

Ans: $x = -5i$ & $x = 2i$

Q. 21. $2x^2 + 3ix + 2 = 0$

Solution: Given:

$$2x^2 + 3ix + 2 = 0$$

$$\Rightarrow 2x^2 + 4ix - ix + 2 = 0$$

$$\Rightarrow 2x(x + 2i) - i\left(x - \frac{2}{i}\right) = 0$$

$$\Rightarrow 2x(x + 2i) - i\left(x - \frac{2 \times i}{i \times i}\right) = 0$$

$$\Rightarrow 2x(x + 2i) - i\left(x - \frac{2 \times i}{-1}\right) = 0$$

$$\Rightarrow 2x(x + 2i) - i(x + 2i) = 0$$

$$\Rightarrow (x + 2i)(2x - i) = 0$$

$$\Rightarrow x + 2i = 0 \text{ \& } 2x - i = 0$$

$$\Rightarrow x = -2i \text{ \& } x = \frac{i}{2}$$

Ans: $x = -2i$ and $x = \frac{i}{2}$

EXERCISE 5F

Q. 1. $\sqrt{5 + 12i}$

Solution: Let, $(a + ib)^2 = 5 + 12i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 5 + 12i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 5 + 12i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 5 \dots \dots \dots \text{eq.1}$$

$$\Rightarrow 2ab = 12 \dots \dots \dots \text{eq.2}$$

$$\Rightarrow a = \frac{6}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(\frac{6}{b}\right)^2 - b^2 = 5$$

$$\Rightarrow 36 - b^4 = 5b^2$$

$$\Rightarrow b^4 + 5b^2 - 36 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -9 \text{ or } b^2 = 4$$

As b is real no. so, $b^2 = 4$

$$b = 2 \text{ or } b = -2$$

Therefore, $a = 3$ or $a = -3$

Hence the square root of the complex no. is $3 + 2i$ and $-3 - 2i$.

Q. 2. $\sqrt{-7 + 24i}$

Solution: Let, $(a + ib)^2 = -7 + 24i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = -7 + 24i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = -7 + 24i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = -7 \dots \dots \dots \text{eq.1}$$

$$\Rightarrow 2ab = 24 \dots \dots \dots \text{eq.2}$$

$$\Rightarrow a = \frac{12}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(\frac{12}{b}\right)^2 - b^2 = -7$$

$$\Rightarrow 144 - b^4 = -7b^2$$

$$\Rightarrow b^4 - 7b^2 - 144 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -9 \text{ or } b^2 = 16$$

As b is real no. so, $b^2 = 16$

$$b = 4 \text{ or } b = -4$$

Therefore, $a = 3$ or $a = -3$

Hence the square root of the complex no. is $3 + 4i$ and $-3 - 4i$.

Q. 3. $\sqrt{-2 + 2\sqrt{3}i}$

Solution: Let, $(a + ib)^2 = -2 + 2\sqrt{3}i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = -2 + 2\sqrt{3}i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = -2 + 2\sqrt{3}i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = -2 \dots \dots \dots \text{eq.1}$$

$$\Rightarrow 2ab = 2\sqrt{3} \dots \dots \dots \text{eq.2}$$

$$\Rightarrow a = \frac{\sqrt{3}}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(\frac{\sqrt{3}}{b}\right)^2 - b^2 = -2$$

$$\Rightarrow 3 - b^4 = -2b^2$$

$$\Rightarrow b^4 - 2b^2 - 3 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -1 \text{ or } b^2 = 3$$

As b is real no. so, $b^2 = 3$

$$b = \sqrt{3} \text{ or } b = -\sqrt{3}$$

Therefore, $a = 1$ or $a = -1$

Hence the square root of the complex no. is $1 + \sqrt{3}i$ and $-1 - \sqrt{3}i$.

Q. 4. $\sqrt{1 + 4\sqrt{-3}}$

Solution: Let, $(a + ib)^2 = 1 + 4\sqrt{3}i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 1 + 4\sqrt{3}i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 1 + 4\sqrt{3}i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 1 \dots\dots\dots \text{eq.1}$$

$$\Rightarrow 2ab = 4\sqrt{3} \dots\dots\dots \text{eq.2}$$

$$\Rightarrow a = \frac{2\sqrt{3}}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(\frac{2\sqrt{3}}{b}\right)^2 - b^2 = 1$$

$$\Rightarrow 12 - b^4 = b^2$$

$$\Rightarrow b^4 + b^2 - 12 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -4 \text{ or } b^2 = 3$$

As b is real no. so, $b^2 = 3$

$$b = \sqrt{3} \text{ or } b = -\sqrt{3}$$

Therefore, $a = 2$ or $a = -2$

Hence the square root of the complex no. is $2 + \sqrt{3}i$ and $-2 - \sqrt{3}i$.

Q. 5. \sqrt{i}

Solution: Let, $(a + ib)^2 = 0 + i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 0 + i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 0 + i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 0 \dots\dots\dots \text{eq.1}$$

$$\Rightarrow 2ab = 1 \dots\dots\dots \text{eq.2}$$

$$\Rightarrow a = \frac{1}{2b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(\frac{1}{2b}\right)^2 - b^2 = 0$$

$$\Rightarrow 1 - 4b^4 = 0$$

$$\Rightarrow 4b^2 = 1$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -\frac{1}{2} \text{ or } b^2 = \frac{1}{2}$$

As b is real no. so, $b^2 = 3$

$$b = \frac{1}{\sqrt{2}} \text{ or } b = -\frac{1}{\sqrt{2}}$$

Therefore, $a = \frac{1}{\sqrt{2}}$ or $a = -\frac{1}{\sqrt{2}}$

Hence the square root of the complex no. is $\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$ and $-\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i$.

Q. 6. $\sqrt{4i}$

Solution: Let, $(a + ib)^2 = 0 + 4i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 0 + 4i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 0 + 4i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 0 \dots\dots\dots \text{eq.1}$$

$$\Rightarrow 2ab = 4 \dots\dots\dots \text{eq.2}$$

$$\Rightarrow a = \frac{2}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(\frac{2}{b}\right)^2 - b^2 = 0$$

$$\Rightarrow 4 - b^4 = 0$$

$$\Rightarrow b^4 = 4$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -2 \text{ or } b^2 = 2$$

As b is real no. so, $b^2 = 2$

$$b = \sqrt{2} \text{ or } b = -\sqrt{2}$$

Therefore, $a = \sqrt{2}$ or $a = -\sqrt{2}$

Hence the square root of the complex no. is $\sqrt{2} + \sqrt{2}i$ and $-\sqrt{2} - \sqrt{2}i$.

Q. 7. $\sqrt{3+4\sqrt{-7}}$

Solution: Let, $(a + ib)^2 = 3 + 4\sqrt{7}i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 3 + 4\sqrt{7}i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 3 + 4\sqrt{7}i$$

now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 3 \dots \dots \dots \text{eq.1}$$

$$\Rightarrow 2ab = 4\sqrt{7} \dots \dots \dots \text{eq.2}$$

$$\Rightarrow a = \frac{2\sqrt{7}}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(\frac{2\sqrt{7}}{b}\right)^2 - b^2 = 3$$

$$\Rightarrow 12 - b^4 = 3b^2$$

$$\Rightarrow b^4 + 3b^2 - 28 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -7 \text{ or } b^2 = 4$$

as b is real no. so, $b^2 = 4$

$$b = 2 \text{ or } b = -2$$

Therefore, $a = \sqrt{7}$ or $a = -\sqrt{7}$

Hence the square root of the complex no. is $\sqrt{7} + 2i$ and $-\sqrt{7} - 2i$.

Q. 8. $\sqrt{16 - 30i}$

Solution: Let, $(a + ib)^2 = 16 - 30i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 16 - 30i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 16 - 30i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 16 \dots\dots\dots \text{eq.1}$$

$$\Rightarrow 2ab = -30 \dots\dots\dots \text{eq.2}$$

$$\Rightarrow a = -\frac{15}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(-\frac{15}{b}\right)^2 - b^2 = 16$$

$$\Rightarrow 225 - b^4 = 16b^2$$

$$\Rightarrow b^4 + 16b^2 - 225 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -25 \text{ or } b^2 = 9$$

As b is real no. so, $b^2 = 9$

$$b = 3 \text{ or } b = -3$$

Therefore, $a = -5$ or $a = 5$

Hence the square root of the complex no. is $-5 + 3i$ and $5 - 3i$.

Q. 9. $\sqrt{-4 - 3i}$

Solution: Let, $(a + ib)^2 = -4 - 3i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = -4 - 3i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = -4 - 3i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = -4 \dots \dots \dots \text{eq.1}$$

$$\Rightarrow 2ab = -3 \dots \dots \dots \text{eq.2}$$

$$\Rightarrow a = -\frac{3}{2b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(-\frac{3}{2b}\right)^2 - b^2 = -4$$

$$\Rightarrow 9 - 4b^4 = -16b^2$$

$$\Rightarrow 4b^4 - 16b^2 - 9 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = \frac{9}{2} \text{ or } b^2 = -2$$

As b is real no. so, $b^2 = \frac{9}{2}$

$$b = \frac{3}{\sqrt{2}} \text{ or } b = -\frac{3}{\sqrt{2}}$$

Therefore, $a = -\frac{1}{\sqrt{2}}$ or $a = \frac{1}{\sqrt{2}}$

Hence the square root of the complex no. is $-\frac{1}{\sqrt{2}} + \frac{3}{\sqrt{2}}i$ and $\frac{1}{\sqrt{2}} - \frac{3}{\sqrt{2}}i$.

Q. 10. $\sqrt{-15 - 8i}$

Solution: Let, $(a + ib)^2 = -15 - 8i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = -15 - 8i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = -15 - 8i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = -15 \dots \dots \dots \text{eq.1}$$

$$\Rightarrow 2ab = -8 \dots\dots\dots \text{eq.2}$$

$$\Rightarrow a = -\frac{4}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(-\frac{4}{b}\right)^2 - b^2 = -15$$

$$\Rightarrow 16 - b^4 = -15b^2$$

$$\Rightarrow b^4 - 15b^2 - 16 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = 16 \text{ or } b^2 = -1$$

As b is real no. so, $b^2 = 16$

$$b = 4 \text{ or } b = -4$$

Therefore, $a = -1$ or $a = 1$

Hence the square root of the complex no. is $-1 + 4i$ and $1 - 4i$.

Q. 11. $\sqrt{-11 - 60i}$

Solution: Let, $(a + ib)^2 = -11 - 60i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = -11 - 60i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = -11 - 60i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = -11 \dots\dots\dots \text{eq.1}$$

$$\Rightarrow 2ab = -60 \dots\dots\dots\text{eq.2}$$

$$\Rightarrow a = -\frac{30}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(-\frac{30}{b}\right)^2 - b^2 = -11$$

$$\Rightarrow 900 - b^4 = -11b^2$$

$$\Rightarrow b^4 - 11b^2 - 900 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = 36 \text{ or } b^2 = -25$$

as b is real no. so, $b^2 = 36$

$$b = 6 \text{ or } b = -6$$

Therefore , $a = -5$ or $a = 5$

Hence the square root of the complex no. is $-5 + 6i$ and $5 - 6i$.

Q. 12. $\sqrt{7 - 30\sqrt{-2}}$

Solution: Let, $(a + ib)^2 = 7 - 30\sqrt{2}i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 7 - 30\sqrt{2}i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 7 - 30\sqrt{2}i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 7 \dots\dots\dots\text{eq.1}$$

$$\Rightarrow 2ab = 30\sqrt{2} \dots \text{eq.2}$$

$$\Rightarrow a = \frac{15\sqrt{2}}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(\frac{15\sqrt{2}}{b}\right)^2 - b^2 = 7$$

$$\Rightarrow 450 - b^4 = 7b^2$$

$$\Rightarrow b^4 + 7b^2 - 450 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -25 \text{ or } b^2 = 18$$

As b is real no. so, $b^2 = 18$

$$b = 3\sqrt{2} \text{ or } b = -3\sqrt{2}$$

Therefore, $a = 5$ or $a = -5$

Hence the square root of the complex no. is $5 + 3\sqrt{2}i$ and $-5 - 3\sqrt{2}i$.

Q. 13. $\sqrt{-8}$

Solution: Let, $(a + ib)^2 = 0 - 8i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 0 - 8i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 0 - 8i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 0 \dots\dots\dots \text{eq.1}$$

$$\Rightarrow 2ab = -8 \dots\dots\dots \text{eq.2}$$

$$\Rightarrow a = -\frac{4}{b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(-\frac{4}{b}\right)^2 - b^2 = 0$$

$$\Rightarrow 16 - b^4 = 0$$

$$\Rightarrow b^4 = 16$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = -4 \text{ or } b^2 = 4$$

As b is real no. so, $b^2 = 4$

$$b = 2 \text{ or } b = -2$$

Therefore, $a = -2$ or $a = 2$

Hence the square root of the complex no. is $-2 + 2i$ and $2 - 2i$.

Q. 14. $\sqrt{1-i}$

Solution: Let, $(a + ib)^2 = 1 - i$

Now using, $(a + b)^2 = a^2 + b^2 + 2ab$

$$\Rightarrow a^2 + (bi)^2 + 2abi = 1 - i$$

Since $i^2 = -1$

$$\Rightarrow a^2 - b^2 + 2abi = 1 - i$$

Now, separating real and complex parts, we get

$$\Rightarrow a^2 - b^2 = 1 \dots\dots\dots \text{eq.1}$$

$$\Rightarrow 2ab = -1 \dots\dots\dots \text{eq.2}$$

$$\Rightarrow a = -\frac{1}{2b}$$

Now, using the value of a in eq.1, we get

$$\Rightarrow \left(-\frac{1}{2b}\right)^2 - b^2 = 1$$

$$\Rightarrow 1 - 4b^4 = 4b^2$$

$$\Rightarrow 4b^4 + 4b^2 - 1 = 0$$

Simplify and get the value of b^2 , we get,

$$\Rightarrow b^2 = \frac{-4 \pm \sqrt{32}}{8}$$

As b is real no. so, $b^2 = \frac{-4 + 4\sqrt{2}}{8}$

$$b^2 = \frac{-1 + \sqrt{2}}{2}$$

$$b = \sqrt{\frac{-1 + \sqrt{2}}{2}} \text{ or } b = -\sqrt{\frac{-1 + \sqrt{2}}{2}}$$

Therefore, $a = -\sqrt{\frac{1 + \sqrt{2}}{2}}$ or $a = \sqrt{\frac{1 + \sqrt{2}}{2}}$

Hence the square root of the complex no. is $-\sqrt{\frac{1 + \sqrt{2}}{2}} + \sqrt{\frac{-1 + \sqrt{2}}{2}} i$

and $\sqrt{\frac{1 + \sqrt{2}}{2}} - \sqrt{\frac{-1 + \sqrt{2}}{2}} i$.

Exercise 5G

Q. 1. Evaluate $i^{\frac{1}{78}}$.

Solution: we have, $\frac{1}{i^{78}}$

$$= \frac{1}{(i^4)^{19} \cdot i^2}$$

We know that, $i^4 = 1$

$$\Rightarrow \frac{1}{1^{19} \cdot i^2}$$

$$\Rightarrow \frac{1}{i^2} = \frac{1}{-1}$$

$$\Rightarrow \frac{1}{i^{78}} = -1$$

Q. 2. Evaluate $(i^{57} + i^{70} + i^{91} + i^{101} + i^{104})$.

Solution: We have, $i^{57} + i^{70} + i^{91} + i^{101} + i^{104}$

$$= (i^4)^{14} \cdot i + (i^4)^{17} \cdot i^2 + (i^4)^{22} \cdot i^3 + (i^4)^{25} \cdot i + (i^4)^{26}$$

We know that, $i^4 = 1$

$$\Rightarrow (1)^{14} \cdot i + (1)^{17} \cdot i^2 + (1)^{22} \cdot i^3 + (1)^{25} \cdot i + (1)^{26}$$

$$= i + i^2 + i^3 + i + 1$$

$$= i - 1 - i + i + 1$$

$$= i$$

Q. 3. Evaluate

$$\left(\frac{i^{180} + i^{178} + i^{176} + i^{174} + i^{172}}{i^{170} + i^{168} + i^{166} + i^{164} + i^{162}} \right)$$

Solution:

$$\begin{aligned}
 \text{We have, } & \left(\frac{i^{180} + i^{178} + i^{176} + i^{174} + i^{172}}{i^{170} + i^{168} + i^{166} + i^{164} + i^{162}} \right) \\
 &= \left(\frac{i^{180} + i^{178} + i^{176} + i^{174} + i^{172}}{i^{170} + i^{168} + i^{166} + i^{164} + i^{162}} \right) \\
 &= \left(\frac{(i^4)^{45} + (i^4)^{44} \cdot i^2 + (i^4)^{44} + (i^4)^{43} \cdot i^2 + (i^4)^{43}}{(i^4)^{42} \cdot i^2 + (i^4)^{42} + (i^4)^{41} \cdot i^2 + (i^4)^{41} + (i^4)^{40} \cdot i^2} \right) \\
 &= \left(\frac{(1)^{45} + (1)^{44} \cdot i^2 + (1)^{44} + (1)^{43} \cdot i^2 + (1)^{43}}{(1)^{42} \cdot i^2 + (1)^{42} + (1)^{41} \cdot i^2 + (1)^{41} + (1)^{40} \cdot i^2} \right) \\
 &= \left(\frac{1 + i^2 + 1 + i^2 + 1}{i^2 + 1 + i^2 + 1 + i^2} \right) \\
 &= \left(\frac{1 - 1 + 1 - 1 + 1}{-1 + 1 - 1 + 1 - 1} \right) \\
 &= \left(\frac{1}{-1} \right) \\
 &= -1
 \end{aligned}$$

Q. 4. Evaluate $(i^{4n+1} - i^{4n-1})$

Solution: We have, $i^{4n+1} - i^{4n-1}$

$$\begin{aligned}
 &= i^{4n} \cdot i - i^{4n} \cdot i^{-1} \\
 &= (i^4)^n \cdot i - (i^4)^n \cdot i^{-1} \\
 &= (1)^n \cdot i - (1)^n \cdot i^{-1} \\
 &= i - i^{-1} \\
 &= i - \frac{1}{i}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{i^2 - 1}{i} \\
 &= \frac{-1 - 1}{i} \\
 &= \frac{-2}{i} \times \frac{i}{i} \\
 &= \frac{-2i}{i^2} = \frac{-2i}{-1} \\
 &= 2i
 \end{aligned}$$

Q. 5. Evaluate $(\sqrt{36} \times \sqrt{-25})$.

Solution: We have, $(\sqrt{36} \times \sqrt{-25})$

$$\begin{aligned}
 &= 6 \times \sqrt{-1 \times 25} \\
 &= 6 \times (\sqrt{-1} \times \sqrt{25}) \\
 &= 6 \times (\sqrt{-1} \times 5) \\
 &= 6 \times 5i = 30i
 \end{aligned}$$

Q. 6. Find the sum $(i^n + i^{n+1} + i^{n+2} + i^{n+3})$, where $n \in \mathbb{N}$.

Solution: We have $i^n + i^{n+1} + i^{n+2} + i^{n+3}$

$$\begin{aligned}
 &= i^n + i^n \cdot i + i^n \cdot i^2 + i^n \cdot i^3 \\
 &= i^n (1 + i + i^2 + i^3) \\
 &= i^n (1 + i - 1 - i) \\
 &= i^n (0) = 0
 \end{aligned}$$

Q. 7. Find the sum $(i + i^2 + i^3 + i^4 + \dots \text{ up to 400 terms})$, where $n \in \mathbb{N}$.

Solution: We have, $i + i^2 + i^3 + i^4 + \dots$ up to 400 terms

We know that given series is GP where $a=i$, $r = i$ and $n = 400$

$$\text{Thus, } S = \frac{a(1-r^n)}{1-r}$$

$$= \frac{i(1 - (i)^{400})}{1 - i}$$

$$= \frac{i(1 - (i^4)^{100})}{1 - i}$$

$$= \frac{i(1 - 1^{100})}{1 - i} [\because i^4 = 1]$$

$$= \frac{i(1 - 1)}{1 - i} = 0$$

Q. 8. Evaluate $(1 + i^{10} + i^{20} + i^{30})$.

Solution: We have, $1 + i^{10} + i^{20} + i^{30}$

$$= 1 + (i^4)^2 \cdot i^2 + (i^4)^5 + (i^4)^7 \cdot i^2$$

We know that, $i^4 = 1$

$$\Rightarrow 1 + (1)^2 \cdot i^2 + (1)^5 + (1)^7 \cdot i^2$$

$$= 1 + i^2 + 1 + i^2$$

$$= 1 - 1 + 1 - 1$$

$$= 0$$

Q. 9. Evaluate: $\left(i^{41} + \frac{1}{i^{71}}\right)$.

Solution: We have, $\left(i^{41} + \frac{1}{i^{71}}\right)$

$$i^{41} = i^{40} \cdot i = i$$

$$i^{71} = i^{68} \cdot i^3 = -i$$

Therefore,

$$\left(i^{41} + \frac{1}{i^{71}}\right) = i - \frac{1}{i} = \frac{i^2 - 1}{i}$$

$$\left(i^{41} + \frac{1}{i^{71}}\right) = -\frac{2}{i} \times \frac{i}{i}$$

$$\left(i^{41} + \frac{1}{i^{71}}\right) = -\frac{2i}{i^2} = 2i$$

Hence, $\left(i^{41} + \frac{1}{i^{71}}\right) = 2i$

Q. 10. Find the least positive integer n for which $\left(\frac{1+i}{1-i}\right)^n = 1$.

Solution: We have, $\left(\frac{1+i}{1-i}\right)^n = 1$

$$\text{Now, } \frac{1+i}{1-i} = \frac{1+i}{1-i} \times \frac{1+i}{1+i}$$

$$= \frac{(1+i)^2}{1^2 - i^2}$$

$$= \frac{1^2 + 2i + i^2}{1 - (-1)}$$

$$= \frac{1 + 2i - 1}{2}$$

$$= i$$

$$\therefore \left(\frac{1+i}{1-i}\right)^n = (i)^n = 1 \Rightarrow n \text{ is multiple of } 4$$

\therefore The least positive integer n is 4

Q. 11. Express $(2 - 3i)^3$ in the form $(a + ib)$.

Solution: We have, $(2 - 3i)^3$
 $= 2^3 - 3 \times 2^2 \times 3i - 3 \times 2 \times (3i)^2 - (3i)^3$
 $= 8 - 36i + 54 + 27i$
 $= 46 - 9i.$

Q. 12. Express $\frac{(3+i\sqrt{5})(3-\sqrt{5})}{(\sqrt{3}+\sqrt{2i})-(\sqrt{3}-\sqrt{2i})}$ **in the form (a + ib).**

Solution: We have, $\frac{(3+i\sqrt{5})(3-i\sqrt{5})}{(\sqrt{3}+\sqrt{2i})-(\sqrt{3}-\sqrt{2i})}$
 $= \frac{(3)^2 - (i\sqrt{5})^2}{\sqrt{3} + \sqrt{2i} - \sqrt{3} + \sqrt{2i}}$ [$\because (a + b)(a - b) = a^2 - b^2$]

$$= \frac{9 + 5}{2\sqrt{2i}} \times \frac{\sqrt{2i}}{\sqrt{2i}}$$

$$= \frac{14\sqrt{2i}}{2(\sqrt{2i})^2}$$

$$= \frac{7\sqrt{2i}}{-2}$$

$$= \frac{-7\sqrt{2i}}{2}$$

Q. 13. Express $\frac{3-\sqrt{-16}}{1-\sqrt{-9}}$ **in the form (a + ib).**

Solution: We have, $\frac{3-\sqrt{-16}}{1-\sqrt{-9}}$

We know that $\sqrt{-1} = i$

Therefore,

$$\frac{3 - \sqrt{-16}}{1 - \sqrt{-9}} = \frac{3 - 4i}{1 - 3i}$$

$$\frac{3 - \sqrt{-16}}{1 - \sqrt{-9}} = \frac{3 - 4i}{1 - 3i} \times \frac{1 + 3i}{1 + 3i}$$

$$\frac{3 - \sqrt{-16}}{1 - \sqrt{-9}} = \frac{3 + 9i - 4i - 12i^2}{(1)^2 - (3i)^2}$$

$$\frac{3 - \sqrt{-16}}{1 - \sqrt{-9}} = \frac{15 + 5i}{1 + 9} = \frac{15}{10} + \frac{5i}{10} = \frac{3}{2} + \frac{1}{2}i$$

Hence,

$$\frac{3 - \sqrt{-16}}{1 - \sqrt{-9}} = \frac{3}{2} + \frac{i}{2}$$

Q. 14. Solve for x: $(1 - i)x + (1 + i)y = 1 - 3i$.

Solution: We have, $(1 - i)x + (1 + i)y = 1 - 3i$

$$\Rightarrow x - ix + y + iy = 1 - 3i$$

$$\Rightarrow (x + y) + i(-x + y) = 1 - 3i$$

On equating the real and imaginary coefficients we get,

$$\Rightarrow x + y = 1 \text{ (i) and } -x + y = -3 \text{ (ii)}$$

From (i) we get

$$x = 1 - y$$

Substituting the value of x in (ii), we get

$$-(1 - y) + y = -3$$

$$\Rightarrow -1 + y + y = -3$$

$$\Rightarrow 2y = -3 + 1$$

$$\Rightarrow y = -1$$

$$\Rightarrow x = 1 - y = 1 - (-1) = 2$$

Hence, $x = 2$ and $y = -1$

Q. 15. Solve for x : $x^2 - 5ix - 6 = 0$.

Solution: We have, $x^2 - 5ix - 6 = 0$

$$\text{Here, } b^2 - 4ac = (-5i)^2 - 4 \times 1 \times -6$$

$$= 25i^2 + 24 = -25 + 24 = -1$$

Therefore, the solutions are given by $x = \frac{-(-5i) \pm \sqrt{-1}}{2 \times 1}$

$$x = \frac{5i \pm i}{2 \times 1}$$

$$x = \frac{5i \pm i}{2}$$

Hence, $x = 3i$ and $x = 2i$

Q. 16. Find the conjugate of $\frac{1}{(3 + 4i)}$.

Solution: Let $z = \frac{1}{3 + 4i}$

$$= \frac{1}{3 + 4i} \times \frac{3 - 4i}{3 - 4i} = \frac{3 - 4i}{9 + 16}$$

$$= \frac{3}{25} - \frac{4}{25}i$$

$$\Rightarrow \bar{z} = \frac{3}{25} + \frac{4}{25}i$$

Q. 17. If $z = (1 - i)$, find z^{-1} .

Solution: We have, $z = (1 - i)$

$$\Rightarrow \bar{z} = 1 + i$$

$$\Rightarrow |z|^2 = (1)^2 + (-1)^2 = 2$$

\therefore The multiplicative inverse of $(1 - i)$,

$$z^{-1} = \frac{\bar{z}}{|z|^2} = \frac{1 + i}{2}$$

$$z^{-1} = \frac{1}{2} + \frac{1}{2}i$$

Q. 18. If $z = (\sqrt{5} + 3i)$, find z^{-1} .

Solution: We have, $z = (\sqrt{5} + 3i)$

$$\Rightarrow \bar{z} = (\sqrt{5} - 3i)$$

$$\Rightarrow |z|^2 = (\sqrt{5})^2 + (3)^2$$

$$= 5 + 9 = 14$$

\therefore The multiplicative inverse of $(\sqrt{5} + 3i)$,

$$z^{-1} = \frac{\bar{z}}{|z|^2} = \frac{\sqrt{5} - 3i}{14}$$

$$z^{-1} = \frac{\sqrt{5}}{14} + \frac{3}{14}i$$

Q. 19. Prove that $\arg(z) + \arg(\bar{z}) = 0$

Solution: Let $z = r(\cos\theta + i \sin\theta)$

$$\Rightarrow \arg(z) = \theta$$

$$\text{Now, } \bar{z} = r(\cos\theta - i \sin\theta) = r(\cos(-\theta) + i \sin(-\theta))$$

$$\Rightarrow \arg(\bar{z}) = -\theta$$

$$\text{Thus, } \arg(z) + \arg(\bar{z}) = \theta - \theta = 0$$

Hence proved.

Q. 20. If $|z| = 6$ and $\arg(z) = \frac{3\pi}{4}$, find z .

Solution: We have, $|z| = 6$ and $\arg(z) = \frac{3\pi}{4}$

$$\text{Let } z = r(\cos\theta + i \sin\theta)$$

$$\text{We know that, } |z| = r = 6$$

$$\text{And } \arg(z) = \theta = \frac{3\pi}{4}$$

$$\text{Thus, } z = r(\cos\theta + i \sin\theta) = 6 \left(\cos\frac{3\pi}{4} + i \sin\frac{3\pi}{4} \right)$$

Q. 21. Find the principal argument of $(-2i)$.

Solution: Let, $z = -2i$

$$\text{Let } 0 = r\cos\theta \text{ and } -2 = r\sin\theta$$

By squaring and adding, we get

$$(0)^2 + (-2)^2 = (r\cos\theta)^2 + (r\sin\theta)^2$$

$$\Rightarrow 0+4 = r^2(\cos^2\theta + \sin^2\theta)$$

$$\Rightarrow 4 = r^2$$

$$\Rightarrow r = 2$$

$$\therefore \cos\theta = 0 \text{ and } \sin\theta = -1$$

Since, θ lies in fourth quadrant, we have

$$\theta = -\frac{\pi}{2}$$

Since, $\theta \in (-\pi, \pi]$ it is principal argument.

Q. 22. Write the principal argument of $(1 + i\sqrt{3})^2$.

Solution: Let, $z = (1 + i\sqrt{3})^2$

$$= (1)^2 + (i\sqrt{3})^2 + 2\sqrt{3}i$$

$$= 1 - 1 + 2\sqrt{3}i$$

$$z = 0 + 2\sqrt{3}i$$

Let $0 = r\cos\theta$ and $2\sqrt{3} = r\sin\theta$

By squaring and adding, we get

$$(0)^2 + (2\sqrt{3})^2 = (r\cos\theta)^2 + (r\sin\theta)^2$$

$$\Rightarrow 0 + (2\sqrt{3})^2 = r^2(\cos^2\theta + \sin^2\theta)$$

$$\Rightarrow (2\sqrt{3})^2 = r^2$$

$$\Rightarrow r = 2\sqrt{3}$$

$$\therefore \cos\theta = 0 \text{ and } \sin\theta = 1$$

Since, θ lies in first quadrant, we have

$$\theta = \frac{\pi}{2}$$

Since, $\theta \in (-\pi, \pi]$ it is principal argument.

Q. 23. Write -9 in polar form.

Solution: We have, $z = -9$

$$\text{Let } -9 = r\cos\theta \text{ and } 0 = r\sin\theta$$

By squaring and adding, we get

$$(-9)^2 + (0)^2 = (r\cos\theta)^2 + (r\sin\theta)^2$$

$$\Rightarrow 81 = r^2(\cos^2\theta + \sin^2\theta)$$

$$\Rightarrow 81 = r^2$$

$$\Rightarrow r = 9$$

$$\therefore \cos\theta = -1 \text{ and } \sin\theta = 0$$

$$\Rightarrow \theta = \pi$$

Thus, the required polar form is $9(\cos \pi + i \sin \pi)$

Q. 24. Write $2i$ in polar form.

Solution: Let, $z = 2i$

$$\text{Let } 0 = r\cos\theta \text{ and } 2 = r\sin\theta$$

By squaring and adding, we get

$$(0)^2 + (2)^2 = (r\cos\theta)^2 + (r\sin\theta)^2$$

$$\Rightarrow 0+4 = r^2(\cos^2\theta + \sin^2\theta)$$

$$\Rightarrow 4 = r^2$$

$$\Rightarrow r = 2$$

$$\therefore \cos\theta = 0 \text{ and } \sin\theta = 1$$

Since, θ lies in first quadrant, we have

$$\theta = \frac{\pi}{2}$$

Thus, the required polar form is $2 \left(\cos \left(\frac{\pi}{2} \right) + i \sin \left(\frac{\pi}{2} \right) \right)$

Q. 25. Write $-3i$ in polar form.

Solution: Let, $z = -3i$

Let $0 = r\cos\theta$ and $-3 = r\sin\theta$

By squaring and adding, we get

$$(0)^2 + (-3)^2 = (r\cos\theta)^2 + (r\sin\theta)^2$$

$$\Rightarrow 0+9 = r^2(\cos^2\theta + \sin^2\theta)$$

$$\Rightarrow 9 = r^2$$

$$\Rightarrow r = 3$$

$$\therefore \cos\theta = 0 \text{ and } \sin\theta = -1$$

Since, θ lies in fourth quadrant, we have

$$\theta = \frac{3\pi}{2}$$

Thus, the required polar form is $3 \left(\cos\left(\frac{3\pi}{2}\right) + i\sin\left(\frac{3\pi}{2}\right) \right)$

Q. 26. Write $z = (1 - i)$ in polar form.

Solution: We have, $z = (1 - i)$

Let $1 = r\cos\theta$ and $-1 = r\sin\theta$

By squaring and adding, we get

$$(1)^2 + (-1)^2 = (r\cos\theta)^2 + (r\sin\theta)^2$$

$$\Rightarrow 1+1 = r^2(\cos^2\theta + \sin^2\theta)$$

$$\Rightarrow 2 = r^2$$

$$\Rightarrow r = \sqrt{2}$$

$$\therefore \cos\theta = \frac{1}{\sqrt{2}} \text{ and } \sin\theta = \frac{-1}{\sqrt{2}}$$

Since, θ lies in fourth quadrant, we have

$$\theta = -\frac{\pi}{4}$$

Thus, the required polar form is $\sqrt{2} \left(\cos \left(-\frac{\pi}{4} \right) + i \sin \left(-\frac{\pi}{4} \right) \right)$

Q. 27. Write $z = (-1 + i\sqrt{3})$ in polar form.

Solution: We have, $z = (-1 + i\sqrt{3})$

Let $-1 = r\cos\theta$ and $\sqrt{3} = r\sin\theta$

By squaring and adding, we get

$$(-1)^2 + (\sqrt{3})^2 = (r\cos\theta)^2 + (r\sin\theta)^2$$

$$\Rightarrow 1+3 = r^2(\cos^2\theta + \sin^2\theta)$$

$$\Rightarrow 4 = r^2$$

$$\Rightarrow r = 2$$

$$\therefore \cos\theta = \frac{-1}{2} \text{ and } \sin\theta = \frac{\sqrt{3}}{2}$$

Since, θ lies in second quadrant, we have

$$\theta = \pi - \frac{\pi}{3} = \frac{2\pi}{3}$$

Thus, the required polar form is $2 \left(\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right)$

Q. 28. If $|z| = 2$ and $\arg(z) = \frac{\pi}{4}$, find z .

Solution: We have, $|z| = 2$ and $\arg(z) = \frac{\pi}{4}$,

Let $z = r(\cos\theta + i \sin\theta)$

We know that, $|z| = r = 2$

And $\arg(z) = \theta = \frac{\pi}{4}$

Thus, $z = r(\cos\theta + i \sin\theta) = 2 \left(\cos\frac{\pi}{4} + i \sin\frac{\pi}{4} \right)$

