Class: XII Session: 2020-21

Subject: Mathematics

Sample Question Paper (Theory)

Time Allowed: 3 Hours Maximum Marks: 80

General Instructions:

1. This question paper contains two **parts A and B**. Each part is compulsory. Part A carries **24** marks and Part B carries **56** marks

- 2. **Part-A** has Objective Type Questions and **Part -B** has Descriptive Type Questions
- 3. Both Part A and Part B have choices.

Part – A:

- 1. It consists of two sections- I and II.
- 2. Section I comprises of 16 very short answer type questions.
- 3. Section II contains 2 case studies. Each case study comprises of 5 case-based MCQs. An examinee is to attempt any 4 out of 5 MCQs.

Part - B:

- 1. It consists of three sections- III, IV and V.
- 2. Section III comprises of 10 questions of 2 marks each.
- 3. Section IV comprises of 7 questions of 3 marks each.
- 4. Section **V** comprises of 3 questions of **5 marks** each.
- 5. Internal choice is provided in **3** questions of Section –III, **2** questions of Section-IV and **3** questions of Section-V. You have to attempt only one of the alternatives in all such questions.

Sr.	Part – A	Mark
No.		S
	Section I	
	All questions are compulsory. In case of internal choices attempt any one.	
1	Check whether the function $f: R \to R$ defined as $f(x) = x^3$ is one-one or not.	1
	OR	

	How many reflexive relations are possible in a set A whose $n(A) = 3$.	1
2	A relation R in $S = \{1,2,3\}$ is defined as $R = \{(1,1),(1,2),(2,2),(3,3)\}$. Which element(s) of relation R be removed to make R an equivalence relation?	1
3	A relation R in the set of real numbers R defined as $R = \{(a, b): \sqrt{a} = b\}$ is a function or not. Justify	1
	OR	
	An equivalence relation R in A divides it into equivalence classes A_1,A_2,A_3 . What is the value of $A_1 \cup A_2 \cup A_3$ and $A_1 \cap A_2 \cap A_3$	1
4	If A and B are matrices of order $3 \times n$ and $m \times 5$ respectively, then find the order of matrix $5A - 3B$, given that it is defined.	1
5	Find the value of A^2 , where A is a 2×2 matrix whose elements are given by $a_{ij} = \begin{cases} 1 & if & i \neq j \\ 0 & if & i = j \end{cases}$	1
	OR	
	Given that A is a square matrix of order 3×3 and A = -4. Find adj A	1
6	Let A = $\begin{bmatrix} a_{ij} \end{bmatrix}$ be a square matrix of order 3×3 and A = -7. Find the value of $a_{11} A_{21} + a_{12} A_{22} + a_{13} A_{23}$	1
	where A_{ij} is the cofactor of element a_{ij}	
7	Find $\int e^x (1 - \cot x + \csc^2 x) dx$	1
	OR _π	1
	Evaluate $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x^2 \sin x dx$	•
8	Find the area bounded by $y = x^2$, the x – axis and the lines $x = -1$ and $x = 1$.	1
9	How many arbitrary constants are there in the particular solution of the differential equation $\frac{dy}{dx} = -4xy^2$; y (0) = 1	1
	OR	
	For what value of n is the following a homogeneous differential equation: $\frac{dy}{dx} = \frac{x^3 - y^n}{x^2y + xy^2}$	1
10	Find a unit vector in the direction opposite to $-\frac{3}{4}\hat{j}$	1
11	Find the area of the triangle whose two sides are represented by the vectors $2\hat{\imath}$ and $-3\hat{\jmath}$.	1

12	Find the angle between the unit vectors \hat{a} and \hat{b} , given that $ \hat{a} + \hat{b} = 1$	1
13	Find the direction cosines of the normal to YZ plane?	1
14	Find the coordinates of the point where the line $\frac{x+3}{3} = \frac{y-1}{-1} = \frac{z-5}{-5}$ cuts the XY plane.	1
15	The probabilities of A and B solving a problem independently are $\frac{1}{3}$ and $\frac{1}{4}$ respectively. If both of them try to solve the problem independently, what is the probability that the problem is solved?	1
16	The probability that it will rain on any particular day is 50%. Find the probability that it rains only on first 4 days of the week.	1
	Section II	
	Both the Case study based questions are compulsory. Attempt any 4 sub parts from each question (17-21) and (22-26). Each question carries 1 mark	
17	An architect designs a building for a multi-national company. The floor consists of a rectangular region with semicircular ends having a perimeter of 200m as shown below:	
	Design of Floor	
	$\begin{array}{c c} A & \begin{array}{c} y \\ \hline z \\ \end{array} \\ \end{array}$	
	Building	
	Based on the above information answer the following:	
	(i) If x and y represents the length and breadth of the rectangular region, then the relation between the variables is	
	a) $x + \pi y = 100$ b) $2x + \pi y = 200$ c) $\pi x + y = 50$ d) $x + y = 100$	
<u> </u>		

(ii)The area of the rectangular region A expressed as a function of x is	1
a) $\frac{2}{\pi} (100 x - x^2)$	
b) $\frac{1}{\pi} (100 x - x^2)$	
c) $\frac{x}{\pi} (100 - x)$	
d) $\pi y^2 + \frac{2}{\pi} \left(100 x - x^2 \right)$	
(iii) The maximum value of area A is	1
a) $\frac{\pi}{3200}m^2$	
b) $\frac{3200}{\pi}m^2$	
c) $\frac{5000}{\pi} m^2$	
$d) \frac{1000}{\pi} m^2$	
(iv) The CEO of the multi-national company is interested in maximizing the area of the whole floor including the semi-circular ends. For this to happen the valve of x should be	1
a) 0 m	
b) 30 m	
c) 50 m	
d) 80 m	
(v) The extra area generated if the area of the whole floor is maximized is :	1
a) $\frac{3000}{\pi}m^2$	
b) $\frac{5000}{\pi}m^2$	
c) $\frac{7000}{\pi} m^2$	
d) No change Both areas are equal	

In an office three employees Vinay, Sonia and Iqbal process incoming copies of a certain form. Vinay process 50% of the forms. Sonia processes 20% and Iqbal 18 the remaining 30% of the forms. Vinay has an error rate of 0.06, Sonia has an error rate of 0.04 and Iqbal has an error rate of 0.03 #!("&\$!^@a&%x#. Based on the above information answer the following: (i) The conditional probability that an error is committed in processing given that Sonia processed the form is: a) 0.0210 b) 0.04 c) 0.47 d) 0.06 (ii)The probability that Sonia processed the form and committed an error is : 1 a) 0.005 b) 0.006 c) 0.008 d) 0.68 (iii)The total probability of committing an error in processing the form is 1 a) 0 b) 0.047 c) 0.234

(iv)The manager of the company wants to do a quality check. During inspection he selects a form at random from the days output of processed forms. If the form selected at random has an error, the probability that the form is NOT processed by Vinay is: a) 1 b) 30/47 c) 20/47 d) 17/47 (v)Let A be the event of committing an error in processing the form and let E_1 , E_2 and E_3 be the events that Vinay, Sonia and Iqbal processed the form. The value of $\sum_{i=1}^3 P(E_i A)$ is a) 0 b) 0.03 c) 0.06 d) 1 Part – B Section III 19 Express $tan^{-1}(\frac{cosx}{1-slinx}), \frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 20 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of $ A $. 21 Find the value(s) of k so that the following function is continuous at $x = 0$			
he selects a form at random from the days output of processed forms. If the form selected at random has an error, the probability that the form is NOT processed by Vinay is : a) 1 b) 30/47 c) 20/47 d) 17/47 (v)Let A be the event of committing an error in processing the form and let E_1 , E_2 and E_3 be the events that Vinay, Sonia and Iqbal processed the form. The value of $\sum_{i=1}^3 P\left(E_i \mid A\right)$ is a) 0 b) 0.03 c) 0.06 d) 1 Part – B Section III 19 Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 20 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of $ A $. OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .		d) 1	
E ₂ and E ₃ be the events that Vinay, Sonia and Iqbal processed the form. The value of $\sum_{i=1}^{3} P\left(E_{i} \mid A\right)$ is a) 0 b) 0.03 c) 0.06 d) 1 Part – B Section III Express $tan^{-1}\left(\frac{cosx}{1-sinx}\right), \frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 2 If A is a square matrix of order 3 such that $A^{2} = 2A$, then find the value of $ A $. OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^{2} - 5A + 71 = 0$. Hence find A^{-1} .		he selects a form at random from the days output of processed forms. If the form selected at random has an error, the probability that the form is NOT processed by Vinay is: a) 1 b) 30/47 c) 20/47	1
value of $\sum_{i=1}^{3} P\left(E_{i} \mid A\right)$ is a) 0 b) 0.03 c) 0.06 d) 1 Part – B Section III Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 2 OR If A is a square matrix of order 3 such that $A^{2} = 2A$, then find the value of $ A $. Part – B Section III 2 Hence find A ⁻¹ .		(v)Let A be the event of committing an error in processing the form and let E_1 ,	1
a) 0 b) 0.03 c) 0.06 d) 1 Part – B Section III		E_2 and E_3 be the events that Vinay, Sonia and Iqbal processed the form. The	
b) 0.03 c) 0.06 d) 1 Part – B Section III		value of $\sum_{i=1}^{3} P(E_i A)$ is	
c) 0.06 d) 1 Part – B Section III Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 2 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of A . OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .		a) 0	
Part – B Section III Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 2 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of $ A $. OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .		b) 0.03	
Part – B Section III Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 2 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of $ A $. OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .		c) 0.06	
Section III 19 Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 20 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of A . OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .		d) 1	
Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 2 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of A . OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .		Part – B	
Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 20 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of A . OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .		Section III	
Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form. 20 If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of A . OR If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .			
$ \text{OR} $ If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .	19	Express $tan^{-1}(\frac{cosx}{1-sinx})$, $\frac{-3\pi}{2} < x < \frac{\pi}{2}$ in the simplest form.	2
If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .	20	If A is a square matrix of order 3 such that $A^2 = 2A$, then find the value of $ A $.	2
If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$. Hence find A^{-1} .		OR	
Hence find A ⁻¹ .			2
		If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = 0$.	2
Find the value(s) of k so that the following function is continuous at $x = 0$		Hence find A ⁻¹ .	
	21	Find the value(s) of k so that the following function is continuous at $x = 0$	2

	$f(x) = \begin{cases} \frac{1-\cos kx}{x\sin x} & \text{if } x \neq 0\\ \frac{1}{2} & \text{if } x = 0 \end{cases}$	
	(2	
22	Find the equation of the normal to the curve	2
	$y = x + \frac{1}{x}$, $x > 0$ perpendicular to the line $3x - 4y = 7$.	
23	Find $\int \frac{1}{\cos^2 x (1 - \tan x)^2} dx$	2
	OR	
	Evaluate $\int_0^1 x(1-x)^n dx$	2
24	Find the area of the region bounded by the parabola $y^2 = 8x$ and the line $x = 2$.	2
25	Solve the following differential equation:	2
	$\frac{dy}{dx} = x^3 \cos c y, given that y(0) = 0.$	
26	Find the area of the parallelogram whose one side and a diagonal are represented by coinitial vectors $\hat{\imath}$ - $\hat{\jmath}$ + \hat{k} and $4\hat{\imath}$ + $5\hat{k}$ respectively	2
27	Find the vector equation of the plane that passes through the point (1,0,0) and contains the line $\vec{r} = \lambda \hat{j}$.	2
28	A refrigerator box contains 2 milk chocolates and 4 dark chocolates. Two chocolates are drawn at random. Find the probability distribution of the number of milk chocolates. What is the most likely outcome?	2
	OR	
	Given that E and F are events such that P(E) = 0.8, P(F) = 0.7, P (E \cap F) = 0.6. Find P ($\bar{E} \mid \bar{F}$)	2
	Section IV	
	All questions are compulsory. In case of internal choices attempt any one.	
29	Check whether the relation R in the set Z of integers defined as R = $\{(a,b): a+b \text{ is "divisible by 2"}\}\$ is reflexive, symmetric or transitive. Write the equivalence class containing 0 i.e. [0].	3
30	If $y = e^{x \sin^2 x} + (\sin x)^x$, find $\frac{dy}{dx}$.	3
31	Prove that the greatest integer function defined by $f(x) = [x]$, $0 < x < 2$ is not differentiable at $x = 1$	3

	OR	
	If $x = a \sec \theta$, $y = b \tan \theta$ find $\frac{d^2y}{dx^2}$ at $x = \frac{\pi}{6}$	3
32	Find the intervals in which the function f given by	3
	$f(x) = \tan x - 4x, x \in \left(0, \frac{\pi}{2}\right)$ is	
	a) strictly increasing b) strictly decreasing	
33	Find $\int \frac{x^2+1}{(x^2+2)(x^2+3)} dx$.	3
	$(x^2+2)(x^2+3)$	
34	Find the area of the region bounded by the curves	3
	$x^2 + y^2 = 4$, $y = \sqrt{3}x$ and $x - axis$ in the first quadrant	
	OR	
	Find the area of the ellipse $x^2 + 9y^2 = 36$ using integration	3
35	Find the general solution of the following differential equation:	3
	$\int x dy - (y + 2x^2) dx = 0$	
	Section V	
	All questions are compulsory. In case of internal choices attempt any	
	one.	
36	If $A = \begin{bmatrix} 1 & 2 & 0 \\ -2 & -1 & -2 \\ 0 & -1 & 1 \end{bmatrix}$, find A^{-1} . Hence	5
	Solve the system of equations;	
	x - 2y = 10	
	2x - y - z = 8	
	-2y + z = 7	
	OR	
	Evaluate the product AB, where	5
	$A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 & 2 & -4 \\ -4 & 2 & -4 \\ 2 & -1 & 5 \end{bmatrix}$	
	Hence solve the system of linear equations	
	x - y = 3	

	2x + 3y + 4z = 17	
	y + 2z = 7	
37	Find the shortest distance between the lines $\vec{r} = 3\hat{\imath} + 2\hat{\jmath} - 4\hat{k} + \lambda(\hat{\imath} + 2\hat{\jmath} + 2\hat{k})$ and $\vec{r} = 5\hat{\imath} - 2\hat{\jmath} + \mu\left(3\hat{\imath} + 2\hat{\jmath} + 6\hat{k}\right)$ If the lines intersect find their point of intersection	5
	OR	
	Find the foot of the perpendicular drawn from the point (-1, 3, -6) to the plane $2x + y - 2z + 5 = 0$. Also find the equation and length of the perpendicular.	5
38	Solve the following linear programming problem (L.P.P) graphically. Maximize $Z = x + 2y$ subject to constraints ; $x + 2y \ge 100$ $2x - y \le 0$ $2x + y \le 200$ $x, y \ge 0$	5
	OR	
	The corner points of the feasible region determined by the system of linear constraints are as shown below: y 11 10 9 8 7 6 5 4 3 2 1 E(4,0) E(4,0)	5
	Ol 1 2 3 4 5 6 7 Answer each of the following:	
	(i) Let $Z = 3x - 4y$ be the objective function. Find the maximum and minimum value of Z and also the corresponding points at which the maximum and minimum value occurs.	

(ii) Let Z = px + qy, where p, q > o be the objective function. Find the condition on p and q so that the maximum value of Z occurs at B(4,10) and C(6,8). Also mention the number of optimal solutions in this case.