




1. In the figure given below, If the areas of the two regions are equal then which of the following is true?



- A. $x = y$
- B. $x = 2y$
- C. $2x = y$
- D. $x = 3y$



$$A_a = x \times 2y + \frac{1}{2}(y + 2y)x = \frac{7}{2}xy$$




$$A_1 = 4 \frac{y^2}{2} + (2x - 2y) \cdot 2y + y^2$$

$$A_1 = A_2$$

$$x = 2y$$

2. Let ABCD be a square of a side length 1, Let P, Q, R, S be points in the interiors of the sides AD, BC, AB, CD, respectively, such that PQ and RS intersect at right angles. If $PQ = \frac{3\sqrt{3}}{4}$, then RS equals

- A. $\frac{2}{\sqrt{3}}$
- B. $\frac{3\sqrt{3}}{4}$
- C. $\frac{\sqrt{2}+1}{2}$
- D. $4 - 2\sqrt{2}$



$$PQ \perp RS$$

$$\Rightarrow c - a = b - d \dots \dots \dots (1)$$

$$\Rightarrow PQ = \frac{3\sqrt{3}}{4}$$

$$\Rightarrow PQ^2 = \frac{27}{16}$$

$$\Rightarrow 1 + (a - c)^2 = \frac{27}{16} \dots \dots \dots (2)$$


$$\Rightarrow RS = \sqrt{(b - d)^2 + 1} \dots \dots \dots (3)$$

By equation (1), (2) and (3)

$$RS = \frac{3\sqrt{3}}{4}$$

3. Let ABCD be a square of side length 1, and Γ a circle passing through B and C, and touching AD. The radius of Γ is

- A. $\frac{3}{8}$
- B. $\frac{1}{2}$
- C. $\frac{1}{\sqrt{2}}$
- D. $\frac{5}{8}$



$$PC=r$$

$$PC^2 = r^2$$

$$(r-1)^2 + \left(\frac{1}{2}-1\right)^2 = r^2$$

$$1 - 2r + \frac{1}{4} = 0$$

$$r = \frac{5}{8}$$


4. Let P be an interior point of a triangle ABC. Let Q and R be the reflections of P in AB and AC, respectively. If Q, A, R are collinear then $\angle A$ equals.

A. 30°

B. 60°

C. 90°

D. 120°



So from Diagram

$$\theta + \theta + \phi + \phi = 180^\circ$$


$$\angle A = \theta + \phi = 90^\circ$$

5. In a triangle ABC, points X and Y are on AB and AC, respectively, such that XY is parallel to BC. Which of the two following always hold(s) good? (Here [PQR] denotes the area of triangle PQR.)

(I) $[BCX] = [BCY]$

(II) $[ACX] \cdot [ABY] = [AXY] \cdot [ABC]$

- A. Neither [I] nor [II]
- B. [I] only
- C. [II] only
- D. both [I] and [II]



$\Delta BCX = \Delta BCY$ (Obvious)

Same base and same height

Now Let $A(\vec{O})$, $\vec{AB} = \vec{b}$, $\vec{AC} = \vec{c}$

So $\vec{AX} = \lambda \vec{b}$, $\vec{AY} = \lambda \vec{c}$

$\Delta ACX = \frac{1}{2} \lambda |\vec{b} \times \vec{c}|$

$\Delta ABY = \frac{1}{2} \lambda |\vec{b} \times \vec{c}|$

$\Delta AXY = \frac{1}{2} \lambda^2 |\vec{b} \times \vec{c}|$

$\Delta ABC = \frac{1}{2} |\vec{b} \times \vec{c}|$

6. The largest non-negative integer k such that 24^k divides $13!$ is.

A. 2

B. 3

C. 4

D. 5

$$24^K \rightarrow (2^3 \times 3)^K$$

Exponent of 2 in $13!$

$$\left[\frac{13}{2} \right] + \left[\frac{13}{2^2} \right] + \left[\frac{13}{2^3} \right] = 10$$


Exponent of 3 in $13!$

$$\left[\frac{13}{3} \right] + \left[\frac{13}{3^2} \right] = 5$$

So $(2^3 \times 3)^3$

Hence, $K=3$

7. In the figure given below, a rectangle of perimeter 76 units is divided into 7 congruent rectangles.



What is the perimeter of each of the smaller rectangles?

- A. 38
- B. 32
- C. 28
- D. 19

Let sides of rectangle are x & y

Then

$$5x + 6y = 76$$

$$4y = 3x$$

After solving we get $y = 6$, $x = 8$

$$\text{Perimeter} = 2(x + y) = 28$$

8. Let x, y, z be positive reals. Which of the following implies $x = y = z$?

(I) $x^3 + y^3 + z^3 = 3xyz$

(II) $x^3 + y^2 z + yz^2 = 3xyz$

(III) $x^3 + y^2 z + z^2 x = 3xyz$

(IV) $(x + y + z)^3 = 27xyz$

- A. I, IV only
- B. I, II, IV only
- C. I, II and III only
- D. All of them

For option (3) if $x = z = 1$ and $y = 2$ then option (3) is right.

So by option (3) we can't say $x = y = z$. Remaining options implies $x = y = z$.

9. If n is the smallest natural number such that $n + 2n + 3n + \dots + 99n$ is a perfect square, then the number of digits in n^2 is

- A. 1
- B. 2
- C. 3
- D. More than 3

$$n + 2n + 3n + \dots + 99n$$

$$n((1 + 2 + 3 + \dots + 99))$$

$$n \frac{(99)(100)}{2} = 9 \times 25 \times 22 \times n \text{ is a perfect square when } n = 22$$

Number of digits in $n = 3$

10. Two distinct polynomial $f(x)$ and $g(x)$ are defined as follows:

$$f(x) = x^2 + ax + 2; g(x) = x^2 + 2x + a$$

If the equation $f(x) = 0$ and $g(x) = 0$ have a common root, then the sum of the roots of the equation $f(x) + g(x) = 0$ is

A. $-\frac{1}{2}$

B. 0

C. $\frac{1}{2}$

D. 1

$$f(x) + g(x) = x^2 + ax + 2 + x^2 + 2x + a$$

Here a common root then

$$\begin{vmatrix} 1 & a \\ 1 & 2 \end{vmatrix} = \begin{vmatrix} a & 2 \\ 2 & a \end{vmatrix} \Rightarrow \begin{vmatrix} 2 & 1 \\ a & 1 \end{vmatrix}^2$$

$$\Rightarrow a = 2, -3$$

$$f(x) + g(x) = 2x^2 + (a + 2)x + a + 2$$

$$\text{Sum of roots} = \frac{-1(a+2)}{2} \text{ if } a = -3 \text{ then sum} = \frac{1}{2}$$

11. Methane is a greenhouse gas because

- A. It absorbs longer wavelengths of the electromagnetic spectrum while transmitting shorter wavelengths.
- B. It absorbs shorter wavelengths of the electromagnetic spectrum while transmitting longer wavelengths.
- C. It absorbs all wavelengths of the electromagnetic spectrum.
- D. It transmits all wavelengths of the electromagnetic spectrum.





Gases that trap heat in the atmosphere are called greenhouse gases. Methane is emitted during the production and transport of coal, natural gas, and oil. Methane emissions also result from livestock and other agricultural practices and by the decay of organic waste in municipal solid waste landfills.

12. If the axis of rotation of the earth were extended into space then it would pass close to

- A. The moon
- B. The sun
- C. The pole star
- D. The centre of mass of all the planets in the solar system.



13. An object at rest at the origin begins to move in the +x direction with a uniform acceleration of 1 m/s^2 for 4 s and then it continues moving with a uniform velocity of 4 m/s in the same direction. The x - t graph for object's motion will be



$$v=0+1\times t$$

$$\frac{dx}{dt}=t$$

$$dx=tdt$$

$$x \propto t^2 \dots\dots\dots\text{Parabolic} \quad (1)$$


$$\frac{dx}{dt}=4$$

$$dx=4dt$$


$$x \propto t \dots\dots\dots\text{Linear} \quad (2)$$

BYJU'S

14. A hollow tilted cylindrical vessel of negligible mass rests on a horizontal plane as shown. The diameter of the base is a and the side of the cylinder makes an angle θ with the horizontal. Water is then slowly poured into the cylinder. The cylinder topples over when the water reaches a certain height h , given by.



- A. $h = 2a \tan \theta$
- B. $h = a \tan^2 \theta$
- C. $h = a \tan \theta$
- D. $h = \frac{a}{2} \tan \theta$



$$\tan \theta = \frac{\frac{h}{2}}{\frac{a}{2}}$$

$$h = a \tan \theta$$

15. In an experiment, mass of an object is measured by applying a known force on it, and then measuring its acceleration. If, in the experiment, the measured values of applied force and the measured acceleration are $F = 10.0 \pm 0.2\text{N}$ and $a = 1.00 \pm 0.01 \text{ m/s}^2$, respectively, the mass of the object is

- A. 10.0 Kg
- B. $10.0 \pm 0.1 \text{ Kg}$
- C. $10.0 \pm 0.3 \text{ Kg}$
- D. $10.0 \pm 0.4\text{Kg}$

$$F=Ma$$

$$M=\frac{f}{a}$$

$$\frac{\Delta M \times 100}{M} = \frac{\Delta f}{f} \times 100 + \frac{\Delta a}{a} \times 100$$

$$= \frac{0.2}{10} + \frac{0.01}{1}$$

$$\Delta M = 0.03 \times 10$$

$$\therefore M = 10 \pm 0.3\text{kg}$$

16. How many ways are there to arrange the letters of the word EDUCATION so that all the following three conditions hold?

- the vowels occur in the same order (EUAIO);
- the consonants occur in the same order(DCTN);
- no two consonants are next to each other.

- A. 15
- B. 24
- C. 72
- D. 120

Vowels can be arranged in one way.

There will be 6 gaps around them. We need 4 gaps as the arrangement can be done in one way.

Therefore, ${}^6C_4 = 15$

17. The number of distinct primes dividing $12! + 13! + 14!$ is

- A. 5
- B. 6
- C. 7
- D. 8

$$12! + 13! + 14!$$

$$12! (1 + 13 + 14 \times 13)$$

$$12! \times 196$$

Which is only divided by possible distinct primes 2, 3, 5, 7, 11

18. Let $n > 1$ be an integer. Which of the following sets of numbers necessarily contains a multiple of 3?

A. $n^{19} - 1, n^{19} + 1$

B. $n^{19}, n^{38} - 1$

C. $n^{38}, n^{38} + 1$

D. $n^{38}, n^{19} - 1$

If $n = 3m$ then n^{19} is multiple of 3

If $n = 3m + 1$ or $3m + 2$ then $n^{38} - 1$ is multiple of 3 by binomial expansion

19. The least positive integer n for which $\sqrt[3]{n+1} - \sqrt[3]{n} < \frac{1}{12}$ is

A. 6

B. 7

C. 8

D. 9

$$(n+1)^{1/3} - (n)^{1/3} < \frac{1}{12}$$

$$(n+1)^{1/3} < (n)^{1/3} + \frac{1}{12}$$

Cube both side are get

$$(n^{1/3}) \left(n^{1/3} + \frac{1}{12} \right) > \frac{1727}{432}$$

So, $n = 8$ only possible least positive integer.


20. A man standing on a railway platform noticed that a train took 21 seconds to cross the platform (this means the time elapsed from the moment the engine enters the platform till the last compartment leaves the platform) which is 88 meters long, and that it took 9 seconds to pass him. Assuming that the train was moving with uniform speed, what is the length of the train in metres?

A. 55

B. 60

C. 66

D. 72



$$9v + 88 = 21v$$

$$12v = 88$$

$$v = \frac{88}{12}$$


$$\text{Required} = 9v = 9 \times \frac{88}{12} = 66$$

21. The international space station is maintained in a nearly circular orbit with a mean altitude of 330 km and a maximum of 410 km. An astronaut is floating in the space station's cabin. The acceleration of astronaut as measured from the earth is.

- A. Zero
- B. Nearly zero and directed towards the earth
- C. Nearly g and directed along the line of travel of the station
- D. Nearly g and directed towards the earth

Earth will be applying the gravitational pull on the astronaut and this pull is producing an acceleration nearly equal to g , which is directed towards the earth.

22. One mole of ideal gas undergoes a linear process as shown in figure below. Its temperature expressed as a function of volume V is.



- A. $\frac{P_0 V_0}{R}$
- B. $\frac{P_0 V}{R}$
- C. $\frac{P_0 V}{R} \left(1 - \frac{V}{V_0}\right)$
- D. $\frac{P_0 V}{R} \left(1 - \left(\frac{V}{V_0}\right)^2\right)$

$$P = \frac{-P_0}{V_0}V + P_0 \dots \dots \dots (1)$$

&

$$PV = nRT \dots \dots \dots (2)$$

$$T = \frac{P_0 V}{R} \left[1 - \frac{V}{V_0}\right]$$

23. A planet is orbiting the sun in an elliptical orbit. Let U denote the potential energy and K denote the kinetic energy of the planet at an arbitrary point on the orbit. Choose the correct statement.

- A. $K < |U|$ always
- B. $K > |U|$ always
- C. $K = |U|$ always
- D. $K = |U|$ for two positions of the planet in the orbit.


Total energy must be less than zero and as potential energy is negative so answer is **A**

24. A ball is launched from the top of Mt. Everest which is at elevation of 9000 m. The ball moves in circular orbit around earth. Acceleration due to gravity near the earth's surface is g . The magnitude of the ball's acceleration while in orbit is

- A. Close to $g/2$
- B. Zero
- C. Much greater than g .
- D. Nearly equal to g .

$$\frac{mv^2}{r} = mg' \text{ (Where } g' \text{ is nearly equal to } g\text{)}$$

25. A light bulb of resistance $r = 16\Omega$ is attached in series with an infinite resistor network with identical resistances r as shown below. A 10 V battery derives current in the circuit. What should be the value of r such that the bulb dissipated about 1 W of power?



- A. 14.8Ω
- B. 29.6Ω
- C. 7.4Ω
- D. 3.7Ω

$$P_{bulb} = \frac{v^2}{R} = i^2 R$$

$$I = \frac{v^2}{16}$$

$$V_B = 4 \text{ V}$$

$$I = i^2 \times 16$$

$$I_B = \frac{1}{4} \text{ Amp.}$$

$$6 = \frac{1}{4} \times r_{eq} \text{ (equivalent of groups of } r)$$

where

$$r_{eq} = r + \frac{r_{eq} \cdot r}{r_{eq} + r}$$

26. A point source of light is moving at a rate of 2 cm-s^{-1} towards a thin convex lens of focal length 10 cm along its optical axis. When the source is 15 cm away from the lens the image is moving at

- A. 4 cm-s^{-1} towards the lens
- B. 8 cm-s^{-1} towards the lens
- C. 4 cm-s^{-1} away from the lens
- D. 8 cm-s^{-1} away from the lens

$$\frac{dv}{dt} = \frac{-v^2 du}{u^2 dt}$$

$$\& \frac{1}{v} = \frac{1}{F} + \frac{1}{u}$$

$$\frac{1}{V} = \frac{1}{10} + \frac{1}{-15}$$

$$\frac{1}{v} = \frac{3-2}{30}$$


$$v = 30 \text{ cm}$$

$$\frac{dv}{dt} = \left(\frac{1}{F}\right)^2 \cdot 2$$

=8 cm/s away from lens

27. An isosceles glass prism with angles 40° is clamped over a tray of water in a position such that the base is just dipped in water. A ray of light incident normally on the inclined face suffers total internal reflection at the base. If the refractive index of water is 1.33 then the condition imposed on the refractive index μ of the glass is


- A. $\mu < 2.07$
- B. $\mu > 2.07$
- C. $\mu < 1.74$
- D. $\mu > 1.74$



$$\begin{aligned}
 1.33 \times \sin 90^\circ &= \mu \sin 40^\circ \\
 &= \frac{1.33 \times 1}{\sin 40^\circ} = \mu \\
 &= \frac{1.33}{3/5} \approx \mu \\
 &= \frac{1.33 \times 5}{3} \approx \mu \\
 &= \mu \approx 2.07 \\
 &= \mu > 2.07 \text{ (For TIR)}
 \end{aligned}$$

28. An optical device is constructed by fixing three identical convex lenses of focal lengths 10 cm each inside a hollow tube at equal spacing of 30 cm each. One end of the device is placed 10 cm away from a point source. How much does the image shift when the device is moved away from the source by another 10 cm?

- A. 0 cm
- B. 5 cm
- C. 15 cm
- D. 45 cm



No Shift in image

29. The beta particles of a radioactive metal originate from.

- A. The free electrons in the metal
- B. The orbiting electrons of the metal atoms
- C. The photons released from the nucleus.
- D. The nucleus of the metal atoms.

The beta particles of a radioactive metal originate from the nucleus of the atom.

30. A parachutist with total weight 75 kg drops vertically onto a sandy ground with a speed of 2 ms^{-1} and comes to a halt over a distance of 0.25m. The average force from the ground on her is close to.

- A. 600 N
- B. 1200 N
- C. 1350 N
- D. 1950 N

$$0^2 = 2^2 - 2a \frac{1}{4}$$

$$a = 8 \text{ m/s}^2$$

$$a_T = (10 + 8) = (18 \text{ m/s}^2)$$


$$f = ma$$

$$= 75 \times 18$$

$$= 1350 \text{ N}$$

31. The functional group present in a molecule having the formula $C_{12}O_9$ is

- A. Carboxylic acid
- B. anhydride
- C. aldehyde
- D. alcohol



32. If Avogadro's number is A_0 , the number of sulphur atoms present in 200 mL of 1N H_2SO_4 is

- A. $A_0/5$
- B. $A_0/2$
- C. $A_0/10$
- D. A_0

Avogadro's number = A_0

Normality = $n_t \times$ Molarity

$1 = 2 \times M$

$M = \frac{1}{2} \text{molL}^{-1}$

Moles of $H_2SO_4 = \frac{1}{2} \times 0.2 = 0.1$ moles


Normality = 1; Volume = 200 ml (0.2 litre)

Moles of hydrogen = 0.2 moles

Moles of sulphur = 0.1 moles


Atoms = $0.1A_0$

33. The major products of the following reaction



Are

- A. ZnO and SO₂
- B. ZnSO₄ and SO₃
- C. ZnSO₄ and SO₂
- D. Zn and SO₂



34. The element which readily forms an ionic bond has the electronic configuration.


- A. 1s²2s²2p³
- B. 1s²2s²2p¹
- C. 1s²2s²2p²
- D. 1s²2s²2p⁶3s¹

Metals form ionic bond as they have low ionization energies.

1s² 2s² 2p⁶ 3s¹ : Sodium metal


35. The gas released when baking soda is mixed with vinegar, is

- A. CO
- B. CO₂
- C. CH₄
- D. O₂




36. Reaction of 2-butane with acidic KMnO_4 gives


- A. CH_3CHO
- B. HCOOH
- C. $\text{CH}_3\text{CH}_2\text{OH}$
- D. CH_3COOH



37. The correct order of acidity of the following compounds is



- A. $1 > 2 > 3$
- B. $1 > 3 > 2$
- C. $3 > 1 > 2$
- D. $3 > 2 > 1$



- (B) OCH_3 exerts +M effect destabilizes the conjugate base of the acid.
- (C) NO_2 exerts - M effect and stabilizes the conjugate base of the acid

38. Maximum number of electrons that can be accommodated in the subshell with azimuthal quantum number $l = 4$, is

A. 10

B. 8

C. 16

D. 18

Total number of electrons = $2(2l+1) = 18$

39. Mendeleev's periodic law states that the properties of elements are a periodic function of their

A. Reactivity of elements

B. atomic size

C. Atomic mass

D. electronic configuration

Mendeleev's periodic law states that properties of elements are periodic function of their atomic masses.

40. The percentage of nitrogen by mass in ammonium sulphate is closest to (atomic masses H = 1, N = 14, O = 16, S = 32)

A. 21%

B. 24%

C. 36%

D. 16%

Ammonium sulphate $(NH_4)_2SO_4 =$

$$\% \text{ of nitrogen} = \frac{28 \times 100}{36 + 96} = \frac{28 \times 100}{132} \cong 21.21\%$$

41. Glycolysis is the breakdown of glucose to pyruvic acid. How many molecules of pyruvic acid are formed from one molecule of glucose?

- A. 1
- B. 2
- C. 3
- D. 4

In glycolysis, one molecule of glucose ($C_6H_{12}O_6$) forms two molecules of pyruvic acid ($CH_3COCOOH$).

42. A person with blood group AB has

- A. antigen A and B on RBCs and both anti-A and anti-B antibodies in plasma
- B. antigen A and B on RBCs but neither anti-A and anti-B antibodies in plasma
- C. no antigen on RBCs but both anti-A and anti-B antibodies present in plasma
- D. antigen A on RBCs and antibodies in plasma

A person with blood group AB have both A and B antigens on membrane of his RBCs but lacks antibodies (a, b) in his plasma.

43. Enzyme X extracted from the digestive system hydrolyses peptide bonds. Which of the following are probable candidate to be enzyme X?

- A. Amylase
- B. Lipase
- C. Trypsin
- D. Maltase

Trypsin is proteolytic enzyme.

44. Restriction endonucleases are enzymes that are used by biotechnologists to

- A. cut DNA at specific base sequences
- B. join fragments of DNA
- C. digest DNA from the 3' end
- D. digest DNA from the 5' end


Restriction endonucleases cut ds DNA at specific base sequences called restriction site.

45. Excess salt inhibits growth of bacteria in pickles by:

- A. endosmosis
- B. exosmosis
- C. oxidation
- D. denaturation

Due to hypertonic solution outside the bacterial cell, bacteria will die by plasmolysis.


46. At constant pressure, the volume of a fixed mass of a gas varies as a function of temperature as shown in the graph.



The volume of the gas at 300°C is larger than that at 0°C by a factor of

- A. 3
- B. 4
- C. 1
- D. 2

From graph.:



$$T_1 = 0^\circ\text{C}$$

$$V_1 = 250$$

$$T_2 = 300^\circ$$

$$V_2 = 500$$

$$\frac{V_2}{V_1} = 2$$


47. The number of electrons required to reduce chromium completely in $\text{Cr}_2\text{O}_7^{2-}$ to Cr^{3+} in acidic medium, is

A. 5

B. 3

C. 6

D. 2




48. The maximum number of isomers ethers with the molecular formula $\text{C}_4\text{H}_{10}\text{O}$ is

A. 2

B. 3

C. 4

D. 5




49. Among Mg, Cu, Fe, Zn, the metal that does not produce hydrogen gas in reaction with hydrochloric acid is.

- A. Cu
- B. Zn
- C. Mg
- D. Fe

Metals having more standard reduction potential than $H^+ / H_2(g)$ can't produce $H_2(g)$ in acidic medium.

50. A sweet smelling compounds formed by reacting acetic acid with ethanol in the presence of sulphuric acid is

- A. $CH_3COOC_2H_5$
- B. C_2H_5COOH
- C. $C_2H_5COOH_3$
- D. CH_3OH



51. Which one of the following is a modified leaf?

- A. Sweet potato
- B. Ginger
- C. Onion
- D. Carrot

Onion is bulb formed by fleshy leaves, sweet potato and carrot are modified root, ginger rhizome is a modified stem.

52. Which of the following pairs are both polysaccharides?

- A. Cellulose and glycogen
- B. Starch and glucose
- C. Cellulose and fructose
- D. Ribose and sucrose

Cellulose is polymer of β , D-glucose and glycogen of α , D-glucose. Glucose, fructose and ribose are monosaccharides.

53. Considering the average molecular mass of a base to be 500 Da, what is the molecular mass of a double stranded DNA of 10 base pairs?

- A. 500 Da
- B. 5kDa
- C. 10 kDa
- D. 1 kDa

1 Base = 500 Da, ds DNA having 10 BP or 20 bases, thus molecular mass of dsDNA = $20 \times 500 = 10 \text{ kDa}$

54. Which, among grass, goat, tiger and vulture, in a food chain, will have the maximum concentration of harmful chemicals in its body due to contamination of pesticides in the soil?

- A. Grass since it grows in the contaminated soil
- B. Goat since it eats the grass
- C. Tiger since it feed on the goat which feeds on the grass
- D. Vulture since it eats the tiger, which in turn eats the goat, which eats the grass.

It is due to biomagnifications. Accumulation of harmful chemicals will be maximum in the organism present at highest level in the food chain.

55. Widal test is prescribed to diagnose.

- A. Typhoid
- B. Pneumonia
- C. Malaria
- D. Filaria

Typhoid is caused by *Salmonella typhi* which is diagnosed by widal test.

56. In which one of the following phyla is the body segmented?

- A. Porifera
- B. Platyhelminthes
- C. Annelida
- D. Echinodermata

Annelids show metameric segmentation.

57. Which one of the following is detrimental to soil fertility?

- A. Saprophytic bacteria
- B. Nitrosomes
- C. Nitrobacter
- D. Pseudomonas

Pseudomonas is denitrifying bacterium.

58. Deficiency of which of the following vitamins can cause impaired blood clotting?

- A. Vitamin B
- B. Vitamin C
- C. Vitamin D
- D. Vitamin K

Vit. K is useful in synthesis of prothrombin and fibrinogen in liver which are necessary for blood clotting.

59. Which one of the following cell types is a part of innate immunity?

- A. Skin epithelial cells
- B. B cells
- C. T lymphocytes
- D. Liver cells

Skin, mucus membranes and phagocytes are part of innate immunity.

60. The process of transfer of electrons from glucose to molecular oxygen in bacteria and mitochondria is known as

- A. TCA cycle
- B. Oxidative phosphorylation
- C. Fermentation
- D. Glycolysis

ETS or electron transport system is also known as oxidative phosphorylation.

61. Stoke's law states that the viscous drag force F experience by a sphere of radius a , moving with a speed v through a fluid with coefficient of viscosity η , is given by $F = 6\pi\eta av$ If this fluid is flowing through a cylindrical pipe of radius r , length l and a pressure difference of P across its two ends, then the volume of water V which flows through the pipe in time t can be written as

$$\frac{V}{t} = k \left(\frac{P}{l} \right)^a \eta^b r^c$$

Where k is a dimensionless constant. Correct values of a , b and c are

- A. $a = 1, b = -1, c = 4$
- B. $a = -1, b = 1, c = 4$
- C. $a = 2, b = -1, c = 3$
- D. $a = 1, b = -2, c = -4$

$$\frac{V}{t} = k \left(\frac{P}{l} \right)^a \eta^b r^c$$

$$L^3 T^{-1} = (M L^{-2} T^{-2})^a L^c (M L^{-1} T^{-1})^b$$

$$a + b = 0$$


$$-2a - b + c = 3$$

$$-2a - b = -1$$

$$c = 4, a = 1, b = -1$$

Answer (A) is correct.

62. A point object is placed 20 cm left of a convex lens of focal length $f = 5$ cm (see the figure). The lens is made to oscillate with small amplitude A along the horizontal axis. The image of the object will also oscillate along the axis with.



- A. amplitude $\frac{A}{9}$, out of phase with the oscillations of the lens
- B. amplitude $\frac{A}{3}$, out of phase with the oscillations of the lens
- C. amplitude $\frac{A}{3}$, in phase with the oscillations of the lens
- D. amplitude $\frac{A}{9}$, in phase with the oscillations of the lens

$$\frac{1}{v} + \frac{1}{20} = \frac{1}{5}$$

$$v = \frac{20}{3} \text{ cm}$$

$$\Delta x_i = \frac{+v^2}{u^2} \Delta x_0$$

$$\Delta x_i = \frac{A}{9} \text{ out of phase with lens}$$

Hence (A) is correct

63. A rigid body in the shape of a "V" has two equal arms made of uniform rods. What must the angle between the two rods be so that when the body is suspended from one end, the other arm is horizontal?

A. $\cos^{-1} \left(\frac{1}{3} \right)$

B. $\cos^{-1} \left(\frac{1}{2} \right)$

C. $\cos^{-1} \left(\frac{1}{4} \right)$

D. $\cos^{-1} \left(\frac{1}{6} \right)$

$$m_1 r_1 + m_2 r_2 = 0$$

$$m r_1 = m r_2$$

$$r_1 = r_2$$

$$\frac{\ell}{2} \cos \theta = \frac{\ell}{2} - \ell \cos \theta$$

$$\frac{3\ell}{2} \cos \theta = \frac{3\ell}{2}$$

$$\cos \theta = \frac{1}{3}$$

64. 1 Kg of ice at -20°C is mixed with 2 Kg of water at 90°C . Assuming that there is no loss of energy to the environment, what will be the final temperature of the mixture? (Assume latent heat of ice = 334.4 KJ/Kg, specific heat of water and ice are 4.18 kJ/(kg.K) and 2.09 kJ/(kg.K), respectively.)

A. 30°C

B. 0°C

C. 80°C


D. 45°C

$$m_i s_i (\Delta T) + m_i L + m_i \cdot s_w (T - 0) = m_w s_w (90 - T)$$

$$1 \times 2.09(20) + 1 \times 334.4 + 1 \times 4.18 \times T = 2 \times 4.18 \times (90 - T)$$

$$T = 60 - 30 = 30^\circ\text{C}$$

65. A girl sees through a circular glass slab (refractive index 1.50) of thickness 20 mm and diameter 60 cm to the bottom of a swimming pool. Refractive index of water is 1.33. The bottom surface of the slab is in contact with the water surface.



The depth of swimming pool is 6m. The area of bottom of swimming pool that can be seen through the slab is approximately.

- A. 100 m²
- B. 160 m²
- C. 190 m²
- D. 220 m²

For maximum possible area

i should be 90°

$$1 \times \sin 90^\circ = \frac{4}{3} \sin r$$


$$\sin r = \frac{3}{4}$$

$$\tan r = \frac{3}{\sqrt{7}}$$

$$\text{Total base area } \pi \left(6 \times \frac{3}{\sqrt{7+0.3}} \right)^2 \approx 160 \text{ m}^2$$

66. The average incomes of the people in two villages are P and Q , respectively. Assume that $P \neq Q$. A person moves from the first village to the second village. The new average incomes are P' and Q' , respectively. Which of the following is not possible?

- A. $P' > P$ and $Q' > Q$
- B. $P' > P$ and $Q' < Q$
- C. $P' = P$ and $Q' = Q$
- D. $P' < P$ and $Q' < Q$



Let number of peoples in two villages be n and m respectively

$$\text{So } nP - P'(n - 1) = (m + 1)Q' - mQ$$

$$n(P - P') + P' = m(Q' - Q) + Q'$$


67. Given are three cylindrical buckets X, Y, Z whose circular bases are of radii 1, 2, 3 units, respectively. Initially, water is filled in these buckets up to the same height. Some water is then transferred from Z to X so that they both have the same volume of water. Some water is then transferred between X and Y so that they both have the same volume of water. If h_y , h_z denote the heights of water at this stage in the buckets Y, Z, respectively, then the ratio $\frac{h_y}{h_z}$ equals.

A. $\frac{4}{9}$

B. 1

C. $\frac{9}{4}$

D. $\frac{81}{40}$



$$V = \pi h \quad \pi 4h \quad \pi 9h \quad \dots \text{stage (1)}$$

$$\pi h \quad 4\pi h \quad 5\pi h \quad \dots \text{stage (2)}$$

$$4.5\pi h \quad 5.4\pi h \quad 5\pi h \quad \dots \text{stage (3)}$$

$$4\pi \cdot h_y = \frac{9\pi h}{2}$$

$$9\pi h_z = 5\pi h$$

$$h_y = \frac{9h}{8}$$


$$h_z = \frac{5h}{9}$$

$$\frac{h_y}{h_z} = \frac{9h}{8} / \frac{5h}{9}$$

$$= \frac{81}{40}$$

68. Let ABCD be a trapezium with AD parallel to BC. Assume there is a point M in the interior of the segment BC such that AB = AM and DC = DM. Then the ratio of the area of the trapezium to the area of triangle AMD is.

- A. 2
- B. 3
- C. 4
- D. not determinable from the data



$$\text{Required} = \frac{\frac{1}{2}[(a+c)+2(a+c)]b}{\frac{1}{2}(a+c)b} = 3$$

69. For a real number x , let $[x]$ denote the largest integer less than or equal to x , and let $\{x\} = x - [x]$. The number of solutions x to be equation $[x]\{x\} = 5$ with is $0 \leq x \leq 2015$ is

- A. 0
- B. 3
- C. 2008
- D. 2009

$$\{x\} = x - [x]$$

$$[x]\{x\} = 5 \quad f \neq 0$$

$$If = 5$$


$$0 < f < 1$$

Possible solutions

$$(6 + \frac{5}{6}), (7 + \frac{5}{7}), \dots, (2014 + \frac{5}{2014}) = 2009$$

70. A triangular corner is cut from a rectangular piece of paper and the resulting pentagon has sides 5, 6, 8, 9, 12 in some order. The ratio of the area of the pentagon to the area of the rectangle is

- A. $\frac{11}{18}$
- B. $\frac{13}{18}$
- C. $\frac{15}{18}$
- D. $\frac{17}{18}$




$$\text{Area of rectangle} = 9 \times 12$$

$$\begin{aligned} \text{Area of pentagon} &= 12 \times 9 - \frac{1}{2} \times 3 \times 4 \\ &= 12 \times 9 - 6 \end{aligned}$$

$$\text{Required} = \frac{12 \times 9 - 6}{12 \times 9} = \frac{17}{18}$$

71. A line is drawn from the exterior of an animal cell to the centre of the nucleus, crossing through one mitochondrion. What is the minimum number of membrane bilayers that the line will cross?

- A. 4
- B. 3
- C. 8
- D. 6







72. What is the advantage of storing glucose as glycogen in animals instead of as monomeric glucose?

- A. Energy obtained from glycogen is more than that from the corresponding glucose monomers.
- B. Glucose present as monomers within the cell exerts more osmotic pressure than a single glycogen molecule, resulting in loss of water from the cells.
- C. Glucose present as monomers within the cell exerts more osmotic pressure than a single glycogen molecule, resulting in excess water within the cells.
- D. Glycogen gives more rigidity to the cells.

Glucose maintains high osmotic pressure inside the cell.

73. A few rabbits are introduced in an un-inhabited island with plenty of food. If these rabbits breed in the absence of any disease, natural calamity and predation, which one of the following graphs best represents their population growth?



In absence of disease, natural calamity and predation growth of rabbit is exponential.

74. If rice is cooked in a pressure cooker on the Siachen glacier, at sea beach, and on Deccan plain, which of the following is correct about the time taken for cooking rice?

- A. Gets cooked faster on the Siachen glacier
- B. Gets cooked faster at sea beach
- C. Gets cooked faster on Deccan plain
- D. Gets cooked at the same time at all the three places.


Rice is cooked faster at sea level than at high altitude.

75. Genomic DNA is digested with Alu I, a restriction enzyme which is a four base - pair cutter. What is the frequency with which it will cut the DNA assuming a random distribution of bases in the genome?


- A. $1/4$
- B. $1/24$
- C. $1/256$
- D. $1/1296$

Alu I is a restriction endonuclease which is a 4 base pair cutter its frequency is $1/256$ BP, while frequency of 6 base pair cutter Bam HI, EcoRI is $1/4096$.


76. The major product of the reaction is :



- A. I
- B. II
- C. III
- D. IV



Mechanism electrophilic addition reaction of alkenes.



77. The amount of Ag (atomic mass = 108) deposited at the cathode when a current of 0.5 amp is passed through a solution of AgNO_3 for 1 hour is closest to :


- A. 2 g
- B. 5 g
- C. 108 g
- D. 11 g

$$W = \frac{E}{96500} \times I \times t$$

$$W = \frac{108}{96500} \times 0.5 \times 3600 = 2gm$$

78. When 22.4 L of C_4H_8 at STP is burnt completely, 89.6 L of CO_2 gas at STP and 72 g of water are produced. The volume of the oxygen gas at STP consumed in the reaction is closest to :

- A. 89.6 L
 B. 112 L
 C. 134.4 L
 D. 22.4 L



22.4 lit 89.6 lit. 72 g

At S.T.P. at S.T.P.

1 mole 4 mole $\frac{72}{18} = 4\text{mole}$

For complete combustion of 1 mole C_4H_8

6 mole O_2 required


$n_{O_2} = 6 \text{ mole}$

$V_{O_2} = 6 \times 22.4$

$V_{O_2} = 134.4 \text{ lit.}$


79. Reaction of ethanol with conc. Sulphuric acid at $170^\circ C$ produces a gas which is then treated with bromine in carbon tetrachloride. The major product obtained in this reaction is :

- A. 1,2-dibromoethane
 B. ethylene glycol
 C. bromoethane
 D. ethyl sulphate



80. When 262 g of xenon (atomic mass = 131) reacted completely with 152 g of fluorine (atomic mass = 19), a mixture of XeF_2 and XeF_6 was produced. The molar ratio $\text{XeF}_2 : \text{XeF}_6$ is :

- A. 1 : 2
- B. 1 : 4
- C. 1 : 1
- D. 1 : 3



$$\frac{262}{131} \quad \frac{152}{38} \quad a \text{ mole} \quad b \text{ mole}$$

$$= 2 \text{ mole} \quad 4 \text{ mole}$$

Let a mole XeF_2 form and b mole XeF_6 form

Apply POAC

$$a \times 1 + b \times 1 = 2 \dots \dots (1)$$

$$2a + 6b = 8 \dots \dots (2)$$

After solving eq. (1) & (2)

$$a = 1 \text{ mole} \ \& \ b = 1 \text{ mole}$$