

# **EXERCISE 9.1**

# **PAGE NO: 9.5**

Solve each of the following equations and also verify your solution:

1. 9  $\frac{1}{4} = y - \frac{1}{1/3}$ **Solution:** We have,  $9\frac{1}{4} = y - \frac{1}{1/3}$ 37/4 = y - 4/3Upon solving we get, y = 37/4 + 4/3By taking LCM for 4 and 3 is 12  $y = (37 \times 3)/12 + (4 \times 4)/12$ = 111/12 + 16/12=(111+16)/12= 127/12:: y = 127/12Verification-RHS = y - 1 1/3= 127/12 - 4/3=(127-16)/12= 111/12= 37/4 $=9\frac{1}{4}$ = LHS 2. 5x/3 + 2/5 = 1**Solution:** We have, 5x/3 + 2/5 = 15x/3 = 1 - 2/5 (by taking LCM) =(5-2)/5By using cross-multiplication we get, 5x/3 = 3/5 $5x = (3 \times 3)/5$  $x = 9/(5 \times 5)$ = 9/25 $\therefore \mathbf{x} = 9/25$ 



Verification-LHS = 5x/3 + 2/5=  $5/3 \times 9/25 + 2/5$ = 3/5 + 2/5= (3 + 2)/5= 5/5= 1 = RHS

3. x/2 + x/3 + x/4 = 13Solution:

We have.

x/2 + x/3 + x/4 = 13let us take LCM for 2, 3 and 4 which is 12  $(x \times 6)/12 + (x \times 4)/12 + (x \times 3)/12 = 13$ 6x/12 + 4x/12 + 3x/12 = 13(6x+4x+3x)/12 = 1313x/12 = 13By using cross-multiplication we get,  $13x = 12 \times 13$ x = 156/13= 12 $\therefore x = 12$ Verification-LHS = x/2 + x/3 + x/4= 12/2 + 12/3 + 12/4= 6 + 4 + 3= 13 = RHS

#### 4. x/2 + x/8 = 1/8 Solution:

We have, x/2 + x/8 = 1/8let us take LCM for 2 and 8 which is 8  $(x\times4)/8 + (x\times1)/8 = 1/8$  4x/8 + x/8 = 1/8 5x/8 = 1/8By using cross-multiplication we get,



5x = 8/85x = 1x = 1/5 $\therefore \mathbf{x} = 1/5$ Verification-LHS = x/2 + x/8=(1/5)/2+(1/5)/8= 1/10 + 1/40= (4 + 1)/40= 5/40= 1/8= RHS5. 2x/3 - 3x/8 = 7/12Solution: We have, 2x/3 - 3x/8 = 7/12

By taking LCM for 3 and 8 is 24  $(2x \times 8)/24 - (3x \times 3)/24 = 7/12$ 16x/24 - 9x/24 = 7/12(16x-9x)/24 = 7/127x/24 = 7/12By using cross-multiplication we get,  $7x \times 12 = 7 \times 24$  $x = (7 \times 24)/(7 \times 12)$ = 24/12= 2 $\therefore \mathbf{x} = 2$ Verification-LHS = 2x/3 - 3x/8= 2(2)/3 - 3(2)/8= 4/3 - 6/8= 4/3 - 3/4=(16-9)/12= 7/12= RHS

6. (x + 2) (x + 3) + (x - 3) (x - 2) - 2x(x + 1) = 0Solution:



We have, (x+2) (x+3) + (x-3) (x-2) - 2x(x+1) = 0Upon expansion we get,  $x^{2} + 5x + 6 + x^{2} - 5x + 6 - 2x^{2} - 2x = 0$ -2x + 12 = 0By dividing the equation using -2 we get, x - 6 = 0 $\mathbf{x} = \mathbf{6}$  $\therefore x = 6$ Verification-LHS = (x + 2) (x + 3) + (x - 3) (x - 2) - 2x(x + 1)= (6+2)(6+3) + (6-3)(6-2) - 2(6)(6+1)=(8)(9)+(3)(4)-12(7)= 72 + 12 - 84= 84 - 84= 0= RHS7. x/2 - 4/5 + x/5 + 3x/10 = 1/5Solution: We have. x/2 - 4/5 + x/5 + 3x/10 = 1/5upon solving we get, x/2 + x/5 + 3x/10 = 1/5 + 4/5by taking LCM for 2, 5 and 10 which is 10  $(x \times 5)/10 + (x \times 2)/10 + (3x \times 1)/10 = 5/5$ 5x/10 + 2x/10 + 3x/10 = 1(5x+2x+3x)/10 = 110x/10 = 1 $\mathbf{x} = 1$  $\therefore \mathbf{x} = 1$ Verification-LHS = x/2 - 4/5 + x/5 + 3x/10 $= \frac{1}{2} - \frac{4}{5} + \frac{1}{5} + \frac{3(1)}{10}$ =(5-8+2+3)/10=(10-8)1/0= 2/10= 1/5= RHS



## 8. 7/x + 35 = 1/10Solution: We have, 7/x + 35 = 1/107/x = 1/10 - 35 $=((1\times 1) - (35\times 10))/10$ =(1-350)/107/x = -349/10By using cross-multiplication we get, x = -70/349 $\therefore x = -70/349$ Verification-LHS = 7/x + 35= 7/(-70/349) + 35 $=(-7 \times 349)/70 + 35$ = -349/10 + 35=(-349+350)/10= 1/10= RHS

# 9. (2x-1)/3 - (6x-2)/5 = 1/3

## Solution:

We have, (2x-1)/3 - (6x-2)/5 = 1/3By taking LCM for 3 and 5 which is 15  $((2x-1)\times 5)/15 - ((6x-2)\times 3)/15 = 1/3$ (10x - 5)/15 - (18x - 6)/15 = 1/3(10x - 5 - 18x + 6)/15 = 1/3(-8x + 1)/15 = 1/3By using cross-multiplication we get, (-8x + 1)3 = 15-24x + 3 = 15-24x = 15 - 3-24x = 12x = -12/24= -1/2:: x = -1/2Verification-LHS = (2x - 1)/3 - (6x - 2)/5



= [2(-1/2) - 1]/3 - [6(-1/2) - 2]/5= (-1 - 1)/3 - (-3 - 2)/5 = -2/3 - (-5/5) = -2/3 + 1 = (-2 + 3)/3 = 1/3 RHS


10. 13(y - 4) - 3(y - 9) - 5(y + 4) = 0**Solution:** We have, 13(y - 4) - 3(y - 9) - 5(y + 4) = 0Upon expansion we get, 13y - 52 - 3y + 27 - 5y - 20 = 013y - 3y - 5y = 52 - 27 + 205y = 45y = 45/5= 9  $\therefore$  y = 9 Verification-LHS = 13(y-4) - 3(y-9) - 5(y+4)= 13 (9 - 4) - 3 (9 - 9) - 5 (9 + 4)= 13(5) - 3(0) - 5(13)= 65 - 0 - 65= 0= RHS

## 11. 2/3(x - 5) - 1/4(x - 2) = 9/2Solution:

We have, 2/3(x - 5) - 1/4(x - 2) = 9/2Upon expansion we get, 2x/3 - 10/3 - x/4 + 2/4 = 9/2 2x/3 - 10/3 - x/4 + 1/2 = 9/2 2x/3 - x/4 = 9/2 + 10/3 - 1/2By taking LCM for (3 and 4 is 12) (2 and 3 is 6)  $(2x \times 4)/12 - (x \times 3)/12 = (9 \times 3)/6 + (10 \times 2)/6 - (1 \times 3)/6$  8x/12 - 3x/12 = 27/6 + 20/6 - 3/6(8x - 3x)/12 = (27 + 20 - 3)/6



5x/12 = 44/6By using cross-multiplication we get,  $5x \times 6 = 44 \times 12$ 30x = 528x = 528/30= 264/15= 88/5Verification-LHS =  $2/3 (x - 5) - \frac{1}{4} (x - 2)$  $= 2/3 [(88/5) - 5] - \frac{1}{4} [(88/5) - 2]$  $= 2/3 [(88 - 25)/5] - \frac{1}{4} [(88 - 10)/5]$  $= 2/3 \times 63/5 - \frac{1}{4} \times 78/5$ =42/5-39/10=(84-39)/10= 45/10= 9/2= RHS





# EXERCISE 9.2

# **PAGE NO: 9.11**

## Solve each of the following equations and also check your results in each case:

1. (2x+5)/3 = 3x - 10Solution: (2x+5)/3 = 3x - 10Let us simplify, (2x+5)/3 - 3x = -10By taking LCM (2x + 5 - 9x)/3 = -10(-7x + 5)/3 = -10By using cross-multiplication we get, -7x + 5 = -30-7x = -30 - 5-7x = -35x = -35/-7= 5

Let us verify the given equation now, (2x+5)/3 = 3x - 10By substituting the value of 'x' we get,  $(2\times5+5)/3 = 3(5) - 10$  (10+5)/3 = 15-10 15/3 = 5 5 = 5Hence, the given equation is verified

#### 2. (a-8)/3 = (a-3)/2 Solution:

(a-8)/3 = (a-3)/2By using cross-multiplication we get, (a-8)2 = (a-3)32a - 16 = 3a - 92a - 3a = -9 + 16-a = 7a = -7

Let us verify the given equation now,



(a-8)/3 = (a-3)/2By substituting the value of 'a' we get, (-7 - 8)/3 = (-7 - 3)/2-15/3 = -10/2-5 = -5Hence, the given equation is verified

### 3. (7y + 2)/5 = (6y - 5)/11Solution: (7y + 2)/5 = (6y - 5)/11

(7y + 2)/5 = (6y - 5)/11By using cross-multiplication we get, (7y + 2)11 = (6y - 5)577y + 22 = 30y - 2577y - 30y = -25 - 2247y = -47y = -47/47y = -1

Let us verify the given equation now, (7y + 2)/5 = (6y - 5)/11By substituting the value of 'y' we get, (7(-1) + 2)/5 = (6(-1) - 5)/11 (-7 + 2)/5 = (-6 - 5)/11 -5/5 = -11/11 -1 = -1Hence, the given equation is verified

### 4. x - 2x + 2 - 16/3x + 5 = 3 - 7/2xSolution:

x - 2x + 2 - 16/3x + 5 = 3 - 7/2xLet us rearrange the equation x - 2x - 16x/3 + 7x/2 = 3 - 2 - 5By taking LCM for 2 and 3 which is 6 (6x - 12x - 32x + 21x)/6 = -4-17x/6 = -4By cross-multiplying  $-17x = -4 \times 6$ -17x = -24x = -24/-17 RD Sharma Solutions for Class 8 Maths Chapter 9 – Linear Equation in One Variable



x = 24/17

Let us verify the given equation now, x - 2x + 2 - 16/3x + 5 = 3 - 7/2xBy substituting the value of 'x' we get, 24/17 - 2(24/17) + 2 - (16/3)(24/17) + 5 = 3 - (7/2)(24/17) 24/17 - 48/17 + 2 - 384/51 + 5 = 3 - 168/34By taking 51 and 17 as the LCM we get, (72 - 144 + 102 - 384 + 255)/51 = (102 - 168)/34 -99/51 = -66/34 -33/17 = -33/17Hence, the given equation is verified

# 5. 1/2x + 7x - 6 = 7x + 1/4

### Solution:

1/2x + 7x - 6 = 7x + 1/4Let us rearrange the equation 1/2x + 7x - 7x = 1/4 + 6 (by taking LCM) 1/2x = (1+24)/41/2x = 25/4By cross-multiplying  $4x = 25 \times 2$ 4x = 50x = 50/4x = 25/2

Let us verify the given equation now, 1/2x + 7x - 6 = 7x + 1/4By substituting the value of 'x' we get, (1/2) (25/2) + 7(25/2) - 6 = 7(25/2) + 1/4 25/4 + 175/2 - 6 = 175/2 + 1/4By taking LCM for 4 and 2 is 4 (25 + 350 - 24)/4 = (350+1)/4 351/4 = 351/4Hence, the given equation is verified

6. 3/4x + 4x = 7/8 + 6x - 6Solution: 3/4x + 4x = 7/8 + 6x - 6



Let us rearrange the equation 3/4x + 4x - 6x = 7/8 - 6By taking 4 and 8 as LCM (3x + 16x - 24x)/4 = (7 - 48)/8 -5x/4 = -41/8By cross-multiplying -5x(8) = -41(4) -40x = -164 x = -164/-40 = 82/20= 41/10

Let us verify the given equation now, 3/4x + 4x = 7/8 + 6x - 6By substituting the value of 'x' we get, (3/4)(41/10) + 4(41/10) = 7/8 + 6(41/10) - 6 123/40 + 164/10 = 7/8 + 246/10 - 6 (123 + 656)/40 = (70 + 1968 - 480)/80 779/40 = 1558/80 779/40 = 779/40Hence, the given equation is verified

#### 7. 7x/2 - 5x/2 = 20x/3 + 10Solution:

7x/2 - 5x/2 = 20x/3 + 10Let us rearrange the equation 7x/2 - 5x/2 - 20x/3 = 10By taking LCM for 2 and 3 is 6 (21x - 15x - 40x)/6 = 10-34x/6 = 10By cross-multiplying -34x = 60x = 60/-34= -30/17

Let us verify the given equation now, 7x/2 - 5x/2 = 20x/3 + 10By substituting the value of 'x' we get, (7-/2)(-30/17) - (5/2)(-30/17) = (20/3)(-30/17) + 10



-210/34 + 150/34 = -600/51 + 10-30/17 = (-600+510)/51= -90/51-30/17 = -30/17Hence, the given equation is verified

#### 8. (6x+1)/2 + 1 = (7x-3)/3Solution:

(6x+1)/2 + 1 = (7x-3)/3 (6x+1+2)/2 = (7x-3)/3By cross-multiplying (6x+3)3 = (7x-3)2 18x + 9 = 14x - 6 18x - 14x = -6 - 9 4x = -15x = -15/4

Let us verify the given equation now, (6x+1)/2 + 1 = (7x-3)/3By substituting the value of 'x' we get, (6(-15/4) + 1)/2 + 1 = (7(-15/4) - 3)/3 (3(-15/2) + 1)/2 + 1 = (-105/4 - 3)/3 (-45/2 + 1)/2 + 1 = (-117/4)/3 (-43/4) + 1 = -117/12 (-43+4)/4 = -39/4 -39/4 = -39/4Hence, the given equation is verified

### 9. (3a-2)/3 + (2a+3)/2 = a + 7/6Solution:

(3a-2)/3 + (2a+3)/2 = a + 7/6Let us rearrange the equation (3a-2)/3 + (2a+3)/2 - a = 7/6By taking LCM for 2 and 3 which is 6 ((3a-2)2 + (2a+3)3 - 6a)/6 = 7/6(6a - 4 + 6a + 9 - 6a)/6 = 7/6(6a + 5)/6 = 7/66a + 5 = 76a = 7-5



6a = 2a = 2/6a = 1/3

Let us verify the given equation now, (3a-2)/3 + (2a+3)/2 = a + 7/6By substituting the value of 'a' we get, (3(1/3)-2)/3 + (2(1/3) + 3)/2 = 1/3 + 7/6 (1-2)/3 + (2/3 + 3)/2 = (2+7)/6 -1/3 + (11/3)/2 = 9/6 -1/3 + 11/6 = 3/2 (-2+11)/6 = 3/2 9/6 = 3/2 3/2 = 3/2Hence, the given equation is verified

# 10. x - (x-1)/2 = 1 - (x-2)/3Solution:

x - (x-1)/2 = 1 - (x-2)/3Let us rearrange the equation x - (x-1)/2 + (x-2)/3 = 1By taking LCM for 2 and 3 which is 6 (6x - (x-1)3 + (x-2)2)/6 = 1(6x - 3x + 3 + 2x - 4)/6 = 1(5x - 1)/6 = 1By cross-multiplying 5x - 1 = 65x = 6 + 1x = 7/5

Let us verify the given equation now, x - (x-1)/2 = 1 - (x-2)/3By substituting the value of 'x' we get, 7/5 - (7/5 - 1)/2 = 1 - (7/5 - 2)/3 7/5 - (2/5)/2 = 1 - (-3/5)/3 7/5 - 2/10 = 1 + 3/15 (14 - 2)/10 = (15+3)/15 12/10 = 18/156/5 = 6/5



Hence, the given equation is verified

# 11. 3x/4 - (x-1)/2 = (x-2)/3Solution:

3x/4 - (x-1)/2 = (x-2)/3Let us rearrange the equation 3x/4 - (x-1)/2 - (x-2)/3 = 0By taking LCM for 4, 2 and 3 which is 12 (9x - (x-1)6 - (x-2)4)/12 = 0(9x - 6x + 6 - 4x + 8)/12 = 0(-x + 14)/12 = 0By cross-multiplying -x + 14 = 0x = 14

Let us verify the given equation now, 3x/4 - (x-1)/2 = (x-2)/3By substituting the value of 'x' we get, 3(14)/4 - (14-1)/2 = (14-2)/3 42/4 - 13/2 = 12/3 (42 - 26)/4 = 4 16/4 = 4Hence, the given equation is verified

## 12. 5x/3 – (x-1)/4 = (x-3)/5 Solution:

5x/3 - (x-1)/4 = (x-3)/5Let us rearrange the equation 5x/3 - (x-1)/4 - (x-3)/5 = 0By taking LCM for 3, 4 and 5 which is 60  $((5x \times 20) - (x-1)15 - (x-3)12)/60 = 0$ (100x - 15x + 15 - 12x + 36)/60 = 0(73x + 51)/60 = 0By cross-multiplying 73x + 51 = 0x = -51/73

Let us verify the given equation now,



5x/3 - (x-1)/4 = (x-3)/5By substituting the value of 'x' we get, (20x - (x-1)3)/12 = (-51/73 - 3)/5(20x - 3x + 3)/12 = (-270/73)/5(17x + 3)/12 = -270/365(17(-51/73) + 3)/12 = -54/73(-867/73 + 3)/12 = -54/73((-867 + 219)/73)/12 = -54/73(-648)/876 = -54/73-54/73 = -54/73 Hence, the given equation is verified

### 13. (3x+1)/16 + (2x-3)/7 = (x+3)/8 + (3x-1)/14Solution:

(3x+1)/16 + (2x-3)/7 = (x+3)/8 + (3x-1)/14Let us rearrange the equation (3x+1)/16 + (2x-3)/7 - (x+3)/8 - (3x-1)/14 = 0By taking LCM for 16, 7, 8 and 14 which is 112 ((3x+1)7 + (2x-3)16 - (x+3)14 - (3x-1)8)/112 = 0(21x + 7 + 32x - 48 - 14x - 42 - 24x + 8)/112 = 0(21x + 32x - 14x - 24x + 7 - 48 - 42 + 8)/112 = 0(15x - 75)/112 = 0By cross-multiplying 15x - 75 = 015x = 75x = 75/15= 5

Let us verify the given equation now, (3x+1)/16 + (2x-3)/7 = (x+3)/8 + (3x-1)/14By substituting the value of 'x' we get, (3(5)+1)/16 + (2(5)-3)/7 = (5+3)/8 + (3(5)-1)/14 (15+1)/16 + (10-3)/7 = 8/8 + (15-1)/14 16/16 + 7/7 = 8/8 + 14/14 1 + 1 = 1 + 12 = 2

Hence, the given equation is verified



14. (1-2x)/7 - (2-3x)/8 = 3/2 + x/4Solution:

(1-2x)/7 - (2-3x)/8 = 3/2 + x/4Let us rearrange the equation (1-2x)/7 - (2-3x)/8 - x/4 = 3/2By taking LCM for 7, 8 and 4 which is 56 ((1-2x)8 - (2-3x)7 - 14x)/56 = 3/2(8 - 16x - 14 + 21x - 14x)/56 = 3/2(-9x - 6)/56 = 3/2By cross-multiplying 2(-9x-6) = 3(56)-18x - 12 = 168-18x = 168+12-18x = 180x = 180/-18x = -10

Let us verify the given equation now, (1-2x)/7 - (2-3x)/8 = 3/2 + x/4By substituting the value of 'x' we get, (1-2(-10))/7 - (2-3(-10))/8 = 3/2 + (-10)/4 (1+20)/7 - (2+30)/8 = 3/2 - 5/2 21/7 - 32/8 = 3/2 - 5/2 3 - 4 = -2/2 -1 = -1Hence, the given equation is verified

### 15. (9x+7)/2 - (x - (x-2)/7) = 36Solution:

(9x+7)/2 - (x - (x-2)/7) = 36Let us simplify the given equation into simple form (9x+7)/2 - (7x-x+2)/7 = 36By taking LCM for 2 and 7 is 14 (7(9x+7) - 2(6x+2))/14 = 36(63x+49 - 12x - 4)/14 = 36(51x + 45)/14 = 36By cross-multiplying 51x + 45 = 36(14)



51x + 45 = 504 51x = 504-45 51x = 459 x = 459/51= 9

Let us verify the given equation now, (9x+7)/2 - (x - (x-2)/7) = 36 (9x+7)/2 - (6x+2)/7 = 36By substituting the value of 'x' we get, (9(9)+7)/2 - (6(9)+2)/7 = 36 (81+7)/2 - (54+2)/7 = 36 88/2 - 56/7 = 36 44 - 8 = 36 36 = 36Hence, the given equation is verified

## 16. 0.18(5x - 4) = 0.5x + 0.8 Solution:

```
0.18(5x - 4) = 0.5x + 0.8
Let us rearrange the equation
0.18(5x - 4) - 0.5x = 0.8
0.90x - 0.72 - 0.5x = 0.8
0.90x - 0.5x = 0.8 + 0.72
0.40x = 1.52
x = 1.52/0.40
= 3.8
```

Let us verify the given equation now, 0.18(5x - 4) = 0.5x + 0.8By substituting the value of 'x' we get, 0.18(5(3.8)-4) = 0.5(3.8) + 0.8 0.18(19-4) = 1.9 + 0.8 2.7 = 2.7Hence, the given equation is verified

17. 2/3x - 3/2x = 1/12Solution: 2/3x - 3/2x = 1/12



By taking LCM for 3x and 2x which is 6x  $((2\times2) - (3\times3))/6x = 1/12$  (4-9)/6x = 1/12 -5/6x = 1/12By cross-multiplying 6x = -60 x = -60/6= -10

Let us verify the given equation now, 2/3x - 3/2x = 1/12By substituting the value of 'x' we get, 2/3(-10) - 3/2(-10) = 1/12 -2/30 + 3/20 = 1/12  $((-2\times2) + (3\times3))/60 = 1/12$  (-4+9)/60 = 1/12 5/60 = 1/12 1/12 = 1/12Hence, the given equation is verified

### 18. 4x/9 + 1/3 + 13x/108 = (8x+19)/18Solution:

4x/9 + 1/3 + 13x/108 = (8x+19)/18Let us rearrange the equation 4x/9 + 13x/108 - (8x+19)/18 = -1/3By taking LCM for 9, 108 and 18 which is 108  $((4x\times12) + 13x\times1 - (8x+19)6)/108 = -1/3$ (48x + 13x - 48x - 114)/108 = -1/3(13x - 114)/108 = -1/3By cross-multiplying (13x - 114)3 = -10839x - 342 = -10839x = -108 + 34239x = 234x = 234/39= 6

Let us verify the given equation now, 4x/9 + 1/3 + 13x/108 = (8x+19)/18



By substituting the value of 'x' we get, 4(6)/9 + 1/3 + 13(6)/108 = (8(6)+19)/18 24/9 + 1/3 + 78/108 = 67/18 8/3 + 1/3 + 13/18 = 67/18  $((8\times6) + (1\times6) + (13\times1))/18 = 67/18$  (48 + 6 + 13)/18 = 67/18 67/18 = 67/18Hence, the given equation is verified

### 19. (45-2x)/15 - (4x+10)/5 = (15-14x)/9Solution:

(45-2x)/15 - (4x+10)/5 = (15-14x)/9By rearranging (45-2x)/15 - (4x+10)/5 - (15-14x)/9 = 0By taking LCM for 15, 5 and 9 which is 45 ((45-2x)3 - (4x+10)9 - (15-14x)5)/45 = 0(135 - 6x - 36x - 90 - 75 + 70x)/45 = 0(28x - 30)/45 = 0By cross-multiplying 28x - 30 = 028x = 30x = 30/28= 15/14

Let us verify the given equation now, (45-2x)/15 - (4x+10)/5 = (15-14x)/9By substituting the value of 'x' we get, (45-2(15/14))/15 - (4(15/14) + 10)/5 = (15 - 14(15/14))/9 (45-15/7)/15 - (30/7 + 10)/5 = (15-15)/9 300/105 - 100/35 = 0 (300-300)/105 = 0 0 = 0Hence, the given equation is verified

## **20.** 5(7x+5)/3 - 23/3 = 13 - (4x-2)/3Solution: 5(7x+5)/3 - 23/3 = 13 - (4x-2)/3

By rearranging (35x + 25)/3 + (4x - 2)/3 = 13 + 23/3



(35x + 25 + 4x - 2)/3 = (39+23)/3(39x + 23)/3 = 62/3 By cross-multiplying (39x + 23)3 = 62(3) 39x + 23 = 62 39x = 62 - 23 39x = 39 x = 1

Let us verify the given equation now, 5(7x+5)/3 - 23/3 = 13 - (4x-2)/3By substituting the value of 'x' we get, (35x + 25)/3 - 23/3 = 13 - (4x-2)/3 (35+25)/3 - 23/3 = 13 - (4-2)/3 60/3 - 23/3 = 13 - 2/3 (60-23)/3 = (39-2)/3 37/3 = 37/3Hence, the given equation is verified

### 21. (7x-1)/4 - 1/3(2x - (1-x)/2) = 10/3Solution:

(7x-1)/4 - 1/3(2x - (1-x)/2) = 10/3Upon expansion (7x-1)/4 - (4x-1+x)/6 = 10/3(7x-1)/4 - (5x-1)/6 = 10/3By taking LCM for 4 and 6 is 24 ((7x-1)6 - (5x-1)4)/24 = 10/3(42x - 6 - 20x + 4)/24 = 10/3(22x - 2)/24 = 10/3By cross-multiplying 22x - 2 = 10(8)22x - 2 = 8022x = 80+222x = 82x = 82/22= 41/11

Let us verify the given equation now, (7x-1)/4 - 1/3(2x - (1-x)/2) = 10/3



By substituting the value of 'x' we get, (7x-1)/4 - (5x-1)/6 = 10/3 (7(41/11)-1)/4 - (5(41/11)-1)/6 = 10/3 (287/11 - 1)/4 - (205/11 - 1)/6 = 10/3 (287-11)/44 - (205-11)/66 = 10/3 (276/44 - 194/66 = 10/3 69/11 - 97/33 = 10/3  $((69\times3) - (97\times1))/33 = 10/3$   $((69\times3) - (97\times1))/33 = 10/3$  (207 - 97)/33 = 10/3 110/33 = 10/3Hence, the given equation is verified

#### 22. 0.5(x-0.4)/0.35 - 0.6(x-2.71)/0.42 = x + 6.1Solution:

 $\begin{array}{l} 0.5(x-0.4)/0.35 - 0.6(x-2.71)/0.42 = x + 6.1\\ \text{Let us simplify}\\ (0.5/0.35)(x-0.4) - (0.6/0.42)(x-2.71) = x + 6.1\\ (x-0.4)/0.7 - (x-2.71)/0.7 = x + 6.1\\ (x-0.4 - x + 2.71)/0.7 = x + 6.1\\ -0.4 + 2.71 = 0.7(x+6.1)\\ 0.7x = 2.71 - 0.4 - 4.27\\ = -1.96\\ x = -1.96/0.7\\ = -2.8\end{array}$ 

Let us verify the given equation now, 0.5(x-0.4)/0.35 - 0.6(x-2.71)/0.42 = x + 6.1By substituting the value of 'x' we get, 0.5(-2.8 - 0.4)/0.35 - 0.6(-2.8 - 2.71)/0.42 = -2.8 + 6.1 -1.6/0.35 + 3.306/0.42 = 3.3 -4.571 + 7.871 = 3.3 3.3 = 3.3Hence, the given equation is verified

#### 23. 6.5x + (19.5x - 32.5)/2 = 6.5x + 13 + (13x - 26)/2Solution: 6.5x + (10.5x - 32.5)/2 = 6.5x + 13 + (13x - 26)/2

6.5x + (19.5x - 32.5)/2 = 6.5x + 13 + (13x - 26)/2By rearranging

B BYJU'S

RD Sharma Solutions for Class 8 Maths Chapter 9 – Linear Equation in One Variable

6.5x + (19.5x - 32.5)/2 - 6.5x - (13x - 26)/2 = 13(19.5x - 32.5)/2 - (13x - 26)/2 = 13 (19.5x - 32.5 - 13x + 26)/2 = 13 (6.5x - 6.5)/2 = 13  $6.5x - 6.5 = 13 \times 2$ 6.5x - 6.5 = 266.5x = 26 + 6.56.5x = 32.5x = 32.5/6.5= 5

Let us verify the given equation now, 6.5x + (19.5x - 32.5)/2 = 6.5x + 13 + (13x - 26)/2By substituting the value of 'x' we get, 6.5(5) + (19.5(5) - 32.5)/2 = 6.5(5) + 13 + (13(5) - 26)/2 32.5 + (97.5 - 32.5)/2 = 32.5 + 13 + (65 - 26)/2 32.5 + 65/2 = 45.5 + 39/2 (65 + 65)/2 = (91+39)/2 130/2 = 130/2 65 = 65Hence, the given equation is verified

24. (3x - 8) (3x + 2) - (4x - 11) (2x + 1) = (x - 3) (x + 7)Solution: (3x - 8) (3x + 2) - (4x - 11) (2x + 1) = (x - 3) (x + 7)Let us simplify  $9x^{2} + 6x - 24x - 16 - 8x^{2} - 4x + 22x + 11 = x^{2} + 7x - 3x - 21$  $9x^{2} + 6x - 24x - 16 - 8x^{2} - 4x + 22x + 11 - x^{2} - 7x + 3x + 21 = 0$  $9x^{2} - 8x^{2} - x^{2} + 6x - 24x - 4x + 22x - 7x + 3x - 16 + 21 + 11 = 0$ -4x + 16 = 0-4x = -16x = 4

Let us verify the given equation now, (3x - 8) (3x + 2) - (4x - 11) (2x + 1) = (x - 3) (x + 7)By substituting the value of 'x' we get, (3(4) - 8) (3(4) + 2) - (4(4) - 11) (2(4) + 1) = (4 - 3) (4 + 7) (12-8) (12+2) - (16-11) (8+1) = 1(11)4 (14) - 5(9) = 11



56 - 45 = 1111 = 11Hence, the given equation is verified

# 25. $[(2x+3) + (x+5)]^2 + [(2x+3) - (x+5)]^2 = 10x^2 + 92$ Solution:

 $[(2x+3) + (x+5)]^{2} + [(2x+3) - (x+5)]^{2} = 10x^{2} + 92$ Let us simplify the given equation  $[3x + 8]^{2} + [x - 2]^{2} = 10x^{2} + 92$ By using the formula  $(a+b)^{2}$  $9x^{2} + 48x + 64 + x^{2} - 4x + 4 = 10x^{2} + 92$ By rearranging  $9x^{2} - 10x^{2} + x^{2} + 48x - 4x = 92 - 64 - 4$ 44x = 24x = 24/44= 6/11

```
Let us verify the given equation now,

[(2x+3) + (x+5)]^{2} + [(2x+3) - (x+5)]^{2} = 10x^{2} + 92
By substituting the value of 'x' we get,

[2(6/11) + 3 + (6/11) + 5]^{2} + [2(6/11) + 3 - (6/11) - 5]^{2} = 10(6/11)^{2} + 92
[(12/11 + 3) + (6/11 + 5)]^{2} + [(12/11 + 3) - (6/11 + 5)]^{2} = 10(6/11)^{2} + 92
[(12+33)/11 + (6+55)/11]^{2} + [(12+33)/11 - (6+55)/11]^{2} = 10(6/11)^{2} + 92
[(45/11) + (61/11)]^{2} + [(45/11) - (61/11)]^{2} = 360/121 + 92
(106/11)^{2} + (-16/11)^{2} = (360 + 11132)/121
11236/121 + 256/121 = 11492/121
Hence, the given equation is verified
```





# **EXERCISE 9.3**

# P&GE NO: 9.17

## Solve the following equations and verify your answer:

1. (2x-3) / (3x+2) = -2/3Solution: We have, (2x-3) / (3x+2) = -2/3Let us perform cross-multiplication we get, 3(2x - 3) = -2(3x + 2)6x - 9 = -6x - 4When rearranged, 6x + 6x = 9 - 412x = 5x = 5/12

Now let us verify the given equation, (2x-3) / (3x+2) = -2/3By substituting the value of 'x' we get, (2(5/12) - 3) / (3(5/12) + 2) = -2/3 ((5/6)-3) / ((5/4) + 2) = -2/3 ((5-18)/6) / ((5+8)/4) = -2/3 (-13/6) / (13/4) = -2/3  $(-13/6) \times (4/13) = -2/3$  -4/6 = -2/3 -2/3 = -2/3Hence, the given equation is verified

# 2. (2-y) / (y+7) = 3/5

#### **Solution:**

We have, (2-y) / (y+7) = 3/5Let us perform cross-multiplication we get, 5(2-y) = 3(y+7) 10 - 5y = 3y + 21When rearranged, 10 - 21 = 3y + 5y 8y = -11y = -11/8



Now let us verify the given equation, (2-y) / (y+7) = 3/5By substituting the value of 'x' we get, (2 - (-11/8)) / ((-11/8) + 7) = 3/5 ((16+11)/8) / ((-11+56)/8) = 3/5 (27/8) / (45/8) = 3/5  $(27/8) \times (8/45) = 3/5$  (27/45 = 3/5)Hence, the given equation is verified

# 3. (5x - 7) / (3x) = 2


## Solution:

We have, (5x - 7) / (3x) = 2Let us perform cross-multiplication we get, 5x - 7 = 2(3x) 5x - 7 = 6x 5x - 6x = 7 -x = 7x = -7

Now let us verify the given equation, (5x - 7) / (3x) = 2By substituting the value of 'x' we get, (5(-7) - 7) / (3(-7)) = 2 (-35 - 7) / -21 = 2 -42/-21 = 2 2 = 2Hence, the given equation is verified

# 4. (3x+5) / (2x + 7) = 4 Solution:

We have, (3x+5)/(2x+7) = 4Let us perform cross-multiplication we get, 3x + 5 = 4(2x+7)3x + 5 = 8x + 28





3x - 8x = 28 - 5-5x = 23x = -23/5

Now let us verify the given equation, (3x+5) / (2x + 7) = 4By substituting the value of 'x' we get, (3(-23/5) + 5) / (2(-23/5) + 7) = 4 (-69/5 + 5) / (-46/5 + 7) = 4 (-69+25)/5 / (-46+35)/5 = 4 -44/5 / -11/5 = 4  $-44/5 \times 5/-11 = 4$  44/11 = 4Hence, the given equation is verified

# 5. (2y + 5) / (y + 4) = 1 Solution:

We have, (2y + 5) / (y + 4) = 1Let us perform cross-multiplication we get, 2y + 5 = y + 4 2y - y = 4 - 5y = -1

Now let us verify the given equation, (2y + 5) / (y + 4) = 1By substituting the value of 'y' we get, (2(-1) + 5) / (-1 + 4) = 1 (-2+5) / 3 = 1 3/3 = 1 1 = 1Hence, the given equation is verified

# 6. (2x + 1) / (3x - 2) = 5/9 Solution:

We have, (2x + 1) / (3x - 2) = 5/9Let us perform cross-multiplication we get,



9(2x + 1) = 5(3x - 2) 18x + 9 = 15x - 10 18x - 15x = -10 - 9 3x = -19x = -19/3

Now let us verify the given equation, (2x + 1) / (3x - 2) = 5/9By substituting the value of 'x' we get, (2(-19/3) + 1) / (3(-19/3) - 2) = 5/9 (-38/3 + 1) / (-57/3 - 2) = 5/9 (-38 + 3)/3 / (-57 - 6)/3 = 5/9  $-35/3 \times 3/-63 = 5/9$  -35/-63 = 5/9 5/9 = 5/9Hence, the given equation is verified

### 7. (1 - 9y) / (19 - 3y) = 5/8Solution: We have, (1 - 9y) / (19 - 3y) = 5/8Let us perform cross-multiplication we get, 8(1 - 9y) = 5(19 - 3y)

8 - 72y = 95 - 15y 8 - 95 = 72y - 15y 57y = -87 y = -87/57= -29/19

Now let us verify the given equation, (1-9y) / (19-3y) = 5/8By substituting the value of 'y' we get, (1-9(-29/19)) / (19-3(-29/19)) = 5/8 (19+261)/19 / (361+87)/19 = 5/8  $280/19 \times 19/448 = 5/8$  280/448 = 5/8 5/8 = 5/8Hence, the given equation is verified



### 8. 2x / (3x + 1) = 1 Solution:

We have, 2x / (3x + 1) = 1Let us perform cross-multiplication we get, 2x = 1(3x + 1) 2x = 3x + 1 2x - 3x = 1 -x = 1x = -1

Now let us verify the given equation, 2x / (3x + 1) = 1By substituting the value of 'x' we get, 2(-1) / (3(-1) + 1) = 1 -2 / (-3+1) = 1 -2/-2 = 1 1 = 1Hence, the given equation is verified

# 9. y - (7 - 8y)/9y - (3 + 4y) = 2/3Solution: We have,

y - (7 - 8y)/9y - (3 + 4y) = 2/3(y - 7 + 8y) / (9y - 3 - 4y) = 2/3 (-7 + 9y) / (5y - 3) = 2/3 Let us perform cross-multiplication we get, 3(-7 + 9y) = 2(5y - 3)-21 + 27y = 10y - 627y - 10y = 21 - 617y = 15y = 15/17

Now let us verify the given equation, y - (7 - 8y)/9y - (3 + 4y) = 2/3By substituting the value of 'y' we get, 15/17 - (7-8(15/17))/9(15/17) - (3 + 4(15/17)) = 2/3 15/17 - (7 - 120/17) / 135/17 - (3 + 60/17) = 2/315/17 - ((119-120)/17) / 135/17 - ((51+60)/17) = 2/3



15/17 - (-1/17) / 135/17 - (111/17) = 2/23((15 + 1)/17) / ((135-111)/17) = 2/316/17 / 24/17 = 2/316/24 = 2/32/3 = 2/3Hence, the given equation is verified

# 10. 6/ 2x - (3 - 4x) = 2/3

## Solution:

We have, 6/2x - (3 - 4x) = 2/3 6/(2x - 3 + 4x) = 2/3 6/(6x - 3) = 2/3Let us perform cross-multiplication we get, 3(6) = 2(6x - 3) 18 = 12x - 6 12x = 18 + 6 12x = 24 x = 24/12= 2

Now let us verify the given equation, 6/2x - (3 - 4x) = 2/3 6/(6x - 3) = 2/3By substituting the value of 'x' we get, 6/(6(2) - 3) = 2/3 6/(12-3) = 2/3 6/9 = 2/3 2/3 = 2/3Hence, the given equation is verified

### 11. 2/3x - 3/2x = 1/12

Solution:

We have, 2/3x - 3/2x = 1/12By taking LCM for 2 and 3 which is 6 4-9/6x = 1/12 -5/6x = 1/12By cross-multiplying we get,





12(-5) = 1 (6x)-60 = 6x x = -60/6= -10

Now let us verify the given equation, 2/3x - 3/2x = 1/12By substituting the value of 'x' we get, 2/3(-10) - 3/2(-10) = 1/12 2/-30 - 3/-20 = 1/12 -4+6/60 = 1/12 5/60 = 1/12 1/12 = 1/12Hence, the given equation is verified

12. (3x + 5)/(4x + 2) = (3x + 4)/(4x + 7)Solution: We have, (3x + 5)/(4x + 2) = (3x + 4)/(4x + 7)(3x + 5)/(4x + 2) - (3x + 4)/(4x + 7) = 0By taking LCM as (4x + 2)(4x + 7)((3x + 5)(4x + 7) - (3x + 4)(4x + 2))/(4x + 2)(4x + 7) = 0

By cross-multiplying we get, (3x + 5) (4x + 7) - (3x + 4) (4x + 2) = 0 (3x + 5) (4x + 7) - (3x + 4) (4x + 2) = 0  $12x^2 + 21x + 20x + 35 - 12x^2 - 6x - 16x - 8 = 0$  19x + 35 - 8 = 0 19x = -27x = -27/19

Now let us verify the given equation, (3x + 5)/(4x + 2) = (3x + 4)/(4x + 7)By substituting the value of 'x' we get, (3(-27/19) + 5)/(4(-27/19) + 2) = (3(-27/19) + 4)/(4(-27/19) + 7) (-81/19 + 5)/(-108/19 + 2) = (-81/19 + 4)/(-108/19 + 7) ((-81+95)/19)/((-108+38)/19) = ((-81+76)/19)/((-108+133)/19) 14/19/-70/19 = -5/19/25/19 -14/70 = -5/25-1/5 = -1/5



Hence, the given equation is verified

# 13. (7x - 2) / (5x - 1) = (7x +3)/(5x + 4) Solution:

We have, (7x - 2) / (5x - 1) = (7x + 3)/(5x + 4) (7x - 2) / (5x - 1) - (7x + 3)/(5x + 4) = 0By taking LCM as (5x - 1) (5x + 4) ((7x-2) (5x+4) - (7x+3)(5x-1)) / (5x - 1) (5x + 4) = 0By cross-multiplying we get, (7x-2) (5x+4) - (7x+3)(5x-1) = 0Upon simplification,  $35x^2 + 28x - 10x - 8 - 35x^2 + 7x - 15x + 3 = 0$  10x - 5 = 0 10x = 5 x = 5/10= 1/2

Now let us verify the given equation, (7x - 2) / (5x - 1) = (7x + 3)/(5x + 4)By substituting the value of 'x' we get, (7(1/2) - 2) / (5(1/2) - 1) = (7(1/2) + 3) / (5(1/2) + 4) (7/2 - 2) / (5/2 - 1) = (7/2 + 3) / (5/2 + 4) ((7-4)/2) / ((5-2)/2) = ((7+6)/2) / ((5+8)/2) (3/2) / (3/2) = (13/2) / (13/2)1 = 1

Hence, the given equation is verified

# 14. $((x+1)/(x+2))^2 = (x+2) / (x+4)$ Solution:

We have,  $((x+1)/(x+2))^{2} = (x+2) / (x + 4)$   $(x+1)^{2} / (x+2)^{2} - (x+2) / (x + 4) = 0$ By taking LCM as  $(x+2)^{2} (x+4)$   $((x+1)^{2} (x+4) - (x+2) (x+2)^{2}) / (x+2)^{2} (x+4) = 0$ By cross-multiplying we get,  $(x+1)^{2} (x+4) - (x+2) (x+2)^{2} = 0$ Let us expand the equation  $(x^{2} + 2x + 1) (x + 4) - (x + 2) (x^{2} + 4x + 4) = 0$ 



$$x^{3} + 2x^{2} + x + 4x^{2} + 8x + 4 - (x^{3} + 4x^{2} + 4x + 2x^{2} + 8x + 8) = 0$$
  

$$x^{3} + 2x^{2} + x + 4x^{2} + 8x + 4 - x^{3} - 4x^{2} - 4x - 2x^{2} - 8x - 8 = 0$$
  

$$-3x - 4 = 0$$
  

$$x = -4/3$$

Now let us verify the given equation,  $((x+1)/(x+2))^2 = (x+2) / (x + 4)$ By substituting the value of 'x' we get,  $(x+1)^2 / (x+2)^2 = (x+2) / (x + 4)$   $(-4/3 + 1)^2 / (-4/3 + 2)^2 = (-4/3 + 2) / (-4/3 + 4)$   $((-4+3)/3)^2 / ((-4+6)/3)^2 = ((-4+6)/3) / ((-4+12)/3)$   $(-1/3)^2 / (2/3)^2 = (2/3) / (8/3)$  1/9 / 4/9 = 2/3 / 8/3 1/4 = 2/8 1/4 = 1/4Hence, the given equation is verified

#### 15. $((x+1)/(x-4))^2 = (x+8)/(x-2)$ Solution: We have

We have,  $((x+1)/(x-4))^2 = (x+8)/(x-2)$   $(x+1)^2/(x-4)^2 - (x+8)/(x-2) = 0$ By taking LCM as  $(x-4)^2 (x-2)$   $((x+1)^2 (x-2) - (x+8) (x-4)^2)/(x-4)^2 (x-2) = 0$ By cross-multiplying we get,  $(x+1)^2 (x-2) - (x+8) (x-4)^2 = 0$ Upon expansion we get,  $(x^2 + 2x + 1) (x-2) - ((x+8) (x^2 - 8x + 16)) = 0$   $x^3 + 2x^2 + x - 2x^2 - 4x - 2 - (x^3 - 8x^2 + 16x + 8x^2 - 64x + 128) = 0$   $x^3 + 2x^2 + x - 2x^2 - 4x - 2 - x^3 + 8x^2 - 16x - 8x^2 + 64x - 128 = 0$  45x - 130 = 0 x = 130/45= 26/9

Now let us verify the given equation,  $((x+1)/(x-4))^2 = (x+8)/(x-2)$   $(x+1)^2/(x-4)^2 = (x+8)/(x-2)$ By substituting the value of 'x' we get,  $(26/9 + 1)^2/(26/9 - 4)^2 = (26/9 + 8)/(26/9 - 2)$ 



 $((26+9)/9)^2 / ((26-36)/9)^2 = ((26+72)/9) / ((26-18)/9)$  $(35/9)^2 / (-10/9)^2 = (98/9) / (8/9)$  $(35/-10)^2 = (98/8)$  $(7/2)^2 = 49/4$ Hence, the given equation is verified

# 16. (9x-7)/(3x+5) = (3x-4)/(x+6)**Solution:** We have, (9x-7)/(3x+5) = (3x-4)/(x+6)(9x-7)/(3x+5) - (3x-4)/(x+6) = 0By taking LCM as (3x+5)(x+6)((9x-7)(x+6) - (3x-4)(3x+5)) / (3x+5)(x+6) = 0By cross-multiplying we get, (9x-7)(x+6) - (3x-4)(3x+5) = 0Upon expansion we get, $9x^{2} + 54x - 7x - 42 - (9x^{2} + 15x - 12x - 20) = 0$ 44x - 22 = 044x = 22x = 22/44= 2/4= 1/2

Now let us verify the given equation, (9x-7)/(3x+5) = (3x-4)/(x+6)By substituting the value of 'x' we get, (9(1/2) - 7) / (3(1/2) + 5) = (3(1/2) - 4) / ((1/2) + 6) (9/2 - 7) / (3/2 + 5) = (3/2 - 4) / (1/2 + 6) ((9-14)/2) / ((3+10)/2) = ((3-8)/2) / ((1+12)/2) -5/2 / 13/2 = -5/2 / 13/2 -5/13 = -5/13Hence, the given equation is verified

# 17. (x+2)/(x+5) = x/(x+6) Solution:

We have, (x+2)/(x+5) = x/(x+6)(x+2)/(x+5) - x/(x+6) = 0



By taking LCM as (x+5) (x+6) ((x+2) (x+6) - x(x+5)) / (x+5) (x+6) = 0By cross-multiplying we get, (x+2) (x+6) - x(x+5) = 0Upon expansion,  $x^{2} + 8x + 12 - x^{2} - 5x = 0$  3x + 12 = 0 3x = -12 x = -12/3= -4

Now let us verify the given equation, (x+2)/(x+5) = x/(x+6)By substituting the value of 'x' we get, (-4+2)/(-4+5) = -4/(-4+6) -2/1 = -4/(2) -2 = -2Hence, the given equation is verified

## 18. 2x - (7-5x) / 9x - (3+4x) = 7/6Solution: We have, 2x - (7-5x) / 9x - (3+4x) = 7/6(2x - 7 + 5x) / (9x - 3 - 4x) = 7/6(7x - 7) / (5x - 3) = 7/6By cross-multiplying we get, 6(7x - 7) = 7(5x - 3)42x - 42 = 35x - 2142x - 35x = -21 + 427x = 21x = 21/7= 3

Now let us verify the given equation, 2x - (7-5x) / 9x - (3+4x) = 7/6 (7x - 7) / (5x - 3) = 7/6By substituting the value of 'x' we get, (7(3) -7) / (5(3) - 3) = 7/6(21-7) / (15-3) = 7/6



14/12 = 7/67/6 = 7/6Hence, the given equation is verified

# 19. (15(2-x) - 5(x+6)) / (1-3x) = 10Solution:

We have, 15(2-x) - 5(x+6) / (1-3x) = 10 (30-15x) - (5x + 30) / (1-3x) = 10By cross-multiplying we get, (30-15x) - (5x + 30) = 10(1-3x) 30-15x - 5x - 30 = 10 - 30x 30-15x - 5x - 30 + 30x = 10 10x = 10 x = 10/10= 1

Now let us verify the given equation, (15(2-x) - 5(x+6)) / (1-3x) = 10By substituting the value of 'x' we get, (15(2-1) - 5(1+6)) / (1-3) = 10 (15 - 5(7))/-2 = 10 (15-35)/-2 = 10 -20/-2 = 10 10 = 10Hence, the given equation is verified

# 20. (x+3)/(x-3) + (x+2)/(x-2) = 2 Solution:

We have, (x+3)/(x-3) + (x+2)/(x-2) = 2By taking LCM as (x-3) (x-2) ((x+3)(x-2) + (x+2) (x-3)) / (x-3) (x-2) = 2By cross-multiplying we get, (x+3)(x-2) + (x+2) (x-3) = 2 ((x-3) (x-2))Upon expansion,  $x^2 + 3x - 2x - 6 + x^2 - 3x + 2x - 6 = 2(x^2 - 3x - 2x + 6)$   $2x^2 - 12 = 2x^2 - 10x + 12$  $2x^2 - 2x^2 + 10x = 12 + 12$ 

RD Sharma Solutions for Class 8 Maths Chapter 9 – Linear Equation in One Variable



10x = 24x = 24/10 = 12/5

Now let us verify the given equation, (x+3)/(x-3) + (x+2)/(x-2) = 2By substituting the value of 'x' we get, (12/5 + 3)/(12/5 - 3) + (12/5 + 2)/(12/5 - 2) = 2 ((12+15)/5)/((12-15)/5) + ((12+10)/5)/((12-10)/5) = 2 (27/5)/(-3/5) + (22/5)/(2/5) = 2 -27/3 + 22/2 = 2  $((-27\times2) + (22\times3))/6 = 2$  (-54 + 66)/6 = 2 12/6 = 2 2 = 2Hence, the given equation is verified

Hence, the given equation is verified

## 21. $((x+2) (2x-3) - 2x^2 + 6)/(x-5) = 2$ Solution: We have, $((x+2) (2x-3) - 2x^2 + 6)/(x-5) = 2$

 $((x+2) (2x-3) - 2x^{2} + 6)/(x-3) = 2$ By cross-multiplying we get,  $(x+2) (2x-3) - 2x^{2} + 6) = 2(x-5)$  $2x^{2} - 3x + 4x - 6 - 2x^{2} + 6 = 2x - 10$ x = 2x - 10x - 2x = -10-x = -10x = 10

Now let us verify the given equation,  $((x+2) (2x-3) - 2x^2 + 6)/(x-5) = 2$ By substituting the value of 'x' we get,  $((10+2) (2(10) - 3) - 2(10)^2 + 6)/(10-5) = 2$  (12(17) - 200 + 6)/5 = 2 (204 - 194)/5 = 2 10/5 = 2 2 = 2Hence, the given equation is verified



22.  $(x^2 - (x+1) (x+2))/(5x+1) = 6$ Solution: We have,  $(x^2 - (x+1) (x+2))/(5x+1) = 6$ By cross-multiplying we get,

 $(x^{2} - (x+1) (x+2)) = 6(5x+1)$   $x^{2} - x^{2} - 2x - x - 2 = 30x + 6$  -3x - 2 = 30x + 6 30x + 3x = -2 - 6 33x = -8x = -8/33

Now let us verify the given equation,  $(x^2 - (x+1) (x+2))/(5x+1) = 6$ By substituting the value of 'x' we get,  $((-8/33)^2 - ((-8/33)+1) (-8/33+2))/(5(-8/33)+1) = 6$  (64/1089 - ((-8+33)/33) ((-8+66)/33)) / (-40+33)/33) = 6 (64/1089 - (25/33) (58/33)) / (-7/33) = 6 (64/1089 - 1450/1089) / (-7/33) = 6 ((64-1450)/1089 / (-7/33)) = 6  $-1386/1089 \times 33/-7 = 6$   $1386 \times 33 / 1089 \times -7 = 6$  6 = 6Hence, the given equation is verified

### 23. ((2x+3) – (5x-7))/(6x+11) = -8/3 Solution:

We have, ((2x+3) - (5x-7))/(6x+11) = -8/3By cross-multiplying we get, 3((2x+3) - (5x-7)) = -8(6x+11) 3(2x + 3 - 5x + 7) = -48x - 88 3(-3x + 10) = -48x - 88 -9x + 30 = -48x - 88 -9x + 48x = -88 - 30 39x = -118x = -118/39

Now let us verify the given equation,



((2x+3) - (5x-7))/(6x+11) = -8/3By substituting the value of 'x' we get, ((2(-118/39) + 3) - (5(-118/39) - 7)) / (6(-118/39) + 11) = -8/3((-336/39 + 3) - (-590/39 - 7)) / (-708/39 + 11) = -8/3(((-336+117)/39) - ((-590-273)/39)) / ((-708+429)/39) = -8/3(-219+863)/39 / (-279)/39 = -8/3644/-279 = -8/3-8/3 = -8/3Hence, the given equation is verified

#### 24. Find the positive value of x for which the given equation is satisfied:

```
(i) (x^2 - 9)/(5 + x^2) = -5/9
Solution:
We have,
(x^2 - 9)/(5 + x^2) = -5/9
By cross-multiplying we get,
9(x^2 - 9) = -5(5 + x^2)
9x^2 - 81 = -25 - 5x^2
9x^2 + 5x^2 = -25 + 81
14x^2 = 56
x^2 = 56/14
x^2 = 4
\mathbf{x} = \sqrt{4}
  = 2
(ii) (y^2 + 4)/(3y^2 + 7) = 1/2
Solution:
We have.
(y^2 + 4)/(3y^2 + 7) = 1/2
By cross-multiplying we get,
2(y^2 + 4) = 1(3y^2 + 7)
2y^2 + 8 = 3y^2 + 7
3y^2 - 2y^2 = 7 - 8
y^2 = -1
y = \sqrt{-1}
  = 1
```



### **EXERCISE 9.4**

### PAGE NO: 9.29

### **1.** Four-fifth of a number is more than three-fourth of the number by **4.** Find the number.

### Solution:

Let us consider the number as 'x' So, Three-fourth of the number is 3x/4Fourth-fifth of the number is 4x/54x/5 - 3x/4 = 4By taking LCM of 5 and 4 is 20 (16x - 15x)/20 = 4By cross-multiplying we get, 16x - 15x = 4(20)x = 80 $\therefore$  The number is 80.

### 2. The difference between the squares of two consecutive numbers is 31. Find the numbers.

### Solution:

Let the two consecutive numbers be x and (x - 1)So now,  $x^{2} - (x-1)^{2} = 31$ By using the formula  $(a-b)^{2} = a^{2} + b^{2} - 2ab$  $x^{2} - (x^{2} - 2x + 1) = 31$  $x^{2} - x^{2} + 2x - 1 = 31$ 2x - 1 = 312x = 31 + 12x = 32x = 32/2= 16Two consecutive numbers are, x and (x-1) : 16 and (16-1) = 15

 $\therefore$  The two consecutive numbers are 16 and 15.

### **3. Find a number whose double is 45 greater than its half. Solution:**

Let us consider the number as 'x' So, 2x - x/2 = 45(4x-x)/2 = 45



By cross-multiplying we get, 3x = 90 x = 90/3 = 30 $\therefore$  The number is 30.

### 4. Find a number such that when 5 is subtracted from 5 times that number, the result is 4 more than twice the number. Solution:

Let us consider the number as 'x' Then, five times the number will be 5x And, two times the number will be 2x So, 5x - 5 = 2x + 4 5x - 2x = 5 + 4 3x = 9 x = 9/3 x = 3 $\therefore$  The number is 3.

### 5. A number whose fifth part increased by 5 is equal to its fourth part diminished by5. Find the number.

### Solution:

Let us consider the number as 'x' So, x/5 + 5 = x/4 - 5x/5 - x/4 = -5 - 5By taking LCM for 5 and 4 which is 20 (4x-5x)/20 = -10By cross-multiplying we get, 4x - 5x = -10(20)-x = -200x = 200 $\therefore$  The number is 200.

## 6. A number consists of two digits whose sum is 9. If 27 is subtracted from the number the digits are reversed. Find the number. Solution:

We know that one of the digit be 'x'



The other digit is 9-x So, the two digit number is 10(9-x) + xThe number obtained after interchanging the digits is 10x + (9-x)10(9-x) + x - 27 = 10x + (9-x)Upon simplification, 90 - 10x + x - 27 = 10x + 9 - x-10x + x - 10x + x = 9 - 90 + 27-18x = -54x = 54/18= 9/3= 3 The two digit number is 10(9-x) + xSubstituting the value of x we get, 10(9-x) + x10(9 - 3) + 310(6) + 360 + 363  $\therefore$  The number is 63.

### 7. Divide 184 into two parts such that one-third of one part may exceed one-seventh of another part by 8.

#### Solution:

Let one of the number be 'x' The other number as 184 - xSo, One-third of one part may exceed one-seventh of another part by 8. x/3 - (184-x)/7 = 8LCM for 3 and 7 is 21 (7x - 552 + 3x)/21 = 8By cross-multiplying we get, (7x - 552 + 3x) = 8(21) 10x - 552 = 168 10x = 168 + 552 10x = 720 x = 720/10 = 72 $\therefore$  One of the number is 72 and other number is 184 - x => 184 - 72 = 112.



## 8. The numerator of a fraction is 6 less than the denominator. If 3 is added to the numerator, the fraction is equal to 2/3. What is the original fraction equal to? Solution:

Let us consider the denominator as x and numerator as (x-6)By using the formula, Fraction = numerator/denominator = (x-6)/x(x - 6 + 3)/x = 2/3(x - 3)/x = 2/3By cross-multiplying 3(x-3) = 2x3x - 9 = 2x3x - 2x = 9x = 9 $\therefore$  The denominator is x = 9, numerator is (x-6) = (9-6) = 3And the fraction = numerator/denominator = (x-6)/x = 3/9 = 1/3

### 9. A sum of Rs 800 is in the form of denominations of Rs 10 and Rs 20. If the total number of notes be 50. Find the number of notes of each type. Solution:

Let the number of 10Rs notes be x Number of 20Rs notes be 50 - xAmount due to 10Rs notes  $= 10 \times x = 10x$ Amount due to 20Rs notes  $= 20 \times (50 - x) = 1000 - 20x$ So the total amount is Rs 800 10x + 1000 - 20x = 800 -10x = 800 - 1000 -10x = -200 x = -200/-10 = 20  $\therefore$  The number of 10Rs notes is 20 Number of 20Rs notes are 50 - 20 = 30

10. Seeta Devi has Rs 9 in fifty-paise and twenty five-paise coins. She has twice as many twenty- five paise coins as she has fifty- paise coins. How many coins of each kind does she have?

#### Solution:

Let the number of fifty paise coins be x Number of twenty five paise coins be 2xAmount due to fifty paise coins =  $(50 \times x)/100 = 0.50x$ 



Amount due to twenty five paise coins =  $(25 \times 2x)/100 = 0.50x$ So the total amount is Rs 9 0.50x + 0.50x = 91x = 9x = 9 $\therefore$  The number of fifty paise coins is x = 9Number of twenty five paise coins,  $2x = 2 \times 9 = 18$ 

11. Sunita is twice as old as Ashima. If six years is subtracted from Ashima's age and four years added to Sunita's age, then Sunita will be four times Ashima's age. How old were they two years ago?

#### Solution:

Let the present age of Ashima be 'x' years The present age of Sunita is 2x years Ashima's new age = (x - 6) years Sunita's new age = (2x + 4) years So, (2x + 4) = 4 (x - 6) 2x + 4 = 4x - 24 2x - 4x = -24 - 4 -2x = -28 x = -28/-2 = 14  $\therefore$  Age of Ashima is x years = 14 years Age of Sunita is 2x years = 2(14) = 28 years Two years ago, age of Ashima is 14 - 2 = 12 years, age of Sunita = 28 - 2 = 26 years

### 12. The ages of Sonu and Monu are in the ratio 7:5 Ten years hence, the ratio of their ages will be 9:7 find their present ages. Solution:

Let the present age of Sonu be 7x years Present age of Monu be 5x years Sonu's age after 10 years = (7x + 10) years Monu's age after 10 years = (5x + 10) years So, (7x + 10) / (5x + 10) = 9/7by using cross-multiplication we get, 7(7x + 10) = 9(5x + 10)49x + 70 = 45x + 9049x - 45x = 90 - 70



4x = 20x = 20/4 = 5  $\therefore$  Present age of Sonu is 7x = 7(5) = 35 years

Present age of Monu is 5x = 5(5) = 25 years


### 13. Five years ago a man was seven times as old as his son. Five years hence, the father will be three times as old as his son. Find their present ages. Solution:

Let the age of son five years ago be x years The age of man five years ago be 7x years After five years, son's age is x + 5 years After five years father's age is 7x + 5 years So, since five years, the relation in their ages are 7x + 5 + 5 = 3(x + 5 + 5) 7x + 10 = 3x + 15 + 15 7x + 10 = 3x + 30 7x - 3x = 30 - 10 4x = 20 x = 5 $\therefore$  Present father's age is 7x + 5 = 7(5) + 5 = 35 + 5 = 40 years

Present son's age is x + 5 = 5 + 5 = 10 years

### 14. I am currently 5 times as old as my son. In 6 years time I will be three times as old as he will be then. What are our ages now? Solution:

Let the present son's age be x years Present father's age be 5x years Son's age after 6 years = (x + 6) years Fathers' age after 6 years = (5x + 6) years So, 5x + 6 = 3(x + 6) 5x + 6 = 3x + 18 5x - 3x = 18 - 6 2x = 12 x = 12/2 = 6  $\therefore$  present son's age is x = 6 years Present father's age is 5x = 5(6) = 30 years





## 15. I have Rs 1000 in ten and five rupee notes. If the number of ten rupee notes that I have is ten more than the number of five rupee notes, how many notes do I have in each denomination?

#### Solution:

Let the number of five rupee notes be x Number of ten rupee notes be (x + 10)Amount due to five rupee notes  $= 5 \times x = 5x$ Amount due to ten rupee notes = 10 (x + 10) = 10x + 100The total amount = Rs 1000 5x + 10x + 100 = 1000 15x = 900 x = 900/15 = 60  $\therefore$  the number of five rupee notes are x = 60Number of ten rupee notes are x + 10 = 60 + 10 = 70

# 16. At a party, colas, squash and fruit juice were offered to guests. A fourth of the guests drank colas, a third drank squash, two fifths drank fruit juice and just three did not drink anything. How many guests were in all? Solution:

Let the number of guests be x The given details are Number of guests who drank colas are x/4 Number of guests who drank squash are x/3 Number of guests who drank fruit juice are 2x/5 Number of guests who did not drink anything are 3 x/4 + x/3 + 2x/5 + 3 = xBy taking LCM for 4, 3 and 5 is 60 (15x+20x+24x-60x)/60 = -3By cross-multiplying we get, (15x+20x+24x-60x) = -3(60) -x = -180 x = 180 $\therefore$  The total number of guests in all were 180

17. There are 180 multiple choice questions in a test. If a candidate gets 4 marks for every correct answer and for every unattempted or wrongly answered question one mark is deducted from the total score of correct answers. If a candidate scored 450 marks in the test, how many questions did he answer correctly?



#### Solution:

Let the number of correct answers be x Number of questions answered wrong are (180 - x)Total score when answered right = 4x Marks deducted when answered wrong = 1(180 - x) = 180 - xSo, 4x - (180 - x) = 450 4x - 180 + x = 450 5x = 450 + 180 5x = 630 x = 630/5 = 126 $\therefore$  126 questions he answered correctly.

# 18. A labourer is engaged for 20 days on the condition that he will receive Rs 60 for each day, he works and he will be fined Rs 5 for each day, he is absent. If he receives Rs 745 in all for how many days he remained absent?

#### Solution:

Let us consider the number of absent days as x So, number of present days is (20 - x)Wage for one day work = Rs 60 Fine for absent day = Rs 5 So, 60(20 - x) - 5x = 7451200 - 60x - 5x = 744-65x = 744-1200-65x = -456x = -456/-65= 7

 $\therefore$  For 7days the labourer was absent.

# 19. Ravish has three boxes whose total weight is 60 ½ Kg. Box B weighs 3 ½ kg more than box A and box C weighs 5 1/3 kg more than box B. Find the weight of box A. Solution:

The given details are total weight of three boxes is  $60 \frac{1}{2} \text{ kg} = 121/2 \text{ kg}$ Let the weight of box A be x kg Weight of box B be x + 7/2 kg Weight of box C be x + 7/2 + 16/3 kg So,



x + x + 7/2 + x + 7/2 + 16/3 = 121/23x = 121/2 - 7/2 - 7/2 - 16/3 By taking LCM for 2 and 3 is 6 3x = (363 - 21 - 21 - 32)/6 3x = 289/6 x = 289/18 ∴ The weight of box A is 289/18 kg

20. The numerator of a rational number is 3 less than the denominator. If the denominator is increased by 5 and the numerator by 2, we get the rational number 1/2. Find the rational number.

#### Solution:

Le the denominator be x and the numerator be (x - 3)

By using the formula

Fraction = numerator/denominator

= (x - 3)/x

So, when numerator is increased by 2 and Denominator is increased by 5, then fraction is  $\frac{1}{2}$ 

(x - 3 + 2)/(x + 5) = 1/2(x - 1)/(x + 5) = 1/2 By using cross-multiplication, we get 2(x - 1) = x + 52x - 2 = x + 52x - x = 2 + 5x = 7

: Denominator is x = 7, numerator is (x - 3) = 7 - 3 = 4And the fraction = numerator/denominator = 4/7

21. In a rational number, twice the numerator is 2 more than the denominator If 3 is added to each, the numerator and the denominator. The new fraction is 2/3. Find the original number.

#### Solution:

Le the numerator be x and the denominator be (2x - 2)By using the formula Fraction = numerator/denominator = x / (2x - 2)So, the numerator and denominator are increased by 3, then fraction is 2/3 (x + 3)/(2x - 2 + 3) = 2/3

(x + 3)/(2x + 1) = 2/3



By cross-multiplying we get, 3(x + 3) = 2(2x + 1) 3x + 9 = 4x + 2 3x - 4x = 2 - 9 -x = -7 x = 7 $\therefore$  The numerator is x = 7, denominator is (2x - 2) = (2(7) - 2) = 14 - 2 = 12

And the fraction is numerator/denominator = 7/12

22. The distance between two stations is 340 km. Two trains start simultaneously from these stations on parallel tracks to cross each other. The speed of one of them is greater than that of the other by 5 km/hr. If the distance between the two trains after 2 hours of their start is 30 km, find the speed of each train. Solution:

Let the speed of one train be x km/hr. Speed of other train be (x + 5) km/hr. Total distance between two stations = 340 kmBy using the formula  $Distance = speed \times time$ So, Distance covered by one train in 2 hrs. Will be  $x \times 2 = 2x$  km Distance covered by other train in 2 hrs. Will be 2(x + 5) = (2x + 10) km Distance between the trains is 30 km 2x + 2x + 10 + 30 = 3404x + 40 = 3404x = 340 - 404x = 300x = 300/4= 75 $\therefore$  The speed of one train is x = 75 km/hr. Speed of other train is (x + 5) = 75 + 5 = 80 km/hr.

23. A steamer goes downstream from one point another in 9 hours. It covers the same distance upstream in 10 hours. If the speed of the stream be 1 km/hr., find the speed of the steamer in still water and the distance between the ports. Solution:

Let the speed of steamer be x km/hr.

Speed of stream = 1 km/hr.

Downstream speed = (x + 1) km/hr.

Upstream speed = (x - 1) km/hr.



By using the formula Distance = speed × time  $= (x + 1) \times 9$  and  $= (x - 1) \times 10$  9x + 9 = 10x - 10 9x - 10x = -10 - 9 -x = -19 x = 19 km/hr.  $\therefore$  The speed of the steamer in still water is 19 km/hr. Distance between the ports is 9(x + 1) = 9(19+1) = 9(20) = 180 km.

## 24. Bhagwanti inherited Rs 12000.00. She invested part of it as 10% and the rest at 12%. Her annual income from these investments is Rs 1280.00 How much did she invest at each rate?

#### **Solution:**

At rate of 10% Let the investment be Rs x At the rate of 12% the investment will be Rs (12000 - x) At 10% of rate the annual income will be  $x \times (10/100) = 10x/100$ At 12% of rate the annual income will be  $(12000 - x) \times 12/100 = (144000 - 12x)/100$ Total investment = 1280 So, 10x/100 + (144000 - 12x)/100 = 1280(10x + 144000 - 12x)/100 = 1280By cross-multiplying we get, 144000 - 2x = 1280(100)-2x = 128000 - 144000-2x = -16000/2= 8000 $\therefore$  At 10% of rate she invested Rs 8000 and at 12% of rate she invested Rs (12000 - x) =

Rs (12000 - 8000) = Rs 4000

25. The length of a rectangle exceeds its breadth by 9 cm. If length and breadth are each increased by 3 cm, the area of the new rectangle will be 84 cm<sup>2</sup> more than that of the given rectangle. Find the length and breadth of the given rectangle. Solution:

Let the breadth of the rectangle be x meter

Length of the rectangle be (x + 9) meter

Area of the rectangle length×breadth =  $x(x + 9) m^2$ 



When length and breadth increased by 3cm then, New length = x + 9 + 3 = x + 12New breadth = x + 3So, Area is (x + 12) (x + 3) = x (x + 9) + 84  $x^2 + 15x + 36 = x^2 + 9x + 84$  15x - 9x = 84 - 36 6x = 48 x = 48/6 = 8 $\therefore$  Length of the rectangle (x + 9) = (8 + 9) = 17cm and breadth of the rectangle is 8cm.

26. The sum of the ages of Anup and his father is 100. When Anup is as old as his father now, he will be five times as old as his son Anuj is now. Anuj will be eight years older than Anup is now, when Anup is as old as his father. What are their ages now?

### Solution:

Let the age of Anup be x years So age of Anup's father will be (100 - x) years The age of Anuj is (100-x)/5 years So, When Anup is as old as his father after (100 - 2x) years, Then Anuj's age = present age of his father (Anup) + 8Present age of Anuj + 100 - 2x = Present age of Anup + 8 (100 - x)/5 + (100 - 2x) = x + 8(100-x)/5 - 3x = 8 - 100(100 - x - 15x)/5 = -92By cross-multiplying we get, 100 - 16x = -460-16x = -460 - 100-16x = -560x = -560/-16= 35: Present age of Anup is 35 years then, Age of Anup's father will be (100-x) = 100-35 =65 years

The age of Anuj is (100-x)/5 = (100 - 35)/5 = 65/5 = 13 years

27. A lady went shopping and spent half of what she had on buying hankies and gave a rupee to a begger waiting outside the shop. She spent half of what was left of what was left on a lunch and followed that up with a two rupee tip. She spent half of



## the remaining amount on a book and three rupees on bus fare. When she reached home, she found that she had exactly one rupee left. How much money did she start with?

### Solution:

Let the amount lady had be Rs x Amount spent for hankies and given to begger is x/2 + 1Remaining amount is x - (x/2 + 1) = x/2 - 1 = (x-2)/2Amount spent for lunch  $(x-2)/2 \times 1/2 = (x-2)/4$ Amount given as tip is Rs 2 Remaining amount after lunch = (x-2)/2 - (x-2)/4 - 2 = (2x - 4 - x + 2 - 8)/4 = (x - 10)/4Amounts spent for books  $=1/2 \times (x-10)/4 = (x-10)/8$ Bus fare is Rs 3 Amount left = (x-10)/4 - (x-10)/8 - 3 = (2x - 20 - x + 10 - 24)/8 = (x-34)/8So from the question we know that the amount left = Rs 1(x-34)/8 = 1By cross-multiplying we get, x - 34 = 8x = 8 + 34= 42 $\therefore$  the lady started with Rs. 42



