


Date : 13th September 2020 **Time** : 02 : 00 pm - 5 : 00 pm

Subject: Chemistry

46. Identify compound X in the following sequences of reactions :









Sol. 1

$$\begin{array}{c|c}
CH_3 & CH \\
\hline
CH_2O \\
\hline
(X) & 373K
\end{array}$$

$$\begin{array}{c|c}
CHO \\
\hline
CHO
\end{array}$$

- 47. Identify a molecule which does not exist.
 - (1) C₂
- $(2) O_2$
- (3) He₂
- (4) Li₂

Sol. 3

Diatomic molecule for which bond order is zero or negative does not exist.

 $\begin{array}{lll} \text{Molecule} & \text{BO} \\ \text{C}_2 & 2 \\ \text{O}_2 & 2 \\ \text{He}_2 & \text{Zero} \\ \text{Li}_2 & 1 \end{array}$

- **48.** Which of the following is a natural polymer?
 - (1) polybutadiene

- (2) poly (Butadiene-acrylonitrile)
- (3) cis-1,4-polyisoprene
- (4) poly (Butadiene-styrene)

Sol. 3



- **49.** An increase in the concentration of the reactants of a reaction leads to change in :
 - (1) threshold energy

(2) collision frequency

(3) activation energy

(4) heat of reaction

Sol. 2

Collision frequency, as number of molecules per unit volume increases.

50. Anisole on cleavage with HI gives :

$$(1) \bigcirc H_{5}I$$

Sol. 3

- Anisole
- **51.** The number of protons, neutrons and electrons in $^{175}_{71}Lu$, respectively, are :
 - (1) 71, 71 and 104

(2) 175, 104 and 71

(3) 71, 104 and 71

(4) 104, 71 and 71

Sol. 3

$$_{(p)=71}^{(n+p)=175}Lu$$

∴ no. of protons

= 71

no. of neutrons

= 175 - 71 = 104

no. of electrons

= 71

- **52.** The calculated spin only magnetic moment of Cr^{2+} ion is :
 - (1) 5.92 BM

(2) 2.84 BM

(3) 3.87 BM

(4) 4.90 BM

Sol. 4

 $Cr^{2+} \rightarrow 1s^2 \ 2s^2 \ 2p^6 \ 3s^2 \ 3p^6 \ 3d^4$

.. n (number of unpaired electrons) = 4

$$\therefore \qquad \mu \qquad = \sqrt{n(n+2)}$$
$$= \sqrt{4(4+2)}$$

= 4.8989

= 4.9 B.M.

13th September 2020 | Chemistry



53. Match the following:

	Oxide		Nature
(a)	CO	(i)	Basic
(b)	BaO	(ii)	Neutral
(c)	Al_2O_3	(iii)	Acidic

(d) Cl₂O₇ (iv) Amphoteric

Which of the following is correct option?

```
(a)
                 (b)
                          (c)
                                  (d)
(1)
        (iii)
                 (iv)
                          (i)
                                  (ii)
(2)
                                   (i)
        (iv)
                 (iii)
                          (ii)
(3)
        (i)
                 (ii)
                          (iii)
                                   (iv)
(4)
        (ii)
                 (i)
                          (iv)
                                   (iii)
```

Sol. 4

CO Neutral BaO Basic Al₂O₃ Amphoteric Cl₂O₇ Acidic

Urea reacts with water to form A which will decompose to form B. B when passed 54. through Cu²⁺ (aq), deep blue colour solution C is formed. What is the formula of C from the following?

```
(1) Cu(OH)<sub>2</sub>
(3) CuSO<sub>4</sub>
```

(2) CuCO₃.Cu(OH)₂ (4) $[Cu(NH_3)_4]^{2+}$

Sol.

$$4NH_3+Cu^{2+}(aq.) \longrightarrow [Cu(NH_3)_4]^{2+}$$
(C)
(deep blue colour solution)

55. Match the following and identify the correct option.

(i)

- $CO(g) + H_2(g)$ (a) Temporary hardness of water (b) B_2H_6
- (c) (d) $H_{2}O_{2}$

(iii)

(4)

(d) (a) (b) (c) (1)(iii) (i) (iv) (ii) (2) (i) (iii) (ii) (iv) (3) (iii) (i) (ii) (iv)

(ii)

- $Mg(HCO_3)_2 + Ca(HCO_3)_2$ (i)
- An electron deficient hydride (ii)
- (iii) Synthesis gas
- Non-planar structure (iv)

(iv)



Sol. 3

 \Rightarrow CO(g) + H₂(g) = Water gas or synthesis gas

⇒ Temporary hardness of water is due to bicarbonates of Ca²⁺ & Mg²⁺

 \Rightarrow B₂H₆ is a electron deficient compound due to presence of banana bond.

 \Rightarrow $H_2^{-}O_2^{-}$ open book like structure which is non-planar

56. The mixture which shows positive deviation from Raoult's law is :

(1) Acetone + Chloroform

(2) Chloroethane + Bromoethane

(3) Ethanol + Acetone

(4) Benzene + Toluene

Sol. 3

Acetone + ethanol shows positive deviation from Raoult's law. Pure ethanol possesses H-bonding and adding acetone to ethanol causes breaking of some H-bonds.

This causes increase in observed vapour pressure.

57. The freezing point depression constant (K_f) of benzene is 5.12 K kg mol⁻¹. The freezing point depression for the solution of molality 0.078 m containing a non-electrolyte solute in benzene is (rounded off upto two decimal places):

(1) 0.40 K

(2) 0.60 K

(3) 0.20 K

(4) 0.80 K

Sol. 1

$$\begin{array}{lll} \Delta T_{_f} = i k_{_f} m \\ \Rightarrow & \Delta T_{_f} & = 1 \times 5.12 \times 0.078 \\ & \Delta T_{_f} & = 0.3993 \\ & \Delta T_{_f} & = 0.40 \text{ K} \end{array}$$

58. Which of the following set of molecules will have zero dipole moment?

(1) Nitrogen trifluoride, beryllium difluoride, water, 1,3-dichlorobenzene

(2) Boron trifluoride, beryllium difluoride, carbon dioxide, 1,4-dichlorobenzene

(3) Ammonia, beryllium difluoride, water 1,4-dichlorobenzene

(4) Boron trifluoride, hydrogen fluoride, carbon dioxide, 1,3-dichlorobenzene

Sol. 2

 $\mu = 0$

Non-polar molecule

Boron trifluoride

F≪Be⇔F

Beryllium difluoride $\mu = 0$

Non-polar molecule

o≌c≌o

Carbon dioxide

 $\mu = 0$

Non-polar molecule

 $\mu = 0$

Non-polar molecule

1,4-dichlorobenzene



- **59.** A tertiary butyl carbocation is more stable than a secondary butyl carbocation because of which of the following ?
 - (1) -R effect of -CH₃ groups
- (2) Hyperconjugation
- (3) –I effect of –CH₃ groups
- (4) + R effect of -CH₃ groups

Sol. 2

CH₃—CH—CH₂—CH₃
sec. butyl carbocation

Tert butyl carbocation stable due to

9 α — H Hyperconjugation

- Stable due to
- $5 \alpha H$ Hyperconjugation
- **60.** Find out the solubility of Ni(OH)₂ in 0.1 M NaOH. Given that the ionic product of Ni(OH)₂ is 2×10^{-15} .
 - (1) $1 \times 10^{-13} \text{ M}$

(2) $1 \times 10^8 \text{ M}$

 $(3) 2 \times 10^{-13} M$

 $(4) 2 \times 10^{-8} M$

Sol. 3

$$Ni(OH)_2$$

- _
- $K_{sp} = [Ni^{2+}][OH^{-}]^{2}$
- $\Rightarrow k_{sp} = s \times (0.1)^2$

$$\Rightarrow \qquad s = \frac{2 \times 10^{-15}}{0.1^2}$$

$$\Rightarrow$$
 s = 2 × 10⁻¹³ M

- **61.** Reaction between acetone and methylmagnesium chloride followed by hydrolysis will give:
 - (1) Tert. butyl alcohol

(2) Isobutyl alcohol

(3) Isopropyl alcohol

(4) Sec. butyl alcohol

Sol. 1

Tertbutylalcohol

62. Which of the following amine will give the carbylamine test?












Sol. 3

Carbylamine reaction is give by 1° amine & aniline.

63. An alkene on ozonolysis gives methanal as one of the product. Its structure is:



Sol.

$$\begin{array}{c} \text{CH}_2\text{-CH=CH}_2 \\ & \xrightarrow{1. \ \text{O}_3} \\ \hline & 2. \ \text{Zn/H}_2\text{O} \end{array} \\ \text{(Methanal)} \end{array}$$

64. A mixture of N_2 and Ar gases in a cylinder contains 7 g of N_2 and 8 g of Ar. If the total pressure of the mixture of the gases in the cylinder is 27 bar, the partial pressure of N_{2}

[Use atomic masses (in g mol⁻¹): N = 14, Ar = 40]

Sol.

According to Dalton's law

$$P_{N_2} = \frac{\left(\frac{7}{28}\right)}{\left(\frac{8}{40} + \frac{7}{28}\right)} \times 27$$

$$P_{N_2} = \left(\frac{0.25}{0.2 + 0.25}\right) \times 27$$

$$P_{N_2} = 15 \text{ bar}$$

Which of the following is the correct order of increasing field strength of ligands to form 65. coordination compounds?

(1)
$$F^- < SCN^- < C_2O_4^{2-} < CN$$

(1)
$$F^- < SCN^- < C_2O_4^{2-} < CN^-$$
 (2) $CN^- < C_2O_4^{2-} < SCN^- < F^-$

(3)
$$SCN^{-} < F^{-} < C_2O_4^{2-} < CN^{-}$$
 (4) $SCN^{-} < F^{-} < CN^{-} < C_2O_4^{2-}$

(4)
$$SCN^- < F^- < CN^- < C_2O_4^2$$

Sol.

Fact from spectrochemical series.



- **66.** Paper chromatography is an example of :
 - (1) Thin layer chromatography
- (2) Column chromatography
- (3) Adsorption chromatography
- (4) Partition chromatography

Sol. 4

It is an example of partition chromatography as mixture of substance are separated by partition.

- **67.** Sucrose on hydrolysis gives:
 - (1) α -D-Glucose + β -D-Fructose
- (2) α -D-Fructose + β -D-Fructose
- (3) β-D-Glucose + α -D-Fructose
- (4) α -D-Glucose + β -D-Glucose

Sol. 1

Sucrose
$$\xrightarrow{H_3O^+}$$
 α - D-Glucose + β - D - Fructose $(C_6H_{12}O_6)$ $(C_6H_{12}O_6)$

- **68.** The rate constant for a first order reaction is $4.606 \times 10^{-3} \text{ s}^{-1}$. The time required to reduce 2.0 g of the reactant to 0.2 g is:
 - (1) 500 s
- (2) 1000 s
- (3) 100 s
- (4) 200 s

Sol. 1

$$t = \frac{1}{K} \ln \frac{C_o}{C_t}$$

$$= \frac{1}{4.606 \times 10^{-3}} \times 2.303 \log \frac{2}{0.2}$$

$$= \frac{1}{2 \times 10^{-3}} \log 10$$


$$= 500 \text{ sec}$$

- **69.** Reaction between benzaldehyde and acetophenone in presence of dilute NaOH is known as:
 - (1) Cross Cannizzaro's reaction
- (2) Cross Aldol condensation

(3) Aldol condensation

(4) Cannizzaro's reaction

Sol. 2



Two different carbonyl compound is treated with base known as cross-aldol condensation.

- **70.** Which of the following is not correct about carbon monoxide?
 - (1) The carboxyhaemoglobin (haemoglobin bound to CO) is less stable than oxyhaemoglobin.
 - (2) It is produced due to incomplete combustion.
 - (3) It forms carboxyhaemoglobin.
 - (4) It reduces oxygen carrying ability of blood.



Sol. 1

> CO when combine with haemoglobin it forms carboxyhaemoglobin which more stable than oxyhaemoglobin thats why it reduces oxygen carring ability of blood.

71. Hydrolysis of sucrose is given by the following reaction.

Sucrose + H₂O ⇌ Glucose + Fructose

If the equilibrium constant (K_c) is 2 \times 10¹³ at 300 K, the value of Δ _cGo at the same temperature will be:

- (1) 8.314 J mol⁻¹ K⁻¹ × 300 K × ln(3 × 10^{13})
- (2) $-8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 300 \text{ K} \times \ln(4 \times 10^{13})$
- (3) $-8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 300 \text{ K} \times \ln(2 \times 10^{13})$
- (4) 8.314 J mol⁻¹ K⁻¹ × 300 K × ln(2 × 10^{13})
- Sol.

 $\Delta G^{\circ} = -RT \ln K$

$$\Rightarrow \Delta G^{\circ} = -8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 300 \text{ K} \times \text{In } (2 \times 10^{13})$$

- HCl was passed through a solution of CaCl₂, MgCl₂ and NaCl. Which of the following 72. compound(s) crystallise(s)?
 - (1) only MgCl₂

- (2) NaCl, MgCl₂ and CaCl₂
- (3) Both MgCl, and CaCl,
- (4) Only NaCl

Sol.

MgCl₂ has least solubility among CaCl₂, MgCl₂ & NaCl thats why it crystallise first on passing HCl.

73. An element has a body centered cubic (bcc) structure with a cell edge of 288 pm. The atomic radius is:

(1)
$$\frac{4}{\sqrt{2}}$$
 × 288 pm

(2)
$$\frac{4}{\sqrt{2}}$$
 × 288 pm

(3)
$$\frac{\sqrt{3}}{4}$$
 × 288 pm

(1)
$$\frac{4}{\sqrt{3}}$$
 × 288 pm (2) $\frac{4}{\sqrt{2}}$ × 288 pm (3) $\frac{\sqrt{3}}{4}$ × 288 pm (4) $\frac{\sqrt{2}}{4}$ × 288 pm

Sol. 3

For bcc

$$\sqrt{3}$$
 a = 4r

$$\Rightarrow r = \frac{\sqrt{3}}{4} \times 288 \text{ pm}$$

- 74. Which of the following oxoacid of sulphur has -O-O- linkage?
 - (1) $H_2S_2O_8$, peroxodisulphuric acid (2) $H_2S_2O_7$, pyrosulphuric acid
 - (3) H₂SO₃, sulphurous acid
- (4) H₂SO₄, sulphuric acid

Sol.



$$\begin{array}{c} O \\ \parallel \\ S \\ OH \end{array} \qquad (H_2SO_3)$$

$$\begin{array}{c}
O \\
\parallel \\
S \\
O \\
OH
\end{array}$$

$$\begin{array}{c}
O \\
(H_2SO_4)
\end{array}$$

- **75.** Identify the incorrect statement.
 - (1) Interstitial compounds are those that are formed when small atoms like H, C or N are trapped inside the crystal lattices of metals.
 - (2) The oxidation states of chromium in CrO_4^{2-} and $Cr_2O_7^{2-}$ are not the same.
 - (3) $Cr^{2+}(d^4)$ is a stronger reducing agent than $Fe^{2+}(d^6)$ in water.
 - (4) The transition metals and their compounds are known for their catalytic activity due to their ability to adopt multiple oxidation states and to form complexes.
- Sol. 2
 - (1) Interstitial compounds are those compounds that are formed when small atoms like H, B, C or N are trapped inside crystal lattices of metals.
 - (2) Oxidation state of chromium in CrO_4^{2-} & $Cr_2O_7^{2-}$ are +6 (same)
 - (3) Fact

Cr²⁺ is stronger reducing agent than Fe²⁺ in water on the basis of standard reduction potential.

- (4) Transition metals are one of the best available catalyst due to their ability to adopt multiple oxidation states & to form complexes.
- **76.** Which of the following is a cationic detergent?
 - (1) Cetyltrimethyl ammonium bromide (2) Sodium dodecylbenzene sulphonate
 - (3) Sodium lauryl sulphate
- (4) Sodium stearate

Sol. 1

Cetyltrimethyl ammonium bromide is a cationic detergent as it's hydrophillic part is a quaternary ammonium ion.

- 77. The correct option for free expansion of an ideal gas under adiabatic condition is:
 - (1) q < 0, $\Delta T = 0$ and w = 0
- (2) q > 0, $\Delta T > 0$ and w > 0
- (3) q = 0, $\Delta T = 0$ and w = 0
- (4) q = 0, $\Delta T < 0$ and w > 0

Sol. 3

For an adiabatic process, q = 0

for free expansion, $P_{ext} = 0$

$$w = 0$$

& from First law of Thermodynamics, $\Delta E = 0$

or $\Delta T = 0$



- **78**. On electrolysis of dil. sulphuric acid using platinum (Pt) electrode, the product obtained at anode will be:
 - (1) H₂S gas
- (2) SO, gas
- (3) Hydrogen gas
- (4) Oxygen gas

Sol.

Anode: $2H_2O \longrightarrow O_2 + 4H^+ + 4e^-$ Cathode: $2H^+ + 2e^- \longrightarrow H_2(g)$

: oxygen gas will be obtained at anode.

- **79.** Identify the correct statement from the following:
 - (1) Vapour phase refining is carried out for Nickel by Van Arkel method.
 - (2) Pig iron can be moulded into a variety of shapes.
 - (3) Wrought iron is impure iron with 4% carbon.
 - (4) Blister copper has blistered appearance due to evolution of CO₂.
- Sol.
 - (1) Vapour phase refining is carried out for nickel by Mond process.
 - (2) Pig iron is molten iron obtained from blast furnace after smelting it can be moulded into variety of shapes.
 - (3) Wrought iron is the purest form of iron which has less than 4% carbon.
 - (4) Blister copper has blistered appearance due to evolution of SO₂.
- Which of the following is a basic amino acid? 80.
 - (1) Tyrosine
- (2) Lysine
- (3) Serine
- (4) Alanine

Sol. 2

$$NH_2$$
 I
 $NH_2-(CH_2)$
 $CH-COOH$ (Lysine)

Basic part is more than acidic part so lysine is basic amino acid.

81. Identify the incorrect match.

Name

IUPAC Official Name

- (a) Unnilunium
- (i) Mendelvium
- (b) Unniltrium
- (ii) Lawrencium
- (c) Unnilhexium
- (iii) Seaborgium
- (d) Unununnium
- (iv) Darmstadtium
- (1) (c), (iii)
- (2) (d), (iv)
- (3) (a), (i) (4) (b), (ii)

- Sol.
 - 101 Unnilunium Mendelevium
 - 103 Unniltrium Lawrencium
 - 106 Unnilhexium Seaborgium
 - 111 Unununium Roentgenium
 - 110 Ununnilium Darmstadtium
- 82. Which of the following alkane cannot be made in good yield by Wurtz reaction?
 - (1) n-Heptane

(2) n-Butane

(3) n-Hexane

((4) 2,3-Dimethylbutane

Sol.

Wurtz reaction is used to prepare symmetrical alkane.

13th September 2020 | Chemistry



- 83. Elimination reaction of 2-Bromo-pentane to form pent-2-ene is:
 - (a) β -Elimination reaction
- (b) Follows Zaitsev rule
- (c) Dehydrohalogenation reaction
- (d) Dehydration reaction

- (1) (b), (c), (d)
- (2) (a), (b), (d) (3) (a), (b), (c)
- (4) (a), (c), (d)

Sol.

Br
$$CH_3-CH-CH_2-CH_2-CH_3 \xrightarrow{Alc-KOH} CH_3-CH=CH-CH_2-CH_3$$

In the reaction (i) Dehydrohalogenation

- (ii) β elimination reaction
- (iii) follows zaitsev rule (highly substituted alkene is formed as major product)
- 84. The number of Faradays (F) required to produce 20g of calcium from molten CaCl, (Atomic mass of $Ca = 40 \text{ g mol}^{-1}$) is:
 - (1) 3
- (2) 4
- (3) 1
- (4)2

Sol.

Equivalent_{Ca} = Equivalent_{current}

$$\Rightarrow \frac{W}{MM} \times nf = No. \text{ of Faradays}$$

⇒ No. of Faradays =
$$\frac{20}{40} \times 2$$

$$\Rightarrow$$
 No. of Faradays = 1

- 85. Which one of the followings has maximum number of atoms?
 - (1) 1g of $O_2(g)$ (Atomic mass of O=16)
 - (2) 1g of Li(s) (Atomic mass of Li=7)
 - (3) 1g of Ag(s) (Atomic mass of Ag = 108)
 - (4) 1g of Mg(s) (Atomic mass of Mg = 24)
- Sol.

(1) no. of atoms =
$$\frac{1}{32} \times 2 \times N_A$$

(2) no. of atoms =
$$\frac{1}{7} \times N_A$$

(3) no. of atoms =
$$\frac{1}{108} \times N_A$$

(4) no. of atoms =
$$\frac{1}{24} \times N_A$$

1 gm of Li is having maximum no. of atoms.

- 86. For the reaction, $2CI(g) \rightarrow CI_2(g)$, the correct option is:
 - (1) $\Delta_r H < 0$ and $\Delta_r S > 0$
- (2) $\Delta_r H < 0$ and $\Delta_r S < 0$
- (3) $\Delta_{r}H > 0$ and $\Delta_{r}S > 0$

(4) $\Delta_{r}H > 0$ and $\Delta_{r}S < 0$



Sol. 2

 $2Cl(g) \longrightarrow Cl_2(g) + Heat$

Due to bond formation stability increases which results in release of heat.

 $\Delta H_r = -ve$ or exothermic process

 $\Delta H_{\perp} < O$

- $\Delta S < O$, because number of Cl atoms decreases in the formation of Cl₂(g) &
- 87. Identify the correct statements from the following:
 - (a) CO₂(g) is used as refrigerant for ice-cream and frozen food.
 - (b) The structure of C_{60} contains twelve six carbon rings and twenty five carbon rings.
 - (c) ZSM-5, a type of zeolite, is used to convert alcohols into gasoline.
 - (d) CO is colourless and odourless gas.
 - (1) (b) and (c) only

- (2) (c) and (d) only
- (3) (a), (b) and (c) only
- (4) (a) and (c) only

- Sol.
 - \Rightarrow CO₂(s) is not used as refrigerant for ice-cream & frozen food.

— six carbon rings – 20 five carbon rings – 12

- ⇒ ZSM-5 zeolite used to convert alcohols into gasoline
- ⇒ CO is colourless & odourless gas
- 88. Measuring Zeta potential is useful in determining which property of colloidal solution?
 - (1) Stability of the colloidal particles
- (2) Size of the colloidal particles

(3) Viscosity

(4) Solubility

Sol.

A large +ve or -ve value of Zeta potential indicate good stability of the colloidal particles.

89. What is the change in oxidation number of carbon in the following reaction?

 $CH_4(g) + 4CI_2(g) \rightarrow CCI_4(I) + 4HCI(g)$

- (1) -4 to +4 (2) 0 to -4
- (3) +4 to +4 (4) 0 to +4

Sol.

$$^{-4}$$
 CH₄(g) + 4Cl₂ \longrightarrow CCl₄(I) + 4HCl(g)

- 90. The following metal ion activates many enzymes, participates in the oxidation of glucose to produce ATP and with Na, is responsible for the transmission of nerve signals.
 - (1) Calcium
- (2) Potassium (3) Iron
- (4) Copper

Sol. 2

Potassium