

**EXERCISE 20.6****PAGE NO: 20.54****1. Insert 6 geometric means between 27 and 1/81.****Solution:**Let the six terms be  $a_1, a_2, a_3, a_4, a_5, a_6$ .

$$A = 27, B = 1/81$$

Now, these 6 terms are between A and B.

So the GP is: A,  $a_1, a_2, a_3, a_4, a_5, a_6, B$ .

So we now have 8 terms in GP with the first term being 27 and eighth being 1/81.

We know that,  $T_n = ar^{n-1}$ Here,  $T_n = 1/81, a = 27$  and

$$1/81 = 27r^{8-1}$$

$$1/(81 \times 27) = r^7$$

$$r = 1/3$$

$$a_1 = Ar = 27 \times 1/3 = 9$$

$$a_2 = Ar^2 = 27 \times 1/9 = 3$$

$$a_3 = Ar^3 = 27 \times 1/27 = 1$$

$$a_4 = Ar^4 = 27 \times 1/81 = 1/3$$

$$a_5 = Ar^5 = 27 \times 1/243 = 1/9$$

$$a_6 = Ar^6 = 27 \times 1/729 = 1/27$$

 $\therefore$  The six GM between 27 and 1/81 are 9, 3, 1, 1/3, 1/9, 1/27**2. Insert 5 geometric means between 16 and 1/4.****Solution:**Let the five terms be  $a_1, a_2, a_3, a_4, a_5$ .

$$A = 16, B = 1/4$$

Now, these 5 terms are between A and B.

So the GP is: A,  $a_1, a_2, a_3, a_4, a_5, B$ .

So we now have 7 terms in GP with the first term being 16 and seventh being 1/4.

We know that,  $T_n = ar^{n-1}$ Here,  $T_n = 1/4, a = 16$  and

$$1/4 = 16r^{7-1}$$

$$1/(4 \times 16) = r^6$$

$$r = 1/2$$

$$a_1 = Ar = 16 \times 1/2 = 8$$

$$a_2 = Ar^2 = 16 \times 1/4 = 4$$

$$a_3 = Ar^3 = 16 \times 1/8 = 2$$

$$a_4 = Ar^4 = 16 \times 1/16 = 1$$

$$a_5 = Ar^5 = 16 \times 1/32 = 1/2$$

∴ The five GM between 16 and  $1/4$  are 8, 4, 2, 1,  $1/2$

### 3. Insert 5 geometric means between $32/9$ and $81/2$ .

**Solution:**

Let the five terms be  $a_1, a_2, a_3, a_4, a_5$ .

$$A = 32/9, B = 81/2$$

Now, these 5 terms are between A and B.

So the GP is: A,  $a_1, a_2, a_3, a_4, a_5, B$ .

So we now have 7 terms in GP with the first term being  $32/9$  and seventh being  $81/2$ .

We know that,  $T_n = ar^{n-1}$

Here,  $T_n = 81/2, a = 32/9$  and

$$81/2 = 32/9r^{7-1}$$

$$(81 \times 9)/(2 \times 32) = r^6$$

$$r = 3/2$$

$$a_1 = Ar = (32/9) \times 3/2 = 16/3$$

$$a_2 = Ar^2 = (32/9) \times 9/4 = 8$$

$$a_3 = Ar^3 = (32/9) \times 27/8 = 12$$

$$a_4 = Ar^4 = (32/9) \times 81/16 = 18$$

$$a_5 = Ar^5 = (32/9) \times 243/32 = 27$$

∴ The five GM between  $32/9$  and  $81/2$  are  $16/3, 8, 12, 18, 27$

### 4. Find the geometric means of the following pairs of numbers:

(i) 2 and 8

(ii)  $a^3b$  and  $ab^3$

(iii) -8 and -2

**Solution:**

(i) 2 and 8

GM between a and b is  $\sqrt{ab}$

Let  $a = 2$  and  $b = 8$

$$GM = \sqrt{2 \times 8}$$

$$= \sqrt{16}$$

$$= 4$$

(ii)  $a^3b$  and  $ab^3$

GM between a and b is  $\sqrt{ab}$

Let  $a = a^3b$  and  $b = ab^3$

$$GM = \sqrt{(a^3b \times ab^3)}$$

$$= \sqrt{a^4b^4}$$

$$= a^2b^2$$

(iii)  $-8$  and  $-2$

GM between  $a$  and  $b$  is  $\sqrt{ab}$

Let  $a = -2$  and  $b = -8$

$$\begin{aligned} \text{GM} &= \sqrt{(-2 \times -8)} \\ &= \sqrt{16} \\ &= 4, -4 \end{aligned}$$

**5. If  $a$  is the G.M. of  $2$  and  $\frac{1}{4}$  find  $a$ .**

**Solution:**

We know that GM between  $a$  and  $b$  is  $\sqrt{ab}$

Let  $a = 2$  and  $b = \frac{1}{4}$

$$\begin{aligned} \text{GM} &= \sqrt{(2 \times \frac{1}{4})} \\ &= \sqrt{\frac{1}{2}} \\ &= \frac{1}{\sqrt{2}} \end{aligned}$$

$\therefore$  value of  $a$  is  $\frac{1}{\sqrt{2}}$

**6. Find the two numbers whose A.M. is  $25$  and GM is  $20$ .**

**Solution:**

Given: A.M =  $25$ , G.M =  $20$ .

$$\text{G.M} = \sqrt{ab}$$

$$\text{A.M} = \frac{(a+b)}{2}$$

So,

$$\sqrt{ab} = 20 \dots\dots (1)$$

$$\frac{(a+b)}{2} = 25 \dots\dots (2)$$

$$a + b = 50$$

$$a = 50 - b$$

Putting the value of ' $a$ ' in equation (1), we get,

$$\sqrt{[(50-b)b]} = 20$$

$$50b - b^2 = 400$$

$$b^2 - 50b + 400 = 0$$

$$b^2 - 40b - 10b + 400 = 0$$

$$b(b - 40) - 10(b - 40) = 0$$

$$b = 40 \text{ or } b = 10$$

If  $b = 40$  then  $a = 10$

If  $b = 10$  then  $a = 40$

$\therefore$  The numbers are  $10$  and  $40$ .

**7. Construct a quadratic equation in  $x$  such that A.M. of its roots is  $A$  and G.M. is  $G$ .**

**Solution:**

Let the root of the quadratic equation be a and b.

So, according to the given condition,

$$\text{A.M} = (a+b)/2 = A$$

$$a + b = 2A \dots (1)$$

$$\text{GM} = \sqrt{ab} = G$$

$$ab = G^2 \dots (2)$$

The quadratic equation is given by,

$$x^2 - x (\text{Sum of roots}) + (\text{Product of roots}) = 0$$

$$x^2 - x (2A) + (G^2) = 0$$

$$x^2 - 2Ax + G^2 = 0 \text{ [Using (1) and (2)]}$$

∴ The required quadratic equation is  $x^2 - 2Ax + G^2 = 0$ .

