

EXERCISE 23.13

PAGE NO: 23.99

1. Find the angles between each of the following pairs of straight lines:

(i) $3x + y + 12 = 0$ and $x + 2y - 1 = 0$

(ii) $3x - y + 5 = 0$ and $x - 3y + 1 = 0$

Solution:

(i) $3x + y + 12 = 0$ and $x + 2y - 1 = 0$

Given:

The equations of the lines are

$3x + y + 12 = 0 \dots (1)$

$x + 2y - 1 = 0 \dots (2)$

Let m_1 and m_2 be the slopes of these lines.

$m_1 = -3, m_2 = -1/2$

Let θ be the angle between the lines.

Then, by using the formula

$$\begin{aligned}\tan \theta &= [(m_1 - m_2) / (1 + m_1 m_2)] \\ &= [(-3 + 1/2) / (1 + 3/2)] \\ &= 1\end{aligned}$$

So,

$\theta = \pi/4 \text{ or } 45^\circ$

 \therefore The acute angle between the lines is 45°

(ii) $3x - y + 5 = 0$ and $x - 3y + 1 = 0$

Given:

The equations of the lines are

$3x - y + 5 = 0 \dots (1)$

$x - 3y + 1 = 0 \dots (2)$

Let m_1 and m_2 be the slopes of these lines.

$m_1 = 3, m_2 = 1/3$

Let θ be the angle between the lines.

Then, by using the formula

$$\begin{aligned}\tan \theta &= [(m_1 - m_2) / (1 + m_1 m_2)] \\ &= [(3 - 1/3) / (1 + 3(1/3))] \\ &= [(9 - 1)/3] / (1 + 1) \\ &= 8/6 \\ &= 4/3\end{aligned}$$

So,

$\theta = \tan^{-1} (4/3)$

 \therefore The acute angle between the lines is $\tan^{-1} (4/3)$.

2. Find the acute angle between the lines $2x - y + 3 = 0$ and $x + y + 2 = 0$.

Solution:

Given:

The equations of the lines are

$$2x - y + 3 = 0 \dots (1)$$

$$x + y + 2 = 0 \dots (2)$$

Let m_1 and m_2 be the slopes of these lines.

$$m_1 = 2, m_2 = -1$$

Let θ be the angle between the lines.

Then, by using the formula

$$\begin{aligned}\tan \theta &= [(m_1 - m_2) / (1 + m_1 m_2)] \\ &= [(2 - (-1)) / (1 + (2)(-1))] \\ &= [3 / (1 - 2)] \\ &= 3\end{aligned}$$

So,

$$\theta = \tan^{-1} (3)$$

\therefore The acute angle between the lines is $\tan^{-1} (3)$.

3. Prove that the points (2, -1), (0, 2), (2, 3) and (4, 0) are the coordinates of the vertices of a parallelogram and find the angle between its diagonals.

Solution:

To prove:

The points (2, -1), (0, 2), (2, 3) and (4, 0) are the coordinates of the vertices of a parallelogram

Let us assume the points, A (2, -1), B (0, 2), C (2, 3) and D (4, 0) be the vertices.

Now, let us find the slopes

$$\begin{aligned}\text{Slope of AB} &= [(2+1) / (0-2)] \\ &= -3/2\end{aligned}$$

$$\begin{aligned}\text{Slope of BC} &= [(3-2) / (2-0)] \\ &= 1/2\end{aligned}$$

$$\begin{aligned}\text{Slope of CD} &= [(0-3) / (4-2)] \\ &= -3/2\end{aligned}$$

$$\begin{aligned}\text{Slope of DA} &= [(-1-0) / (2-4)] \\ &= 1/2\end{aligned}$$

Thus, AB is parallel to CD and BC is parallel to DA.

Hence proved, the given points are the vertices of a parallelogram.

Now, let us find the angle between the diagonals AC and BD.

Let m_1 and m_2 be the slopes of AC and BD, respectively.

$$m_1 = [(3+1) / (2-2)] \\ = \infty$$

$$m_2 = [(0-2) / (4-0)] \\ = -1/2$$

Thus, the diagonal AC is parallel to the y-axis.

$$\angle ODB = \tan^{-1} (1/2)$$

In triangle MND,

$$\angle DMN = \pi/2 - \tan^{-1} (1/2)$$

\therefore The angle between the diagonals is $\pi/2 - \tan^{-1} (1/2)$.

4. Find the angle between the line joining the points (2, 0), (0, 3) and the line $x + y = 1$.

Solution:

Given:

Points (2, 0), (0, 3) and the line $x + y = 1$.

Let us assume A (2, 0), B (0, 3) be the given points.

Now, let us find the slopes

$$\text{Slope of AB} = m_1 \\ = [(3-0) / (0-2)] \\ = -3/2$$

Slope of the line $x + y = 1$ is -1

$$\therefore m_2 = -1$$

Let θ be the angle between the line joining the points (2, 0), (0, 3) and the line $x + y = 1$

$$\tan \theta = |[(m_1 - m_2) / (1 + m_1 m_2)]| \\ = [(-3/2 + 1) / (1 + 3/2)] \\ = 1/5$$

$$\theta = \tan^{-1} (1/5)$$

\therefore The acute angle between the line joining the points (2, 0), (0, 3) and the line $x + y = 1$ is $\tan^{-1} (1/5)$.


5. If θ is the angle which the straight line joining the points (x_1, y_1) and (x_2, y_2)

subtends at the origin, prove that $\tan \theta = \frac{x_2 y_1 - x_1 y_2}{x_1 x_2 + y_1 y_2}$ **and** $\cos \theta = \frac{x_1 y_2 + y_1 y_2}{\sqrt{x_1^2 + y_1^2} \sqrt{x_2^2 + y_2^2}}$

Solution:

We need to prove:

$$\tan \theta = \frac{x_2 y_1 - x_1 y_2}{x_1 x_2 + y_1 y_2} \text{ and } \cos \theta = \frac{x_1 x_2 + y_1 y_2}{\sqrt{x_1^2 + y_1^2} \sqrt{x_2^2 + y_2^2}}.$$



Let us assume A (x_1, y_1) and B (x_2, y_2) be the given points and O be the origin.

Slope of OA = $m_1 = y_1/x_1$

Slope of OB = $m_2 = y_2/x_2$

It is given that θ is the angle between lines OA and OB.

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

Now, substitute the values, we get

$$= \frac{\frac{y_1}{x_1} - \frac{y_2}{x_2}}{1 + \frac{y_1}{x_1} \times \frac{y_2}{x_2}}$$

$$\tan \theta = \frac{x_2 y_1 - x_1 y_2}{x_1 x_2 + y_1 y_2}$$

Now,

As we know that $\cos \theta = \frac{1}{\sqrt{1 + \tan^2 \theta}}$

Now, substitute the values, we get

$$\cos \theta = \frac{x_1 x_2 + y_1 y_2}{\sqrt{(x_2 y_1 - x_1 y_2)^2 + (x_1 x_2 + y_1 y_2)^2}}$$

$$\cos \theta = \frac{x_1 x_2 + y_1 y_2}{\sqrt{x_1^2 y_1^2 + x_1^2 y_2^2 + x_1^2 x_2^2 + y_1^2 y_2^2}}$$

$$\cos \theta = \frac{x_1 x_2 + y_1 y_2}{\sqrt{x_1^2 + y_1^2} \sqrt{x_2^2 + y_2^2}}$$

Hence proved.