

Exercise 11(A)

Page No: 152

1. Find which of the following sequence form a G.P.:

(i) 8, 24, 72, 216,

(ii) $1/8, 1/24, 1/72, 1/216, \dots$

(iii) 9, 12, 16, 24,

Solution:

(i) Given sequence: 8, 24, 72, 216,

Since,

$$24/8 = 3, 72/24 = 3, 216/72 = 3$$

$$\Rightarrow 24/8 = 72/24 = 216/72 = \dots = 3$$

Therefore 8, 24, 72, 216, is a G.P. with a common ratio 3.

(ii) Given sequence: $1/8, 1/24, 1/72, 1/216, \dots$

Since,

$$(1/24)/(1/8) = 1/3, (1/72)/(1/24) = 1/3, (1/216)/(1/72) = 1/3$$

$$\Rightarrow (1/24)/(1/8) = (1/72)/(1/24) = (1/216)/(1/72) = \dots = 1/3$$

Therefore $1/8, 1/24, 1/72, 1/216, \dots$ is a G.P. with a common ratio $1/3$.

(iii) Given sequence: 9, 12, 16, 24,

Since,

$$12/9 = 4/3; 16/12 = 4/3; 24/16 = 3/2$$

$$12/9 = 16/12 \neq 24/16$$

Therefore, 9, 12, 16, 24 is not a G.P.

2. Find the 9th term of the series: 1, 4, 16, 64,

Solution:

It's seen that, the first term is $(a) = 1$

And, common ratio $(r) = 4/1 = 4$

We know that, the general term is

$$t_n = ar^{n-1}$$

Thus,

$$t_9 = (1)(4)^{9-1} = 4^8 = 65536$$

3. Find the seventh term of the G.P: 1, $\sqrt{3}$, 3, $3\sqrt{3}$,

Solution:

It's seen that, the first term is $(a) = 1$

And, common ratio $(r) = \sqrt{3}/1 = \sqrt{3}$

We know that, the general term is

$$t_n = ar^{n-1}$$

Thus,

$$t_7 = (1)(\sqrt{3})^{7-1} = (\sqrt{3})^6 = 27$$

4. Find the 8th term of the sequence:

$$\frac{3}{4}, 1\frac{1}{2}, 3, \dots$$

Solution:

The given sequence can be rewritten as,
 $\frac{3}{4}, \frac{3}{2}, 3, \dots$

It's seen that, the first term is $(a) = \frac{3}{4}$

And, common ratio $(r) = \frac{3/2}{3/4} = 2$

We know that, the general term is

$$t_n = ar^{n-1}$$

Thus,

$$t_8 = \left(\frac{3}{4}\right)(2)^{8-1} = \left(\frac{3}{4}\right)(2)^7 = 3 \times 2^5 = 3 \times 32 = 96$$

5. Find the 10th term of the G.P. :

$$12, 4, 1\frac{1}{3}, \dots$$

Solution:

The given sequence can be rewritten as,
 $12, 4, \frac{4}{3}, \dots$

It's seen that, the first term is $(a) = 12$

And, common ratio $(r) = \frac{4}{12} = \frac{1}{3}$

We know that, the general term is

$$t_n = ar^{n-1}$$

Thus,

$$t_{10} = (12)\left(\frac{1}{3}\right)^{10-1} = (12)\left(\frac{1}{3}\right)^9 = 12 \times \frac{1}{19683} = \frac{4}{6561}$$

6. Find the n th term of the series:

$$1, 2, 4, 8, \dots$$

Solution:

It's seen that, the first term is $(a) = 1$

And, common ratio $(r) = \frac{2}{1} = 2$

We know that, the general term is

$$t_n = ar^{n-1}$$

Thus,

$$t_n = (1)(2)^{n-1} = 2^{n-1}$$

Exercise 11(B)

Page No: 154

1. Which term of the G.P. :

$$-10, \frac{5}{\sqrt{3}}, -\frac{5}{6}, \dots, \text{is } -\frac{5}{72}?$$

Solution:

In the given G.P.

First term, $a = -10$

Common ratio, $r = (5/\sqrt{3}) / (-10) = 1/(-2\sqrt{3})$

We know that, the general term is

$$t_n = ar^{n-1}$$

So,

$$t_n = (-10) \left(\frac{1}{-2\sqrt{3}} \right)^{n-1} = -5/72$$

$$-\frac{5}{72} = -10 \times \left(\frac{1}{-2\sqrt{3}} \right)^{n-1}$$

$$\frac{1}{144} = \left(\frac{-1}{2\sqrt{3}} \right)^{n-1}$$

$$\frac{-1}{2 \times 2 \times 2 \times 2 \times \sqrt{3} \times \sqrt{3} \times \sqrt{3} \times \sqrt{3}} = \left(\frac{-1}{2\sqrt{3}} \right)^{n-1}$$

$$\left(\frac{-1}{2\sqrt{3}} \right)^4 = \left(\frac{-1}{2\sqrt{3}} \right)^{n-1}$$

Now, equating the exponents we have

$$n - 1 = 4$$

$$n = 5$$

Thus, the 5th of the given G.P. is $-5/72$

2. The fifth term of a G.P. is 81 and its second term is 24. Find the geometric progression.

Solution:

Given,

$$t_5 = 81 \text{ and } t_2 = 24$$

We know that, the general term is

$$t_n = ar^{n-1}$$

So,

$$t_5 = ar^{5-1} = ar^4 = 81 \dots (1)$$

And,

$$t_2 = ar^{2-1} = ar^1 = 24 \dots (2)$$

Dividing (1) by (2), we have

$$ar^4 / ar = 81 / 24$$

$$r^3 = 27 / 8$$

$$r = 3/2$$

Using r in (2), we get

$$a(3/2) = 24$$

$$a = 16$$

Hence, the G.P. is

$$\text{G.P.} = a, ar, ar^2, ar^3, \dots$$

$$= 16, 16 \times (3/2), 16 \times (3/2)^2, 16 \times (3/2)^3$$

$$= 16, 24, 36, 54, \dots$$

3. Fourth and seventh terms of a G.P. are $1/18$ and $-1/486$ respectively. Find the G.P.

Solution:

Given,

$$t_4 = 1/18 \text{ and } t_7 = -1/486$$

We know that, the general term is

$$t_n = ar^{n-1}$$

So,

$$t_4 = ar^{4-1} = ar^3 = 1/18 \dots (1)$$

And,

$$t_7 = ar^{7-1} = ar^6 = -1/486 \dots (2)$$

Dividing (2) by (1), we have

$$ar^6 / ar^3 = (-1/486) / (1/18)$$

$$r^3 = -1/27$$

$$r = -1/3$$

Using r in (1), we get

$$a(-1/3)^3 = 1/18$$

$$a = -27/18 = -3/2$$

Hence, the G.P. is

$$\text{G.P.} = a, ar, ar^2, ar^3, \dots$$

$$= -3/2, -3/2(-1/3), -3/2(-1/3)^2, -3/2(-1/3)^3, \dots$$

$$= -3/2, 1/2, -1/6, 1/18, \dots$$

4. If the first and the third terms of a G.P are 2 and 8 respectively, find its second term.

Solution:

Given,

$$t_1 = 2 \text{ and } t_3 = 8$$

We know that, the general term is

$$t_n = ar^{n-1}$$

So,

$$t_1 = ar^{1-1} = a = 2 \dots (1)$$

And,

$$t_3 = ar^{3-1} = ar^2 = 8 \dots (2)$$

Dividing (2) by (1), we have

$$ar^2 / a = 8 / 2$$

$$r^2 = 4$$

$$r = \pm 2$$

Hence, the 2nd term of the G.P. is

When $a = 2$ and $r = 2$ is $2(2) = 4$
Or when $a = 2$ and $r = -2$ is $2(-2) = -4$

5. The product of 3rd and 8th terms of a G.P. is 243. If its 4th term is 3, find its 7th term
Solution:

Given,

Product of 3rd and 8th terms of a G.P. is 243

The general term of a G.P. with first term a and common ratio r is given by,

$$t_n = ar^{n-1}$$

So,

$$t_3 \times t_8 = ar^{3-1} \times ar^{8-1} = ar^2 \times ar^7 = a^2r^9 = 243$$

Also given,

$$t_4 = ar^{4-1} = ar^3 = 3$$

Now,

$$a^2r^9 = (ar^3) ar^6 = 243$$

Substituting the value of ar^3 in the above equation, we get,

$$(3) ar^6 = 243$$

$$ar^6 = 81$$

$$ar^{7-1} = 81 = t_7$$

Thus, the 7th term of the G.P is 81.

Exercise 11(C)

Page No: 156

1. Find the seventh term from the end of the series: $\sqrt{2}, 2, 2\sqrt{2}, \dots, 32$

Solution:

Given series: $\sqrt{2}, 2, 2\sqrt{2}, \dots, 32$

Here,

$$a = \sqrt{2}$$

$$r = 2 / \sqrt{2} = \sqrt{2}$$

And, the last term (l) = 32

$$l = t_n = ar^{n-1} = 32$$

$$(\sqrt{2})(\sqrt{2})^{n-1} = 32$$

$$(\sqrt{2})^n = 32$$

$$(\sqrt{2})^n = (2)^5 = (\sqrt{2})^{10}$$

Equating the exponents, we have

$$n = 10$$

So, the 7th term from the end is $(10 - 7 + 1)$ th term.

i.e. 4th term of the G.P

Hence,

$$t_4 = (\sqrt{2})(\sqrt{2})^{4-1} = (\sqrt{2})(\sqrt{2})^3 = (\sqrt{2}) \times 2\sqrt{2} = 4$$

2. Find the third term from the end of the G.P.

$2/27, 2/9, 2/3, \dots, 162$

Solution:

Given series: $2/27, 2/9, 2/3, \dots, 162$

Here,

$$a = 2/27$$

$$r = (2/9) / (2/27)$$

$$r = 3$$

And, the last term (l) = 162

$$l = t_n = ar^{n-1} = 162$$

$$(2/27)(3)^{n-1} = 162$$

$$(3)^{n-1} = 162 \times (27/2)$$

$$(3)^{n-1} = 2187$$

$$(3)^{n-1} = (3)^7$$

$$n - 1 = 7$$

$$n = 7 + 1$$

$$n = 8$$

So, the third term from the end is $(8 - 3 + 1)$ th term

i.e. 6th term of the G.P. = t_6

Hence,

$$t_6 = ar^{6-1}$$

$$t_6 = (2/27)(3)^{6-1}$$

$$t_6 = (2/27)(3)^5$$

$$t_6 = 2 \times 3^2$$

$$t_6 = 18$$

3. Find the G.P. $1/27, 1/9, 1/3, \dots, 81$; find the product of fourth term from the beginning and the fourth term from the end.

Solution:

Given G.P. $1/27, 1/9, 1/3, \dots, 81$

Here, $a = 1/27$, common ratio (r) = $(1/9) / (1/27) = 3$ and $l = 81$

We know that,

$$l = t_n = ar^{n-1} = 81$$

$$(1/27)(3)^{n-1} = 81$$

$$3^{n-1} = 81 \times 27 = 2187$$

$$3^{n-1} = 3^7$$

$$n - 1 = 7$$

$$n = 8$$

Hence, there are 8 terms in the given G.P.

Now,

4^{th} term from the beginning is t_4 and the 4^{th} term from the end is $(8 - 4 + 1) = 5^{\text{th}}$ term (t_5)

Thus,

$$\text{the product of } t_4 \text{ and } t_5 = ar^{4-1} \times ar^{5-1} = ar^3 \times ar^4 = a^2r^7 = (1/27)^2(3)^7 = 3$$

4. If for a G.P., p^{th} , q^{th} and r^{th} terms are a , b and c respectively; prove that:

$$(q - r) \log a + (r - p) \log b + (p - q) \log c = 0$$

Solution:

Let's take the first term of the G.P. be A and its common ratio be R .

Then,

$$p^{\text{th}} \text{ term} = a \Rightarrow AR^{p-1} = a$$

$$q^{\text{th}} \text{ term} = b \Rightarrow AR^{q-1} = b$$

$$r^{\text{th}} \text{ term} = c \Rightarrow AR^{r-1} = c$$

Now,

$$\begin{aligned} a^{q-r} \times b^{r-p} \times c^{p-q} &= (AR^{p-1})^{q-r} \times (AR^{q-1})^{r-p} \times (AR^{r-1})^{p-q} \\ &= A^{q-r} \cdot R^{(p-1)(q-r)} \times A^{r-p} \cdot R^{(q-1)(r-p)} \times A^{p-q} \cdot R^{(r-1)(p-q)} \\ &= A^{q-r+r-p+p-q} \times R^{(p-1)(q-r)+(q-1)(r-p)+(r-1)(p-q)} \\ &= A^0 \times R^0 \\ &= 1 \end{aligned}$$

On taking log on both the sides, we get

$$\log(a^{q-r} \times b^{r-p} \times c^{p-q}) = \log 1$$

$$\Rightarrow (q - r)\log a + (r - p)\log b + (p - q)\log c = 0$$

- Hence Proved

Exercise 11(D)

Page No: 156

1. Find the sum of G.P.:

(i) $1 + 3 + 9 + 27 + \dots$ to 12 terms

(ii) $0.3 + 0.03 + 0.003 + 0.0003 + \dots$ to 8 terms.

(iii) $1 - 1/2 + 1/4 - 1/8 + \dots$ to 9 terms

(iv) $1 - 1/3 + 1/3^2 - 1/3^3 + \dots$ to n terms

(v) $\frac{x+y}{x-y} + 1 + \frac{x-y}{x+y} + \dots$ upto n terms

(vi) $\sqrt{3} + \frac{1}{\sqrt{3}} + \frac{1}{3\sqrt{3}} + \dots$ to n terms.

Solution:

(i) Given G.P: $1 + 3 + 9 + 27 + \dots$ to 12 terms

Here,

$$a = 1 \text{ and } r = 3/1 = 3 \text{ (} r > 1 \text{)}$$

Number of terms, $n = 12$

Hence,

$$S_n = a(r^n - 1) / r - 1$$

$$\Rightarrow S_{12} = (1)((3)^{12} - 1) / 3 - 1$$

$$= (3^{12} - 1) / 2$$

$$= (531441 - 1) / 2$$

$$= 531440 / 2$$

$$= 265720$$

(ii) Given G.P: $0.3 + 0.03 + 0.003 + 0.0003 + \dots$ to 8 terms

Here,

$$a = 0.3 \text{ and } r = 0.03/0.3 = 0.1 \text{ (} r < 1 \text{)}$$

Number of terms, $n = 8$

Hence,

$$S_n = a(1 - r^n) / 1 - r$$

$$\Rightarrow S_8 = (0.3)(1 - 0.1^8) / (1 - 0.1)$$

$$= 0.3(1 - 0.1^8) / 0.9$$

$$= (1 - 0.1^8) / 3$$

$$= 1/3(1 - (1/10)^8)$$

(iii) Given G.P: $1 - 1/2 + 1/4 - 1/8 + \dots$ to 9 terms

Here,

$$a = 1 \text{ and } r = (-1/2) / 1 = -1/2 \text{ (} |r| < 1 \text{)}$$

Number of terms, $n = 9$

Hence,

$$S_n = a(1 - r^n) / 1 - r$$

$$\Rightarrow S_9 = (1)(1 - (-1/2)^9) / (1 - (-1/2))$$

$$= (1 + (1/2)^9) / (3/2)$$

$$= 2/3 \times (1 + 1/512)$$

$$= \frac{2}{3} \times \left(\frac{513}{512}\right)$$

$$= \frac{171}{256}$$

(iv) Given G.P: $1 - \frac{1}{3} + \frac{1}{3^2} - \frac{1}{3^3} + \dots$ to n terms

Here,

$$a = 1 \text{ and } r = \left(-\frac{1}{3}\right) / 1 = -\frac{1}{3} \quad (|r| < 1)$$

Number of terms is n

Hence,

$$S_n = \frac{a(1 - r^n)}{1 - r}$$

$$S_n = \frac{1 \left(1 - \left(-\frac{1}{3}\right)^n \right)}{1 - \left(-\frac{1}{3}\right)}$$

$$= \frac{1 \left(1 - \left(-\frac{1}{3}\right)^n \right)}{1 + \frac{1}{3}}$$

$$= \frac{\left[1 - \left(-\frac{1}{3}\right)^n \right]}{\frac{4}{3}}$$

$$= \frac{3}{4} \left[1 - \left(-\frac{1}{3}\right)^n \right]$$

(v) Given G.P:

$$\frac{x+y}{x-y} + 1 + \frac{x-y}{x+y} + \dots \text{ upto } n \text{ terms}$$

Here,

$$a = \frac{(x+y)}{(x-y)} \text{ and } r = \frac{1}{\left[\frac{(x+y)}{(x-y)}\right]} = \frac{(x-y)}{(x+y)} \quad (|r| < 1)$$

Number of terms = n

Hence,

$$S_n = \frac{a(1 - r^n)}{1 - r}$$

$$\begin{aligned}
 S_n &= \frac{\frac{x+y}{x-y} \left(1 - \left(\frac{x-y}{x+y} \right)^n \right)}{1 - \left(\frac{x-y}{x+y} \right)} \\
 &= \frac{\frac{x+y}{x-y} \left(1 - \left(\frac{x-y}{x+y} \right)^n \right)}{\frac{x+y-x+y}{x+y}} \\
 &= \frac{\frac{x+y}{x-y} \left(1 - \left(\frac{x-y}{x+y} \right)^n \right)}{\frac{2y}{x+y}} \\
 &= \frac{(x+y)^2 \left(1 - \left(\frac{x-y}{x+y} \right)^n \right)}{2y(x-y)}
 \end{aligned}$$

(vi) Given G.P:

$$\sqrt{3} + \frac{1}{\sqrt{3}} + \frac{1}{3\sqrt{3}} + \dots \text{to } n \text{ terms.}$$

Here,

$$a = \sqrt{3} \text{ and } r = \frac{1/\sqrt{3}}{\sqrt{3}} = \frac{1}{3} \quad (|r| < 1)$$

Number of terms = n

Hence,

$$S_n = a(1 - r^n) / 1 - r$$

$$S_n = \frac{\sqrt{3} \left(1 - \left(\frac{1}{3} \right)^n \right)}{1 - \frac{1}{3}} = \frac{\sqrt{3} \left(1 - \frac{1}{3^n} \right)}{\frac{2}{3}}$$

$$= \frac{3\sqrt{3}}{2} \left(1 - \frac{1}{3^n} \right)$$

2. How many terms of the geometric progression $1 + 4 + 16 + 64 + \dots$ must be added to get sum equal to 5461?

Solution:

Given G.P: $1 + 4 + 16 + 64 + \dots$

Here,

$$a = 1 \text{ and } r = 4/1 = 4 \quad (r > 1)$$

And,

$$S_n = 5461$$

We know that,

$$S_n = a(r^n - 1) / r - 1$$

$$\Rightarrow S_n = (1)((4)^n - 1) / 4 - 1$$

$$= (4^n - 1) / 3$$

$$5461 = (4^n - 1) / 3$$

$$16383 = 4^n - 1$$

$$4^n = 16384$$

$$4^n = 4^7$$

$$n = 7$$

Therefore, 7 terms of the G.P must be added to get a sum of 5461.

3. The first term of a G.P. is 27 and its 8th term is 1/81. Find the sum of its first 10 terms.

Solution:

Given,

First term (a) of a G.P = 27

And, 8th term = $t_8 = ar^{8-1} = 1/81$

$$(27)r^7 = 1/81$$

$$r^7 = 1/(81 \times 27)$$

$$r^7 = (1/3)^7$$

$$r = 1/3 \quad (r < 1)$$

$$S_n = a(1 - r^n) / 1 - r$$

Now,

Sum of first 10 terms = S_{10}

$$S_{10} = \frac{27 \left(1 - \left(\frac{1}{3} \right)^{10} \right)}{1 - \frac{1}{3}} = \frac{27 \left(1 - \frac{1}{3^{10}} \right)}{\frac{2}{3}}$$

$$= \frac{81}{2} \left(1 - \frac{1}{3^{10}} \right)$$

4. A boy spends Rs.10 on first day, Rs.20 on second day, Rs.40 on third day and so on. Find how much, in all, will he spend in 12 days?

Solution:

Given,

Amount spent on 1st day = Rs 10

Amount spent on 2nd day = Rs 20

And amount spent on 3rd day = Rs 40

It's seen that,

10, 20, 40, forms a G.P with first term, $a = 10$ and common ratio, $r = 20/10 = 2$ ($r > 1$)

The number of days, $n = 12$

Hence, the sum of money spend in 12 days is the sum of 12 terms of the G.P.

$$S_n = a(r^n - 1) / r - 1$$

$$S_{12} = (10)(2^{12} - 1) / 2 - 1 = 10 (2^{12} - 1) = 10 (4096 - 1) = 10 \times 4095 = 40950$$

Therefore, the amount spent by him in 12 days is Rs 40950

5. The 4th term and the 7th term of a G.P. are 1/27 and 1/729 respectively. Find the sum of n terms of the G.P.

Solution:

Given,

$$t_4 = 1/27 \text{ and } t_7 = 1/729$$

We know that,

$$t_n = ar^{n-1}$$

So,

$$t_4 = ar^{4-1} = ar^3 = 1/27 \dots (1)$$

$$t_7 = ar^{7-1} = ar^6 = 1/729 \dots (2)$$

Dividing (2) by (1) we get,

$$ar^6 / ar^3 = (1/729) / (1/27)$$

$$r^3 = (1/3)^3$$

$$r = 1/3 \text{ (} r < 1 \text{)}$$

In (1)

$$a \times 1/27 = 1/27$$

$$a = 1$$

Hence,

$$S_n = a(1 - r^n) / 1 - r$$

$$S_n = (1 - (1/3)^n) / 1 - (1/3)$$

$$= (1 - (1/3)^n) / (2/3)$$

$$= 3/2 (1 - (1/3)^n)$$

6. A geometric progression has common ratio = 3 and last term = 486. If the sum of its terms is 728; find its first term.

Solution:

Given,

For a G.P.,

$$r = 3, l = 486 \text{ and } S_n = 728$$

$$\frac{l r - a}{r - 1} = 728$$

$$\frac{486 \times 3 - a}{3 - 1} = 728$$

$$\frac{1458 - a}{2} = 728$$

$$1458 - a = 728 \times 2 = 1456$$

$$\text{Thus, } a = 2$$

7. Find the sum of G.P.: 3, 6, 12,, 1536.

Solution:

Given G.P: 3, 6, 12,, 1536

Here,

$$a = 3, l = 1536 \text{ and } r = 6/3 = 2$$

So,

$$\begin{aligned} \text{The sum of terms} &= (lr - a) / (r - 1) \\ &= (1536 \times 2 - 3) / (2 - 1) \\ &= 3072 - 3 \\ &= 3069 \end{aligned}$$

8. How many terms of the series $2 + 6 + 18 + \dots$ must be taken to make the sum equal to 728?

Solution:

Given G.P: $2 + 6 + 18 + \dots$

Here,

$$a = 2 \text{ and } r = 6/2 = 3$$

Also given,

$$S_n = 728$$

$$S_n = a(r^n - 1) / r - 1$$

$$728 = (2)(3^n - 1) / 3 - 1 = 3^n - 1$$

$$729 = 3^n$$

$$3^6 = 3^n$$

$$n = 6$$

Therefore, 6 terms must be taken to make the sum equal to 728.

9. In a G.P., the ratio between the sum of first three terms and that of the first six terms is 125:152.

Find its common ratio.

Solution:

Given,

$$\frac{a(r^3 - 1)}{r - 1} : \frac{a(r^6 - 1)}{r - 1} = 125 : 152$$

$$\frac{a(r^3 - 1)}{r - 1} = \frac{125}{152}$$

$$\frac{(r^3 - 1)}{(r^6 - 1)} = \frac{125}{152}$$

$$\frac{r^3 - 1}{(r^3)^2 - (1)^2} = \frac{125}{152}$$

$$\frac{r^3 - 1}{(r^3 - 1)(r^3 + 1)} = \frac{125}{152}$$

$$\frac{1}{r^3 + 1} = \frac{125}{152}$$

$$r^3 + 1 = \frac{152}{125}$$

$$r^3 = \frac{152}{125} - 1 = \frac{152 - 125}{125} = \frac{27}{125}$$

$$r^3 = \left(\frac{3}{5}\right)^3$$

$$r = 3/5$$

Therefore, the common ratio is 3/5.