



## Section A


1. The correct order of the following compounds showing increasing tendency towards nucleophilic substitution reaction is:




(i)



(ii)



(iii)



(iv)

- a. (iv) < (i) < (iii) < (ii)  
c. (i) < (ii) < (iii) < (iv)


- b. (iv) < (i) < (ii) < (iii)  
d. (iv) < (iii) < (ii) < (i)

**Ans (c)**

**Solution:**




(i)




(- M & -I)

(ii)



(- M & -I)

(iii)



(- M & -I)

(iv)

Reactivity  $\propto$  - M group present at o/p position.

2. Match List-I with List-II

List-I

(Metal)

(a) Aluminum

(b) Iron

(c) Copper

(d) Zinc

List-II

(Ores)

(i) Siderite

(ii) Calamine

(iii) Kaolinite

(iv) Malachite

# JEE MAIN 24<sup>th</sup> Feb. 2021 Shift-2 (Chemistry)



- a. (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i)
- b. (a)-(i), (b)-(ii), (c)-(iii), (d)-(iv)
- c. (a)-(iii), (b)-(i), (c)-(iv), (d)-(ii)
- d. (a)-(ii), (b)-(iv), (c)-(i), (d)-(iii)

**Ans (c)**

Solution:

Siderite       $\text{FeCO}_3$

Calamine       $\text{ZnCO}_3$

Kaolinite       $\text{Si}_2\text{Al}_2\text{O}_5(\text{OH})_4$  or  $\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot 2\text{H}_2\text{O}$

Malachite       $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$

### 3. Match List-I with List-II

List- I	List-II
(Salt)	(Flame colour wavelength)
(a) LiCl	(i) 455.5 nm
(b) NaCl	(ii) 970.8 nm
(c) RbCl	(iii) 780.0 nm
(d) CsCl	(iv) 589.2 nm

Choose the correct answer from the options given below:

- a. (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)
- b. (a)-(ii), (b)-(iv), (c)-(iii), (d)-(i)
- c. (a)-(iv), (b)-(ii), (c)-(iii), (d)-(i)
- d. (a)-(i), (b)-(iv), (c)-(ii), (d)-(iii)

**Ans (b)**

Solution:

Range of visible region: -  
390 nm – 760 nm

VIBGYOR

Violet - Red

LiCl      Crimson Red

NaCl      Golden yellow

RbCl      Violet

CsCl      Blue

## JEE MAIN 24<sup>th</sup> Feb. 2021 Shift-2 (Chemistry)



So, LiCl which is crimson have wave length closed to red in the spectrum of visible region which is as per given data.

4. Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R.

Assertion A: Hydrogen is the most abundant element in the Universe, but it is not the most abundant gas in the troposphere.

Reason R: Hydrogen is the lightest element.

In the light of the above statements, choose the correct answer from the given below

- (1) A is false but R is true
- (2) Both A and R are true and R is the correct explanation of A
- (3) A is true but R is false
- (4) Both A and R are true but R is NOT the correct explanation of A

- a. A is false but R is true
- b. Both A and R are true and R is the correct explanation of A
- c. A is true but R is false
- d. Both A and R are true but R is NOT the correct explanation of A

**Ans (b)**

**Solution:**

Hydrogen is most abundant element in universe because all luminous body of universe i.e. stars & nebulae are made up of hydrogen which acts as nuclear fuel & fusion reaction is responsible for their light.

5. Given below are two statements:

Statement I: The value of the parameter "Biochemical Oxygen Demand (BOD)" is important for survival of aquatic life.

Statement II: The optimum value of BOD is 6.5 ppm.

In the light of the above statements, choose the most appropriate answer from the options given below.





- a. Both Statement I and Statement II are false
- b. Statement I is false but Statement II is true
- c. Statement I is true but Statement II is false
- d. Both Statement I and Statement II are true

**Ans (c)**

**Solution:**

For survival of aquatic life dissolved oxygen is responsible its optimum limit 6.5 ppm and optimum limit of BOD ranges from 10-20 ppm & BOD stands for biochemical oxygen demand.



6. Which one of the following carbonyl compounds cannot be prepared by addition of water on an alkyne in the presence of  $\text{HgSO}_4$  and  $\text{H}_2\text{SO}_4$ ?



**Ans (a)**

**Solution:**

Reaction of Alkyne with  $\text{HgSO}_4$  &  $\text{H}_2\text{SO}_4$  follow as



Hence, by this process preparation of  $\text{CH}_3\text{CH}_2\text{CHO}$  can't be possible.

7. Which one of the following compounds is non-aromatic?


a.




b.



c.




d.



**Ans (b)**

Solution:



Hence, it is non-aromatic.

8. The incorrect statement among the following is:

- a.  $\text{VOSO}_4$  is a reducing agent
- b. Red color of ruby is due to the presence of  $\text{CO}^{3+}$
- c.  $\text{Cr}_2\text{O}_3$  is an amphoteric oxide
- d.  $\text{RuO}_4$  is an oxidizing agent

Ans (b)

Solution:

Red color of ruby is due to presence of  $\text{CrO}_3$  or  $\text{Cr}^{+6}$  not  $\text{CO}^{3+}$

9. According to Bohr's atomic theory:

- (a) Kinetic energy of electron is  $\propto \frac{Z^2}{n^2}$
- (b) The product of velocity (v) of electron and principal quantum number (n). ' $v_n$ '  $\propto Z^2$
- (c) Frequency of revolution of electron in an orbit is  $\propto \frac{Z^3}{n^3}$
- (d) Coulombic force of attraction on the electron is  $\propto \frac{Z^3}{n^4}$

Choose the most appropriate answer from the options given below:

- a. (c) only
- b. (a) and (d) only
- c. (a) only
- d. (a), (c) and (d) only

Ans (b)

Solution:

(a)  $\text{KE} = -\text{TE} = 13.6 \times \frac{Z^2}{n^2} \text{ eV}$

$$\text{KE} \propto \frac{Z^2}{n^2}$$



$$(b) V = 2.188 \times 10^6 \times \frac{Z}{n} \text{ m/s}$$

$$\text{So, } V_n \propto Z$$

$$\text{Frequency} = \frac{V}{2\pi r}$$

$$F \propto \frac{Z^2}{n^3} \left[ \therefore r \propto \frac{n^2}{Z} \text{ and } v \propto \frac{Z}{n} \right]$$

$$(d) \text{ Force} \propto \frac{Z^2}{r^2}$$

$$\text{So, } F \propto \frac{Z^3}{n^4}$$

So, only statement (A) is correct.

10. Match List-I with List-II

List-I	List-II
(a) Valium	(i) Antifertility drug
(b) Morphine	(ii) Pernicious anaemia
(c) Norethindrone	(iii) Analgesic
(d) Vitamin B12	(iv) Tranquilizer

a. (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i)

b. (a)-(i), (b)-(iii), (c)-(iv), (d)-(ii)

c. (a)-(ii), (b)-(iv), (c)-(iii), (d)-(i)

d. (a)-(iv), (b)-(iii), (c)-(i), (d)-(ii)

**Ans (d)**

**Solution:**

(a) Valium	(iv) Tranquilizer
(b) Morphine	(iii) Analgesic
(c) Norethindrone	(i) Antifertility drug
(d) Vitamin B12	(ii) Pernicious anemia

11. The Correct set from the following in which both pairs are in correct order of melting point is

a.  $\text{LiF} > \text{LiCl}$  ;  $\text{NaCl} > \text{MgO}$

b.  $\text{LiF} > \text{LiCl}$  ;  $\text{MgO} > \text{NaCl}$

c.  $\text{LiCl} > \text{LiF}$  ;  $\text{NaCl} > \text{MgO}$

d.  $\text{LiCl} > \text{LiF}$  ;  $\text{MgO} > \text{NaCl}$

**Ans (b)**

Solution:

Generally

$$\text{M.P.} \propto \text{Lattice energy} = \frac{KQ_1Q_2}{r^+ + r^-}$$

$\propto$  (packing efficiency)

12. The calculated magnetic moments (spin only value) for species  $[\text{FeCl}_4]^{2-}$ ,  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$  and  $\text{MnO}_4^{2-}$  respectively are:
- |                        |                        |
|------------------------|------------------------|
| a. 5.92, 4.90 and 0 BM | b. 5.82, 0 and 0 BM    |
| c. 4.90, 0 and 1.73 BM | d. 4.90, 0 and 2.83 BM |

**Ans (c)**

Solution:




$\text{Fe}^{2+} 3d^6 \rightarrow 4$  unpaired electrons. as  $\text{Cl}^-$  in a weak field ligand.

$$\mu_{\text{spin}} = \sqrt{24} \text{ BM}$$

$$= 4.9 \text{ BM}$$

13.




Which of the following reagent is suitable for the preparation of the product in the above reaction?

- a. Red P +  $\text{Cl}_2$
- b.  $\text{NH}_2\text{-NH}_2 / \text{C}_2\text{H}_5\text{O}^-\text{Na}^+$
- c.  $\text{Ni}/\text{H}_2$
- d.  $\text{NaBH}_4$

**Ans: (b)**


Solution:




It is wolff-kishner reduction of carbonyl compounds.

14. The diazonium salt of which of the following compounds will form a coloured dye on reaction with  $\beta$ -Naphthol in NaOH?


a.




b.



c.




d.




Ans: (c)

Solution:






15. What is the correct sequence of reagents used for converting nitrobenzene into m-dibromobenzene?



- a.  $\xrightarrow{\text{Sn/HCl}}$  /  $\xrightarrow{\text{Br}_2}$  /  $\xrightarrow{\text{NaNO}_2}$  /  $\xrightarrow{\text{NaBr}}$
- b.  $\xrightarrow{\text{Sn/HCl}}$  /  $\xrightarrow{\text{KBr}}$  /  $\xrightarrow{\text{Br}_2}$  /  $\xrightarrow{\text{H}^+}$
- c.  $\xrightarrow{\text{NaNO}_2}$  /  $\xrightarrow{\text{HCl}}$  /  $\xrightarrow{\text{KE}}$
- d.  $\xrightarrow{\text{Br}_2/\text{Fe}}$  /  $\xrightarrow{\text{Sn/HCl}}$  /  $\xrightarrow{\text{NaNO}_2/\text{HCl}}$  /  $\xrightarrow{\text{CuBr/HBr}}$

Ans: (d)

Solution:




16. The correct shape and I-I-I bond angles respectively in  $I_3^-$  ion are:
- Trigonal planar;  $120^\circ$
  - Distorted trigonal planar;  $135^\circ$  and  $90^\circ$
  - Linear;  $180^\circ$
  - T-shaped;  $180^\circ$  and  $90^\circ$

Ans: (c)

Solution:

$I_3^-$  has  $sp^3d$  hybridization (2 BP + 3 LP) and linear geometry.



17. What is the correct order of the following elements with respect to their density?

- a.  $Cr < Fe < Co < Cu < Zn$   
c.  $Zn < Cu < Co < Fe < Cr$

- b.  $Cr < Zn < Co < Cu < Fe$   
d.  $Zn < Cr < Fe < Co < Cu$

Ans: (d)





Solution:

Fact Based

Density depends on many factors like atomic mass, atomic radius and packing efficiency.

18. Match List-I and List-II.

List - I



List-II

(i)  $Br_2/NaOH$

(ii)  $H_2/Pd-BaSO_4$

(iii)  $Zn(Hg)/Conc. HCl$


(iv)  $Cl_2/Red P, H_2O$




- a. (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)
- b. (a)-(iii), (b)-(iv), (c)-(i), (d)-(ii)
- c. (a)-(iii), (b)-(i), (c)-(iv), (d)-(ii)
- d. (a)-(ii), (b)-(iv), (c)-(i), (d)-(iii)

**Ans:** (d)


**Solution:**




(b)



(c)



(d)



19. In polymer Buna-S: 'S' stands for :

- a. Styrene
- b. Sulphur
- c. Strength
- d. Sulphonation

**Ans:** (a)

**Solution:**

Buna-S is the co-polymer of buta-1,3-diene & styrene

20. Most suitable salt which can be used for efficient clotting of blood will be:

- |                                |                    |
|--------------------------------|--------------------|
| a. $\text{Mg}(\text{HCO}_3)_2$ | b. $\text{FeSO}_4$ |
| c. $\text{NaHCO}_3$            | d. $\text{FeCl}_3$ |

**Ans:** (d)

**Solution:**

Blood is a negative sol, according to Hardy-Schulz's rule, the cation with high charge has high coagulation power. Hence,  $\text{FeCl}_3$  can be used for clotting blood.

## Section B


1. The magnitude of the change in oxidising power of the  $\text{MnO}_4^-/\text{Mn}^{2+}$  couple is  $x \times 10^{-4}$  V, if the  $\text{H}^+$  concentration is decreased from 1M to  $10^{-4}$  M at  $25^\circ\text{C}$ . (Assume concentration of  $\text{MnO}_4^-$  and  $\text{Mn}^{2+}$  to be same on change in  $\text{H}^+$  concentration). The value of x is \_\_\_\_.

(Rounded off to the nearest integer)

$$\left[ \text{Given : } \frac{2303RT}{F} = 0.059 \right]$$

**Ans:** 3776

**Solution:**



$$Q = \frac{[\text{Mn}^{2+}]}{[\text{H}^+]^8 [\text{MnO}_4^-]} \quad E_1 = E^\circ - \frac{0.059}{5} \log(Q_1)$$

$$E_2 = E^\circ - \frac{0.059}{5} \log(Q_2) \quad E_2 - E_1 = \frac{0.059}{5} \log\left(\frac{Q_1}{Q_2}\right)$$

$$\frac{0.059}{5} \log\left\{\frac{[\text{H}^+]_{\text{II}}}{[\text{H}^+]_{\text{I}}}\right\}^8 = \frac{0.059}{5} \log\left(\frac{10^{-4}}{1}\right)^8$$

$$(E_2 - E_1) = \frac{0.059}{5} \times (-32) \quad |(E_2 - E_1)| = 32 \times \frac{0.059}{5} = x \times 10^{-4}$$

$$\frac{32 \times 590}{5} \times 10^{-4} = x \times 10^{-4}$$

$$= 3776 \times 10^{-4} \quad \text{so, } x = 3776$$

2. Among the following allotropic forms of sulphur, the number of allotropic forms, which will show paramagnetism is \_\_\_\_.

(1)  $\alpha$ -sulphur

(2)  $\beta$ -sulphur

(3)  $S_2$ -form

**Ans: 1**

**Solution:**

$S_2$  is like  $O_2$  i.e. paramagnetic as per molecular orbital theory.

3.  $C_6H_6$  freezes at  $5.5^\circ C$ . The temperature at which a solution of 10 g of  $C_4H_{10}$  in 200 g of  $C_6H_6$  freeze is \_\_\_\_  $^\circ C$ . (The molal freezing point depression constant of  $C_6H_6$  is  $5.12^\circ C/m$ ).

**Ans: 1**

**Solution:**

$$\Delta T_f = i \times K_f \times m$$
$$= 1 \times 5.12 \times \frac{10/58}{200} \times 1000$$

$$\Delta T_f = \frac{5.12 \times 50}{58} = 4.414$$

$$T_{f(\text{solution})} = T_{K(\text{solvent})} - \Delta T_f = 5.5 - 4.414 = 1.086^\circ C$$
$$\approx 1.09^\circ C = 1 \text{ (nearest integer)}$$

4. The volume occupied by 4.75 g of acetylene gas at  $50^\circ C$  and 740 mmHg pressure is \_\_\_\_ L.

(Rounded off to the nearest integer)

(Given  $R = 0.0826 \text{ L atm K}^{-1} \text{ mol}^{-1}$ )

**Ans: 5**

**Solution:**

$$T = 50^\circ C = 323.15 \text{ K}$$

$$P = 740 \text{ mm of Hg} = \frac{740}{760} \text{ atm}$$

$$V = ?$$

$$\text{moles (n)} = \frac{4.75}{26} \text{ atm}$$

$$V = \frac{4.75}{26} \times \frac{0.0821 \times 323.15}{740} \times 760$$

$$V = 4.97 \approx 5 \text{ Lit}$$

5. The solubility product of  $PbI_2$  is  $8.0 \times 10^{-9}$ . The solubility of lead iodide in 0.1 molar solution of lead nitrate is  $x \times 10^{-6}$  mol/L. The value of  $x$  is \_\_\_\_\_ (Rounded off to the nearest integer)

Given  $\sqrt{2} = 1.41$

**Ans:** 141

Solution:  $PbI_2(s) \rightleftharpoons Pb^{2+}(aq) + 2I^{-}(aq)$

$S+0.1 \qquad 2s$

$$K_{SP}(PbI_2) = 8 \times 10^{-9}$$

$$K_{SP} = [Pb^{+2}][I^{-}]^2$$

$$8 \times 10^{-9} = (S + 0.1)(2S)^2 \Rightarrow (8 \times 10^{-9} + 0.1) \times 4S^2$$


$$\Rightarrow S^2 = 2 \times 10^{-8}$$

$$S = 1.414 \times 10^{-4} \text{ mol/Lit}$$

$$= x \times 10^{-6} \text{ mol/Lit} \quad \therefore x = 141.4 \approx 141$$


6. The total number of amines among the following which can be synthesized by Gabriel synthesis is \_\_\_\_\_.

1.




2.  $\text{CH}_3\text{CH}_2\text{NH}_2$

3.



4.



**Ans: 3**


**Solution:**

Only 1° amines can be prepared by Gabriel synthesis.

7. 1.86 g of aniline completely reacts to form acetanilide. 10% of the product is lost during purification. Amount of acetanilide obtained after purification (in g) is \_\_\_\_  $\times 10^{-2}$ .

**Ans: 243**

**Solution:**



Molar mass = 93    Molar mass = 135

93 g Aniline produce 135 g acetanilide

$$1.86 \text{ g produce } \frac{135 \times 1.86}{93} = 2.70 \text{ g}$$


At 10% loss, 90% product will be formed after purification.

$$\therefore \text{Amount of product obtained} = \frac{2.70 \times 90}{100} = 2.43 \text{ g} = 243 \times 10^{-2} \text{ g}$$

8. The formula of a gaseous hydrocarbon which requires 6 times of its own volume of  $\text{O}_2$  for complete oxidation and produces 4 times its own volume of  $\text{CO}_2$  is  $\text{C}_x\text{H}_y$ . The value of y is .....

**Ans: 8**

**Solution:**



Applying POAC on 'O' atoms

$$6 \times 2 = 4 \times 2 + \frac{y}{2} \times 1$$

$$\frac{y}{2} = 4 \Rightarrow y = 8$$

9. Sucrose hydrolyses in acid solution into glucose and fructose following first order rate law with a half-life of 3.33 h at 25°C. After 9h, the fraction of sucrose remaining is f. The value of  $\log_{10} \frac{1}{f}$  is \_\_\_\_\_  $\times 10^{-2}$  (Rounded off to the nearest integer)

[Assume:  $\ln 10 = 2.303$ ,  $\ln 2 = 0.693$ ]

**Ans : 81**

**Solution:**

Sucrose  $\xrightarrow{\text{Hydrolysis}}$  Glucose + Fructose

$$t_{1/2} = 3.33\text{h} = \frac{10}{3}\text{h}$$

$$C_t = \frac{C_o}{2^{t/t_{1/2}}}$$

$$\text{Fraction of sucrose remaining} = f = \frac{C_t}{C_o} = \frac{1}{2^{\frac{t}{t_{1/2}}}}$$

$$\frac{1}{f} = 2^{t/t_{1/2}}$$

$$\log(1/f) = \log(2^{t/t_{1/2}}) = \frac{t}{t_{1/2}} \log(2)$$

$$\frac{9}{10/3} \times 0.3 = \frac{8.1}{10} = 0.81$$

$$= x \times 10^{-2} \quad x = 81$$

10. Assuming ideal behavior, the magnitude of  $\log K$  for the following reaction at 25°C is  $x \times 10^{-1}$ . The value of x is \_\_\_\_\_. (Integer answer)



[Given:  $\Delta_f G^\circ(\text{HC} \equiv \text{CH}) = -2.04 \times 10^5 \text{ J mol}^{-1}$ ;  $\Delta_f G^\circ(\text{C}_6\text{H}_6) = -1.24 \times 10^5 \text{ J mol}^{-1}$ ;

$R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$ ]

**Ans: 855**

**Solution:**

$$\Delta G_r^\circ = \Delta G_f^\circ[\text{C}_6\text{H}_6(\text{l})] - 3 \times \Delta G_f^\circ[\text{HC} \equiv \text{CH}]$$

$$= [-1.24 \times 10^5 - 3 \times (-2.04 \times 10^5)]$$

$$= 4.88 \times 10^5 \text{ J/mol}$$

$$\Delta G_r^\circ = -RT \ln(K_{\text{eq}})$$

$$\log(K_{\text{eq}}) = \frac{-\Delta G_r^\circ}{2.303RT}$$

$$\frac{-4.88 \times 10^5}{2.303 \times 8.314 \times 298} = -8.55 \times 101 = 855 \times 10^{-1}$$