

1. Let  $a, b \in \mathbf{R}$ . If the mirror image of the point  $P(a, 6, 9)$  with respect to the line  $\frac{x-3}{7} = \frac{y-2}{5} = \frac{z-1}{-9}$  is  $(20, b, -a-9)$ , then  $|a+b|$  is equal to :
- (1) 86                      (2) 88                      (3) 84                      (4) 90

**Ans. (2)**

Sol.  $P(a, 6, 9)$ ,  $Q(20, b, -a-9)$

Mid point of  $PQ = \left(\frac{a+20}{2}, \frac{b+6}{2}, -\frac{a}{2}\right)$  lie on the line.

$$\frac{\frac{a+20}{2}-3}{7} = \frac{\frac{b+6}{2}-2}{5} = \frac{-\frac{a}{2}-1}{-9}$$

$$\Rightarrow \frac{a+20-6}{14} = \frac{b+6-4}{10} = \frac{-a-2}{-18}$$

$$\Rightarrow \frac{a+14}{14} = \frac{a+2}{18}$$

$$\Rightarrow 18a + 252 = 14a + 28$$

$$\Rightarrow 4a = -224$$

$$\boxed{a = -56}$$

$$\frac{b+2}{10} = \frac{a+2}{18}$$

$$\Rightarrow \frac{b+2}{10} = \frac{-54}{18}$$

$$\Rightarrow \frac{b+2}{10} = -3 \Rightarrow b = -32$$

$$|a+b| = |-56-32| = 88$$

2. Let  $f$  be a twice differentiable function defined on  $\mathbf{R}$  such that  $f(0) = 1, f'(0) = 2$  and  $f'(x) \neq 0$  for all  $x \in \mathbf{R}$ . If  $\left| \frac{f(x)}{f'(x)} - \frac{f'(x)}{f''(x)} \right| = 0$ , for all  $x \in \mathbf{R}$ , then the value of  $f(1)$  lies in the interval :
- (1) (9, 12)                      (2) (6, 9)                      (3) (3, 6)                      (4) (0, 3)

**Ans. (2)**

Sol. Given  $f(x)f''(x) - (f'(x))^2 = 0$

$$\text{Let } h(x) = \frac{f(x)}{f'(x)}$$

$$\text{Then } h'(x) = 0 \Rightarrow h(x) = k$$

$$\Rightarrow \frac{f(x)}{f'(x)} = k \Rightarrow f(x) = kf'(x)$$

$$\Rightarrow f(0) = kf'(0)$$

$$\Rightarrow k = \frac{1}{2}$$

$$\text{Now, } f(x) = \frac{1}{2}f'(x)$$

# JEE Main 2021 24 Feb Shift 2



$$\Rightarrow \int 2 dx = \int \frac{f'(x)}{f(x)} dx$$

$$\Rightarrow 2x = \ln|f(x)| + C$$

$$\text{As } f(0) = 1 \Rightarrow C = 0$$

$$\Rightarrow 2x = \ln|f(x)|$$

$$\Rightarrow f(x) = \pm e^{2x}$$

$$\text{As } f(0) = 1 \Rightarrow f(x) = e^{2x}$$

$$\therefore f(1) = e^2 \approx 7.38$$

3. A possible value of  $\tan\left(\frac{1}{4}\sin^{-1}\frac{\sqrt{63}}{8}\right)$  is :

(1)  $\frac{1}{2\sqrt{2}}$

(2)  $\frac{1}{\sqrt{7}}$


(3)  $\sqrt{7} - 1$

(4)  $2\sqrt{2} - 1$

Ans. (2)

Sol.  $\tan\left(\frac{1}{4}\sin^{-1}\frac{\sqrt{63}}{8}\right)$

$$\text{Let } \sin^{-1}\left(\frac{\sqrt{63}}{8}\right) = \theta \quad \sin \theta = \frac{\sqrt{63}}{8}$$



$$\cos \theta = \frac{1}{8}$$

$$2 \cos^2 \frac{\theta}{2} - 1 = \frac{1}{8}$$

$$\Rightarrow \cos^2 \frac{\theta}{2} = \frac{9}{16}$$

$$\cos \frac{\theta}{2} = \frac{3}{4}$$

$$\Rightarrow \frac{1 - \tan^2 \frac{\theta}{4}}{1 + \tan^2 \frac{\theta}{4}} = \frac{3}{4}$$

$$\tan \frac{\theta}{4} = \frac{1}{\sqrt{7}}$$

4. The probability that two randomly selected subsets of the set  $\{1, 2, 3, 4, 5\}$  have exactly two elements in their intersection, is :

(1)  $\frac{65}{2^7}$

(2)  $\frac{135}{2^9}$

(3)  $\frac{65}{2^8}$

(4)  $\frac{35}{2^7}$

**Ans. (2)**

**Sol.** Let  $A$  and  $B$  be two subsets.

For each  $x \in \{1, 2, 3, 4, 5\}$ , there are four possibilities :

$$x \in A \cap B, x \in A' \cap B, x \in A \cap B', x \in A' \cap B'$$

So, the number of elements in sample space =  $4^5$

Required probability

$$\begin{aligned} &= \frac{{}^5C_2 \times 3^3}{4^5} \\ &= \frac{10 \times 27}{2^{10}} = \frac{135}{2^9} \end{aligned}$$

5. The vector equation of the plane passing through the intersection of the planes

$$\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 1 \text{ and } \vec{r} \cdot (\hat{i} - 2\hat{j}) = -2, \text{ and the point } (1, 0, 2) \text{ is :}$$

$$(1) \vec{r} \cdot (\hat{i} - 7\hat{j} + 3\hat{k}) = \frac{7}{3}$$

$$(2) \vec{r} \cdot (\hat{i} + 7\hat{j} + 3\hat{k}) = 7$$

$$(3) \vec{r} \cdot (3\hat{i} + 7\hat{j} + 3\hat{k}) = 7$$

$$(4) \vec{r} \cdot (\hat{i} + 7\hat{j} + 3\hat{k}) = \frac{7}{3}$$

**Ans. (2)**

**Sol.** Family of planes passing through intersection of planes is

$$\{\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) - 1\} + \lambda\{\vec{r} \cdot (\hat{i} - 2\hat{j}) + 2\} = 0$$

The above curve passes through  $\hat{i} + 2\hat{k}$ ,

$$(3 - 1) + \lambda(1 + 2) = 0 \Rightarrow \lambda = -\frac{2}{3}$$

Hence, equation of plane is

$$3\{\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) - 1\} - 2\{\vec{r} \cdot (\hat{i} - 2\hat{j}) + 2\} = 0$$

$$\Rightarrow \vec{r} \cdot (\hat{i} + 7\hat{j} + 3\hat{k}) = 7$$

**TRICK:** Only option (2) satisfies the point (1, 0, 2)

6. If P is a point on the parabola  $y = x^2 + 4$  which is closest to the straight line  $y = 4x - 1$ , then the co-ordinates of P are :

$$(1) (-2, 8)$$

$$(2) (1, 5)$$

$$(3) (3, 13)$$

$$(4) (2, 8)$$


**Ans. (4)**

**Sol.** Tangent at P is parallel to the given line.

$$\frac{dy}{dx} \Big|_P = 4$$

$$\Rightarrow 2x_1 = 4$$

# JEE Main 2021 24 Feb Shift 2



$\Rightarrow x_1 = 2$

Required point is (2, 8)

7. Let a, b, c be in arithmetic progression. Let the centroid of the triangle with vertices (a, c), (2, b) and (a, b) be  $(\frac{10}{3}, \frac{7}{3})$ . If  $\alpha, \beta$  are the roots of the equation  $ax^2 + bx + 1 = 0$ , then the value of  $\alpha^2 + \beta^2 - \alpha\beta$  is :
- (1)  $\frac{71}{256}$                       (2)  $-\frac{69}{256}$                       (3)  $\frac{69}{256}$                       (4)  $-\frac{71}{256}$

**Ans. (4)**

Sol.  $2b = a + c$   
 $\frac{2a+2}{3} = \frac{10}{3}$  and  $\frac{2b+c}{3} = \frac{7}{3}$   
 $\Rightarrow a = 4$      $\left. \begin{matrix} 2b + c = 7 \\ 2b - c = 4 \end{matrix} \right\}$  solving  
 $b = \frac{11}{4}$  and  $c = \frac{3}{2}$   
 $\therefore$  Quadratic equation is  $4x^2 + \frac{11}{4}x + 1 = 0$   
 $\therefore$  The value of  $(\alpha + \beta)^2 - 3\alpha\beta = \frac{121}{256} - \frac{3}{4} = -\frac{71}{256}$

8. The value of the integral,  $\int_1^3 [x^2 - 2x - 2] dx$ , where [x] denotes the greatest integer less than or equal to x, is :
- (1) -4                      (2) -5                      (3)  $-\sqrt{2} - \sqrt{3} - 1$                       (4)  $-\sqrt{2} - \sqrt{3} + 1$

**Ans. (3)**

Sol.  $I = \int_1^3 -3dx + \int_1^3 [(x - 1)^2] dx$   
 Put  $x - 1 = t$  ;  $dx = dt$   
 $I = (-6) + \int_0^2 [t^2] dt$   
 $I = -6 + \int_0^1 0dt + \int_1^{\sqrt{2}} 1dt + \int_{\sqrt{2}}^{\sqrt{3}} 2dt + \int_{\sqrt{3}}^2 3dt$

# JEE Main 2021 24 Feb Shift 2

B

$$I = -6 + (\sqrt{2} - 1) + 2\sqrt{3} - 2\sqrt{2} + 6 - 3\sqrt{3}$$
$$I = -1 - \sqrt{2} - \sqrt{3}$$

9. Let  $f : \mathbf{R} \rightarrow \mathbf{R}$  be defined as

$$f(x) = \begin{cases} -55x, & \text{if } x < -5 \\ 2x^3 - 3x^2 - 120x, & \text{if } -5 \leq x \leq 4 \\ 2x^3 - 3x^2 - 36x - 336, & \text{if } x > 4 \end{cases}$$

Let  $A = \{x \in \mathbf{R} : f \text{ is increasing}\}$ . Then  $A$  is equal to :

- (1)  $(-5, -4) \cup (4, \infty)$                                       (2)  $(-5, \infty)$   
(3)  $(-\infty, -5) \cup (4, \infty)$                                       (4)  $(-\infty, -5) \cup (-4, \infty)$

Ans. (1)

Sol. 
$$f'(x) = \begin{cases} -55 & ; x < -5 \\ 6(x^2 - x - 20) & ; -5 < x < 4 \\ 6(x^2 - x - 6) & ; x > 4 \end{cases}$$
$$\Rightarrow f'(x) = \begin{cases} -55 & ; x < -5 \\ 6(x-5)(x+4) & ; -5 < x < 4 \\ 6(x-3)(x+2) & ; x > 4 \end{cases}$$

Hence,  $f(x)$  is monotonically increasing in  $(-5, -4) \cup (4, \infty)$

10. If the curve  $y = ax^2 + bx + c$ ,  $x \in \mathbf{R}$  passes through the point  $(1, 2)$  and the tangent line to this curve at origin is  $y = x$ , then the possible values of  $a, b, c$  are :

- (1)  $a = 1, b = 1, c = 0$                                       (2)  $a = -1, b = 1, c = 1$   
(3)  $a = 1, b = 0, c = 1$                                       (4)  $a = \frac{1}{2}, b = \frac{1}{2}, c = 1$

Ans. (1)

Sol.  $2 = a + b + c$   
 $\frac{dy}{dx} = 2ax + b, \left(\frac{dy}{dx}\right)_{(0,0)} = 1$   
 $\Rightarrow b = 1 \text{ and } a + c = 1$   
Since  $(0, 0)$  lies on curve,  
 $\therefore c = 0, a = 1$

TRICK :  $(0,0)$  lies on the curve. Only option (1) has  $c = 0$

11. The negation of the statement  $\sim p \wedge (p \vee q)$  is :

- (1)  $\sim p \wedge q$                                       (2)  $p \wedge \sim q$                                       (3)  $\sim p \vee q$                                       (4)  $p \vee \sim q$

# JEE Main 2021 24 Feb Shift 2



**Ans. (4)**

Sol. Negation of  $\sim p \wedge (p \vee q)$  is

$$\begin{aligned} & \sim [\sim p \wedge (p \vee q)] \\ & \equiv p \vee \sim (p \vee q) \\ & \equiv p \vee (\sim p \wedge \sim q) \\ & \equiv (p \vee \sim p) \wedge (p \vee \sim q) \\ & \equiv T \wedge (p \vee \sim q), \text{ where } T \text{ is tautology.} \\ & \equiv p \vee \sim q \end{aligned}$$

**12.** For the system of linear equations :

$$x - 2y = 1, x - y + kz = -2, ky + 4z = 6, k \in \mathbf{R}$$

consider the following statements :

- (A) The system has unique solution if  $k \neq 2, k \neq -2$ .
- (B) The system has unique solution if  $k = -2$ .
- (C) The system has unique solution if  $k = 2$ .
- (D) The system has no-solution if  $k = 2$ .
- (E) The system has infinite number of solutions if  $k \neq -2$ .

Which of the following statements are **correct** ?

- (1) (B) and (E) only
- (2) (C) and (D) only
- (3) (A) and (D) only
- (4) (A) and (E) only

**Ans. (3)**

Sol.  $x - 2y + 0.z = 1$

$$x - y + kz = -2$$

$$0.x + ky + 4z = 6$$

$$\Delta = \begin{vmatrix} 1 & -2 & 0 \\ 1 & -1 & k \\ 0 & k & 4 \end{vmatrix} = 4 - k^2$$

For unique solution,  $4 - k^2 \neq 0$

$$\boxed{k \neq \pm 2}$$

For  $k = 2$ ,

$$x - 2y + 0.z = 1$$

$$x - y + 2z = -2$$

# JEE Main 2021 24 Feb Shift 2



$$0. x + 2y + 4z = 6$$

$$\Delta_x = \begin{vmatrix} 1 & -2 & 0 \\ -2 & -1 & 2 \\ 6 & 2 & 4 \end{vmatrix} = (-8) + 2[-20]$$

$$\Rightarrow \Delta_x = -48 \neq 0$$

For  $k = 2, \Delta_x \neq 0$

So, for  $k = 2$ , the system has no solution.

13. For which of the following curves, the line  $x + \sqrt{3}y = 2\sqrt{3}$  is the tangent at the point  $\left(\frac{3\sqrt{3}}{2}, \frac{1}{2}\right)$ ?

(1)  $x^2 + 9y^2 = 9$

(2)  $2x^2 - 18y^2 = 9$

(3)  $y^2 = \frac{1}{6\sqrt{3}}x$

(4)  $x^2 + y^2 = 7$

Ans. (1)

Sol. Tangent to  $x^2 + 9y^2 = 9$  at point  $\left(\frac{3\sqrt{3}}{2}, \frac{1}{2}\right)$  is  $x\left(\frac{3\sqrt{3}}{2}\right) + 9y\left(\frac{1}{2}\right) = 9$

$$\Rightarrow 3\sqrt{3}x + 9y = 18 \Rightarrow x + \sqrt{3}y = 2\sqrt{3}$$

$\Rightarrow$  Option (1) is true.

14. The angle of elevation of a jet plane from a point A on the ground is  $60^\circ$ . After a flight of 20 seconds at the speed of 432 km/hour, the angle of elevation changes to  $30^\circ$ . If the jet plane is flying at a constant height, then its height is :

(1)  $1200\sqrt{3}$  m


(2)  $1800\sqrt{3}$  m

(3)  $3600\sqrt{3}$  m

(4)  $2400\sqrt{3}$  m

Ans. (1)

Sol.



$$v = 432 \times \frac{1000}{60 \times 60} \text{ m/sec} = 120 \text{ m/sec}$$

$$\text{Distance PQ} = v \times 20 = 2400 \text{ m}$$

In  $\Delta PAC$

$$\tan 60^\circ = \frac{h}{AC} \Rightarrow AC = \frac{h}{\sqrt{3}}$$

In  $\Delta AQD$

$$\tan 30^\circ = \frac{h}{AD} \Rightarrow AD = \sqrt{3}h$$

$$AD = AC + CD$$

$$\Rightarrow \sqrt{3}h = \frac{h}{\sqrt{3}} + 2400 \Rightarrow \frac{2h}{\sqrt{3}} = 2400$$

$$\Rightarrow h = 1200\sqrt{3} \text{ m}$$

15. For the statements  $p$  and  $q$ , consider the following compound statements :

(a)  $(\sim q \wedge (p \rightarrow q)) \rightarrow \sim p$

(b)  $((p \vee q) \wedge \sim p) \rightarrow q$

Then which of the following statements is **correct** ?

(1) (a) is a tautology but not (b)

(2) (a) and (b) both are not tautologies.

(3) (a) and (b) both are tautologies.

(4) (b) is a tautology but not (a).



# JEE Main 2021 24 Feb Shift 2



**Ans. (3)**

	$p$	$q$	$\sim q$	$p \rightarrow q$	$\sim q \wedge (p \rightarrow q)$	$\sim p$	$(\sim q) \wedge (p \rightarrow q) \rightarrow \sim p$
	$T$	$T$	$F$	$T$	$F$	$F$	$T$
Sol. (a)	$T$	$F$	$T$	$F$	$F$	$F$	$T$
	$F$	$T$	$F$	$T$	$F$	$T$	$T$
	$F$	$F$	$T$	$T$	$T$	$T$	$T$

(a) is tautology.

	$p$	$q$	$p \vee q$	$\sim p$	$(p \vee q) \wedge \sim p$	$((p \vee q) \wedge \sim p) \rightarrow q$
	$T$	$T$	$T$	$F$	$F$	$T$
(b)	$T$	$F$	$T$	$F$	$F$	$T$
	$F$	$T$	$T$	$T$	$T$	$T$
	$F$	$F$	$F$	$T$	$F$	$T$

(b) is tautology.

$\therefore$  (a) and (b) both are tautologies.

**16.** Let A and B be  $3 \times 3$  real matrices such that A is symmetric matrix and B is skew-symmetric matrix. Then the system of linear equations  $(A^2 B^2 - B^2 A^2)X = O$ , where X is a  $3 \times 1$  column matrix of unknown variables and O is a  $3 \times 1$  null matrix, has :

- |                               |                           |
|-------------------------------|---------------------------|
| (1) a unique solution         | (2) exactly two solutions |
| (3) infinitely many solutions | (4) no solution           |

**Ans. (3)**

Sol.  $A^T = A, B^T = -B$

Let  $A^2 B^2 - B^2 A^2 = P$

$$P^T = (A^2 B^2 - B^2 A^2)^T = (A^2 B^2)^T - (B^2 A^2)^T$$

$$= (B^2)^T (A^2)^T - (A^2)^T (B^2)^T$$

$$= B^2 A^2 - A^2 B^2$$

$\Rightarrow P$  is a skew-symmetric matrix.

$$\begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\therefore ay + bz = 0 \quad \dots(1)$$

$$-ax + cz = 0 \quad \dots(2)$$

$$-bx - cy = 0 \quad \dots(3)$$

From equation (1), (2), (3)

$$\Delta = 0 \text{ and } \Delta_1 = \Delta_2 = \Delta_3 = 0$$

$\therefore$  System of equations has infinite number of solutions.

# JEE Main 2021 24 Feb Shift 2



17. If  $n \geq 2$  is a positive integer, then the sum of the series

${}^{n+1}C_2 + 2({}^2C_2 + {}^3C_2 + {}^4C_2 + \dots + {}^nC_2)$  is :

(1)  $\frac{n(n+1)^2(n+2)}{12}$

(2)  $\frac{n(n-1)(2n+1)}{6}$

(3)  $\frac{n(n+1)(2n+1)}{6}$

(4)  $\frac{n(2n+1)(3n+1)}{6}$

**Ans. (3)**

Sol.  ${}^2C_2 = {}^3C_3$

Let  $S = {}^3C_3 + {}^3C_2 + \dots + {}^nC_2 = {}^{n+1}C_3$  ( $\because {}^nC_r + {}^nC_{r-1} = {}^{n+1}C_r$ )

$\therefore {}^{n+1}C_2 + {}^{n+1}C_3 + {}^{n+1}C_3$

$= {}^{n+2}C_3 + {}^{n+1}C_3$

$= \frac{(n+2)!}{3!(n-1)!} + \frac{(n+1)!}{3!(n-2)!}$

$= \frac{(n+2)(n+1)n}{6} + \frac{(n+1)(n)(n-1)}{6} = \frac{n(n+1)(2n+1)}{6}$

**TRICK :** Put  $n = 2$  and verify the options.

18. If a curve  $y = f(x)$  passes through the point  $(1, 2)$  and satisfies  $x \frac{dy}{dx} + y = bx^4$ , then for

what value of  $b, \int_1^2 f(x) dx = \frac{62}{5}$  ?

(1) 5

(2)  $\frac{62}{5}$

(3)  $\frac{31}{5}$

(4) 10

**Ans. (4)**

Sol.  $\frac{dy}{dx} + \frac{y}{x} = bx^3$

I.F. =  $e^{\int \frac{dx}{x}} = x$

$\therefore yx = \int bx^4 dx = \frac{bx^5}{5} + c$

Above curve passes through  $(1,2)$ .

$2 = \frac{b}{5} + c$

Also,  $\int_1^2 \left( \frac{bx^4}{5} + \frac{c}{x} \right) dx = \frac{62}{5}$

$\Rightarrow \frac{b}{25} \times 32 + c \ln 2 - \frac{b}{25} = \frac{62}{5}$

$\Rightarrow c = 0$  and  $b = 10$

# JEE Main 2021 24 Feb Shift 2




19. The area of the region :  $R = \{(x, y) : 5x^2 \leq y \leq 2x^2 + 9\}$  is :

- (1)  $9\sqrt{3}$  square units                      (2)  $12\sqrt{3}$  square units  
 (3)  $11\sqrt{3}$  square units                      (4)  $6\sqrt{3}$  square units

**Ans. (2)**

Sol.



Required area

$$\begin{aligned} &= 2 \int_0^{\sqrt{3}} (2x^2 + 9 - 5x^2) dx \\ &= 2 \int_0^{\sqrt{3}} (9 - 3x^2) dx \\ &= 2 [9x - x^3]_0^{\sqrt{3}} = 12\sqrt{3} \end{aligned}$$

20. Let  $f(x)$  be a differentiable function defined on  $[0, 2]$  such that  $f'(x) = f'(2 - x)$  for all  $x \in (0, 2)$ ,  $f(0) = 1$  and  $f(2) = e^2$ . Then the value of  $\int_0^2 f(x) dx$  is :

- (1)  $1 + e^2$                       (2)  $1 - e^2$                       (3)  $2(1 - e^2)$                       (4)  $2(1 + e^2)$

**Ans. (1)**

Sol.  $f'(x) = f'(2 - x)$

On integrating both sides, we get

$$f(x) = -f(2 - x) + c$$

Put  $x = 0$

$$f(0) + f(2) = c$$

$$\Rightarrow c = 1 + e^2$$

$$\Rightarrow f(x) + f(2 - x) = 1 + e^2$$

# JEE Main 2021 24 Feb Shift 2



$$I = \int_0^2 f(x)dx = \int_0^1 \{f(x) + f(2-x)\}dx = 1 + e^2$$

## Section B

1. The number of the real roots of the equation  $(x+1)^2 + |x-5| = \frac{27}{4}$  is \_\_\_\_\_.

**Ans.** 2

**Sol.** For  $x \geq 5$ ,

$$(x+1)^2 + (x-5) = \frac{27}{4}$$

$$\Rightarrow x^2 + 3x - 4 = \frac{27}{4}$$

$$\Rightarrow x^2 + 3x - \frac{43}{4} = 0$$

$$\Rightarrow 4x^2 + 12x - 43 = 0$$

$$x = \frac{-12 \pm \sqrt{144 + 688}}{8}$$

$$x = \frac{-12 \pm \sqrt{832}}{8} = \frac{-12 \pm 28.8}{8}$$

$$= \frac{-3 \pm 7.2}{2}$$

$$= \frac{-3+7.2}{2}, \frac{-3-7.2}{2} \quad (\text{therefore, no solution})$$

For  $x < 5$ ,

$$(x+1)^2 - (x-5) = \frac{27}{4}$$

$$\Rightarrow x^2 + x + 6 - \frac{27}{4} = 0$$

$$\Rightarrow 4x^2 + 4x - 3 = 0$$

$$x = \frac{-4 \pm \sqrt{16+48}}{8}$$

$$x = \frac{-4 \pm 8}{8} \Rightarrow x = -\frac{12}{8}, \frac{4}{8}$$

$\therefore$  2 real roots.

2. The students  $S_1, S_2, \dots, S_{10}$  are to be divided into 3 groups A, B and C such that each group has at least one student and the group C has at most 3 students. Then the total number of possibilities of forming such groups is \_\_\_\_\_.



Ans. 31650

Sol.

$$C \rightarrow 1 \quad 9 \begin{cases} A \\ B \end{cases}$$

$$C \rightarrow 2 \quad 8 \begin{cases} A \\ B \end{cases}$$

$$C \rightarrow 3 \quad 7 \begin{cases} A \\ B \end{cases}$$

Number of ways

$$\begin{aligned} &= {}^{10}C_1 [2^9 - 2] + {}^{10}C_2 [2^8 - 2] + {}^{10}C_3 [2^7 - 2] \\ &= 2^7 [{}^{10}C_1 \times 4 + {}^{10}C_2 \times 2 + {}^{10}C_3] - 20 - 90 - 240 \\ &= 128 [40 + 90 + 120] - 350 \\ &= (128 \times 250) - 350 \\ &= 10[3165] = 31650 \end{aligned}$$

3. If  $a + \alpha = 1, b + \beta = 2$  and  $af(x) + \alpha f\left(\frac{1}{x}\right) = bx + \frac{\beta}{x}, x \neq 0$ , then the value of the expression

$$\frac{f(x) + f\left(\frac{1}{x}\right)}{x + \frac{1}{x}} \text{ is } \underline{\hspace{2cm}}.$$

Ans. 2

Sol.  $af(x) + \alpha f\left(\frac{1}{x}\right) = bx + \frac{\beta}{x} \quad \dots(i)$

Replace  $x$  by  $\frac{1}{x}$

$$af\left(\frac{1}{x}\right) + \alpha f(x) = \frac{b}{x} + \beta x \quad \dots(ii)$$

(i) + (ii)

$$(a + \alpha) \left[ f(x) + f\left(\frac{1}{x}\right) \right] = \left( x + \frac{1}{x} \right) (b + \beta)$$

$$\Rightarrow \frac{f(x) + f\left(\frac{1}{x}\right)}{x + \frac{1}{x}} = \frac{2}{1} = 2$$

# JEE Main 2021 24 Feb Shift 2



4. If the variance of 10 natural numbers  $1, 1, 1, \dots, 1, k$  is less than 10, then the maximum possible value of  $k$  is \_\_\_\_\_.

**Ans. 11**

Sol. 
$$\sigma^2 = \frac{\Sigma x^2}{n} - \left(\frac{\Sigma x}{n}\right)^2$$

$$\Rightarrow \sigma^2 = \frac{(9+k^2)}{10} - \left(\frac{9+k}{10}\right)^2 < 10$$

$$\Rightarrow 10(9+k^2) - (81+k^2+18k) < 1000$$

$$\Rightarrow 90+10k^2-k^2-18k-81 < 1000$$

$$\Rightarrow 9k^2-18k+9 < 1000$$

$$\Rightarrow (k-1)^2 < \frac{1000}{9} \Rightarrow k-1 < \frac{10\sqrt{10}}{3}$$

$$\Rightarrow k < \frac{10\sqrt{10}}{3} + 1$$

Maximum possible integral value of  $k$  is 11.

5. Let  $\lambda$  be an integer. If the shortest distance between the lines  $x - \lambda = 2y - 1 = -2z$  and

$x = y + 2\lambda = z - \lambda$  is  $\frac{\sqrt{7}}{2\sqrt{2}}$ , then the value of  $|\lambda|$  is \_\_\_\_\_.

**Ans. 1**

Sol. 
$$\frac{x-\lambda}{1} = \frac{y-\frac{1}{2}}{\frac{1}{2}} = \frac{z}{-\frac{1}{2}}$$

$$\frac{x-\lambda}{2} = \frac{y-\frac{1}{2}}{1} = \frac{z}{-1} \quad \dots(1) \quad \text{Point on line} = \left(\lambda, \frac{1}{2}, 0\right)$$

$$\frac{x}{1} = \frac{y+2\lambda}{1} = \frac{z-\lambda}{1} \quad \dots(2) \quad \text{Point on line} = (0, -2\lambda, \lambda)$$

Distance between skew lines =  $\frac{[\vec{a}_2 - \vec{a}_1 \quad \vec{b}_1 \quad \vec{b}_2]}{|\vec{b}_1 \times \vec{b}_2|}$

$$\frac{\begin{vmatrix} \lambda & \frac{1}{2}+2\lambda & -\lambda \\ 2 & 1 & -1 \\ 1 & 1 & 1 \end{vmatrix}}{\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -1 \\ 1 & 1 & 1 \end{vmatrix}}$$

$$= \frac{|-5\lambda - \frac{3}{2}|}{\sqrt{14}} = \frac{\sqrt{7}}{2\sqrt{2}} \quad (\text{Given})$$

$\Rightarrow |10\lambda + 3| = 7 \Rightarrow \lambda = -1$  as  $\lambda$  is an integer.

$\Rightarrow |\lambda| = 1$

# JEE Main 2021 24 Feb Shift 2



6. Let  $i = \sqrt{-1}$ . If  $\frac{(-1+i\sqrt{3})^{21}}{(1-i)^{24}} + \frac{(1+i\sqrt{3})^{21}}{(1+i)^{24}} = k$ , and  $n = [|k|]$  be the greatest integral part of  $|k|$ . Then  $\sum_{j=0}^{n+5} (j+5)^2 - \sum_{j=0}^{n+5} (j+5)$  is equal to \_\_\_\_.

**Ans. 310**

Sol. 
$$\frac{(2e^{i\frac{2\pi}{3}})^{21}}{(\sqrt{2}e^{-i\frac{\pi}{4}})^{24}} + \frac{(2e^{i\frac{\pi}{3}})^{21}}{(\sqrt{2}e^{i\frac{\pi}{4}})^{24}}$$
$$= \frac{2^{21} \cdot e^{i14\pi}}{2^{12} \cdot e^{-i6\pi}} + \frac{2^{21} (e^{i7\pi})}{2^{12} (e^{i6\pi})}$$
$$= 2^9 e^{i(20\pi)} + 2^9 e^{i\pi}$$
$$= 2^9 + 2^9(-1) = 0 = k$$
$$\therefore n = 0$$

$$\sum_{j=0}^5 (j+5)^2 - \sum_{j=0}^5 (j+5)$$
$$= [5^2 + 6^2 + 7^2 + 8^2 + 9^2 + 10^2] - [5 + 6 + 7 + 8 + 9 + 10]$$
$$= [(1^2 + 2^2 + \dots + 10^2) - (1^2 + 2^2 + 3^2 + 4^2)] - [(1 + 2 + 3 + \dots + 10) - (1 + 2 + 3 + 4)]$$
$$= (385 - 30) - [55 - 10]$$
$$= 355 - 45 = 310$$

7. Let a point P be such that its distance from the point (5, 0) is thrice the distance of P from the point (-5, 0). If the locus of the point P is a circle of radius r, then  $4r^2$  is equal to \_\_\_\_.

**Ans. 56.25**

Sol. Let P be (h, k), A(5, 0) and B(-5, 0)

Given PA = 3PB

$$\Rightarrow PA^2 = 9PB^2$$

$$\Rightarrow (h-5)^2 + k^2 = 9[(h+5)^2 + k^2]$$

$$\Rightarrow 8h^2 + 8k^2 + 100h + 200 = 0$$

$$\therefore \text{Locus of } P \text{ is } x^2 + y^2 + \left(\frac{25}{2}\right)x + 25 = 0$$

$$\text{Centre} \equiv \left(\frac{-25}{4}, 0\right)$$

$$\therefore r^2 = \left(\frac{-25}{4}\right)^2 - 25$$

# JEE Main 2021 24 Feb Shift 2



$$= \frac{625}{16} - 25$$

$$= \frac{225}{16}$$

$$\therefore 4r^2 = 4 \times \frac{225}{16} = \frac{225}{4} = 56.25$$

8. For integers n and r, let  $\binom{n}{r} = \begin{cases} {}^n C_r, & \text{if } n \geq r \geq 0 \\ 0, & \text{otherwise} \end{cases}$

The maximum value of k for which the sum

$$\sum_{i=0}^k \binom{10}{i} \binom{15}{k-i} + \sum_{i=0}^{k+1} \binom{12}{i} \binom{13}{k+1-i}$$
 exists, is equal to \_\_\_\_\_.

**Ans.** \*

**Sol.** BONUS

$$(1+x)^{10} = {}^{10}C_0 + {}^{10}C_1x + {}^{10}C_2x^2 + \dots + {}^{10}C_{10}x^{10}$$

$$(1+x)^{15} = {}^{15}C_0 + {}^{15}C_1x + \dots + {}^{15}C_{k-1}x^{k-1} + {}^{15}C_kx^k + {}^{15}C_{k+1}x^{k+1} + \dots + {}^{15}C_{15}x^{15}$$

$$\sum_{i=0}^k ({}^{10}C_i)({}^{15}C_{k-i}) = {}^{10}C_0 \cdot {}^{15}C_k + {}^{10}C_1 \cdot {}^{15}C_{k-1} + \dots + {}^{10}C_k \cdot {}^{15}C_0$$

Coefficient of  $x_k$  in  $(1+x)^{25}$

$$= {}^{25}C_k$$

$$\sum_{i=0}^{k+1} ({}^{12}C_i)({}^{13}C_{k+1-i}) = {}^{12}C_0 \cdot {}^{13}C_{k+1} + {}^{12}C_1 \cdot {}^{13}C_k + \dots + {}^{12}C_{k+1} \cdot {}^{13}C_0$$

Coefficient of  $x^{k+1}$  in  $(1+x)^{25}$

$$= {}^{25}C_{k+1}$$

$${}^{25}C_k + {}^{25}C_{k+1} = {}^{26}C_{k+1}$$

By the given definition of  $\binom{n}{r}$ , k can be as large as possible.

9. The sum of first four terms of a geometric progression (G.P.) is  $\frac{65}{12}$  and the sum of their

respective reciprocals is  $\frac{65}{18}$ . If the product of first three terms of the G.P. is 1, and the third

term is  $\alpha$ , then  $2\alpha$  is \_\_\_\_\_.

**Ans.** 3

**Sol.** a, ar, ar<sup>2</sup>, ar<sup>3</sup>

$$a + ar + ar^2 + ar^3 = \frac{65}{12} \quad \dots\dots(1)$$

$$\frac{1}{a} + \frac{1}{ar} + \frac{1}{ar^2} + \frac{1}{ar^3} = \frac{65}{18}$$



# JEE Main 2021 24 Feb Shift 2



$$\Rightarrow \frac{1}{a} \left( \frac{r^3 + r^2 + r + 1}{r^3} \right) = \frac{65}{18} \quad \dots\dots(2)$$

$\frac{(1)}{(2)}$ , we get

$$a^2 r^3 = \frac{18}{12} = \frac{3}{2}$$

$$\text{Also, } a^3 r^3 = 1 \Rightarrow a \left( \frac{3}{2} \right) = 1 \Rightarrow a = \frac{2}{3}$$

$$\frac{4}{9} r^3 = \frac{3}{2} \Rightarrow r^3 = \frac{3^3}{2^3} \Rightarrow r = \frac{3}{2}$$


$$\alpha = ar^2 = \frac{2}{3} \cdot \left( \frac{3}{2} \right)^2 = \frac{3}{2}$$

$$\therefore 2\alpha = 3$$

10. If the area of the triangle formed by the positive x-axis, the normal and the tangent to the circle  $(x-2)^2 + (y-3)^2 = 25$  at the point  $(5, 7)$  is  $A$ , then  $24A$  is equal to \_\_\_\_\_.

Ans.

Sol.



Equation of normal at P is

$$(y - 7) = \left( \frac{7-3}{5-2} \right) (x - 5)$$

$$\Rightarrow 3y - 21 = 4x - 20$$

$$\Rightarrow 4x - 3y + 1 = 0$$

$$\Rightarrow M \text{ is } \left( -\frac{1}{4}, 0 \right)$$

Equation of tangent at P is

$$(y - 7) = -\frac{3}{4}(x - 5)$$

$$\Rightarrow 4y - 28 = -3x + 15$$

$$\Rightarrow 3x + 4y = 43$$

$$\Rightarrow N \text{ is } \left( \frac{43}{3}, 0 \right)$$

The question is **wrong**. The normal cuts at a point on the negative axis.



BYJU'S