

SECTION – A

1. Match List-I with List-II.

List – I

Chemical Compound

- (a) Sucralose
- (b) Glyceryl ester of stearic acid
- (c) Sodium benzoate
- (d) Bithionol

List – II

Used as

- (i) Synthetic detergent
- (ii) Artificial sweetener
- (iii) Antiseptic
- (iv) Food preservative

Choose the correct match:



- a. (a)-(i), (b)-(ii), (c)-(iv), (d)-(iii)
- b. (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)
- c. (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i)
- d. (a)-(iii), (b)-(ii), (c)-(iv), (d)-(i)

Ans: (b)

Solution:

- (a) Sucralose → Artificial sweetener
- (b) Glyceryl ester of stearic acid → Synthetic detergent
- (c) Sodium benzoate → Food preservative
- (d) Bithionol → Antiseptic

2.




In the above reactions, the enzyme A and enzyme B respectively are:

- a. Invertase and Amylase
- b. Amylase and Invertase
- c. Invertase and Zymase
- d. Zymase and Invertase

Ans: (c)

Solution:



3. The correct pair(s) of the ambident nucleophiles is (are):


- (a) AgCN/KCN
- (b) RCOOAg/RCOOK
- (c) AgNO₂/KNO₂
- (d) AgI/KI

- a. (A) and (C) only
- c. (B) and (C) only

- b. (B) only
- d. (A) only

Ans: (a)

Solution:



More than one electron donating side

4. During which of the following processes, does entropy decrease?

- (a) Freezing of water to ice at 0°C
- (b) Freezing of water to ice at -10°C
- (c) N₂(g) + 3H₂(g) → 2NH₃(g)
- (d) Adsorption of CO(g) on lead surface.
- (e) Dissolution of NaCl in water

- a. (A), (B), (C) and (D) only
- c. (A) and (E) only

- b. (A), (C) and (E) only
- d. (B) and (C) only

Ans: (a)

Solution:

A, B → Freezing of water will decrease entropy as particles will move closer and forces of attraction will increase. This leads to decrease in randomness. So, entropy decrease.

C → No. of molecules decreasing




D → Adsorption will lead to decrease in randomness of gaseous particles.

E → NaCl(s) → Na⁺(aq) + Cl⁻(aq); ΔS > 0

So, (A, B, C, D) decreases entropy.

5. Match List-I with List-II:

List-I



List-II

(i) Linkage isomerism

(ii) Solvate isomerism

(iii) Co-ordination isomerism

(iv) Optical isomerism

Choose the correct answer from the options given below:

a. (a)-(iii), (b)-(i), (c)-(ii), (d)-(iv)

b. (a)-(i), (b)-(ii), (c)-(iii), (d)-(iv)

c. (a)-(ii), (b)-(i), (c)-(iii), (d)-(iv)

d. (a)-(iv), (b)-(ii), (c)-(iii), (d)-(i)

Ans: (a)

Solution:

Theory based

6. The common positive oxidation states for an element with atomic number 24, are:

a. +1 and +3

b. +1 to +6

c. +1 and +3 to +6

d. +2 to +6

Ans: (d)

Solution:

Fact

7. The set of elements that differ in mutual relationship from those of the other sets is:

a. Be – Al

b. Li – Na

c. B – Si

d. Li – Mg

Ans: (b)

Solution:

Li and Na does not have diagonal relationship.

8. Given below are two statements:

Statement I: 2-methylbutane on oxidation with KMnO_4 gives 2-methylbutan-2-ol.



Statement II: n-alkanes can be easily oxidized to corresponding alcohols with KMnO_4 .

Choose the correct option:

- a. Both statement I and statement II are incorrect
- b. Statement I is correct but statement II is incorrect
- c. Both statement I and statement II are correct
- d. Statement I is incorrect but statement II is correct

Ans: (b)

Solution:




9. Amongst the following, the linear species is:


- a. N_3^-
- b. Cl_2O
- c. O_3
- d. NO_2

Ans: (a)


Solution:




b)



c)



d)



10. For the coagulation of a negative sol, the species below, that has the highest flocculating power is:

- | | |
|----------------|----------------|
| a. SO_4^{2-} | b. Na^+ |
| c. Ba^{2+} | d. PO_4^{3-} |

Ans: (c)

Solution:

For a negative sol, positive ion is required for flocculation.

Greater the valence of the flocculating ion added, the greater is its power to cause precipitation. This is called Hardy-Schulz law.

So, Ba^{+2} has highest flocculating power.

11. The functional groups that are responsible for the ion-exchange property of cation and anion exchange resins, respectively, are:


- | | |
|-------------------------|-------------------------|
| a. $-SO_3H$ and $-COOH$ | b. $-SO_3H$ and $-NH_2$ |
| c. $-NH_2$ and $-SO_3H$ | d. $-NH_2$ and $-COOH$ |

Ans: (b)

Solution:

$-SO_3H$ and $-COOH$ are cation exchange resin and $-NH_2$ is anion exchange resin.

12. Choose the correct statement regarding the formation of carbocations A and B given.




- a. Carbocation A is more stable and formed relatively at faster rate
- b. Carbocation B is more stable and formed relatively at faster rate
- c. Carbocation A is more stable and formed relatively at slow rate
- d. Carbocation B is more stable and formed relatively at slow rate

Ans: (b)

Solution:


B carbocation is more stable due to more hyperconjugation & it form relatively faster rate compared to A.

13.




In the above reaction, the structural formula of (A), "X" and "Y" respectively are:


a.




b.



c.




d.



Ans: (b)

Solution:



14. Fructose is an example of:

- Heptose
- Aldohexose
- Pyranose
- Ketohexose

Ans: (d)

Solution:

Fructose is an example of Keto-hexose.

15. Which of the following statement(s) is (are) incorrect reason for eutrophication?

- (a) excess usage of fertilisers
- (b) excess usage of detergents
- (c) dense plant population in water bodies
- (d) lack of nutrients in water bodies that prevent plant growth

Choose the most appropriate answer from the option given below:

- a. (D) only
- b. (C) only
- c. (B) and (D) only
- d. (A) only

Ans: (a)

Solution:



Lack of nutrients in water bodies that prevent plant growth.


16. Primary, secondary and tertiary amines can be separated using:

- a. Para-Toluene sulphonyl chloride
- b. Chloroform and KOH
- c. Acetyl amide
- d. Benzene sulphonic acid

Ans: (a)

Solution:





17. Match List-I with List-II

List-I	List-II
(A) Haematite	(i) $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}$
(B) Bauxite	(ii) Fe_2O_3
(C) Magnetite	(iii) $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$
(D) Malachite	(iv) Fe_3O_4

Choose the correct answer from the options given below:

- a. (a)-(ii), (b)-(iii), (c)-(i), (d)-(iv) b. (a)-(iv), (b)-(i), (c)-(ii), (d)-(iii)
 c. (a)-(i), (b)-(iii), (c)-(ii), (d)-(iv) d. (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)

Ans: (d)

Solution:

Fact

18. The set that represents the pair of neutral oxides of nitrogen is:

- a. NO and N_2O b. NO and NO_2
 c. N_2O and NO_2 d. N_2O and N_2O_3


Ans: (a)

Solution:


NO and N_2O are neutral oxides and N_2O_3 , NO_2 and N_2O_5 are acidic oxides.

19. Nitrogen can be estimated by Kjeldahl's method for which of the following compound?


a.




b.



c.




d.



Ans: (d)

Solution:

If nitrogen atom is a part of the ring or in the form of the diazonium or NO_2 , then it is not estimated with the Kjeldahl's method.



20. One of the by-products formed during the recovery of NH_3 from Solvay process is:


- a. NaHCO_3
- c. CaCl_2

- b. Ca(OH)_2
- d. NH_4Cl

Ans: (c)

Solution:

Recovery of NH₃



SECTION - B

1. The reaction $2\text{A} + \text{B}_2 \rightarrow 2\text{AB}$ is an elementary reaction. For a certain quantity of reactants, if the volume of the reaction vessel is reduced by a factor of 3, the rate of the reaction increases by a factor of (Round off to the Nearest Integer).

Ans: 27

Solution:

For elementary reaction,

$$\text{Rate of reaction} = k[\text{A}]^2 [\text{B}_2]$$

$$\text{Initial rate} = k \left(\frac{n_{\text{A}}}{V_0} \right)^2 \left(\frac{n_{\text{B}}}{V_0} \right)$$


$$\text{Final rate} = k \left(\frac{n_{\text{A}}}{V_0/3} \right)^2 \left(\frac{n_{\text{B}}}{V_0/3} \right) = 27 k \left(\frac{n_{\text{A}}}{V_0} \right) \left(\frac{n_{\text{B}}}{V_0} \right)$$

$$\text{Final rate} = 27 \times \text{Initial rate}$$

2. In the ground state of atomic Fe ($Z = 26$), the spin-only magnetic moment is $\times 10^{-1}$ BM. (Round off to the Nearest Integer).
[Given: $\sqrt{3} = 1.73$, $\sqrt{2} = 1.41$]

Ans: 49


Solution:



In $3d^6$, no. of unpaired $e^- = 4$

$$\begin{aligned} \text{Spin only magnetic moment} &= \sqrt{n(n+2)} \text{ BM} \\ &= \sqrt{4(4+2)} \\ &= \sqrt{24} \\ &= 4.9 \\ &= 49 \times 10^{-1} \end{aligned}$$


3.



Consider the above reaction. The percentage yield of amide product is
 (Round off to the Nearest Integer)
 [Given: Atomic mass: C : 12.0 u, H : 1.0 u, N : 14.0, O : 16.0 u, Cl : 35.5 u]

Ans: 77

Solution:



Stoichiometric moles of amide = 10^{-3} mol

Actual moles of amide = 7.69×10^{-4} mol

$$\% \text{ yield} = \frac{7.69 \times 10^{-4}}{10^{-3}} \times 100$$


$$= 76.9\%$$

$$\approx 77\%$$

4. On complete reaction of FeCl_3 with oxalic acid in aqueous solution containing KOH, resulted in the formation of product A. The secondary valency of Fe in the product A is
 (Round off to the Nearest Integer)

Ans: 6

Solution:



(A)

CN = 6

S.V. = C.N

5. Consider the reaction $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$. The temperature at which $K_c = 20.4$ and $K_p = 600.1$, is K. (Round off to the Nearest Integer).
[Assume all gases are ideal and $R = 0.0831 \text{ L bar K}^{-1} \text{ mol}^{-1}$]

Ans: 354

Solution:

$$K_p = K_c (RT)^{\Delta n_g}, \Delta n_g = 1 \text{ (for given reaction)}$$

$$600.1 = 20.4 (RT)^1$$

$$T \approx 354 \text{ K}$$

6. A KCl solution of conductivity 0.14 S m^{-1} shows a resistance of 4.19Ω in a conductivity cell. If the same cell is filled with an HCl solution, the resistance drops 1.03Ω . The conductivity of the HCl solution is $\times 10^{-2} \text{ S m}^{-1}$. (Round off to the Nearest Integer).

Ans: 56

Solution:

For KCl solution,

$$R = \left(\frac{1}{K}\right)\left(\frac{\ell}{A}\right) \Rightarrow \frac{\ell}{A} = R \times K = 4.19 \times 0.14$$

$$= 0.58 \text{ ohm}$$

For HCl solution,

$$R = \left(\frac{1}{K}\right)\left(\frac{\ell}{A}\right)$$

$$K = \frac{(\ell / A)}{R} = \frac{0.58}{1.03} = 0.56 = 56 \times 10^{-2} \text{ Sm}^{-1}$$

Ans = 56

7. A 1 molal $\text{K}_4\text{Fe}(\text{CN})_6$ solution has a degree of dissociation of 0.4. Its boiling point is equal to that of another solution which contains 18.1 weight percent of a non-electrolytic solute A. The molar mass of A is g/mol. (Round off to the Nearest Integer).

Ans: 85

Solution:

Since boiling point is same,
elevation in boiling point is also same for both solution.

$$(\Delta T_B)_{K_4[Fe(CN)_6]} = (\Delta T_B)_A$$

$$(ik_b m)_{K_4[Fe(CN)_6]} = (ik_b m)_A$$

$$(1 + 4\alpha) \times 1 = \frac{1 \times \frac{18.1}{M} \times 1000}{100 - 18.1}$$

$$2.6 = \frac{18.1}{M} \times \frac{1000}{81.9}$$

$$M = 85$$

8. The number of chlorine atoms in 20 mL of chlorine gas at STP is 10^{21} . (Round off to the Nearest Integer).

[Assume chlorine is an ideal gas at STP

$R=0.083 \text{ L bar mol}^{-1} \text{ K}^{-1}$, $N_A = 6.023 \times 10^{23}$]

Ans: 1

Solution:

$$n = \frac{PV}{RT} = \frac{1 \times 20 \times 10^{-3}}{0.083 \times 273}$$

$$\text{No. of atoms} = \frac{1 \times 20 \times 10^{-3}}{0.083 \times 273} \times 2 \times 6.023 \times 10^{23} = 1.06 \times 10^{21}$$

9. KBr is doped with 10^{-5} mole percent of SrBr_2 . The number of cationic vacancies in 1 g of KBr crystal is 10^{14} . (Round off to the Nearest Integer).

[Atomic Mass: K = 39.1 u, Br = 79.9 u

$N_A = 6.023 \times 10^{23}$]

Ans: 5

Solution:

For every Sr^{+2} ion, 1 cationic vacancy is created. Hence, no. of Sr^{+2} ion = Number of cationic vacancies

Since mole percentage of SrBr_2 doped is 10^{-5} to that of total moles of KBr.

Hence,

$$\text{No. of cationic vacancy} = \frac{10^{-5}}{100} \times \frac{1}{119} \times N_A$$

$$= \frac{1}{119} \times 10^{-7} \times 6.022 \times 10^{23} = 5 \times 10^{-2} \times 10^{-7} \times 10^{23} = 5 \times 10^{14}$$

Ans. 5

10. The total number of C-C sigma bond/s in mesityl oxide (C₆H₁₀O) is (Round off to the Nearest Integer).

Ans: 5

Solution:

