



JEE Main 2021(July) Paper

(Memory Based)

Date of Exam: 20th July 2021

Time: 9:00 a.m.-12 a.m.

Subject: Mathematics

- The mean of 6 numbers is 6.5 and its variance is 10.25. If 4 numbers are 2, 4, 5 and 7, then find the other two.

Solution:

Let a and b be the other two numbers.

$$\bar{x} = 6.5$$

$$\Rightarrow \frac{a+b+2+4+5+7}{6} = 6.5$$

$$\Rightarrow a+b = 21 \quad \dots (i)$$

$$\sigma^2 = 10.25$$

$$\Rightarrow \frac{a^2+b^2+4^2+16+25+49}{6} - (6.5)^2 = 10.25$$

$$\Rightarrow a^2+b^2 = 221 \quad \dots (ii)$$

On solving (i) and (ii),

$$a = 10, b = 11 \text{ or } a = 11, b = 10$$

- Find the coefficient of x^{256} in $(1-x)^{101}(x^2+x+1)^{100}$

Solution:

$$(1-x)^{101}(x^2+x+1)^{100}$$

$$= (1-x)(1-x)^{100}(x^2+x+1)^{100}$$

$$= (1-x)(1-x^3)^{100}$$

$$= (1-x^3)^{100} - x(1-x^3)^{100}$$

First term will not have x^{256} .

Coefficient of x^{256} is $-1 \times$ coefficient of x^{255} in $(1-x^3)^{100}$

$$= -1(-{}^{100}C_{85}) = {}^{100}C_{85}$$

- If $\vec{a}, \vec{b}, \vec{c}$ are three mutually perpendicular vectors equally inclined to $\vec{a} + \vec{b} + \vec{c}$ at angle θ , then find the value of $36 \cos^2 2\theta$

Solution:

$$\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$$

$$(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = |\vec{a} + \vec{b} + \vec{c}|^2$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 = |\vec{a} + \vec{b} + \vec{c}|^2$$

$$\text{Now, } \vec{a} \cdot (\vec{a} + \vec{b} + \vec{c}) = |\vec{a}| |\vec{a} + \vec{b} + \vec{c}| \cos \theta$$

$$\Rightarrow |\vec{a}|^2 = |\vec{a}| |\vec{a} + \vec{b} + \vec{c}| \cos \theta$$

$$\Rightarrow |\vec{a}| = |\vec{a} + \vec{b} + \vec{c}| \cos \theta$$

$$\text{Similarly, } |\vec{b}| = |\vec{a} + \vec{b} + \vec{c}| \cos \theta, |\vec{c}| = |\vec{a} + \vec{b} + \vec{c}| \cos \theta$$

$$\therefore 3|\vec{a} + \vec{b} + \vec{c}|^2 \cos^2 \theta = |\vec{a} + \vec{b} + \vec{c}|^2$$

$$\Rightarrow \cos^2 \theta = \frac{1}{3} \Rightarrow 36 \cos^2 2\theta = 36 \cdot \left(2 \times \frac{1}{3} - 1\right)^2 = 4$$

JEE Main 2021(July) Paper

4. If $f(x) = \begin{cases} \sin x - e^x, & x < 0 \\ a + [-x], & 0 \leq x < 1 \\ 2x - b, & x \geq 1 \end{cases}$ is continuous in $(-\infty, 1]$, find the value of $a + b$.

Solution:

$$f(x) = \begin{cases} \sin x - e^x, & x < 0 \\ a + [-x], & 0 \leq x < 1 \\ 2x - b, & x \geq 1 \end{cases}$$

$$f(1^-) = a + (-1) = a - 1$$

$$f(1^+) = 2 - b$$

$$f(1^-) = f(1^+)$$

$$\Rightarrow a - 1 = 2 - b$$

$$\Rightarrow a + b = 3$$

5. The number of solutions of $\tan^{-1} \sqrt{x(x+1)} + \sin^{-1} \sqrt{x^2+x+1} = \frac{\pi}{2}$ is

Solution:

$$\tan^{-1} \sqrt{x(x+1)} + \sin^{-1} \sqrt{x^2+x+1} = \frac{\pi}{2}$$

$$\Rightarrow \tan^{-1} \sqrt{x(x+1)} = \frac{\pi}{2} - \sin^{-1} \sqrt{x^2+x+1}$$

$$\Rightarrow \tan^{-1} \sqrt{x(x+1)} = \cos^{-1} \sqrt{x^2+x+1}$$


$$\Rightarrow \cos^{-1} \frac{1}{\sqrt{x^2+x+1}} = \cos^{-1} \sqrt{x^2+x+1}$$

$$\Rightarrow \sqrt{x^2+x+1} = \frac{1}{\sqrt{x^2+x+1}}$$

$$\Rightarrow x^2 + x + 1 = 1$$

$$\Rightarrow x(x+1) = 0 \Rightarrow x = 0, -1$$

∴ Number of solutions is 2.



6. There are 6 batsmen, 7 bowlers and 2 wicket keepers. Find the number of ways of forming a team of 11 players having at least 4 batsmen, at least 5 bowlers and at least 1 wicket keeper.

- a. 567
c. 462

- b. 525
d. 777

Solution:

Team Size = 11 players

Case 1: 4 batsmen & 5 bowlers & 2 wicket keepers

$$\Rightarrow {}^6C_4 \times {}^7C_5 \times {}^2C_2 = \frac{6 \times 5}{2} \times \frac{7 \times 6}{2} \times 1 = 315$$

Case 2: 4 batsmen & 6 bowlers & 1 wicket keeper

$$\Rightarrow {}^6C_4 \times {}^7C_6 \times {}^2C_1 = \frac{6 \times 5}{2} \times 7 \times 2 = 210$$

Case 3: 5 batsmen & 5 bowlers & 1 wicket keeper

$$\Rightarrow {}^6C_5 \times {}^7C_5 \times {}^2C_1 = 6 \times \frac{7 \times 6}{2} \times 2 = 252$$

So total number of ways = $315 + 210 + 252 = 777$

JEE Main 2021(July) Paper

7. All the letters of the word EXAMINATION are arranged. Find the probability that 'M' is at the 4th position.

a. $\frac{2}{11}$
c. $\frac{4}{11}$

b. $\frac{1}{11}$
d. $\frac{8}{11}$

Solution:

EXAMINATION

Letters: AA II NN E T O X M

Total possible words

$$n(S) = \frac{11!}{(2!)^3}$$

Now when M is at 4th position

$$n(E) = \frac{10!}{(2!)^3}$$

$$P(M \text{ is at } 4^{\text{th}} \text{ position}) = \frac{n(E)}{n(S)} = \frac{1}{11}$$

8. Find the number of integral terms in the expansion of $(4^{\frac{1}{4}} + 5^{\frac{1}{6}})^{120}$

- a. 1
c. 20

- b. 21
d. 11

Solution:

$$\begin{aligned} T_{r+1} &= {}^{120}C_r 4^{\frac{120-r}{4}} \cdot 5^{\frac{r}{6}} \\ &= {}^{120}C_r 2^{\frac{120-r}{2}} \cdot 5^{\frac{r}{6}} \end{aligned}$$

For integral terms, r should be even and multiple of 6
i.e., r should be a multiple of 6.

r can be 0, 6, 12, ..., 120

\therefore Total number of integral terms = 21

9. If $\lim_{x \rightarrow 0} \{2 - \cos x \sqrt{\cos 2x}\}^{\frac{x+2}{x^2}} = e^\alpha$, then $\alpha = ?$

Solution:

$$\lim_{x \rightarrow 0} \{2 - \cos x \sqrt{\cos 2x}\}^{\frac{x+2}{x^2}} = e^\alpha \quad (1^\infty \text{ form})$$

$$\Rightarrow e^{\lim_{x \rightarrow 0} \{1 - \cos x \sqrt{\cos 2x}\} \times \left(\frac{x+2}{x^2}\right)} = e^\alpha$$

$$\Rightarrow e^{\lim_{x \rightarrow 0} \left\{ \frac{1 - \cos^2 x (\cos 2x)}{1 + \cos x \sqrt{\cos 2x}} \right\} \times \left(\frac{x+2}{x^2}\right)} = e^\alpha$$

$$\Rightarrow e^{\lim_{x \rightarrow 0} \left\{ \frac{1 - \cos^2 x (1 - 2 \sin^2 x)}{1 + \cos x \sqrt{\cos 2x}} \right\} \times \left(\frac{x+2}{x^2}\right)} = e^\alpha$$

$$\Rightarrow e^{\lim_{x \rightarrow 0} \left\{ \frac{1 - \cos^2 x + 2 \sin^2 x \cos^2 x}{1 + \cos x \sqrt{\cos 2x}} \right\} \times \left(\frac{x+2}{x^2}\right)} = e^\alpha$$

$$\Rightarrow e^{\lim_{x \rightarrow 0} \left\{ \frac{\sin^2 x (2 \cos^2 x + 1)}{1 + \cos x \sqrt{\cos 2x}} \right\} \times \left(\frac{x+2}{x^2}\right)} = e^\alpha$$

$$\Rightarrow e^{\lim_{x \rightarrow 0} \left\{ \frac{\sin^2 x \times (2 \cos^2 x + 1)}{x^2} \times \frac{(2 \cos^2 x + 1)}{1 + \cos x \sqrt{\cos 2x}} \right\} \times (x+2)} = e^\alpha$$

$$\Rightarrow e^{\frac{2+1}{1+1} \times 2} = e^\alpha \Rightarrow e^3 = e^\alpha \Rightarrow \alpha = 3$$

JEE Main 2021(July) Paper



10. Evaluate $\int_{-1}^1 \log(\sqrt{1-x} + \sqrt{1+x}) dx$

Solution:

$$f(-x) = f(x)$$

$\Rightarrow f$ is even.

$$\therefore I = 2 \int_0^1 \log(\sqrt{1-x} + \sqrt{1+x}) dx \quad \dots (1)$$

We will solve by integration by parts.

$$\text{Let } u = \ln(\sqrt{1-x} + \sqrt{1+x})$$

$$\begin{aligned} du &= \frac{\frac{1}{2} \left(\frac{-1}{\sqrt{1-x}} + \frac{1}{\sqrt{1+x}} \right)}{\sqrt{1-x} + \sqrt{1+x}} dx = \frac{1}{2} \frac{(\sqrt{1-x} - \sqrt{1+x})}{\sqrt{1-x^2} (\sqrt{1-x} + \sqrt{1+x})} dx \\ &= \frac{-1}{4x} \frac{(\sqrt{1-x} - \sqrt{1+x})^2}{\sqrt{1-x^2}} dx = \frac{1}{2x} \frac{(\sqrt{1-x^2} - 1)}{\sqrt{1-x^2}} dx = \frac{1}{2x} \left(1 - \frac{1}{\sqrt{1-x^2}} \right) dx \end{aligned}$$

$$I = 2 \int_0^1 \log(\sqrt{1-x} + \sqrt{1+x}) dx$$

$$I = 2x \ln(\sqrt{1-x} + \sqrt{1+x}) \Big|_0^1 - \int_0^1 \left(1 - \frac{1}{\sqrt{1-x^2}} \right) dx$$

$$I = 2x \ln(\sqrt{1-x} + \sqrt{1+x}) \Big|_0^1 - (x - \sin^{-1} x) \Big|_0^1$$

$$I = 2 \ln(2) - \left(1 - \frac{\pi}{2} \right)$$

11. The logical equivalence of the Boolean expression $\sim(p \wedge \sim q) \vee (q \vee \sim p)$ is

Solution:

$$\sim(p \wedge \sim q) \vee (q \vee \sim p)$$

$$\equiv (\sim p \vee q) \vee (q \vee \sim p)$$

$$\equiv \sim p \vee q$$

$$\equiv p \rightarrow q$$

12. Consider the curve $y^2 = 2x$ and point $A(2, 2)$. If the normal at A intersects the curve again at point B and the tangent at A intersects the x -axis at C , then area of ΔABC is

Solution:

Equation of tangent at A

$$yy_1 = x + x_1$$

$$\Rightarrow 2y = x + 2$$

Tangent at A intersects the x -axis at C

$$C(-2, 0)$$


Equation of normal at A

$$y + 2x = 6$$

Solving normal equation with $y^2 = 2x$, we get $B\left(\frac{9}{2}, -3\right)$

So, area will be $\frac{1}{2} \times AC \times AB$

$$= \frac{1}{2} \times 2\sqrt{5} \times \sqrt{\frac{25}{4} + 25} = \frac{25}{2}$$



JEE Main 2021(July) Paper

13. If for vectors A and B , $A \cdot B = |A \times B|$, then $|A - B|$ is

- a) $\sqrt{A^2 + B^2 + \sqrt{2(AB)}}$
- b) $\sqrt{A^2 + B^2 - \sqrt{2(AB)}}$
- c) $\sqrt{(A^2 + B^2 - \sqrt{2} AB)}$
- d) $\sqrt{(A^2 + B^2 + \sqrt{2} AB)}$

Solution:

$$A \cdot B = |A \times B| \Rightarrow \cos \theta = \sin \theta$$

$$\Rightarrow \tan \theta = 1$$

$$\theta = \frac{\pi}{4}$$

$$|A - B|^2 = A^2 + B^2 - 2A \cdot B$$

$$= A^2 + B^2 - 2AB \cos\left(\frac{\pi}{4}\right)$$

$$= A^2 + B^2 - \sqrt{2}AB$$

$$\Rightarrow |A - B| = \sqrt{(A^2 + B^2 - \sqrt{2} AB)}$$

14. If the roots of the quadratic equation $x^2 + 3^{1/4}x + 3^{1/2} = 0$ are α and β , then the value of

$$\alpha^{96}(\alpha^{12} - 1) + \beta^{96}(\beta^{12} - 1)$$

$$a) 50 \cdot 3^{24}$$

$$c) 52 \cdot 3^{24}$$

$$b) 51 \cdot 3^{24}$$

$$d) 104 \cdot 3^{24}$$

Solution:

$$\text{Sol. } x^2 + 3^{1/2} = -3^{1/4}x$$

$$\Rightarrow x^4 + 3 = -3^{1/2}x^2$$

$$\Rightarrow x^8 + 9 + 6x^4 = 3x^4$$

$$\Rightarrow x^8 + 9 + 3x^4 = 0$$

$$\therefore \alpha^8 = -9 - 3\alpha^4$$

$$\text{and } \alpha^{12} = -9\alpha^4 - 3\alpha^8 = -9\alpha^4 - 3(-9 - 3\alpha^4) = 27$$

$$\text{Similarly, } \beta^{12} = 27$$

$$\alpha^{96}(\alpha^{12} - 1) + \beta^{96}(\beta^{12} - 1) = (27)^8 \cdot 26 + (27)^8 \cdot 26 = 52 \cdot 3^{24}$$

15. In $\triangle ABC$, if $AB=5$, $\angle B = \cos^{-1}\left(\frac{3}{5}\right)$ and the radius of circumcircle of triangle is 5. Then the area of $\triangle ABC$ is

$$a) 6 + 8\sqrt{3}$$

$$b) 3 + 4\sqrt{3}$$

$$c) 3 + 8\sqrt{3}$$

$$d) 6 + 4\sqrt{3}$$

Solution:

$$\cos B = \frac{3}{5} \Rightarrow \sin B = \frac{4}{5}, R = 5$$

$$b = 2R \sin B = 8$$

$$AB = c = 5$$

JEE Main 2021(July) Paper

By cosine rule,

$$\begin{aligned}\cos B &= \frac{a^2 + c^2 - b^2}{2ac} = \frac{3}{5} \\ \Rightarrow \frac{a^2 + 25 - 64}{2a(5)} &= \frac{3}{5} \\ \Rightarrow a^2 - 39 &= 6a \\ \Rightarrow a^2 - 6a - 39 &= 0 \\ \Rightarrow a &= \frac{(6+8\sqrt{3})}{2} \Rightarrow a = 3 + 4\sqrt{3} \\ \Delta &= \frac{abc}{4R} = 6 + 8\sqrt{3}\end{aligned}$$

16. If the shortest distance between the lines $r_1 = \alpha\hat{i} + 2\hat{j} + 2\hat{k} + \lambda(\hat{i} - 2\hat{j} + 2\hat{k})$, $\lambda \in \mathbb{R}, \alpha > 0$ and $r_2 = -4\hat{i} - \hat{k} + \mu(3\hat{i} - 2\hat{j} - 2\hat{k})$, $\mu \in \mathbb{R}$ is 9, then the value of α is

- a. 2
- b. 4
- c. 6
- d. $\sqrt{6}$

Solution:

$$\begin{aligned}\text{Shortest distance} &= \left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_2 \times \vec{b}_1)}{|\vec{b}_2 \times \vec{b}_1|} \right| \\ 9 &= \left| \frac{((\alpha+4)\hat{i}+2\hat{j}+3\hat{k}) \cdot (8\hat{i}+8\hat{j}+4\hat{k})}{\sqrt{64+64+16}} \right| \\ \Rightarrow |(\alpha+4) \times 8 + 16 + 12| &= 108 \\ \Rightarrow (\alpha+4) \times 8 &= 80 \quad (\because \alpha > 0) \\ \Rightarrow \alpha &= 6\end{aligned}$$

17. Let $y = mx + c, m > 0$ be the focal chord of $y^2 = -64x$ which is tangent to $(x + 10)^2 + y^2 = 4$. Then the value of $4\sqrt{2}(m + c)$ is equal to

Solution:

Focus of parabola is $(-16, 0)$

$$\text{So, } -16m + c = 0 \Rightarrow c = 16m \quad \dots (i)$$

Now slope form of tangent to the circle is given by

$$y = m(x + 10) \pm 2\sqrt{1 + m^2}$$

$$\therefore c = 10m \pm 2\sqrt{1 + m^2} \quad \dots (ii)$$

So, from (i) and (ii)

$$16m = 10m \pm 2\sqrt{1 + m^2}$$

$$\Rightarrow 9m^2 = 1 + m^2 \Rightarrow m = \frac{1}{2\sqrt{2}}$$

$$\Rightarrow c = 16m = \frac{8}{2\sqrt{2}}$$

$$\therefore 4\sqrt{2}(m + c) = 34$$

JEE Main 2021(July) Paper


18. A continuous differentiable function $f(x)$ is increasing in $(-\infty, \frac{3}{2})$ and decreasing in $(\frac{3}{2}, \infty)$. Then

$x = \frac{3}{2}$ is

- a) point of local maxima
- b) point of local minima
- c) point of inflection
- d) None of these

Solution:

Roughly graph of $f(x)$ can be drawn as



Thus $x = \frac{3}{2}$ is a point of local maxima.

19. If z and ω are complex numbers such that $|z\omega| = 1$, $\arg(z) - \arg(\omega) = \frac{3\pi}{2}$. then find $\arg\left(\frac{1-2\bar{z}\omega}{1+3\bar{z}\omega}\right)$

- a) $\frac{\pi}{4}$
- b) $-\frac{\pi}{4}$
- c) $\frac{3\pi}{4}$
- d) $-\frac{3\pi}{4}$

Solution.

Let $z = re^{i\theta}$ and $\omega = \frac{1}{r}e^{i(\theta - \frac{3\pi}{2})}$

$$\text{Then } \frac{1-2\bar{z}\omega}{1+3\bar{z}\omega} = \frac{1-2re^{-i\theta} \cdot \frac{1}{r}e^{i(\theta - \frac{3\pi}{2})}}{1+3re^{-i\theta} \cdot \frac{1}{r}e^{i(\theta - \frac{3\pi}{2})}}$$

$$= \frac{1-2e^{i(-\frac{3\pi}{2})}}{1+3e^{i(-\frac{3\pi}{2})}} = \frac{1-2i}{1+3i}$$

$$= -\frac{1}{2} - \frac{1}{2}i$$

$$\arg\left(-\frac{1}{2} - \frac{1}{2}i\right) = -\frac{3\pi}{4}$$

20. If an invertible function $f(x)$ is defined as $f(x) = 3x - 2$, $g(x)$ is also an invertible function such that $f^{-1}(g^{-1}(x)) = x - 2$, then $g(x)$ is

- a) $\frac{x-8}{3}$
- b) $\frac{x+8}{3}$
- c) $\frac{x-3}{8}$
- d) $\frac{x+3}{8}$

Solution:

$$f^{-1}(g^{-1}(x)) = x - 2$$

$$\Rightarrow f(x-2) = g^{-1}(x)$$

$$\Rightarrow 3(x-2) - 2 = g^{-1}(x)$$

$$\Rightarrow 3x - 8 = g^{-1}(x)$$

$$\Rightarrow g^{-1}(x) = 3x - 8$$

$$\text{or } x = 3g(x) - 8$$

$$\therefore g(x) = \frac{x+8}{3}$$

JEE Main 2021(July) Paper

21. The probability of selecting integers $a \in [-5, 30]$, such that $x^2 + 2(a+4)x - 5a + 64 > 0$ for all $x \in \mathbf{R}$ is:

Solution:

$$x^2 + 2(a+4)x - (5a - 64) > 0$$

$$D < 0$$

$$\therefore 4(a+4)^2 + 4(5a - 64) < 0$$

$$\Rightarrow a^2 + 13a - 48 < 0$$

$$\Rightarrow (a+16)(a-3) < 0$$

$$\text{So, } a \in (-16, 3)$$

Total integers are 18

$$\therefore \text{Probability} = \frac{18}{36} = \frac{1}{2}$$

22. If $\int_0^a e^{x-[x]} dx = 10e - 9$, then the value of 'a' is (where $[.]$ is greatest integer function)

a) $9 + \ln 2$

c) 10

b) $10 + \ln 2$

d) 9

Solution:

$$\text{Let } a = K + \lambda, 0 \leq \lambda < 1$$

$$\int_0^a e^{\{x\}} dx = \int_0^K e^{\{x\}} dx + \int_K^{K+\lambda} e^{\{x\}} dx$$

$$= K \int_0^1 e^x dx + \int_0^\lambda e^x dx$$

$$= K(e-1) + (e^\lambda - 1)$$

$$= (Ke + e^\lambda) - (K + 1)$$

$$\text{Given, } \int_0^a e^{x-[x]} dx = 10e - 9$$

$$\Rightarrow (Ke + e^\lambda) - (K + 1) = 10e - 9$$

$$\Rightarrow K = 10 \text{ and } e^\lambda = 11 - 9 = 2$$

$$\Rightarrow \lambda = \ln 2$$

$$\therefore a = 10 + \ln 2$$

23. $a_{ij} = \begin{cases} 1, & i = j \\ -x, & |i - j| = 1 \\ 2x + 1, & \text{otherwise} \end{cases}$ $A = [a_{ij}]_{3 \times 3}$. $f(x) = \text{Det}(A)$. Then find the sum of local maximum and minimum values of $f(x)$.

a) $\frac{20}{27}$

b) $-\frac{20}{27}$

c) $\frac{88}{27}$

d) $-\frac{88}{27}$

Solution:

$$f(x) = \begin{vmatrix} 1 & -x & 2x+1 \\ -x & 1 & -x \\ 2x+1 & -x & 1 \end{vmatrix} = 4x^3 - 4x^2 - 4x$$

JEE Main 2021(July) Paper

$$f'(x) = 4(3x^2 - 2x - 1)$$

$$f'(x) = 4(x - 1)\left(x + \frac{1}{3}\right)$$

$$\therefore f(1) = 4 - 4 - 4 = -4$$

$$f\left(-\frac{1}{3}\right) = \frac{-4}{27} - \frac{4}{9} + \frac{4}{3} = \frac{-4-12+36}{27} = \frac{20}{27}$$

$$\frac{20}{27} - 4 = \frac{20-108}{27} = -\frac{88}{27}$$

24. Find the coefficient of $a^3b^4c^5$ in $(ab + bc + ca)^6$.

- a) 60
c) 40

- b) 45
d) 90

Solution:

$$(ab + bc + ca)^6 = \sum_{p+q+r=6} \frac{6!}{p! q! r!} (ab)^p (bc)^q (ca)^r \\ = \sum_{p+q+r=6} \frac{6!}{p! q! r!} a^{p+r} b^{p+q} c^{q+r}$$

For $a^3b^4c^5$, we need

$$p + r = 3$$

$$p + q = 4$$

$$q + r = 5$$

Solving we get, $p = 1, q = 3, r = 2$

\therefore Co-efficient of $a^3b^4c^5$ in $(ab + bc + ca)^6$ is $\frac{6!}{1!3!2!} = 60$.

25. $x \frac{dy}{dx} \cdot \tan \frac{y}{x} = y \tan \frac{y}{x} + x$, $y\left(\frac{1}{2}\right) = \frac{\pi}{6}$. Area bounded by $x = 0, x = \frac{1}{\sqrt{2}}, y = y(x)$.

Solution:

$$x \frac{dy}{dx} \cdot \tan \frac{y}{x} = y \tan \frac{y}{x} + x$$

$$\therefore \frac{dy}{dx} \tan \left(\frac{y}{x}\right) = \frac{y}{x} \tan \left(\frac{y}{x}\right) + 1$$

Let $y = vx$

$$\frac{dy}{dx} = v + x \frac{dv}{dx}$$

$$\left(v + x \frac{dv}{dx}\right) \tan v = v \tan v + 1$$

$$\Rightarrow v + x \frac{dv}{dx} = v + \cot v$$

JEE Main 2021(July) Paper

$$\Rightarrow \frac{dv}{\cot v} = \frac{dx}{x}$$

$$\Rightarrow \tan v \, dv = \frac{dx}{x}$$

$$-\log|\cos v| = \log |cx|$$

$$-\cos v = cx$$

$$\therefore y = x \cos^{-1} x$$

$$\int_0^{1/\sqrt{2}} x \cos^{-1} x = \frac{1}{8}(\pi - 1)$$



Aakash
+ BYJU'S