




Grade 10

Mathematics Chapter Notes



Triangles



Topics



1. Similar Triangles


2. Criteria of Similarity of Triangles

3. Pythagoras Theorem

4. Basic Proportionality Theorem




SSS




SAS

$$a^2 + b^2 = c^2$$



AAA

Similar Triangles



Relation between Corresponding Sides and Angles

- ★ Two triangles are similar, if
 - ★ Their corresponding angles are equal.

$$\angle A = \angle P$$

$$\angle B = \angle Q$$

$$\angle C = \angle R$$

- ★ Their corresponding sides are in the same ratio.


$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{CA}{RP} = k$$

Criteria for Similarity of Triangles


Side-Side-Side (SSS)

★ Corresponding sides are proportional.

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{CA}{RP}$$



Angle-Angle-Angle (AAA) / Angle-Angle (AA)



★ Corresponding angles are equal.


★ Triangles are similar even if a pair of corresponding angles are equal.

Side-Angle-Side (SAS)


★ Pair of adjacent corresponding sides are proportional and one angle is equal.

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{2}{3}$$

$$\angle B = \angle Q$$



Ratio of Areas of Similar Triangles



Ratio of Area of Similar Triangles

$$\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle PQR)} = \left(\frac{AB}{PQ}\right)^2 = \left(\frac{BC}{QR}\right)^2 = \left(\frac{CA}{RP}\right)^2$$


Properties of Right-Angled Triangles

Similarity of triangles when a perpendicular is drawn from the vertex of the right angle.

$$\triangle ABC \sim \triangle ADC \sim \triangle ADB \text{ (AA Similarity)}$$

All the three triangles have:

- ★ A right-angle.
- ★ A common angle.



Basic Proportionality Theorem



If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, then the other two sides are divided in the same ratio.

Proof:

$$\text{Area of } \triangle APQ = \frac{1}{2} \times AP \times QN$$

$$\text{Area of } \triangle PBQ = \frac{1}{2} \times PB \times QN$$

$$\text{Area of } \triangle APQ = \frac{1}{2} \times AQ \times PM$$

$$\text{Area of } \triangle QCP = \frac{1}{2} \times QC \times PM$$

Now,

$$\frac{\text{Area of } \triangle APQ}{\text{Area of } \triangle PBQ} = \frac{\frac{1}{2} \times AP \times QN}{\frac{1}{2} \times PB \times QN} = \frac{AP}{PB} \dots\dots\dots(1)$$

Similarly,


$$\frac{\text{Area of } \triangle APQ}{\text{Area of } \triangle QCP} = \frac{\frac{1}{2} \times AQ \times PM}{\frac{1}{2} \times QC \times PM} = \frac{AQ}{QC} \dots\dots\dots(2)$$

The triangles drawn between the same parallel lines and on the same base have equal areas.

$$\therefore \text{Area of } \triangle PBQ = \text{Area of } \triangle QCP \dots\dots\dots(3)$$

From (1), (2) and (3)

$$\frac{AP}{PB} = \frac{AQ}{QC}$$



Converse of Basic Proportionality Theorem



If a line divides any two sides of a triangle in the same ratio, then the line is parallel to the third side.

Proof:

If $\frac{AD}{DB} = \frac{AE}{EC}$, then $DE \parallel BC$.

Suppose a line DE , intersects the two sides of a triangle AB and AC at D and E , such that:

$$\frac{AD}{DB} = \frac{AE}{EC} \dots\dots(1)$$

Assume DE is not parallel to BC . Now, draw a line DE' parallel to BC . Hence, by Basic Proportionality Theorem,

$$\frac{AD}{DB} = \frac{AE'}{E'C} \dots\dots(2)$$

From eq. 1 and 2, we get

$$\frac{AE}{EC} = \frac{AE'}{E'C}$$

Adding 1 on both the sides


$$\frac{AE}{EC} + 1 = \frac{AE'}{E'C} + 1 \Rightarrow \frac{AE+EC}{EC} = \frac{AE'+E'C}{E'C}$$

$$\frac{AC}{EC} = \frac{AC}{E'C} \Rightarrow \text{So, } EC = E'C$$

This is possible only when E and E' coincides.

But $DE' \parallel BC$

$\therefore DE \parallel BC$.



Properties of Right-Angled Triangles

Pythagoras Theorem




In a right-angled triangle, the square of the hypotenuse side is equal to the sum of squares of the other two sides.

Proof:

$$\triangle ADB \sim \triangle ABC$$

$$\therefore \frac{AD}{AB} = \frac{AB}{AC} \text{ (corresponding sides of similar triangles)}$$

$$AB^2 = AD \times AC \text{(1)}$$



$$\text{Also, } \triangle ADC \sim \triangle ABC$$

$$\therefore \frac{CD}{BC} = \frac{BC}{AC} \text{ (corresponding sides of similar triangles)}$$

$$BC^2 = CD \times AC \text{(2)}$$

$$(1) + (2)$$

$$AB^2 + BC^2 = AD \times AC + CD \times AC$$

$$AB^2 + BC^2 = AC (AD + CD)$$

$$\text{Since, } AD + CD = AC$$

$$\therefore AC^2 = AB^2 + BC^2$$

Converse of Pythagoras Theorem




If the square of the length of the longest side of a triangle is equal to the sum of the squares of the other two sides, then the triangle is a right triangle.

Proof:

Construct another triangle, $\triangle EGF$, such as
 $AC = EG$ and $BC = FG$.

In $\triangle EGF$, by Pythagoras Theorem:

$$EF^2 = EG^2 + FG^2 = b^2 + a^2 \dots\dots(1)$$



In $\triangle ABC$, by Pythagoras Theorem:

$$AB^2 = AC^2 + BC^2 = b^2 + a^2 \dots\dots(2)$$

From (1) and (2)


$$EF^2 = AB^2$$

$$EF = AB$$

$$\Rightarrow \triangle ACB \cong \triangle EGF \text{ (By SSS)}$$

$$\Rightarrow \angle C \text{ is right angle}$$



$$\therefore \triangle ABC \text{ is a right triangle.}$$







Similarity of Triangles



SSS



AAA/AA




SAS



Pythagoras Theorem

★ In a right-angled triangle, the square of the hypotenuse side is equal to the sum of squares of the other two sides.


$$a^2 + b^2 = c^2$$



Basic Proportionality Theorem

★ If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, then the other two sides are divided in the same ratio.

$$PQ \parallel BC, \frac{AP}{PB} = \frac{AQ}{QC}$$





Mind Map

